*Acta Mathematica Academiae Paedagogicae Nyíregyháziensis*
**24** (2008), 115–123
www.emis.de/journals
ISSN 1786-0091

# ON NONLINEAR CONNECTIONS IN HIGHER ORDER LAGRANGE SPACES

MARCEL ROMAN

Abstract. Considering a Lagrangian of order $k$, we determine a nonlinear connection $N$ on $T^kM$ such that the horizontal and vertical distributions to be Lagrangian subbundles for the presymplectic structure given by the Cartan-Poincaré two-form $\omega_L^k$.

## 1. Introduction

We denote by $(T^kM, \pi^k, M)$, $k \geq 1$, the space of tangent bundle of order $k$ over a smooth, real, $n$-dimensional manifold $M$, [5]. Local coordinates $(x^i)$ on $M$ induce local coordinates $(x^i, y^{(1)i}, \ldots, y^{(k)i})$ on $T^kM$, where for a $k$-jet $j_0^k\rho \in T^kM$, the coordinate functions are defined as follows

$$y^{(\alpha)i}(j_0^k\rho) = \frac{1}{\alpha!}\frac{d^\alpha(x^i \circ \rho)}{dt^\alpha}\bigg|_{t=0}, \ \alpha \in \{1, \ldots, k\}.$$

The tangent structure of order $k, J$ is defined as follows, [6],

$$J = \frac{\partial}{\partial y^{(1)i}} \otimes dx^i + \frac{\partial}{\partial y^{(2)i}} \otimes dy^{(1)i} + \cdots + \frac{\partial}{\partial y^{(k)i}} \otimes dy^{(k-1)i}.$$

The foliated structure of $T^kM$ allows for $k$ regular, integrable, vertical distributions, $V_{k-\alpha+1} = \operatorname{Ker} J^\alpha = \operatorname{Im} J^{k-\alpha+1}$, $\alpha \in \{1, \ldots, k\}$.

2000 *Mathematics Subject Classification.* 53B40, 53C20, 53C60.
*Key words and phrases.* higher order Lagrange spaces, nonlinear connection, Cartan-Poincaré forms.
This work has been supported by CEEX Grant 31742/2006 and by Grant CNCSIS 1158/2006 from the Romanian Ministry of Education.

The following $k$ vertical vector fields are globally defined on $T^kM$ and they are called Liouville vector fields:

$$\Gamma_k = y^{(1)i}\frac{\partial}{\partial y^{(1)i}} + 2y^{(2)i}\frac{\partial}{\partial y^{(2)i}} + \cdots + ky^{(k)i}\frac{\partial}{\partial y^{(k)i}},$$
$$\Gamma_\alpha = J^{k-\alpha}\left(\Gamma_k\right),\ \alpha \in \{1, 2, \ldots, k\}.$$

A semispray is a globally defined vector field $S$ on $T^kM$ that satisfies the equation $JS = \Gamma_k$. Therefore, a semispray $S$, which is a vector field of order $k+1$, which can be expressed as follows

$$S = y^{(1)i}\frac{\partial}{\partial x^i} + 2y^{(2)i}\frac{\partial}{\partial y^{(1)i}} + \cdots + ky^{(k)i}\frac{\partial}{\partial y^{(k-1)i}} - \tag{1.1}$$
$$- (k+1)G^i(x, y^{(1)}, \ldots, y^{(k)})\frac{\partial}{\partial y^{(k)i}}$$

and it is perfectly determined by its coefficient functions $G^i(x, y^{(1)}, \ldots, y^{(k)})$.

A nonlinear connection, or a horizontal distribution on $T^kM$ is a regular distribution $N\colon u \in T^kM \mapsto N(u) \subset T_uT^kM$ such that the following direct sum holds true:

$$T_u(T^kM) = N(u) \oplus N_1(u) \oplus \cdots \oplus N_{k-1}(u) \oplus V_k(u), \tag{1.2}$$

where $N_1 = J(N), N_{\alpha-1} = J^{\alpha-1}(N),\ \alpha \in \{3, \ldots, k\}$. The adapted basis to this decomposition is given by

$$\left\{\frac{\delta}{\delta x^i}, \frac{\delta}{\delta y^{(1)i}}, \ldots, \frac{\delta}{\delta y^{(k-1)i}}, \frac{\partial}{\partial y^{(k)i}}\right\},$$

where, [7]:

$$\begin{aligned}
\frac{\delta}{\delta x^i} &= \frac{\partial}{\partial x^i} - \underset{(1)}{N^j_i}\frac{\partial}{\partial y^{(1)j}} - \cdots - \underset{(k)}{N^j_i}\frac{\partial}{\partial y^{(k)j}},\\
\frac{\delta}{\delta y^{(1)i}} &= \frac{\partial}{\partial y^{(1)i}} - \underset{(1)}{N^j_i}\frac{\partial}{\partial y^{(2)j}} - \cdots - \underset{(k-1)}{N^j_i}\frac{\partial}{\partial y^{(k)j}},\\
&\vdots\\
\frac{\delta}{\delta y^{(k-1)i}} &= \frac{\partial}{\partial y^{(k-1)i}} - \underset{(1)}{N^j_i}\frac{\partial}{\partial y^{(k)j}}.
\end{aligned} \tag{1.3}$$

The functions $\underset{(1)}{N^j_i}, \underset{(2)}{N^j_i}, \ldots, \underset{(k)}{N^j_i}$ are called the local coefficients of the nonlinear connection $N$.

The dual basis of the previous adapted basis is given by $\left\{dx^i, \delta y^{(1)i}, \ldots, \delta y^{(k)i}\right\}$, where:

$$
\begin{aligned}
\delta y^{(1)i} &= dy^{(1)i} + \underset{(1)}{M^i}_j dx^j, \\
\delta y^{(2)i} &= dy^{(2)i} + \underset{(1)}{M^i}_j dy^{(1)j} + \underset{(2)}{M^i}_j dx^j, \\
&\vdots \\
\delta y^{(k)i} &= dy^{(k)i} + \underset{(1)}{M^i}_j dy^{(k-1)j} + \cdots + \underset{(k)}{M^i}_j dx^j.
\end{aligned} \tag{1.4}
$$

The functions $\underset{(1)}{M^j}_i, \underset{(2)}{M^j}_i, \ldots, \underset{(k)}{M^j}_i$ are called the dual coefficients of the nonlinear connection $N$.

## 2. Cartan-Poincaré forms for a higher order Lagrangian

Let us consider a regular Lagrangian of order $k$, $(k > 1)$, $L(x^i, y^{(1)i}, \ldots, y^{(k)i})$. The metric tensor is given by the symmetric $d$-tensor field:

$$g_{ij} = \frac{1}{2} \frac{\partial^2 L}{\partial y^{(k)i} \partial y^{(k)j}}, \tag{2.1}$$

which has maximal rank on $T^k M$, $\operatorname{rank} ||g_{ij}|| = n$.

For a regular Lagrangian of order $k$, we consider the following Cartan-Poincaré one-forms, [4]:

$$\theta_L^\alpha = d_{J^\alpha} L = \frac{\partial L}{\partial y^{(\alpha)i}} dx^i + \cdots + \frac{\partial L}{\partial y^{(k-\alpha)i}} dy^{(\alpha)i}, \quad \alpha \in \{1, \ldots, k\}. \tag{2.2}$$

We consider also the following Cartan-Poincaré two-forms:

$$
\begin{aligned}
&\omega_L^\alpha = d\theta_L^\alpha = d\left(\frac{\partial L}{\partial y^{(\alpha)i}}\right) \wedge dx^i + \cdots + d\left(\frac{\partial L}{\partial y^{(k-\alpha)i}}\right) \wedge dy^{(\alpha)i}, \\
&\alpha \in \{1, \ldots, k\}.
\end{aligned} \tag{2.3}
$$

We remark here that for $k > 1$, the regularity of the Lagrangian $L$ implies the fact that $\operatorname{rank}(\omega_L^k) = 2n < (k+1)n = \dim(T^k M)$. We refer to $\omega_L^k$ as to the canonical presymplectic structure of the Lagrangian $L$.

## 3. Nonlinear connection

Now, our question is: Can we find an adapted basis such that the horizontal distribution to be Lagrangian subbundle for the presymplectic structure $\omega_L^k$? Taking into account (1.3) this is equivalent with the determination of the coefficients of a nonlinear connection $N$.

As we know, [1], in the $k = 1$ case, we consider the canonical nonlinear connection and we have $\omega_L = 2g_{ji}\delta y^j \wedge dx^i$. It follows that in the basis $(dx^i, \delta y^i)$ of the cotangent bundle, the matrix of $\omega_L$ is

$$\begin{pmatrix} 0 & 2g_{ji} \\ -2g_{ji} & 0 \end{pmatrix}.$$

For $k > 1$, we consider the Cartan-Poincaré two-form:

$$\begin{aligned} \omega_L^k &= d\left(\frac{\partial L}{\partial y^{(k)i}}\right) \wedge dx^i \\ &= \frac{1}{2}\left[\frac{\delta}{\delta x^j}\left(\frac{\partial L}{\partial y^{(k)i}}\right) - \frac{\delta}{\delta x^i}\left(\frac{\partial L}{\partial y^{(k)j}}\right)\right] dx^j \wedge dx^i \\ &\quad + \frac{\delta}{\delta y^{(1)j}}\left(\frac{\partial L}{\partial y^{(k)i}}\right) \delta y^{(1)j} \wedge dx^i + \cdots \\ &\quad + \frac{\delta}{\delta y^{(k-1)j}}\left(\frac{\partial L}{\partial y^{(k)i}}\right) \delta y^{(k-1)j} \wedge dx^i \\ &\quad + 2g_{ji}\delta y^{(k)j} \wedge dx^i. \end{aligned} \tag{3.1}$$

We are looking for a nonlinear connection on $T^kM$ such that the presymplectic structure of the lagrangian $L$ to be: $\omega_L^k = 2g_{ji}\delta y^{(k)j} \wedge dx^i$, i.e. to have the following matrix:

$$\begin{pmatrix} 0 & 0 \cdots 0 & 2g_{ji} \\ 0 & & \\ \vdots & O_{kn} & \\ -2g_{ji} & & \end{pmatrix}_{(k+1)n \times (k+1)n}.$$

This is equivalent with the vanishes of all coefficients from (3.1), except the last one. We will obtain the coefficients of the nonlinear connection N.

We have:

$$\frac{\delta}{\delta y^{(k-1)j}}\left(\frac{\partial L}{\partial y^{(k)i}}\right) = 0 \quad \Longrightarrow \quad \frac{\partial^2 L}{\partial y^{(k-1)j}\partial y^{(k)i}} - \underset{(1)}{N_j^m} 2g_{mi} = 0$$

Therefore, we find the first coefficient for the nonlinear connection:

$$\underset{(1)}{N_j^m} = \frac{1}{2}g^{mi}\frac{\partial^2 L}{\partial y^{(k-1)j}\partial y^{(k)i}}. \tag{3.2}$$

Now, we take $\dfrac{\delta}{\delta y^{(k-2)j}}\left(\dfrac{\partial L}{\partial y^{(k)i}}\right) = 0$ and we obtain the second coefficient of the nonlinear connection:

$$\underset{(2)}{N_j^m} = \frac{1}{2}g^{mi}\frac{\partial^2 L}{\partial y^{(k-2)j}\partial y^{(k)i}} - \underset{(1)}{N_j^r}\frac{1}{2}g^{mi}\frac{\partial^2 L}{\partial y^{(k-1)r}\partial y^{(k)i}}. \tag{3.3}$$

Finally, from $\dfrac{\delta}{\delta y^{(1)j}}\left(\dfrac{\partial L}{\partial y^{(k)i}}\right)=0$ we obtain the $(k-1)^{th}$ coefficient of the nonlinear connection $N$:

$$\begin{aligned}\underset{(k-1)}{N^m_j} &= \frac{1}{2}g^{mi}\frac{\partial^2 L}{\partial y^{(1)j}\partial y^{(k)i}} - \underset{(1)}{N^r_j}\frac{1}{2}g^{mi}\frac{\partial^2 L}{\partial y^{(2)r}\partial y^{(k)i}}\\ &\quad -\cdots-\underset{(k-2)}{N^r_j}\frac{1}{2}g^{mi}\frac{\partial^2 L}{\partial y^{(k-1)r}\partial y^{(k)i}}.\end{aligned}\tag{3.4}$$

Consequently, the coefficients $\underset{(1)}{N^m_j}, \underset{(2)}{N^m_j}, \ldots, \underset{(k-1)}{N^m_j}$ are unique determined.

We have:

**Theorem 3.1.** *With respect to a transformation of the local coordinates* $\left(x^i, y^{(1)i}, \ldots, y^{(k)i}\right) \longrightarrow \left(\widetilde{x}^i, \widetilde{y}^{(1)i}, \ldots, \widetilde{y}^{(k)i}\right)$ *on* $T^kM$, *the coefficient of the nonlinear connection* $N$ *are transformed by the rule:*

$$\begin{aligned}
\underset{(1)}{\widetilde{N}^i_m}\frac{\partial \widetilde{x}^m}{\partial x^j} &= \frac{\partial \widetilde{x}^i}{\partial x^m}\underset{(1)}{N^m_j} - \frac{\partial \widetilde{y}^{(1)i}}{\partial x^j},\\
\underset{(2)}{\widetilde{N}^i_m}\frac{\partial \widetilde{x}^m}{\partial x^j} &= \frac{\partial \widetilde{x}^i}{\partial x^m}\underset{(2)}{N^m_j} + \frac{\partial \widetilde{y}^{(1)i}}{\partial x^m}\underset{(1)}{N^m_j} - \frac{\partial \widetilde{y}^{(2)i}}{\partial x^j},\\
&\vdots\\
\underset{(k-1)}{\widetilde{N}^i_m}\frac{\partial \widetilde{x}^m}{\partial x^j} &= \frac{\partial \widetilde{x}^i}{\partial x^m}\underset{(k-1)}{N^m_j} + \frac{\partial \widetilde{y}^{(1)i}}{\partial x^m}\underset{(k-2)}{N^m_j} + \cdots + \frac{\partial \widetilde{y}^{(k-2)i}}{\partial x^m}\underset{(1)}{N^m_j} - \frac{\partial \widetilde{y}^{(k-1)i}}{\partial x^j}.
\end{aligned}\tag{3.5}$$

*Proof.* A transformation of local coordinates

$$\left(x^i, y^{(1)i}, \ldots, y^{(k)i}\right) \longrightarrow \left(\widetilde{x}^i, \widetilde{y}^{(1)i}, \ldots, \widetilde{y}^{(k)i}\right)$$

on the manifold $T^kM$ is given by, [7]:

$$\begin{aligned}
\widetilde{x}^i &= \widetilde{x}^i(x^1, \ldots, x^n),\ \operatorname{rank}\|\frac{\partial \widetilde{x}^i}{\partial x^j}\| = n,\\
\widetilde{y}^{(1)i} &= \frac{\partial \widetilde{x}^i}{\partial x^j}\, y^{(1)j},\\
2\widetilde{y}^{(2)i} &= \frac{\partial \widetilde{y}^{(1)i}}{\partial x^j}\, y^{(1)j} + 2\frac{\partial \widetilde{y}^{(1)i}}{\partial y^{(1)j}}\, y^{(2)j},\\
&\vdots\\
k y^{(k)i} &= \frac{\partial \widetilde{y}^{(k-1)i}}{\partial x^j}\, y^{(1)j} + 2\frac{\partial \widetilde{y}^{(k-1)i}}{\partial y^{(1)j}}\, y^{(2)j} + \cdots + k\frac{\partial \widetilde{y}^{(k-1)i}}{\partial y^{(k-1)j}}\, y^{(k)j}.
\end{aligned}$$

Also, we have the identities, [7]:

$$\frac{\partial \widetilde{y}^{(\alpha)i}}{\partial x^j}=\frac{\partial \widetilde{y}^{(\alpha+1)i}}{\partial y^{(1)j}}=\cdots=\frac{\partial \widetilde{y}^{(k)i}}{\partial y^{(k-\alpha)j}},\ (\alpha=0,\dots,k-1; y^{(0)i}=x^i).$$

With respect to the previous transformation of local coordinates, the natural basis is changed by the following rule:

$$\begin{aligned}
\frac{\partial}{\partial x^i}&=\frac{\partial \widetilde{x}^j}{\partial x^i}\frac{\partial}{\partial \widetilde{x}^j}+\frac{\partial \widetilde{y}^{(1)j}}{\partial x^i}\frac{\partial}{\partial \widetilde{y}^{(1)j}}+\cdots+\frac{\partial \widetilde{y}^{(k)j}}{\partial x^i}\frac{\partial}{\partial \widetilde{y}^{(k)j}},\\
\frac{\partial}{\partial y^{(1)i}}&=\frac{\partial \widetilde{y}^{(1)j}}{\partial y^{(1)i}}\frac{\partial}{\partial \widetilde{y}^{(1)j}}+\cdots+\frac{\partial \widetilde{y}^{(k)j}}{\partial y^{(1)i}}\frac{\partial}{\partial \widetilde{y}^{(k)j}},\\
&\vdots\\
\frac{\partial}{\partial y^{(k)i}}&=\frac{\partial \widetilde{y}^{(k)j}}{\partial y^{(k)i}}\frac{\partial}{\partial \widetilde{y}^{(k)j}}.
\end{aligned}$$

Taking into account last formulas, we have

$$\frac{\partial L}{\partial y^{(k)i}}=\frac{\partial \widetilde{y}^{(k)j}}{\partial y^{(k)i}}\frac{\partial L}{\partial \widetilde{y}^{(k)j}}$$

and consequently, we obtain

$$\frac{\partial^2 L}{\partial y^{(k)i}\partial y^{(k-1)m}}=\frac{\partial \widetilde{y}^{(k)j}}{\partial y^{(k)i}}\frac{\partial \widetilde{y}^{(k-1)r}}{\partial y^{(k-1)m}}\frac{\partial^2 L}{\partial \widetilde{y}^{(k)j}\partial \widetilde{y}^{(k-1)r}}+\frac{\partial \widetilde{y}^{(k)r}}{\partial y^{(k-1)m}}\frac{\partial \widetilde{y}^{(k)j}}{\partial y^{(k)i}}\frac{\partial^2 L}{\partial \widetilde{y}^{(k)j}\partial \widetilde{y}^{(k)r}}.$$

But,

$$\frac{\partial \widetilde{y}^{(k)r}}{\partial y^{(k-1)m}}=\frac{\partial \widetilde{y}^{(1)r}}{\partial x^m}\ \text{and}\ 2g_{jr}=\frac{\partial^2 L}{\partial \widetilde{y}^{(k)j}\partial \widetilde{y}^{(k)r}}.$$

Now, contracting by $\frac{1}{2}g^{ij}$, we obtain

$$\underset{(1)}{N}{}^j_m=\frac{\partial \widetilde{x}^j}{\partial x^i}\frac{\partial \widetilde{x}^r}{\partial x^m}\underset{(1)}{\widetilde{N}}{}^i_r+\frac{\partial \widetilde{y}^{(1)r}}{\partial x^m}\frac{\partial \widetilde{x}^j}{\partial x^r}$$

and finally,

$$\underset{(1)}{\widetilde{N}}{}^r_i\frac{\partial \widetilde{x}^i}{\partial x^m}=\underset{(1)}{N}{}^j_m\frac{\partial \widetilde{x}^r}{\partial x^j}-\frac{\partial \widetilde{y}^{(1)r}}{\partial x^m}.$$

Similarly, in order to check the second relation from (3.5), we calculate $\frac{\partial^2 L}{\partial y^{(k)i}\partial y^{(k-2)m}}$, for the third relation we calculate $\frac{\partial^2 L}{\partial y^{(k)i}\partial y^{(k-3)m}}$ and so on, for the last relation we calculate $\frac{\partial^2 L}{\partial y^{(k)i}\partial y^{(1)m}}$. □

Now, considering the following notations:

$$\underset{(\alpha)}{M}{}^r_j=\frac{1}{2}g^{ri}\frac{\partial^2 L}{\partial y^{(k-\alpha)j}\partial y^{(k)i}},\quad \alpha\in\{1,\dots,k-1\},\tag{3.6}$$

we obtain:

**Proposition 3.1.** *The relationship between the coefficients* $\underset{(1)}{N^m_i}, \underset{(2)}{N^m_i}, \ldots, \underset{(k-1)}{N^m_i}$ *of the nonlinear connection N and the coefficents given in (3.6) is expressed by:*

$$
\begin{aligned}
&\underset{(1)}{N^m_i} = \underset{(1)}{M^m_i} \\
&\underset{(2)}{N^m_i} = \underset{(2)}{M^m_i} - \underset{(1)}{M^m_r}\,\underset{(1)}{N^r_i} \\
&\qquad \vdots \\
&\underset{(k-1)}{N^m_i} = \underset{(k-1)}{M^m_i} - \underset{(k-2)}{M^m_r}\,\underset{(1)}{N^r_i} - \cdots - \underset{(1)}{M^m_r}\,\underset{(k-2)}{N^r_i}\,.
\end{aligned}
\tag{3.7}
$$

Indeed, replacing (3.6) in (3.2),(3.3) and (3.4) we have the conclusion. Therefore, the system of functions $\left\{ \underset{(1)}{M^m_i}, \underset{(2)}{M^m_i}, \ldots, \underset{(k-1)}{M^m_i} \right\}$ is the system of dual coefficients of the nonlinear connection $N$.

*Example* 3.1. Let $\mathcal{R} = (M, g_{ij}(x))$ be a Riemannian space and $\mathrm{Prol}^2\mathcal{R}^n$ its prolongation of order 2, [8].

We consider the Liouville $d$-vector fields, [7]:

$$
\begin{aligned}
&z^{(1)m} = y^{(1)m}, \\
&z^{(2)m} = \frac{1}{2}\left[\Gamma z^{(1)m} + \gamma^m_{ij} z^{(1)i} z^{(1)j}\right],
\end{aligned}
\tag{3.8}
$$

where $\gamma^m_{ij}(x)$ are the Christoffel symbols and the operator $\Gamma$ is given by:

$$\Gamma = y^{(1)i}\frac{\partial}{\partial x^i} + 2y^{(2)i}\frac{\partial}{\partial y^{(1)i}}.$$

Considering the Lagrangian $L = g_{ij}(x) z^{(k)i} z^{(k)j}$, we remark that the first coefficient $\underset{(1)}{N^m_i}$ of our nonlinear connection is the same with the first coefficient from nonlinear connections determined by I. Bucataru, [2], M. de Leon, [4], and R. Miron, [7], i.e. is expressed by:

$$\underset{(1)}{N^i_{\ j}} = \underset{(1)}{M^i_{\ j}} = \gamma^i_{jh}(x) y^{(1)h}.$$

Now, for the last coefficient of the nonlinear connection $\underset{(k)}{N^i_{\ j}}$, we have:

**Theorem 3.2.** *The skew symmetric part of the coefficient* $N^i_{\ j\ (k)}$ *is expressed as follows:*

$$
\begin{aligned}
N_{\ [ji]\ (k)} &= \frac{1}{4}\left(\frac{\partial^2 L}{\partial x^i \partial y^{(k)j}} - \frac{\partial^2 L}{\partial x^j \partial y^{(k)i}}\right) \\
&\quad - \frac{1}{2} g^{sm} \sum_{\alpha=1}^{k-1} \left( N_{\ js\ (\alpha)} M_{\ im\ (k-\alpha)} - N_{\ is\ (\alpha)} M_{\ jm\ (k-\alpha)} \right).
\end{aligned} \tag{3.9}
$$

*where* $M_{\ ji\ (k)} = g_{im} M^m_{j\ (k)}$.

*Proof.* For the last coefficient of the nonlinear connection $N^i_{\ j\ (k)}$ we have to consider the first coefficient from (3.1):

$$
\frac{1}{2}\left[\frac{\delta}{\delta x^j}\left(\frac{\partial L}{\partial y^{(k)i}}\right) - \frac{\delta}{\delta x^i}\left(\frac{\partial L}{\partial y^{(k)j}}\right)\right] = 0 \tag{3.10}
$$

We obtain:

$$
\begin{aligned}
N_{\ [ji]\ (k)} &= \frac{1}{2}\left(N_{\ ji\ (k)} - N_{\ ij\ (k)}\right) \\
&= \frac{1}{4}\left(\frac{\partial^2 L}{\partial x^i \partial y^{(k)j}} - \frac{\partial^2 L}{\partial x^j \partial y^{(k)i}}\right) \\
&\quad - \frac{1}{2}\left(N^m_{j\ (1)} \frac{\partial^2 L}{\partial y^{(1)m} \partial y^{(k)i}} - N^m_{i\ (1)} \frac{\partial^2 L}{\partial y^{(1)m} \partial y^{(k)j}}\right) \\
&\quad - \cdots - \frac{1}{2}\left(N^m_{j\ (k-1)} \frac{\partial^2 L}{\partial y^{(k-1)m} \partial y^{(k)i}} - N^m_{i\ (k-1)} \frac{\partial^2 L}{\partial y^{(k-1)m} \partial y^{(k)j}}\right),
\end{aligned} \tag{3.11}
$$

where $N_{\ ji\ (k)} = g_{im} N^m_{j\ (k)}$.

The condition (3.10) determines uniquely the skew symmetric part of the coefficient $N^m_{j\ (k)}$, only. □

For the symmetric part we need a supplementary condition. In the $k = 1$ case, I. Bucataru proved that the symmetric part is uniquely determined by a metric condition, [3].

So far, we have a whole family of nonlinear connections that are determined by the presymplectic structure $\omega^k_L$. These connections are derived directly from the Lagrangian $L$ and does not use the canonical semispray.

## References

[1] M. Anastasiei. Symplectic structures and Lagrange geometry. In *Finsler and Lagrange geometries (Iaşi, 2001)*, pages 9–16. Kluwer Acad. Publ., Dordrecht, 2003.

[2] I. Bucătaru. Sprays and homogeneous connections in the higher order geometry. *Stud. Cerc. Mat.*, 50(5-6):307–315, 1998.

[3] I. Bucataru. Metric nonlinear connections. *Differential Geom. Appl.*, 25(3):335–343, 2007.

[4] M. de León and P. R. Rodrigues. *Generalized classical mechanics and field theory*, volume 112 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1985. A geometrical approach of Lagrangian and Hamiltonian formalisms involving higher order derivatives, Notes on Pure Mathematics, 102.

[5] C. Ehresmann. Les prolongements d'une variété différentiable. I. Calcul des jets, prolongement principal. *C. R. Acad. Sci. Paris*, 233:598–600, 1951.

[6] H. A. Eliopoulos. On the general theory of differentiable manifolds with almost tangent structure. *Canad. Math. Bull.*, 8:721–748, 1965.

[7] R. Miron. *The geometry of higher-order Lagrange spaces*, volume 82 of *Fundamental Theories of Physics*. Kluwer Academic Publishers Group, Dordrecht, 1997. Applications to mechanics and physics.

[8] R. Miron and G. Atanasiu. Prolongation of Riemannian, Finslerian and Lagrangian structures. *Rev. Roumaine Math. Pures Appl.*, 41(3-4):237–249, 1996.

[9] W. M. Tulczyjew. The Lagrange differential. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.*, 24(12):1089–1096, 1976.

Marcel Roman,
Technical University "Gh. Asachi",
Department of Mathematics,
Iaşi, 700506, Romania
*E-mail address*: marcelroman@gmail.com