Acta Mathematica Academiae Paedagogicae Nyíregyháziensis
32 (2016), 135–139
www.emis.de/journals
ISSN 1786-0091

ON A FINSLER SPACE WITH A SPECIAL METRICAL CONNECTION

M. K. GUPTA AND ANIL K. GUPTA

Abstract. In this paper, we consider a Finsler space with a special metrical connection and find necessary and sufficient condition when the (v)hv-torsion tensor ${}^*P^i_{jk}$ with respect to the special metrical connection coincides with the (v)hv-torsion P^i_{jk} with respect to general Finsler connection. The relation in hv-curvature tensor, h-curvature tensor and v(h)-torsion tensor with respect to these two connection are also obtained.

1. Introduction

Let $F^n = (M^n, L)$ be an n-dimensional Finsler space equipped with the fundamental function $L(x, y)$. The metric tensor, the angular metric tensor and Cartan tensor are defined by

$$g_{ij} = \frac{1}{2}\dot\partial_i\dot\partial_j L^2, \quad h_{ij} = L\dot\partial_i\dot\partial_j L \quad \text{and} \quad C_{ijk} = \frac{1}{2}\dot\partial_k g_{ij}$$

respectively, where $\dot\partial_i = \partial/\partial y^i$.

A Finsler connection is a triad $F\Gamma = (\Gamma^i_{jk}, N^i_j, C^i_{jk})$, where Γ^i_{jk} are connection coefficients of h-connection, N^i_j are connection coefficients of non-linear connection and C^i_{jk} are connection coefficients of v-connection. For a given connection, the h- and v-covariant derivatives of any vector X^i are given by

$$X^i_{|k} = \delta_k X^i + X^r\Gamma^i_{rk},$$

and

$$X^i|_k = \dot\partial_k X^i + X^r C^i_{rk},$$

where $\delta_k = \partial_k - N^r_k\dot\partial_r$ and $\partial_k = \frac{\partial}{\partial x^k}$.

2010 *Mathematics Subject Classification.* 53B40.
Key words and phrases. Finsler space, Finsler connection, metrical connection.
The first author is supported by UGC, Government of India.

In 2006, H. S. Park et.al. [4] defined a new non-linear connection $\overline{N}^i_j$ with the help of given non-linear connection N^i_j for an (α, β)-metric as

$$\overline{N}^i_j = N^i_j + \nabla_j L\frac{y^i}{L}, \tag{1.1}$$

where '∇_j' denotes the covariant derivative with respect to the associated Riemannian connection.

In 2008, H. G. Nagaraja [3] defined a new non-linear connection ${}^*N^i_j$ with the help of given non-linear connection for Randers space as

$${}^*N^i_j = N^i_j + \frac{L_{|j}y^i}{L}, \tag{1.2}$$

where '$_{|j}$' denote the covariant derivative with respect to Finsler connection $F\Gamma$, and find a new Finsler connection ${}^*F\Gamma = ({}^*\Gamma^i_{jk}, {}^*N^i_j, C^i_{jk})$.

In this paper, we consider a Finsler space F^n admitting the Finsler connection ${}^*F\Gamma$ and we find a relation between $v(hv)$-torsion tensors with respect to these two Finsler connection connections $F\Gamma$ and ${}^*F\Gamma$. We obtain necessary and sufficient condition that two $(v)hv$-torsions coincides. We also find relation in hv-curvature tensor, h-curvature tensor and v(h) torsion tensor with respect to these two Finsler connections.

The Terminology and notion are referred to [2, 5].

2. A special metrical connection

Let $F^n = (M^n, L)$ be an n-dimensional Finsler space and $F\Gamma = (\Gamma^i_{jk}, N^i_j, C^i_{jk})$ be a Finsler connection. Let the Finsler space $F^n = (M^n, L)$ admits a new Finsler connection ${}^*F\Gamma = ({}^*\Gamma^i_{jk}, {}^*N^i_j, C^i_{jk})$, which is $h(h)$-torsion free and non-linear coefficients ${}^*N^i_j$ are given by (1.4). Then we have

$${}^*\delta_k = \partial_k - {}^*N^r_k\dot{\partial}_r = \partial_k - N^r_k\dot{\partial}_r - \frac{L_{|k}y^r}{L}\dot{\partial}_r = \delta_k - \frac{L_{|k}y^r}{L}\dot{\partial}_r. \tag{2.1}$$

The h-covariant derivative of L with respect to ${}^*F\Gamma$ is given by

$$L_{*|k} = {}^*\delta_k L = \delta_k L - \frac{L_{|k}y^r}{L}\dot{\partial}_r L = \delta_k L - L_{|k} = 0. \tag{2.2}$$

Therefore the Finsler connection ${}^*F\Gamma$ is h-metrical. Since ${}^*F\Gamma$ is h-metrical and $h(h)$- torsion ${}^*T^i_{jk}$ is zero, the linear connection coefficients ${}^*\Gamma^i_{jk}$ of ${}^*F\Gamma$ are given in [1] by

$$\Gamma^i_{jk} = g^{ir}\Gamma_{jrk} = \frac{1}{2}g^{ir}[\delta_j g_{rk} + \delta_k g_{rj} - \delta_r g_{jk}]. \tag{2.3}$$

Using (2.1) and $(\dot{\partial}_j g_{ik})y^j = 0$ in (2.3), we have

$${}^*\Gamma^i_{jk} = \Gamma^i_{jk}. \tag{2.4}$$

Thus, the new connection ${}^*F\Gamma = ({}^*\Gamma^i_{jk}, {}^*N^i_j, C^i_{jk})$ reduces to ${}^*F\Gamma = (\Gamma^i_{jk}, {}^*N^i_j, C^i_{jk})$.

The (v)hv-torsion tensor. The (v)hv-torsion tensor P^i_{jk} of a Finsler space with respect to connection $F\Gamma$ is defined by

$$P^i_{jk} = \dot{\partial}_k N^i_j - \Gamma^i_{jk}.$$

Therefore the (v)hv-torsion tensor ${}^*P^i_{jk}$ of a Finsler space with respect to the connection ${}^*F\Gamma$ is given by

$${}^*P^i_{jk} = \dot{\partial}^*_k N^i_j - {}^*\Gamma^i_{jk}.$$

Using (1.2) and (2.4), we get

$$ {}^*P^i_{jk} = P^i_{jk} + l_{k|j}l^i + L_{|j}\frac{h^i_k}{L}. \tag{2.5}$$

Let us suppose ${}^*P^i_{jk} = P^i_{jk}$. Then equation (2.5) implies that $l_{k|j}l^i + L_{|j}\frac{h^i_k}{L} = 0$. Transvecting by y_i and using $h^i_k y_i = 0$, we get $Ll_{k|j} = 0$, which gives $l_{k|j} = 0$.

Conversely, let $l_{k|j} = 0$. Also let us assume that deflection tensor D^i_j for the Finsler connection $F\Gamma$ is zero, i.e. $D^i_j = 0$. Then we get $y^i_{|j} = 0$. Again $l_{k|j} = 0$ and $y^i_{|j} = 0$, implies that $L_{|j} = 0$, and hence equation (2.5) yields ${}^*P^i_{jk} = P^i_{jk}$. Thus, we have:

Theorem 2.1. *Let the Finsler space F^n admits a Finsler connection $F\Gamma$ with zero deflection tensor and a Finsler connection ${}^*F\Gamma$. Then ${}^*P^i_{jk} = P^i_{jk}$ if and only if $l_{k|j} = 0$.*

Rephrasing, the theorem concludes that (v)hv-torsion tensors with respect to Finsler connections $F\Gamma$ and ${}^*F\Gamma$ are equal if and only if covariant differentiation of directional derivative of Fundamental metric function L vanishes.

The hv-curvature tensor. The hv-curvature tensor P^i_{hjk} of a Finsler space with respect to connection $F\Gamma$ is defined by

$$P^i_{hjk} = \dot{\partial}_k \Gamma^i_{hj} - C^i_{hk|j} + C^i_{hm}P^m_{jk}.$$

Therefore the hv-torsion tensor ${}^*P^i_{jk}$ of a Finsler space with respect to connection ${}^*F\Gamma$ is given by

$${}^*P^i_{hjk} = \dot{\partial}^*_k \Gamma^i_{hj} - C^i_{hk|j} + C^i_{hm}\,{}^*P^m_{jk}.$$

Using (2.4) and (2.5) we get

$$ {}^*P^i_{hjk} = P^i_{hjk} + C^i_{hm}h^m_k\frac{L_{|j}}{L}. \tag{2.6}$$

Thus, we have:

Theorem 2.2. *Let the Finsler space F^n admits the Finsler connections $F\Gamma$ and ${}^*F\Gamma$. Then expression for ${}^*P^i_{hjk}$ is given by* (2.6).

Let us suppose that ${}^*P^i_{hjk} = P^i_{hjk}$. Then equation (2.6) implies that

$$C^i_{hm} h^m_k \frac{L_{|j}}{L} = 0,$$

which gives $C^i_{hm} = 0$, i.e. the space is Riemannian. Thus, we have:

Theorem 2.3. *If hv-curvature tensor with respect to Finsler connections $F\Gamma$ and ${}^*F\Gamma$ coincides, then the space will be Riemannian.*

The v(h)-torsion tensor. The v(h)-torsion tensor R^i_{jk} of a Finsler space with respect to connection $F\Gamma$ is defined by

$$R^i_{jk} = \delta_k N^i_j - \delta_j N^i_k.$$

Therefore the v(h)-torsion tensor ${}^*R^i_{jk}$ of a Finsler space with respect to connection ${}^*F\Gamma$ is given by

$${}^*R^i_{jk} = {}^*\delta_k {}^*N^i_j - {}^*\delta_j {}^*N^i_k.$$

Using (1.2) and (2.1) and simplification gives us

$$ {}^*R^i_{jk} = R^i_{jk} + [L_{|j|k} l^i + L_{|j} l^i_{|k} - L_{|k} l^r l_{r|j} l^i - j/k], \tag{2.7}$$

where $-j/k$ denotes interchange of j and k and subtract the terms within the bracket.

Theorem 2.4. *Let the Finsler space F^n admits the Finsler connections $F\Gamma$ and ${}^*F\Gamma$. Then expression for ${}^*R^i_{jk}$ is given by* (2.7).

Equation (2.7) can be re-written as

$$ {}^*R^i_{jk} = R^i_{jk} + [(L_{|j|k} - L_{|k|j}) l^i + L_{|j} \frac{y^i_{|k}}{L} - L_{|k} \frac{y^i_{|j}}{L} + L_{|k} \frac{y^r_{|j}}{L} l^r l^i - L_{|j} \frac{y^r_{|k}}{L} l^r l^i]. \tag{2.8}$$

Let us assume that deflection tensor D^i_j for the Finsler connection $F\Gamma$ is zero i.e. $D^i_j = 0$. Then we get $y^i_{|j} = 0$ and then equation (2.8) becomes

$$ {}^*R^i_{jk} = R^i_{jk} + (L_{|j|k} - L_{|k|j}) l^i. \tag{2.9}$$

Thus, we have:

Corollary 1. *Let the Finsler space F^n admits the Finsler connection $F\Gamma$ with zero deflection tensor and a Finsler connection ${}^*F\Gamma$. Then v(h)-torsion tensor ${}^*R^i_{jk}$ for the connection ${}^*F\Gamma$ is given by* (2.9).

The h-curvature tensor. The h-curvature tensor R^i_{hjk} of a Finsler space with respect to connection $F\Gamma$ is defined by

$$R^i_{hjk} = C^i_{hm} R^m_{jk} + \delta_k \Gamma^i_{hj} + \Gamma^m_{hj} \Gamma^i_{mk} - \delta_j \Gamma^i_{hk} - \Gamma^m_{hk} \Gamma^i_{mj}.$$

Therefore the h-curvature tensor ${}^*R^i_{jk}$ of a Finsler space with respect to connection ${}^*F\Gamma$ is given by

$${}^*R^i_{hjk} = C^i_{hm} {}^*R^m_{jk} + {}^*\delta_k \Gamma^i_{hj} + \Gamma^m_{hj} \Gamma^i_{mk} - {}^*\delta_j \Gamma^i_{hk} - \Gamma^m_{hk} \Gamma^i_{mj}.$$

Using (2.1) and (2.7) and simplifying, we obtain

$$(2.10) \qquad {}^*R^i_{hjk} = R^i_{hjk} - (L_{|k}l^r\dot\partial_r\Gamma^i_{hj} - L_{|j}l^r\dot\partial_r\Gamma^i_{hk}) + C^i_{hm}[L_{|j}l^m_{|k} - L_{|k}l^m_{|j}].$$

Thus, we have:

Theorem 2.5. *Let the Finsler space F^n admits Finsler connections $F\Gamma$ and ${}^*F\Gamma$. Then expression for ${}^*R^i_{hjk}$ is given by* (2.10).

Equation (2.10) can be re-written as

$$(2.11) \qquad {}^*R^i_{hjk} = R^i_{hjk} - L_{|k}l^r\dot\partial_r\Gamma^i_{hj} + L_{|j}l^r\dot\partial_r\Gamma^i_{hk} + C^i_{hm}[L_{|j}\frac{y^m_{|k}}{L} - L_{|k}\frac{y^m_{|j}}{L}].$$

Also let us assume that deflection tensor D^i_j for the Finsler connection $F\Gamma$ is zero, i.e. $D^i_j = 0$. Then we get $y^i_{|j} = 0$ and the equation (2.11) yields

$$(2.12) \qquad {}^*R^i_{hjk} = R^i_{hjk} - (L_{|k}l^r\dot\partial_r\Gamma^i_{hj}) + (L_{|j}l^r\dot\partial_r\Gamma^i_{hk}).$$

Thus, we have:

Corollary 2. *Let the Finsler space F^n admits the Finsler connections $F\Gamma$ with deflection zero and ${}^*F\Gamma$. Then expression for ${}^*R^i_{hjk}$ is given by* (2.12).

References

[1] P. L. Antonelli, R. S. Ingarden, and M. Matsumoto. *The theory of sprays and Finsler spaces with applications in physics and biology*, volume 58 of *Fundamental Theories of Physics*. Kluwer Academic Publishers Group, Dordrecht, 1993.

[2] M. Matsumoto. *Foundations of Finsler geometry and special Finsler spaces*. Kaiseisha Press, Shigaken, 1986.

[3] H. G. Nagaraja. Randers space with special nonlinear connection. *Lobachevskii J. Math.*, 29(1):27–31, 2008.

[4] H. S. Park, H. Y. Park, and B. D. Kim. On a Finsler space with (α, β)-metric and certain metrical non-linear connection. *Commun. Korean Math. Soc.*, 21(1):177–183, 2006.

[5] H. Rund. *The differential geometry of Finsler spaces*. Die Grundlehren der Mathematischen Wissenschaften, Bd. 101. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1959.

Received September 9, 2015.

M. K. Gupta (corresponding author),
Department of Pure and Applied Mathematics,
Guru Ghasidas Vishwavidyalaya,
Bilaspur (C.G.), India
E-mail address: mkgiaps@gmail.com

Anil K. Gupta,
Department of Pure and Applied Mathematics,
Guru Ghasidas Vishwavidyalaya,
Bilaspur (C.G.), India
E-mail address: gupta.anil409@gmail.com