

A certain complete space-like hypersurface in Lorentz manifolds

Hyang Sook Kim and Young-Mi Kim

Abstract

In this paper, we find an upper bound of the squared norm of the second fundamental tensor of a complete space-like hypersurface in a Lorentz space form $M_1^m(c)$ satisfying some curvature conditions. Then it gives naturally an extension of some theorems of Cheng and Nakagawa ([3]), Ishihara ([7]), Li ([8]) and Nishikawa ([9]).

Mathematics Subject Classification: 53C50, 53C25, 53C42.

Key words: space-like hypersurface, constant mean curvature, locally symmetric, maximal space-like, totally geodesic.

1 Introduction

In connection with the negative settlement of the Berstein problem by Calabi ([2]), Cheng-Yau ([4]) and Chouque-Bruhat et al.([6]) proved the following famous theorem independently.

Theorem A *Let M be a complete space-like Lorentz space form $M_1^{n+1}(c)$, $c > 0$. If M is maximal, then it is totally geodesic.*

On the other hand, complete space-like hypersurface with constant mean curvature in a Lorentz space form $M_1^m(c)$ are investigated by many differential geometers in various view points ; for example Akutagawa ([1]), Cheng and Nakagawa ([3]), Li ([8]), Nishikawa([9]) and Ramanathan ([11]). In this paper, we'll give an upper bound of the squared norm of the second fundamental form, of complete space-like hypersurface with constant mean curvature in a Lorentz space form $M_1^m(c)$. Namely, the following assertion is our main theorem.

Main Theorem *Let M' be an $(n + 1)$ -dimensional Lorentz manifold which satisfies the condition (*) and M be a complete space-like hypersurface with constant mean curvature. If M is not maximal and if it satisfies*

$$2nc_2 + c_1 > 0,$$

then there exist a positive constant a_1 depending on c_1, c_2, c_3, h and n such that $h_2 \leq a_1$, where the condition $(*)$ means (5.1), (5.2) and (5.3), h_2 denotes the square norm of the second fundamental form.

As an application of this result, we able to make a generalization of some theorem which are investigated by Cheng and Nakagawa ([3]), Ishihara ([7]), Li ([8]), and Nishikawa ([9]) in new different view point.

2 Definitions

Let $M' = (M', g')$ be a Lorentz manifold with a Lorentz metric g' of signature $(-, +, \dots, +)$. M' has uniquely defined torsion-free affine connection ∇' compatible with the metric g' . M' is called *locally symmetric* if the curvature tensor R' of M' is parallel, that is, $\nabla' R' = 0$. Let M be a hypersurface immersed in M' . M is said to be *space-like* if the Lorentz metric g' of M' induces a Riemannian metric g on M . For a space-like hypersurface M there is naturally defined the second fundamental form (the extrinsic curvature) α of M . M is called *maximal space-like* if the mean(extrinsic) curvature $H = \text{Tr } \alpha$, the trace of α , of M vanishes identically. M is maximal space-like if and only if it is extreme under the variations, with compact support through space-like hypersurfaces, for the induced volume. M is said to be *totally geodesic* (a moment of time symmetry) if the second fundamental form α vanishes identically.

3 Preliminaries

Let M be a space-like hypersurface in a Lorentz $(n + 1)$ -manifold $M' = (M', g')$. We choose a local field of Lorentz orthonormal frames e_0, \dots, e_n are tangent to M' such that, restricted to M , the vectors e_1, \dots, e_n are tangent to M . Here and in the sequel the following convention on the range of indices used throughout this paper, unless otherwise stated:

$$i, j, k, \dots = 1, 2, \dots, n \quad \alpha, \beta, \dots = 0, 1, 2 \dots n$$

Let ω_α be its dual frame field so that the Lorentz metric g' can be written as $g' = -\omega_0^2 + \sum_i \omega_i^2$. then the connection forms $\omega_{\alpha\beta}$ of M' are characterized by the equations

$$(3.1) \quad \begin{aligned} d\omega_i &= -\sum_k \omega_{ik} \wedge \omega_k + \omega_{i0} \wedge \omega_0, \\ d\omega_0 &= -\sum_k \omega_{0k} \wedge \omega_k, \omega_{\alpha\beta} + \omega_{\beta\alpha} = 0. \end{aligned}$$

The curvature forms $\Omega'_{\alpha\beta}$ of M' are given by

$$(3.2) \quad \begin{aligned} \Omega'_{ij} &= d\omega_{ij} + \sum_k \omega_{ik} \wedge \omega_{kj} - \omega_{i0} \wedge \omega_{0j}, \\ \Omega'_{0i} &= d\omega_{0i} + \sum_k \omega_{0k} \wedge \omega_{ki}, \end{aligned}$$

and we have

$$(3.3) \quad \Omega'_{\alpha\beta} = -\frac{1}{2} \sum_{\gamma,\delta} R'_{\alpha\beta\gamma\delta} \omega_\gamma \wedge \omega_\delta,$$

where $R'_{\alpha\beta\gamma\delta}$ are components of the curvature tensor R' of M' . We restrict these forms to M .

Then

$$(3.4) \quad \omega_0 = 0,$$

and the induced Riemannian metric g of M is written as $g = \sum_i \omega_i^2$. From formulas (3.1) \sim (3.4), we obtain the structure equations of M

$$(3.5) \quad \begin{aligned} d\omega_i &= -\sum_k \omega_{ik} \wedge \omega_k, \quad \omega_{ij} + \omega_{ji} = 0, \\ d\omega_{ij} &= -\sum_k \omega_{ik} \wedge \omega_{kj} + \omega_{i0} \wedge \omega_{0j} + \Omega'_{ij}, \\ \Omega_{ij} &= d\omega_{ij} + \sum_k \omega_{ik} \wedge \omega_{kj} = -\frac{1}{2} \sum_{k,l} R_{ijkl} \omega_k \wedge \omega_l, \end{aligned}$$

where Ω_{ij} and R_{ijkl} denote the curvature forms and the components of the curvature tensor R of M , respectively. We can also write

$$(3.6) \quad \omega_{i0} = \sum_j h_{ij} \omega_j,$$

where h_{ij} are components of the second fundamental form $\alpha = \sum_{i,j} h_{ij} \omega_i \otimes \omega_j$ of M . Using (3.6) in (3.5) then gives the Gauss formula

$$(3.7) \quad R_{ijkl} = R'_{ijkl} - (h_{ik}h_{jl} - h_{il}h_{jk}).$$

Let h_{ijk} denote the covariant derivative of h_{ij} so that

$$(3.8) \quad \sum_k h_{ijk} \omega_k = dh_{ij} - \sum_k h_{kj} \omega_{ki} - \sum_k h_{ik} \omega_{kj}.$$

Then, by exterior differentiating (3.6), we obtain the Codazzi equation

$$(3.9) \quad h_{ijk} - h_{ikj} = R'_{0ijk}.$$

From the exterior derivative of (3.8), we define the second covariant derivative of h_{ij} by

$$\sum_l h_{ijkl}\omega_l = dh_{ijk} - \sum_l h_{ijk}\omega_{li} - \sum_l h_{ilk}\omega_{lj} - \sum_l h_{ijl}\omega_{lk}.$$

Then we obtain the Ricci formula

$$(3.10) \quad h_{ijkl} - h_{ijlk} = \sum_m h_{mj}R_{mikl} + \sum_m h_{im}R_{mjkl}.$$

The components of the Ricci tensor S and the scalar curvature r of M are given by

$$\begin{aligned} S_{ij} &= \sum_k R'_{kikj} - hh_{ij} + h_{ij}^2, \\ r &= \sum_{j,k} R_{kjkj} - h^2 + h_2, \end{aligned}$$

where $h = \sum_i h_{ii}$, $h_{ij}^2 = \sum_r h_{ir}h_{rj}$ and $h_2 = \sum_j h_{jj}^2$.

Let us now denote the covariant derivative of $R'_{\alpha\beta\gamma\delta}$, as a curvature tensor of M' , by $R'_{\alpha\beta\gamma\delta;\eta}$. Then restricting on M , $R'_{0ijk;l}$ is given by

$$(3.11) \quad R'_{0ijk;l} = R'_{0ijkl} - R'_{0i0k}h_{jl} - R'_{0ij0}h_{kl} - \sum_m R'_{mijk}h_{ml},$$

where R'_{0ijkl} denotes the covariant derivative of R'_{0ijk} as a tensor on M so that

$$\sum_l R'_{0ijkl}\omega_l = dR'_{0ijk} - \sum_l R'_{0ljk}\omega_{li} - \sum_l R'_{0ilk}\omega_{lj} - \sum_l R'_{0ijl}\omega_{lk}.$$

For the sake of brevity, a tensor h_{ij}^{2m} and a function h_{2m} on M , for any integer $m(\geq 2)$, are introduced as follows:

$$\begin{aligned} h_{ij}^{2m} &= \sum_{i_1, \dots, i_{m-1}} h_{ii_1}^2 h_{i_1 i_2}^2 \cdots h_{i_{m-1} j}^2, \\ h_{2m} &= \sum_i h_{ii}^{2m}. \end{aligned}$$

First of all, let us introduce a fundamental property for the generalized maximal principle due to Omori ([10]) and Yau([13]).

Theorem 3.1 ([10], [13]) *Let M be a complete Riemannian manifold whose Ricci curvature is bounded from below on M . Let F be a C^2 -function bounded from below on M , then for any $\epsilon > 0$, there exists a point p such that*

$$|\nabla F(p)| < \epsilon, \quad \Delta F(p) > -\epsilon \quad \text{and} \quad \inf F + \epsilon > F(p).$$

We also know the following result ([5]).

Theorem 3.2 ([5]) *Let M be a complete Riemannian manifold whose Ricci curvature is bounded from below. Let F be a polynomial of the variable f with constant coefficients such that*

$$(3.12) \quad F(f) = c_0 f^n + c_1 f^{n-1} + \cdots + c_k f^{n-k} + c_{k+1},$$

where $n > 1$, $1 \geq n - k \geq 0$ and $c_0 > c_{k+1}$. If a C^2 -positive function f satisfies

$$\Delta f \geq F(f),$$

then we have

$$F(f_1) \leq 0,$$

where f_1 denotes the supremum of the given function f .

4 The Laplacian operator

Let M be a space-like hypersurface of an $(n + 1)$ -dimensional Lorentz manifold M' . Then the Laplacian Δh_{ij} of the components h_{ij} of α is defined by

$$\Delta h_{ij} = \sum_k h_{ijkk}.$$

From (3.9) we have

$$(4.1) \quad \Delta h_{ij} = \sum_k h_{kijj} + \sum_k R'_{0ijkk},$$

and from (3.10) it follows that

$$(4.2) \quad h_{kijj} = h_{kikj} + \sum_m h_{mi} R_{mkjk} + \sum_m h_{km} R_{mijk}.$$

By using (3.9), replace h_{kikj} in (4.2) by $h_{kkij} + R'_{0kikj}$ and substitute the right hand side of (4.2) into h_{kijj} in (4.1). Then we get

$$(4.3) \quad \begin{aligned} \Delta h_{ij} &= \sum_k (h_{kkij} + R'_{0kikj} + R'_{0ijkk}) \\ &+ \sum_k \left(\sum_m h_{mi} R_{mkjk} + \sum_m h_{km} R_{mijk} \right). \end{aligned}$$

From (3.7), (3.11) and (4.3) we have

$$\begin{aligned}
 (4.4) \quad \Delta h_{ij} &= \sum_k h_{kki j} + \sum_k R'_{0kik;j} + \sum_k R'_{0ijk;k} \\
 &+ \sum_k (h_{kk} R'_{0ij0} + h_{ij} R'_{0k0k}) \\
 &+ \sum_{m,k} (h_{mj} R'_{mki k} + 2h_{mk} R'_{mijk} + h_{mi} R'_{mkjk}) \\
 &- \sum_{m,k} (h_{mi} h_{mj} h_{kk} + h_{km} h_{mj} h_{ik} - h_{km} h_{mk} h_{ij} - h_{mi} h_{mk} h_{kj}).
 \end{aligned}$$

5 Some curvature conditions

Let M' be an $(n+1)$ -dimensional Lorentz manifold and let M be a space-like hypersurface of M' . For a point x in M let $\{e_0, e_1, \dots, e_n\}$ be a local field of orthonormal frames of M' around of x in such a way that, restricted to M , the vectors e_1, \dots, e_n are tangent to M and the other is normal to M . Accordingly, e_1, \dots, e_n are space-like vectors and e_0 is a time-like one. For linearly independent vectors u and v in the tangent space $T_x M'$, by which the non-degenerate plane section is spanned, we denote by $K'(u, v)$ the sectional curvature of the plane section in M' and by R' or $Ric'(u, u)$ the Riemannian curvature tensor on M or the Ricci curvature in the direction of u in M' , respectively. Let us denote by ∇' the Riemannian connection on M' . We assume that the ambient space M' satisfies the following three conditions : For some constants c_1, c_2 and c_3

$$(5.1) \quad K'(u, v) = \frac{c_1}{n},$$

for any space-like vector u and time-like vector v ,

$$(5.2) \quad K'(u, v) \geq c_2,$$

for any space-like vectors u and v

$$(5.3) \quad |\nabla' R'| \leq \frac{c_3}{n}.$$

When M' satisfies the above conditions (5.1), (5.2) and (5.3), it is said simply for M' to satisfy the (*) condition.

Remark 5.1 *It can be easily seen that $c_3=0$, then the ambient space M' is locally symmetric.*

Remark 5.2 *If M is maximal, then the condition (5.1) can be replaced by*

$$(5.4) \quad Ric'(v, v) \geq c_1$$

for any time-like vector v .

If M' satisfies the conditions (5.4), (5.2) and (5.3), it is said simply for M' to satisfy the condition $(*)'$.

Remark 5.3 *If M' is a Lorentz space form $M_1^{n+1}(c)$ of index 1 and of constant curvature c , then it satisfies the condition $(*)'$, where $-\frac{c_1}{n}=c_2=c$.*

Now we assume that the ambient space M' satisfies the condition $(*)$ and the mean curvature of the hypersurface M is constant. Then the Laplacian of the squared norm h_2 of the second fundamental form α of M is given by

$$\begin{aligned}\Delta h_2 &= \Delta\left(\sum_{i,j} h_{ij}h_{ij}\right) = 2\sum_{i,j,k} (h_{ijk}h_{ij})_k = 2\sum_{i,j,k} (h_{ijk}h_{ij} + h_{ijk}h_{ij}) \\ &= 2|\nabla\alpha|^2 + 2\sum_{i,j,k} h_{ijk}h_{ij} = 2|\nabla\alpha|^2 + 2\sum_{i,j} (\Delta h_{ij})h_{ij},\end{aligned}$$

where $\nabla\alpha$ is the covariant derivative of the second fundamental form α and $|\nabla\alpha|$ is the norm of $\nabla\alpha$ which is defined by $\sum_{i,j,k} h_{ijk}h_{ijk}$. Hence, by (4.4) and the assumption $\sum_k h_{kkj}=0$, we have

$$\begin{aligned}\Delta h_2 &= 2|\nabla\alpha|^2 + 2\sum_{i,j} \left\{ \sum_k (R'_{0kik;j} + R'_{0ijk;k}) + \sum_k (h_{kk}R'_{0ij0} + h_{ij}R'_{0k0k}) \right. \\ &\quad \left. + \sum_{k,m} (2h_{km}R'_{mijk} + h_{mj}R'_{mki} + h_{mi}R'_{mkjk}) - hh_{ij}^2 + h_2h_{ij} \right\} h_{ij}.\end{aligned}$$

Thus we get

$$\begin{aligned}(5.5) \quad \Delta h_2 &= 2|\nabla\alpha|^2 + 2\sum_{i,j,k} h_{ij}(R'_{0kik;j} + R'_{0ijk;k}) \\ &\quad + 2\left(\sum_{i,j} hh_{ij}R'_{0ij0} + h_2\sum_k R'_{0k0k}\right) \\ &\quad + 4\left(\sum_{i,j,k,m} h_{ij}h_{km}R'_{mijk} + \sum_{j,k,m} h_{mj}^2R'_{mkjk}\right) - 2(hh_3 - h_2^2),\end{aligned}$$

where we have denoted by $h_{ij}^3 = \sum h_{ir}h_{rj}^2$ and $h_3 = \sum h_{ii}^3$. Since the matrix $H=(h_{ij})$ can be diagonalized, the component of h_{ij} of H can be expressed by

$$(5.6) \quad h_{ij} = \lambda_i\delta_{ij},$$

where λ_i is the principle curvature on M . By definition, we see

$$\lambda_i^2 \leq h_2 = \sum_i \lambda_i^2,$$

and hence we have

$$(5.7) \quad -\sqrt{h_2} \leq \lambda_i \leq \sqrt{h_2},$$

$$(5.8) \quad -h_2 \leq \lambda_i \lambda_j \leq h_2.$$

Now, we estimate (5.5) from above. First, we treat with the second term of (5.5). It is seen that we have

$$\begin{aligned} -2 \sum_{i,j,k} (R'_{0kik;j} + R'_{0ij;k;k}) h_{ij} &= -2 \sum_{j,k} \lambda_j (R'_{0kj;k;j} + R'_{0jj;k;k}) \\ &\leq 2 \sum_{j,k} |\lambda_j| (|R'_{0kj;k;j}| + |R'_{0jj;k;k}|). \end{aligned}$$

So by (5.3) and (5.7) we have

$$(5.9) \quad \text{the second term of (5.5)} \geq -4c_3 \sqrt{h_2}.$$

Next, we consider the third term of (5.4). It is estimated as follows:

$$\begin{aligned} 2 \left(\sum_{i,j} h h_{ij} R'_{0ij0} + h_2 \sum_k R'_{0k0k} \right) &= 2 \sum_k (h \lambda_k R'_{0k0k0} + h_2 \sum_k R'_{0k0k}) \\ &= 2 \sum_k (h_2 - h \lambda_k) R'_{0k0k} = 2 \sum_k (h_2 - h \lambda_k) \frac{c_1}{n}, \end{aligned}$$

where we have used (5.1). Hence we have

$$(5.10) \quad \text{the third term of (5.5)} = \frac{2c_1(nh_2 - h^2)}{n}.$$

It is evident that if the ambient space M' is a Lorentz space form $M_1^{n+1}(c)$ of constant curvature c and if the hypersurface M is *maximal*, then it also holds under (5.4), namely if M' satisfies the condition $(*)'$, then *the third term of (5.5)* $\geq 2c_1 h_2$. Last we estimate the fourth term of (5.5). We have by (5.2)

$$\begin{aligned} 4 \left(\sum_{i,j,k,m} h_{ij} h_{km} R'_{mijk} + \sum_{j,k,m} h_{mj}^2 R'_{mkjk} \right) &= 4 \sum_{j,k} (\lambda_j \lambda_k R'_{kjjk} + \lambda_j^2 R'_{kjkj}) \\ &= 4 \sum_{j,k} (\lambda_j^2 - \lambda_j \lambda_k) R'_{kjkj} = 2 \sum_{j,k} (\lambda_j - \lambda_k)^2 R'_{kjkj} \geq 2c_2 \sum_{j,k} (\lambda_j - \lambda_k)^2. \end{aligned}$$

Accordingly, we obtain

$$(5.11) \quad \text{the fourth term of (5.5)} \geq 4c_2(nh_2 - h^2),$$

where we have used the formula

$$\sum_{j,k} (\lambda_j - \lambda_k)^2 = 2n \sum_j \lambda_j^2 - 2 \sum_{j,k} \lambda_j \lambda_k = 2n \sum_j \lambda_j^2 - 2 \left(\sum_j \lambda_j \right)^2$$

and the definitions of $h_2 = \sum_j \lambda_j^2$ and $h^2 = \left(\sum_j \lambda_j \right)^2$. Thus, substituting (5.9), (5.10) and (5.11) into (5.5), we can prove the following.

Lemma 5.4 *Let M' be an $(n+1)$ -dimensional Lorentz manifold satisfying the condition (*) and M a space-like hypersurface of M' . If its mean curvature is constant, then we have*

$$(5.12) \quad \Delta h_2 \geq -4c_3 \sqrt{h_2} + \frac{2(2nc_2 + c_1)(nh_2 - h^2)}{n} - 2(hh_3 - h_2^2).$$

In particular, if M is maximal, we have

$$\Delta h_2 \geq -4c_3 \sqrt{h_2} + 2(2nc_2 + c_1)h_2 + 2h_2^2.$$

Also, if $M' = M_1^{n+1}(c)$, then we obtain

$$\Delta h_2 \geq 2c(nh_2 - h^2) - 2(hh_3 - h_2^2).$$

6 Proof of Main Theorem

Let M' be an $(n+1)$ -dimensional Lorentz manifold and let M be a complete hypersurface of M' with constant mean curvature. Assume that the ambient space satisfies the condition (*). The condition (*) is defined by (5.1), (5.2) and (5.3). Now, by (5.12) in Lemma 5.1 the function h_2 satisfies

$$\Delta h_2 \geq -4c_3 \sqrt{h_2} + \frac{2(2nc_2 + c_1)(nh_2 - h^2)}{n} - 2(hh_3 - h_2^2).$$

Moreover, we obtain

$$(6.1) \quad -2hh_3 = -2h \sum_i h_{ii}^3 = -2h \sum_j \lambda_j^3 \geq -2h \sum_j \sqrt{h_2}^3 = -2nhh_2 \sqrt{h_2},$$

from which together with (5.12) it follows that

$$(6.2) \quad \Delta h_2 \geq -4c_3 \sqrt{h_2} + 2(2nc_2 + c_1) \left(h_2 - \frac{h^2}{n} \right) - 2nhh_2 \sqrt{h_2} + 2h_2^2.$$

Now we define a non-negative function f by $f^2 = h_2$. Then it turns out to be

$$(6.3) \quad \Delta f^2 \geq 2[f^4 - nhf^3 + (2nc_2 + c_1)f^2 - 2c_3f - \frac{h^2}{n}(2nc_2 + c_1)].$$

Proof of the Main Theorem

Let $\lambda_1, \dots, \lambda_n$ be principal curvatures on M . The Ricci tensor S_{ij} is expressed by

$$S_{ij} = \sum_k (R'_{kikj} - h_{ij}h_{kk} + h_{ik}h_{jk}).$$

So we have

$$S_{jj} \geq (n-1)c_2 - h\lambda_j + \lambda_j^2 \geq (n-1)c_2 - \frac{h^2}{4},$$

which yields the Ricci curvature of M is bounded from below. For the function f defined by $f^2 = h_2$, by (6.3) we have

$$\Delta f^2 \geq F(f^2),$$

where the function $F(x)$ is defined by

$$F(x) = 2[x^2 - nhx^{\frac{3}{2}} + (2nc_2 + c_1)x - 2c_3x^{\frac{1}{2}} - \frac{h^2}{n}(2nc_2 + c_1)].$$

By comparing with (3.12), we get

$n = 2$, $n - k = \frac{1}{2}$, $c_0 = 2$, $c_{k+1} = -\frac{2h^2(2nc_2+c_1)}{n}$, where we have used $2nc_2 + c_1 > 0$. Now we are able to apply Theorem 3.2 to the function f^2 . Then we obtain

$$(6.4) \quad F(f_1^2) \leq 0,$$

where f_1^2 denotes the supremum of the given function f^2 .

We define the function $y = y(x)$ of the variable x by

$$y = y(x) = x^4 - nhx^3 + (2nc_2 + c_1)x^2 - 2c_3x - \frac{h^2}{n}(2nc_2 + c_1).$$

By the assumption $2nc_2 + c_1 > 0$ and the fact that the hypersurface is not maximal, the algebraic equation $y(x) = 0$ with constant coefficients has positive roots because $y(0) < 0$ and it converges to infinity as x tends to infinity. We denote by $\sqrt{a_1}$ ($a_1 > 0$) the minimal root among the positive roots. So it depends only on the constant coefficients, namely, it depends on c_1, c_2, c_3, h and n , and by definition we see that

$$y|_{[0, \sqrt{a_1})} < 0.$$

From the above equation together with (6.4) it follows that we have $0 \leq f_1 \leq \sqrt{a_1}$. Since the squared norm h_2 of the second fundamental form is given by $h_2 = f^2$, we have

$$\sup h_2 = f_1^2 \leq a_1.$$

So we get the conclusion. \square

If the hypersurface M is maximal, then we have by (6.3)

$$\Delta f^2 \geq 2\{f^4 + (2nc_2 + c_1)f^2 - 2c_3f\} = F(f^2),$$

where a non-negative function f is defined by $f^2 = h_2$. By a similar method to the proof of our Main Theorem, we have

$$(6.5) \quad F(f_1^2) \leq 0,$$

where f_1 denotes the supremum of the function f . We define a function y of the variable x by

$$y = y(x) = x\{x^3 + (2nc_2 + c_1)x - 2c_3\}.$$

By the direct calculus, there exists a unique positive root of the equation $y(x) = 0$, say $\sqrt{a_1}$, if $c_3 > 0$.

Corollary 6.1 *Let M' be an $(n+1)$ -dimensional Lorentz manifold which satisfies the condition $(*)$ and let M be a complete space-like maximal hypersurface. If M' is not locally symmetric, then there exists a positive constant a_1 depending on $c_1, c_2, c_3 (> 0), h$ and n such that $h_2 \leq a_1$.*

Remark 6.2 *Corollary 6.1 was proved by Li ([8]) under the additional condition $c_3^2 + \frac{(2nc_2+c_1)^3}{27} < 0$.*

Remark 6.3 *In the case where the ambient space is locally symmetric and it satisfies the condition $(*)'$, the constant a_1 is the positive root of the algebraic equation*

$$F(x^2) = x^2\{x^2 + (2nc_2 + c_1)\} = 0,$$

which yields that if $2nc_2 + c_1 \geq 0$, then $F|(0, \infty) > 0$, which means that we have no positive roots. In the case where $2nc_2 + c_1 < 0$ there exists a unique positive root of the equation $y(x) = 0$, say $\sqrt{a_1}$. In the first case, considering (6.5) we have $f_1 = 0$. By definition of $f_1 = \sup f$ for the non-negative function f , we see that f vanishes identically on M . It yields that M is totally geodesic. So if it satisfies $2nc_2 + c_1 < 0$, then we have $a_1 = -(2nc_2 + c_1)$. This result was derived by Li ([8]). The first assertion of Corollary 6.1 was also proved by Nishikawa([9]). In particular, when $M' = H_1^{n+1}(c)$, this reduces to Ishihara's theorem ([7]).

Acknowledgement.

The first named author was supported by LG Yonam Foundation 2002.

References

- [1] K. Akutagawa, *On space-like hypersurfaces with constant mean curvature in the de sitter space*, Math. Z., 196(1987), 12-19.
- [2] E. Calabi, *Examples of Bernstein problems for some nonlinear equations*, Proc.Pure and Appl. Math., 15(1970), 223-230.
- [3] Q. M. Cheng and H. Nakagawa, *Totally umbilic hypersurfaces*, Hiroshima Math. J., 20(1990), 1-10.

- [4] S. Y. Cheng and S. T. Yau, *Maximal space-like hypersurfaces in the Lorentz-Minkowski spaces*, Ann. of Math., 104(1976), 223-230.
- [5] S. M. Choi, J. H. Kwon and Y. J. Suh, *A Liouville type theorem for complete Riemannian manifolds*, Bull. Korean Math.Soc., 35(1998), 301-309.
- [6] Y. Chouque-Bruhat, A. E. Fisher and J. E. Marsdan, *Maximal hypersurfaces and positive mass*, Proc.of the E.Fermi Summer School of the Italian Physical Society, J.Ehlers ed.North-holland, 1979.
- [7] T.Ishihara, *Maximal space-like submanifolds of pseudo-hyperbolic space with second fundamental form of maximal length*, Preprint.
- [8] H. Li, *On complete maximal space-like hypersurfaces in a Lorentzian manifold*,Soochow J. Math., 23(1997), 79-89.
- [9] S. Nishikawa, *On maximal space-like hypersurfaces in a Lorentzian manifold*,Nagoya Math. J.,95(1984), 117-124.
- [10] H. Omori, *Isometric immersions of Riemannian manifolds*, J. Math. Soc. Japan, 19(1967), 205-211.
- [11] J. Ramanathan, *Complete space-like hypersurfaces of constant mean curvature in de Sitter space*, Indiana Univ. Math. J., 36(1987), 349-359.
- [12] J. A. Wolf, *Spaces of Constant Curvature*, McGrew-Hill, New-York ,1967.
- [13] S. T. Yau, *Harmonic functions on complete Riemannian manifolds*, Comm. Pure and Appl. Math.,28(1975),201-228 Springer-Verlag 1987.

Hyang Sook Kim

Computational Mathematics, School of Computer Aided Science, Inje University,
Kimhae 621-749, Korea.

e-mail address: mathkim@inje.ac.kr

Young-Mi Kim

Department of Mathematics, Silla University, Busan 617-736, Korea.

e-mail address: ymkim@silla.ac.kr