

**ON SOME ENTIRE MODULAR FORMS OF WEIGHTS 5
 AND 6 FOR THE CONGRUENCE GROUP $\Gamma_0(4N)$**

G. LOMADZE

ABSTRACT. Two entire modular forms of weight 5 and two of weight 6 for the congruence subgroup $\Gamma_0(4N)$ are constructed, which will be useful for revealing the arithmetical sense of additional terms in formulas for the number of representations of positive integers by quadratic forms in 10 and 12 variables.

In this paper N, a, k, n, r, s, t denote positive integers; u are odd positive integers; $H, c, g, h, j, m, \alpha, \beta, \gamma, \delta, \xi, \eta$ are integers; A, B, C, D are complex numbers, and $z, \tau (\operatorname{Im} \tau > 0)$ are complex variables. Further, $(\frac{h}{u})$ is the generalized Jacobi symbol, $\binom{n}{t}$ a binomial coefficient, $\varphi(k)$ Euler's function, and $e(z) = \exp 2\pi iz$, $\eta(\gamma) = 1$ if $\gamma \geq 0$ and $\eta(\gamma) = -1$ if $\gamma < 0$.

Let

$$\begin{aligned}\Gamma &= \left\{ \frac{\alpha\tau + \beta}{\gamma\tau + \delta} \mid \alpha\delta - \beta\gamma = 1 \right\}, \\ \Gamma_0(4N) &= \left\{ \frac{\alpha\tau + \beta}{\gamma\tau + \delta} \in \Gamma \mid \gamma \equiv 0 \pmod{4N} \right\}.\end{aligned}$$

Definition. We shall say that a function F defined on $\mathcal{H} = \{\tau \in \mathbb{C} \mid \operatorname{Im} \tau > 0\}$ is an entire modular form of weight r and character $\chi(\delta)$ for the congruence subgroup $\Gamma_0(4N)$ if

- 1) F is regular on \mathcal{H} ,
- 2) for all substitutions from $\Gamma_0(4N)$ and all $\tau \in \mathcal{H}$

$$F\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) = \chi(\delta)(\gamma\tau + \delta)^r F(\tau),$$

1991 *Mathematics Subject Classification.* 11F11, 11F03, 11F27.

3) in the neighborhood of the point $\tau = i\infty$

$$F(\tau) = \sum_{m=0}^{\infty} A_m e(m\tau),$$

4) for all substitutions from Γ , in the neighborhood of each rational point $\tau = -\frac{\delta}{\gamma}$ ($\gamma \neq 0$, $(\gamma, \delta) = 1$)

$$(\gamma\tau + \delta)^r F(\tau) = \sum_{m=0}^{\infty} A'_m e\left(\frac{n}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right).$$

For $\eta \neq 0$, ξ, g, h, N with $\xi g + \eta h + \xi\eta N \equiv 0 \pmod{2}$, put

$$S_{gh} \left(\begin{array}{c} \xi \\ \eta \end{array}; c, N \right) = \sum_{\substack{m \pmod{N|\eta|} \\ m \equiv c \pmod{N}}} (-1)^{h(m-c)/N} e\left(\frac{\xi}{2N\eta} \left(m + \frac{g}{2}\right)^2\right).$$

It is known ([2] p. 323, formula (2.4)–(2.6)) that

$$\begin{aligned} S_{g+2j,h} \left(\begin{array}{c} \xi \\ \eta \end{array}; c, N \right) &= S_{gh} \left(\begin{array}{c} \xi \\ \eta \end{array}; c+j, N \right), \\ S_{g,h+2j} \left(\begin{array}{c} \xi \\ \eta \end{array}; c, N \right) &= S_{gh} \left(\begin{array}{c} \xi \\ \eta \end{array}; c, N \right), \\ S_{gh} \left(\begin{array}{c} \xi \\ \eta \end{array}; c+N_j, N \right) &= (-1)^{hj} S_{gh} \left(\begin{array}{c} \xi \\ \eta \end{array}; c, N \right). \end{aligned} \quad (1)$$

Let

$$\begin{aligned} \vartheta_{gh}(z|\tau; c, N) &= \\ &= \sum_{m \equiv c \pmod{N}} (-1)^{h(m-c)/N} e\left(\frac{1}{2N} \left(m + \frac{g}{2}\right)^2 \tau\right) e\left(\left(m + \frac{g}{2}\right) z\right), \end{aligned} \quad (2)$$

hence

$$\begin{aligned} \frac{\partial^n}{\partial z^n} \vartheta_{gh}(z|\tau; c, N) &= (\pi i)^n \sum_{m \equiv c \pmod{N}} (-1)^{h(m-c)/N} (2m+g)^n \times \\ &\quad \times e\left(\frac{1}{2N} \left(m + \frac{g}{2}\right)^2 \tau\right) e\left(\left(m + \frac{g}{2}\right) z\right). \end{aligned} \quad (3)$$

Put

$$\begin{aligned} \vartheta_{gh}^{(n)}(\tau; c, N) &= \frac{\partial^n}{\partial z^n} \vartheta_{gh}(z|\tau; c, N) \Big|_{z=0}, \\ \vartheta_{gh}^{(0)}(\tau; c, N) &= \vartheta_{gh}(\tau; c, N) = \vartheta_{gh}(0|\tau; c, N). \end{aligned} \quad (4)$$

It is known ([2], p.318, formula (1.3); p. 321, formula (1.12); p.327, formulas (3.9), (3.5), (3.3), (3.7); p.324, formula (2.16); p.327, formulas (3.10), (3.11)) that

$$\vartheta_{g,h+2j}(z|\tau; c, N) = \vartheta_{gh}(z|\tau; c, N); \quad (5)$$

$$\begin{aligned} \vartheta_{gh}\left(\frac{z}{\tau} \mid -\frac{1}{\tau}; c, N\right) &= \left(-\frac{i\tau}{N}\right)^{1/2} e\left(\frac{Nz^2}{2\tau}\right) \times \\ &\times \sum_{H \bmod N} e\left(-\frac{1}{N}\left(c + \frac{g}{2}\right)\left(H + \frac{h}{2}\right)\right) \vartheta_{hg}(z|\tau; H, N); \end{aligned} \quad (6)$$

$$\begin{aligned} \vartheta_{gh}\left(\frac{z}{\gamma\tau + \delta} \mid \frac{\alpha\tau + \beta}{\gamma\tau + \delta}; c, N\right) &= \left(\frac{-i(\gamma\tau + \delta) \operatorname{sgn} \gamma}{N|\gamma|}\right)^{\frac{1}{2}} e\left(\frac{N\gamma z^2}{2(\gamma\tau + \delta)}\right) \times \\ &\times \sum_{H \bmod N} \varphi_{g'gh}(c, H; N) \vartheta_{g'h'}(z|\tau; H, N) \quad (\gamma \neq 0), \end{aligned} \quad (7)$$

where

$$g' = \alpha g + \gamma h + \alpha\gamma N, \quad h' = \beta g + \delta h + \beta\delta N, \quad (8)$$

$$\begin{aligned} \varphi_{g'gh}(c, H; N) &= e\left(-\frac{\beta\delta}{2N}\left(H + \frac{g'}{2}\right)^2\right) e\left(\frac{\beta}{N}\left(c + \frac{g}{2}\right)\left(H + \frac{g'}{2}\right)\right) \times \\ &\times S_{g-\delta g', h+\beta g'}\left(\frac{\alpha}{\gamma}; c - \delta H, N\right); \end{aligned} \quad (9)$$

$$\begin{aligned} \vartheta_{gh}(z|\tau + \beta; c, N) &= e\left(\frac{\beta}{2N}\left(c + \frac{g}{2}\right)^2\right) \vartheta_{g,h+\beta g+\beta N}(z|\tau; c, N), \\ \vartheta_{gh}(-z|\tau - \beta; c, N) &= \\ &= e\left(-\frac{\beta}{2N}\left(c + \frac{g}{2}\right)^2\right) \vartheta_{-g,-h+\beta g-\beta N}(z|\tau; -c, N). \end{aligned} \quad (10)$$

From (5) and (10), according to the notations (4), it follows that

$$\begin{aligned} \vartheta_{g,h+2j}(\tau; c, N) &= \vartheta_{gh}(\tau; c, N), \\ \vartheta_{g,h+2j}^{(n)}(\tau; c, N) &= \vartheta_{gh}^{(n)}(\tau; c, N); \end{aligned} \quad (11)$$

$$\begin{aligned} \vartheta_{gh}^{(n)}(\tau + \beta; c, N) &= e\left(\frac{\beta}{2N}\left(c + \frac{g}{2}\right)^2\right) \vartheta_{g,h+\beta g+\beta N}^{(n)}(\tau; c, N), \\ \vartheta_{gh}^{(n)}(\tau - \beta; c, N) &= \\ &= (-1)^n e\left(-\frac{\beta}{2N}\left(c + \frac{g}{2}\right)^2\right) \vartheta_{-g,-h+\beta g-\beta N}^{(n)}(\tau; -c, N). \end{aligned} \quad (12)$$

From (2) and (3), according to the notations (4), in particular, it follows

that

$$\begin{aligned} \vartheta_{gh}(\tau; 0, N) &= \sum_{m=-\infty}^{\infty} (-1)^{hm} e\left(\frac{1}{2N}\left(Nm + \frac{g}{2}\right)^2 \tau\right), \\ \vartheta_{gh}^{(n)}(\tau; 0, N) &= \\ &= (\pi i)^n \sum_{m=-\infty}^{\infty} (-1)^{hm} (2Nm + g)^n e\left(\frac{1}{2N}\left(Nm + \frac{g}{2}\right)^2 \tau\right). \end{aligned} \quad (13)$$

1.

Lemma 1. *For $n \geq 0$*

$$\begin{aligned} \vartheta_{gh}^{(n)}\left(-\frac{1}{\tau}; c, N\right) &= (Ni)^n \left(-\frac{i\tau}{N}\right)^{(2n+1)/2} \times \\ &\times \sum_{H \bmod N} e\left(-\frac{1}{N}(c + \frac{g}{2})(H + \frac{h}{2})\right) \times \\ &\times \left\{ \vartheta_{hg}^{(n)}(\tau; H, N) + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{hg}^{(n-t)}(\tau; H, N) \right\}, \end{aligned}$$

where

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)! \left(\frac{N\pi i}{\tau}\right)^k & \text{if } t = 2k \\ 0 & \text{if } t \neq 2k \end{cases} \quad (t = 1, 2, \dots, n; k = 1, 2, \dots, t). \quad (1.1)$$

Proof. From (6), by Leibniz's formula, we obtain

$$\begin{aligned} \frac{\partial^n}{\partial z^n} \vartheta_{gh}\left(\frac{z}{\tau} - \frac{1}{\tau}; c, N\right) &= \tau^n \left(\frac{-i\tau}{N}\right)^{1/2} \frac{\partial^n}{\partial z^n} \left\{ e\left(\frac{Nz^2}{2\tau}\right) \times \right. \\ &\times \sum_{H \bmod N} e\left(-\frac{1}{N}(c + \frac{g}{2})(H + \frac{h}{2})\right) \vartheta_{hg}(z|\tau; H, N) \Big\} = \\ &= (Ni)^n \left(\frac{-i\tau}{N}\right)^{(2n+1)/2} \times \\ &\times \sum_{H \bmod N} e\left(-\frac{1}{N}(c + \frac{g}{2})(H + \frac{h}{2})\right) \left\{ e\left(\frac{Nz^2}{2\tau}\right) \frac{\partial^n}{\partial z^n} \vartheta_{hg}(z|\tau; H, N) \right\} + \\ &+ \sum_{t=1}^n \binom{n}{t} \frac{\partial^n}{\partial z^n} e\left(\frac{Nz^2}{2\tau}\right) \frac{\partial^{n-t}}{\partial z^{n-t}} \vartheta_{hg}(z|\tau; H, N). \end{aligned} \quad (1.2)$$

According to formulae (a) and (b) of [1], p. 37¹

$$\frac{\partial^t}{\partial z^t} e\left(\frac{Nz^2}{2\tau}\right) = A_{t1} e\left(\frac{Nz^2}{2\tau}\right) + \frac{A_{t2}}{2!} e\left(\frac{Nz^2}{2\tau}\right) + \cdots + \frac{A_{tt}}{t!} e\left(\frac{Nz^2}{2\tau}\right)$$

$$(t = 1, 2, \dots, n),$$

where

$$A_{tk} = \frac{\partial^t}{\partial z^t} \left(\frac{N\pi iz^2}{\tau} \right)^k - k \left(\frac{N\pi iz^2}{\tau} \right) \frac{\partial^t}{\partial z^t} \left(\frac{N\pi iz^2}{\tau} \right)^{k-1} +$$

$$+ \frac{k(k-1)}{2!} \left(\frac{N\pi iz^2}{\tau} \right)^2 \frac{\partial^t}{\partial z^t} \left(\frac{N\pi iz^2}{\tau} \right)^{k-2} +$$

$$+ \cdots + (-1)^{k-1} k \left(\frac{N\pi iz^2}{\tau} \right)^{k-1} \frac{\partial^t}{\partial z^t} \left(\frac{N\pi iz^2}{\tau} \right) \quad (k = 1, 2, \dots, t),$$

hence

$$\frac{\partial^t}{\partial z^t} e\left(\frac{Nz^2}{2\tau}\right) \Big|_{z=0} = \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \quad (t = 1, 2, \dots, n) \quad (1.3)$$

and

$$A_{tk} \Big|_{z=0} = \frac{\partial^t}{\partial z^t} \left(\frac{N\pi iz^2}{\tau} \right)^k \Big|_{z=0} \quad (k = 1, 2, \dots, t). \quad (1.4)$$

Thus, in view of notations (4), the lemma follows from (1.2)–(1.4). \square

Lemma 2. *If $\gamma \neq 0$, then for $n \geq 0$*

$$\vartheta_{gh}^{(n)}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; c, N\right) = (N|\gamma|i \operatorname{sgn} \gamma)^n \left(-i(\gamma\tau + \delta)\frac{\operatorname{sgn} \gamma}{N|\gamma|}\right)^{(2n+1)/2} \times$$

$$\times \sum_{H \bmod N} \varphi_{g'gh}(c, H; N) \left\{ \vartheta_{g'h'}^{(n)}(\tau; H, N) + \right.$$

$$\left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{g'h'}^{(n-t)}(\tau; H, N) \right\},$$

where $g' h'$ and $\varphi_{g'gh}(c, H; N)$ are defined by the formulas (8) and (9),

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)! \left(\frac{N\gamma\pi i}{\gamma\tau+\delta} \right)^k & \text{if } t = 2k \\ 0 & \text{if } t \neq 2k \end{cases}$$

$$(t = 1, 2, \dots, n; k = 1, 2, \dots, t).$$

¹Page 75 in the Russian version of [1] published in 1933.

Proof. From (7), according to Leibnitz's formula, we obtain

$$\begin{aligned}
& \frac{\partial^n}{\partial z^n} \vartheta_{gh} \left(\frac{z}{\gamma\tau + \delta} \middle| \frac{\alpha\tau + \beta}{\gamma\tau + \delta}; c, N \right) = (\gamma\tau + \delta)^n \left(-i(\gamma\tau + \delta) \frac{\operatorname{sgn} \gamma}{N|\gamma|} \right)^{1/2} \times \\
& \times \frac{\partial^n}{\partial z^n} \left\{ e \left(\frac{N\gamma z^2}{2(\gamma\tau + \delta)} \right) \sum_{H \bmod N} \varphi_{g'gh}(c, H; N) \vartheta_{g'h'}(z|\tau; H, N) \right\} = \\
& = (N|\gamma|i \operatorname{sgn} \gamma)^n \left(-i(\gamma\tau + \delta) \frac{\operatorname{sgn} \gamma}{N|\gamma|} \right)^{(2n+1)/2} \times \\
& \times \sum_{H \bmod N} \varphi_{g'gh}(c, H; N) \left\{ e \left(\frac{N\gamma z^2}{2(\gamma\tau + \delta)} \right) \frac{\partial^n}{\partial z^n} \vartheta_{g'h'}(z|\tau; H, N) + \right. \\
& \left. + \sum_{t=1}^n \binom{n}{t} \frac{\partial^t}{\partial z^t} e \left(\frac{N\gamma z^2}{2(\gamma\tau + \delta)} \right) \frac{\partial^{n-t}}{\partial z^{n-t}} \vartheta_{g'h'}(z|\tau; H, N) \right\}. \quad (1.5)
\end{aligned}$$

As in Lemma 1, but with $e \left(\frac{N\gamma z^2}{2(\gamma\tau + \delta)} \right)$ instead of $e \left(\frac{Nz^2}{2\tau} \right)$, we have

$$\frac{\partial^t}{\partial z^t} e \left(\frac{N\gamma z^2}{2(\gamma\tau + \delta)} \right) \Big|_{z=0} = \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \quad (t = 1, 2, \dots, n) \quad (1.6)$$

and

$$A_{tk} \Big|_{z=0} = \frac{\partial^t}{\partial z^t} \left(\frac{N\gamma\pi iz}{\gamma\tau + \delta} \right)^{2k} \Big|_{z=0} \quad (k = 1, 2, \dots, t). \quad (1.7)$$

Thus, according to the notations (4), the lemma follows from (1.5)-(1.7). \square

Lemma 3. *If g is even, then for $n \geq 0$ and all substitutions from $\Gamma_0(4N)$ we have*

$$\begin{aligned}
& \vartheta_{gh}^{(n)} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N \right) = \\
& = (\operatorname{sgn} \delta)^n i^{(2n+1)\eta(\gamma)(\operatorname{sgn} \delta - 1)/2} i^{(1-|\delta|)/2} \left(\frac{2\beta N \operatorname{sgn} \delta}{|\delta|} \right) \times \\
& \times (\gamma\tau + \delta)^{(2n+1)/2} e \left(-\frac{\alpha\gamma\delta^2 h^2}{16N} \right) e \left(\frac{\beta\delta g^2}{4} \frac{\delta^{2\varphi(2N)-2}}{4N} \right) \vartheta_{\alpha g, h}^{(n)}(\tau; 0, 2N).
\end{aligned}$$

Proof. 1) Let $\gamma \neq 0$. In [4] (p.18, formula (5.1)) it is shown that

$$\begin{aligned}
& S_{g0} \left(\frac{\beta}{\delta}; 0, 2N \right) = \\
& = e \left(\frac{\beta\delta g^2}{4} \frac{\delta^{2\varphi(2n)-2}}{4N} \right) i^{(1-|\delta|)/2} \left(\frac{2\beta N \operatorname{sgn} \delta}{|\delta|} \right) |\delta|^{1/2}. \quad (1.8)
\end{aligned}$$

Replacing $\alpha, \beta, \gamma, \delta, \tau, c, N$ by $\beta, -\alpha, \delta, -\gamma, \tau', 0, 2N$ in Lemma 2, we obtain

$$\begin{aligned} \vartheta_{gh}^{(n)}\left(\frac{\beta\tau' - \alpha}{\delta\tau' - \gamma}; 0, 2N\right) &= (2N|\delta|i \operatorname{sgn} \delta)^n \left(-i(\delta\tau' - \gamma)\frac{\operatorname{sgn} \delta}{2N|\delta|}\right)^{(2n+1)/2} \times \\ &\quad \times \sum_{H \bmod 2N} \varphi_{h'gh}(0, H; 2N) \left\{ \vartheta_{h', \alpha g}^{(n)}(\tau'; H, 2N) + \right. \\ &\quad \left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{h', \alpha g}^{(n-t)}(\tau'; H, 2N) \right\}, \end{aligned} \quad (1.9)$$

where by (8),(9),(11) and (1)

$$h' = \beta g + \delta h + 2\beta\delta N, \quad (1.10)$$

$$\varphi_{h'gh}(0, H; 2N) = e\left(-\frac{\alpha\gamma h'^2}{16N}\right) e\left(-\frac{\alpha g}{4N}\left(H + \frac{h'}{2}\right)\right) S_{g0}\left(\frac{\beta}{\delta}; 0, 2N\right), \quad (1.11)$$

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)! \left(\frac{2N\delta\pi i}{\delta\tau' - \gamma}\right)^k & \text{if } t = 2k \\ 0 & \text{if } t \neq 2k \end{cases} \quad (k = 1, 2, \dots, t). \quad (1.12)$$

Writing $-\frac{1}{\tau}$ instead of τ' in (1.9) and (1.12), according to (1.11), we obtain

$$\begin{aligned} \vartheta_{gh}^{(n)}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N\right) &= (2Ni)^n (|\delta| \operatorname{sgn} \delta)^n \left(\frac{-i(-\frac{\delta}{\tau} - \gamma) \operatorname{sgn} \delta}{2N|\delta|}\right)^{(2n+1)/2} \times \\ &\quad \times e\left(-\frac{\alpha\gamma h'^2}{16N}\right) e\left(\frac{\beta\delta g^2}{4} \frac{\delta^{2\varphi(2N)-2}}{4N}\right) i^{(1-|\delta|)/2} \left(\frac{2\beta N \operatorname{sgn} \delta}{|\delta|}\right) |\delta|^{1/2} \times \\ &\quad \times \sum_{H \bmod 2N} e\left(-\frac{\alpha g}{4N}\left(H + \frac{h'}{2}\right)\right) \left\{ \vartheta_{h', \alpha g}^{(n)}\left(-\frac{1}{\tau}; H, 2N\right) + \right. \\ &\quad \left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{h', \alpha g}^{(n-t)}\left(-\frac{1}{\tau}; H, 2N\right) \right\}, \end{aligned} \quad (1.13)$$

where

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)! \left(\frac{2N\delta\pi i\tau}{\gamma\tau + \delta}\right)^k & \text{if } t = 2k \\ 0 & \text{if } t \neq 2k \end{cases} \quad (k = 1, 2, \dots, t).$$

Writing $\alpha g, h', -\frac{1}{\tau}, 0, 2N$ instead of g, h, τ, c, N in Lemma 1, we obtain

$$\begin{aligned} \vartheta_{\alpha g, h'}^{(n)}(\tau; 0, 2N) &= (2Ni)^n \left(\frac{i}{2N\tau} \right)^{(2n+1)/2} \sum_{H \bmod 2N} e\left(-\frac{\alpha g}{4N}\left(H + \frac{h'}{2}\right)\right) \times \\ &\quad \times \left\{ \vartheta_{h', \alpha g}^{(n)}\left(-\frac{1}{\tau}; H, 2N\right) + \right. \\ &\quad \left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{h', \alpha g}^{(n-t)}\left(-\frac{1}{\tau}; H, 2N\right) \right\}, \end{aligned} \quad (1.14)$$

where

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)!(-2N\pi i\tau)^k & \text{if } t = 2k \\ 0 & \text{if } t \neq 2k \end{cases} \quad (k = 1, 2, \dots, t).$$

From (1.14) it follows that

$$\begin{aligned} &\sum_{H \bmod 2N} e\left(-\frac{\alpha g}{4N}\left(H + \frac{h'}{2}\right)\right) \left\{ \vartheta_{h', \alpha g}^{(n)}\left(-\frac{1}{\tau}; H, 2N\right) + \right. \\ &\quad \left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{h', \alpha g}^{(n-t)}\left(-\frac{1}{\tau}; H, 2N\right) \right\} = \\ &= (2\pi i)^{-n} (-2N\pi i\tau)^{(2n+1)/2} \vartheta_{\alpha g, h'}^{(n)}(\tau; 0, 2N). \end{aligned} \quad (1.15)$$

From (1.13) and (1.15) we obtain

$$\begin{aligned} \vartheta_{gh}^{(n)}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N\right) &= (-2N\pi i\tau)^{(2n+1)/2} (|\delta| \operatorname{sgn} \delta)^n \times \\ &\quad \times \left(\frac{-i(-\frac{\delta}{\tau} - \gamma) \operatorname{sgn} \delta}{2N|\delta|} \right)^{(2n+1)/2} i^{(1-|\delta|)/2} e\left(-\frac{\alpha\gamma h'^2}{16N}\right) \times \\ &\quad \times e\left(\frac{\beta\delta g^2}{4} \frac{\delta^{2\varphi(2N)-2}}{4N}\right) \left(\frac{2\beta N \operatorname{sgn} \delta}{|\delta|} \right) |\delta|^{1/2} \vartheta_{\alpha g, h'}^{(n)}(\tau; 0, 2N). \end{aligned} \quad (1.16)$$

In [3] (p.85, formula (6.16) with $2N$ instead of N) it is shown that

$$\begin{aligned} &\left(\frac{-i(-\frac{\delta}{\tau} - \gamma) \operatorname{sgn} \delta}{2N|\delta|} \right)^{1/2} (-2N\pi i\tau)^{1/2} = \\ &= i^{\operatorname{sgn} \gamma (\operatorname{sgn} \delta - 1)/2} \left(\frac{\gamma\tau + \delta}{|\delta|} \right)^{1/2}. \end{aligned} \quad (1.17)$$

From (1.16) and (1.17) it follows that

$$\begin{aligned} & \vartheta_{gh}^{(n)}\left(\frac{\alpha\tau+\beta}{\gamma\tau+\delta}; 0, 2N\right) = \\ &= (|\delta| \operatorname{sgn} \delta)^n \left(i^{\operatorname{sgn} \gamma (\operatorname{sgn} \delta - 1)} \frac{\gamma\tau+\delta}{|\delta|}\right)^{(2n+1)/2} i^{(1-|\delta|)/2} \times \\ & \times e\left(-\frac{\alpha\gamma h'^2}{16N}\right) e\left(\frac{\beta\delta g^2}{4} \frac{\delta^{2\varphi(2N)-2}}{4N}\right) \left(\frac{2\beta N \operatorname{sgn} \delta}{|\delta|}\right) |\delta|^{1/2} \vartheta_{\alpha g, h'}^{(n)}(\tau; 0, 2N). \end{aligned}$$

Thus, in view of (1.10) and (1.11), the lemma is proved for $\gamma \neq 0$.

2) Let $\gamma = 0$. Then $\alpha = \delta = 1$ or $\alpha = \delta = -1$ in (12). Putting $c = 0$ in (12) and writing $2N$ instead of N , by (11), we obtain

$$\begin{aligned} \vartheta_{gh}^{(n)}(\tau + \beta; 0, 2N) &= e\left(\frac{\beta g^2}{16N}\right) \vartheta_{gh}^{(n)}(\tau; 0, 2N), \\ \vartheta_{gh}^{(n)}(\tau - \beta; 0, 2N) &= (-1)^n e\left(-\frac{\beta g^2}{16N}\right) \vartheta_{-g, h}^{(n)}(\tau; 0, 2N), \end{aligned}$$

i.e., the lemma is also proved for $\gamma = 0$. \square

Lemma 4. *If $\gamma \neq 0$, then for $n \geq 0$*

$$\begin{aligned} & (\gamma\tau + \delta)^{(2n+1)/2} \vartheta_{gh}^{(n)}(\tau; 0, 2N) = \\ &= e\left((2n+1) \operatorname{sgn} \gamma / 8\right) (2N|\gamma|)^{-1/2} (-i \operatorname{sgn} \gamma)^n \times \\ & \times \sum_{H \bmod 2N} \varphi_{g'gh}(0, H; 2N) \left\{ \vartheta_{g'h'}^{(n)}\left(\frac{\alpha\tau+\beta}{\gamma\tau+\delta}; H, 2N\right) + \right. \\ & \left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{g'h'}^{(n-t)}\left(\frac{\alpha\tau+\beta}{\gamma\tau+\delta}; H, 2N\right) \right\}, \end{aligned}$$

where

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)! (-2N\gamma\pi i(\gamma\tau + \delta))^k & \text{if } t = 2k \\ 0 & \text{if } t \neq 2k \end{cases} \quad (t = 1, 2, \dots; k = 1, 2, \dots, t). \quad (1.18)$$

Remark. From (1.18) it follows that

$$\sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} = \begin{cases} \frac{A_{t,t/2}}{(t/2)!} \Big|_{z=0} & \text{if } 2|t \\ 0 & \text{if } 2\nmid t. \end{cases} \quad (1.19)$$

Proof. Replacing $\alpha, \beta, \gamma, \delta, \tau, c, N$ by $\delta, -\beta, -\gamma, \alpha, \tau', 0, 2N$ in Lemma 2, we obtain

$$\begin{aligned} & \vartheta_{gh}^{(n)}\left(\frac{\delta\tau' - \beta}{-\gamma\tau' + \alpha}; 0, 2N\right) = \\ & = (-2N|\gamma|i \operatorname{sgn} \gamma)^n \left(i(-\gamma\tau' + \alpha)\frac{\operatorname{sgn} \gamma}{2N|\gamma|}\right)^{(2n+1)/2} \times \\ & \quad \times \sum_{H \bmod 2N} \varphi_{g'gh}(0, H; 2N) \left\{ \vartheta_{g'h'}^{(n)}(\tau'; H, 2N) + \right. \\ & \quad \left. + \sum_{t=1}^n \binom{n}{t} \sum_{k=1}^t \frac{A_{tk}}{k!} \Big|_{z=0} \cdot \vartheta_{g'h'}^{(n-t)}(\tau'; H, 2N) \right\}, \end{aligned} \quad (1.20)$$

where

$$g' = \delta g - \gamma h - 2\gamma\delta N, \quad h' = -\beta g + \alpha h - 2\alpha\beta N, \quad (1.21)$$

$$\begin{aligned} \varphi_{g'gh}(0, H; 2N) & = e\left(-\frac{\alpha\beta}{4N}(H + \frac{g'}{2})^2\right) e\left(-\frac{\beta g}{4N}(H + \frac{g'}{2})\right) \times \\ & \quad \times S_{g-\alpha g', h-\beta g'}\left(\frac{\delta}{-\gamma}; -\alpha H, 2N\right), \end{aligned} \quad (1.22)$$

$$A_{tk} \Big|_{z=0} = \begin{cases} (2k)! \left(\frac{2N\gamma\pi i}{\gamma\tau' - \alpha}\right)^k & \text{if } t = 2k, \\ 0 & \text{if } t \neq 2k. \end{cases} \quad (1.23)$$

In [3, p.87, formula (6.23)] (with $2N$ instead of N) it is shown that

$$\left(\frac{i \operatorname{sgn} \gamma}{2N|\gamma|(\gamma\tau + \delta)}\right)^{1/2} = \frac{e(\operatorname{sgn} \gamma/8)}{(2N|\gamma|)^{1/2}(\gamma\tau + \delta)^{1/2}}. \quad (1.24)$$

Taking $\frac{\alpha\tau + \beta}{\gamma\tau + \delta}$ instead of τ' in (1.20) and (1.23) and using (1.24), we complete the proof of the lemma. \square

2.

Lemma 5. *For a given N let*

$$\begin{aligned} \Psi_1(\tau) & = \Psi_1(\tau; g_1, g_2, h_1, h_2, c_1, c_2, N_1, N_2) = \\ & = \frac{1}{N_1} \vartheta_{g_1 h_1}'''(\tau; c_1, 2N_1) \vartheta_{g_2 h_2}'(\tau; c_2, 2N_2) - \\ & \quad - \frac{1}{N_2} \vartheta_{g_2 h_2}'''(\tau; c_2, 2N_2) \vartheta_{g_1 h_1}'(\tau; c_1, 2N_1) \end{aligned} \quad (2.1)$$

and

$$\Psi_2(\tau) = \Psi_2(\tau; g_1, g_2, h_1, h_2, c_1, c_2, N_1, N_2) =$$

$$\begin{aligned}
&= \frac{1}{N_1^2} \vartheta_{g_1 h_1}^{(4)}(\tau; c_1, 2N_1) \vartheta_{g_2 h_2}(\tau; c_2, 2N_2) + \\
&\quad + \frac{1}{N_2^2} \vartheta_{g_2 h_2}^{(4)}(\tau; c_2, 2N_2) \vartheta_{g_1 h_1}(\tau; c_1, 2N_1) - \\
&\quad - \frac{6}{N_1 N_2} \vartheta_{g_1 h_1}''(\tau; c_1, 2N_1) \vartheta_{g_2 h_2}''(\tau; c_2, 2N_2),
\end{aligned} \tag{2.2}$$

where

$$2|g_1, 2|g_2, N_1|N, N_2|N, 4|N\left(\frac{h_1}{N_1} + \frac{h_2}{N_2}\right). \tag{2.3}$$

For all substitutions from Γ in the neighborhood of each rational point $\tau = -\frac{\delta}{\gamma}$ ($\gamma \neq 0, (\gamma, \delta) = 1$), we then have

$$\begin{aligned}
&(\gamma\tau + \delta)^5 \Psi_j(\tau; g_1, g_2, h_1, h_2, 0, 0, N_1, N_2) = \\
&= \sum_{n=0}^{\infty} C_n^{(j)} e\left(\frac{n}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \quad (j = 1, 2).
\end{aligned} \tag{2.4}$$

Proof. I. From Lemma 4 for $n = 3$ (with $g_1, h_1, N_1, g'_1, h'_1, H_1$ instead of g, h, N, g', h', H) and $n = 1$ (with $g_2, h_2, N_2, g'_2, h'_2, H_2$ instead of g, h, N, g', h', H), according to (1.19), it follows that

$$\begin{aligned}
&\frac{1}{N_1} (\gamma\tau + \delta)^5 \vartheta_{g_1 h_1}'''(\tau; 0, 2N_1) \vartheta_{g_2 h_2}'(\tau; 0, 2N_2) = \\
&= \frac{1}{N_1} e\left(\frac{5}{4} \operatorname{sgn} \gamma\right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\
&\times \sum_{H_1 \bmod 2N_1} \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \left\{ \vartheta_{g'_1 h'_1}''\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1\right) + \right. \\
&\quad \left. + \frac{3 \cdot 2}{2!} A_{21} \Big|_{z=0} \cdot \vartheta'_{g'_1 h'_1}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1\right) \right\} \times \\
&\times \sum_{H_2 \bmod 2N_2} \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \vartheta'_{g'_2 h'_2}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2\right) = \\
&= e\left(\frac{5}{4} \operatorname{sgn} \gamma\right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\
&\times \sum_{\substack{H_1 \bmod 2N_1 \\ H_2 \bmod 2N_2}} \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \times \\
&\times \left\{ \frac{1}{N_1} \vartheta_{g'_1 h'_1}'''(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1) \vartheta'_{g'_2 h'_2}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2\right) - \right. \\
&\quad \left. - 12\gamma\pi i(\gamma\tau + \delta) \vartheta'_{g'_1 h'_1}\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1\right) \times \right.
\end{aligned}$$

$$\times \vartheta'_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \}. \quad (2.5)$$

Replacing $N_1, g_1, h_1, H_1, g'_1, h'_1$ by $N_2, g_2, h_2, H_2, g'_2, h'_2$ in (2.5) and vice versa, we obtain

$$\begin{aligned} & \frac{1}{N_2} (\gamma\tau + \delta)^5 \vartheta'''_{g'_2 h'_2} (\tau; 0, 2N_2) \vartheta'_{g'_1 h'_1} (\tau; 0, 2N_1) = \\ &= e \left(\frac{5}{4} \operatorname{sgn} \gamma \right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\ & \times \sum_{\substack{H_2 \bmod 2N_2 \\ H_1 \bmod 2N_1}} \varphi_{g'_2 g_2 h_2} (0, H_2; 2N_2) \varphi_{g'_1 g_1 h_1} (\tau; 0, H_1; 2N_1) \times \\ & \times \left\{ \frac{1}{N_2} \vartheta'''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \vartheta'_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - \right. \\ & \quad \left. - 12\gamma\pi i(\gamma\tau + \delta) \vartheta'_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \times \right. \\ & \quad \left. \times \vartheta'_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right\}. \end{aligned} \quad (2.6)$$

Subtracting (2.6) from (2.5), according to (2.1), we obtain

$$\begin{aligned} & (\gamma\tau + \delta)^5 \Psi_1 (\tau; g_1, g_2, h_1, h_2, 0, 0, N_1, N_2) = \\ &= e \left(\frac{5}{4} \operatorname{sgn} \gamma \right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\ & \times \sum_{\substack{H_1 \bmod 2N_1 \\ H_2 \bmod 2N_2}} \varphi_{g'_1 g_1 h_1} (0, H_1; 2N_1) \varphi_{g'_2 g_2 h_2} (0, H_2; 2N_2) \times \\ & \times \Psi_1 \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; g'_1, g'_2, h'_1, h'_2, H_1, H_2, N_1, N_2 \right). \end{aligned} \quad (2.7)$$

In (2.7) let γ be even. Then, by (1.21) and (2.3), g'_1 and g'_2 are also even. Therefore, according to (3) and the notations (4), we have

$$\begin{aligned} & \vartheta'''_{g'_r h'_r} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_r, 2N_r \right) \vartheta'_{g'_s h'_s} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_s, 2N_s \right) = \\ &= \sum_{n_1=0}^{\infty} B_{n_1} e \left(\frac{N n_1 / N_r}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta} \right) \sum_{n_2=0}^{\infty} B_{n_2} e \left(\frac{N n_2 / N_s}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta} \right) = \\ &= \sum_{n=0}^{\infty} B_n e \left(\frac{n}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta} \right) \\ & \text{for } r = 1, s = 2 \text{ and } r = 2, s = 1. \end{aligned} \quad (2.8)$$

Hence, for even γ , (2.4) follows from (2.1), (2.7), and (2.8) if $j = 1$.

Now let γ in (2.7) be odd. If h_1 and h_2 are both even, then by (1.21) and (2.3), g'_1 and g'_2 are also even, and we obtain the same result. But if h_r is odd, then by (1.21) g'_r will also be odd, and in (3) we have

$$(m + \frac{g'_r}{2})^2 = (m + \frac{1}{2}(g'_r - 1))^2 + (m + \frac{1}{2}(g'_r - 1)) + \frac{1}{4},$$

hence

$$\begin{aligned} \vartheta'_{g'_r h'_r}''\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_r, 2N_r\right) &= -\pi^3 ie\left(\frac{1}{16N_r} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \times \\ &\quad \times \sum_{m \equiv H_r \pmod{2N_r}} (-1)^{h'_r(m-H_r)/2N_r} (2m + g'_r)^3 \times \\ &\quad \times e\left\{\frac{1}{4N_r} \left((m + \frac{1}{2}(g'_r - 1))^2 + (m + \frac{1}{2}(g'_r - 1))\right) \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right\} = \\ &= e\left(\frac{h_r}{16N_r} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \sum_{n_1=0}^{\infty} B'_{n_1} e\left(\frac{n_1}{4N_r} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right), \end{aligned}$$

since (5) implies that $h_r = 1$. Analogously,

$$\vartheta'_{g'_s h'_s}''\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_s, 2N_s\right) = e\left(\frac{h_s}{16N_s} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \sum_{n_2=0}^{\infty} B'_{n_2} e\left(\frac{n_2}{4N_s} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right),$$

which implies that $h_s = 1$ if h_s is odd and $h_s = 0$ if h_s is even. Thus, if among h_1 and h_2 at least one is odd, then we have, for $r = 1, s = 2$ and $r = 2, s = 1$,

$$\begin{aligned} \vartheta'_{g'_r h'_r}''\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_r, 2N_r\right) \vartheta'_{g'_s h'_s}''\left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_s, 2N_s\right) &= \\ &= e\left(\frac{N/4(h_r/N_r + h_s/N_s)}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \sum_{n_1=0}^{\infty} B'_{n_1} e\left(\frac{n_1}{4N_r} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \times \\ &\quad \times \sum_{n_2=0}^{\infty} B'_{n_2} e\left(\frac{n_2}{4N_s} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) = \sum_{n=0}^{\infty} B'_n e\left(\frac{n}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right), \end{aligned} \tag{2.9}$$

since, by (2.3), $\frac{N}{4}(\frac{h_r}{N_r} + \frac{h_s}{N_s})$ is an integer. Hence for odd γ (2.4) follows from (2.1), (2.7), and (2.9) if $j = 1$.

II. From Lemma 4 for $n = 4$ and $n = 0$, as in I, it follows that

$$\begin{aligned} \frac{1}{N_2^2} (\gamma\tau + \delta)^5 \vartheta_{g_1 h_1}^{(4)}(\tau; 0, 2N_1) \vartheta_{g_2 h_2}(\tau; 0, 2N_2) &= \\ &= \frac{1}{N_1^2} e\left(\frac{5}{4} \operatorname{sgn} \gamma\right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \end{aligned}$$

$$\begin{aligned}
& \times \sum_{H_1 \bmod 2N_1} \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \left\{ \vartheta_{g'_1 h'_1}^{(4)} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) + \right. \\
& \quad + \frac{4 \cdot 3}{2!} A_{21} \Big|_{z=0} \cdot \vartheta''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) + \\
& \quad \left. + \frac{A_{42}}{2!} \Big|_{z=0} \cdot \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right\} \times \\
& \times \sum_{H_2 \bmod 2N_2} \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) = \\
& = e \left(\frac{5}{4} \operatorname{sgn} \gamma \right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\
& \times \sum_{\substack{H_1 \bmod 2N_1 \\ H_2 \bmod 2N_2}} \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \times \\
& \quad \times \left\{ \frac{1}{N_1^2} \vartheta_{g'_1 h'_1}^{(4)} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \times \right. \\
& \quad \times \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - \frac{24\gamma\pi i}{N_1} (\gamma\tau + \delta) \vartheta''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \times \\
& \quad \left. \times \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - 48\gamma^2\pi^2(\gamma\tau + \delta)^2 \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \times \right. \\
& \quad \left. \times \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \right\}. \tag{2.10}
\end{aligned}$$

Replacing $N_1, g_1, h_1, H_1, g'_1, h'_1$ by $N_2, g_2, H_2, g'_2, h'_2$ in (2.10) and vice versa, we obtain

$$\begin{aligned}
& \frac{1}{N_1^2} (\gamma\tau + \delta)^5 \vartheta_{g_2 h_2}^{(4)}(\tau; 0, 2N_2) \vartheta_{g_1 h_1}(\tau; 0, 2N_1) = \\
& = e \left(\frac{5}{4} \operatorname{sgn} \gamma \right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\
& \times \sum_{\substack{H_2 \bmod 2N_2 \\ H_1 \bmod 2N_1}} \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \times \\
& \quad \times \left\{ \frac{1}{N_2^2} \vartheta_{g'_2 h'_2}^{(4)} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \times \right. \\
& \quad \times \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - \frac{24\gamma\pi i}{N_2} (\gamma\tau + \delta) \vartheta''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \times \\
& \quad \left. \times \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - 48\gamma^2\pi^2(\gamma\tau + \delta)^2 \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \times \right. \\
& \quad \left. \times \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right\}
\end{aligned}$$

$$\times \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \}. \quad (2.11)$$

Analogously, from Lemma 4 for $n = 2$ it follows that

$$\begin{aligned} & \frac{6}{N_1 N_2} (\gamma\tau + \delta)^5 \vartheta''_{g_1 h_1}(\tau; 0, 2N_1) \vartheta''_{g_2 h_2}(\tau; 0, 2N_2) = \\ &= e\left(\frac{5}{4} \operatorname{sgn} \gamma\right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\ & \times \sum_{\substack{H_1 \bmod 2N_1 \\ H_2 \bmod 2N_2}} \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \times \\ & \times \left\{ \frac{6}{N_1 N_2} \vartheta''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \vartheta''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - \right. \\ & - \frac{24\gamma\pi i}{N_2} (\gamma\tau + \delta) \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - \\ & - \frac{24\gamma\pi i}{N_1} (\gamma\tau + \delta) \vartheta''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \vartheta''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - \\ & \left. - 96\gamma^2\pi^2 (\gamma\tau + \delta)^2 \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \times \right. \\ & \left. \times \vartheta_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \right\}. \end{aligned} \quad (2.12)$$

Subtracting (2.12) from the sum of (2.10) and (2.11), according to (2.2), we obtain

$$\begin{aligned} & (\gamma\tau + \delta)^5 \Psi_2(\tau; g_1, g_2, h_1, h_2, 0, 0, N_1, N_2) = \\ &= e\left(\frac{5}{4} \operatorname{sgn} \gamma\right) (2|\gamma|(N_1 N_2)^{1/2})^{-1} \times \\ & \times \sum_{\substack{H_1 \bmod 2N_1 \\ H_2 \bmod 2N_2}} \varphi_{g'_1 g_1 h_1}(0, H_1; 2N_1) \varphi_{g'_2 g_2 h_2}(0, H_2; 2N_2) \times \\ & \times \Psi_2 \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; g'_1, g'_2, h'_1, h'_2, H_1, H_2, N_1, N_2 \right). \end{aligned} \quad (2.13)$$

Further, reasoning as in I, from (2.2) and (2.13) we obtain (2.4) if $j = 2$. \square

Theorem 1. *For a given N the functions $\Psi_1(\tau)$ and $\Psi_2(\tau)$ with $c_1 = c_2 = 0$ are entire modular forms of weight 10 and character $\chi(\delta) = \operatorname{sgn} \delta \left(\frac{-\Delta}{|\delta|} \right)$ (Δ is the determinant of an arbitrary positive quadratic form in 5 variables)*

for the congruence group $\Gamma_0(4N)$ if the following conditions hold:

$$1) \quad 2|g_1, 2|g_2, N_1|N, N_2|N, \quad (2.14)$$

$$2) \quad 4|N\left(\frac{h_1^2}{N_1} + \frac{h_2^2}{N_2}\right), 4|\frac{g_1^2}{4N_1} + \frac{g_2^2}{4N_2}, \quad (2.15)$$

3) for all α and δ with $\alpha \equiv \delta \equiv 1 \pmod{4N}$

$$\begin{aligned} & \left(\frac{N_1 N_2}{|\delta|}\right) \Psi_j(\tau; \alpha g_1, \alpha g_2, h_1, h_2, 0, 0, N_1, N_2) = \\ & = \left(\frac{\Delta}{|\delta|}\right) \Psi_j(\tau; g_1, g_2, h_1, h_2, 0, 0, N_1, N_2) \quad (j = 1, 2). \end{aligned} \quad (2.16)$$

Proof.

I. It is wellknown that the thetaseries (2)-(3) are regular on \mathcal{H} , hence the functions $\Psi_1(\tau)$ and $\Psi_2(\tau)$ satisfy condition 1 of the definition.

II. From (2.15), since $\delta^2 \equiv 1 \pmod{4}$, it follows that

$$4|N\delta^2\left(\frac{h_1^2}{N_1} + \frac{h_2^2}{N_2}\right), 4|\frac{g_1^2}{4N_1}\delta^{2\varphi(2N_1)-2} + \frac{g_2^2}{4N_2}\delta^{2\varphi(2N_2)-2}. \quad (2.17)$$

By Lemma 3 for $n = 3$ and $n = 1$ (with g_r, h_r, N_r and g_s, h_s, N_s instead of g, h, N), according to (2.14) and (2.17), for each substitution from $\Gamma_0(4N)$, we have

$$\begin{aligned} & \vartheta''_{g_r h_r} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_r \right) \vartheta'_{g_s h_s} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_s \right) = i^{\eta(\gamma)(sgn\delta-1)} i^{1-|\delta|} \times \\ & \times (\gamma\tau + \delta)^5 \left(\frac{N_r N_s}{|\delta|} \right) \vartheta'''_{\alpha g_r, h_r}(\tau; 0, 2N_r) \vartheta'_{\alpha g_s, h_s}(\tau; 0, 2N_s) = \\ & = \operatorname{sgn} \delta \left(\frac{-N_r N_s}{|\delta|} \right) (\gamma\tau + \delta)^5 \times \\ & \times \vartheta'''_{\alpha g_r, h_r}(\tau; 0, 2N_r) \vartheta'_{\alpha g_s, h_s}(\tau; 0, 2N_s) \end{aligned} \quad (2.18)$$

for $r = 1, s = 2$ and $r = 2, s = 1$.

Analogously, by Lemma 3, we have:

1) for $n = 4$ and $n = 0$

$$\begin{aligned} & \vartheta^{(4)}_{g_r h_r} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_r \right) \vartheta_{g_s h_s} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_s \right) = i^{\eta(\gamma)(sgn\delta-1)} i^{1-|\delta|} \times \\ & \times (\gamma\tau + \delta)^5 \left(\frac{N_r N_s}{|\delta|} \right) \vartheta^{(4)}_{\alpha g_r, h_r}(\tau; 0, 2N_r) \vartheta_{\alpha g_s, h_s}(\tau; 0, 2N_s) = \\ & = \operatorname{sgn} \delta \left(\frac{-N_r N_s}{|\delta|} \right) (\gamma\tau + \delta)^5 \times \\ & \times \vartheta^{(4)}_{\alpha g_r, h_r}(\tau; 0, 2N_r) \vartheta'_{\alpha g_s, h_s}(\tau; 0, 2N_s) \end{aligned} \quad (2.19)$$

if $r = 1, s = 2$ and $r = 2, s = 1$;

2) for $n = 2$

$$\begin{aligned} & \vartheta''_{g_1 h_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_1 \right) \vartheta''_{g_2 h_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_2 \right) = \\ &= \operatorname{sgn} \delta \left(\frac{-N_1 N_2}{|\delta|} \right) (\gamma\tau + \delta)^5 \times \\ & \quad \times \vartheta''_{\alpha g_1, h_1}(\tau; 0, 2N_1) \vartheta''_{\alpha g_2, h_2}(\tau; 0, 2N_2). \end{aligned} \quad (2.20)$$

Hence, according to (2.16), for all α and δ with $\alpha\delta \equiv 1 \pmod{4N}$, we have

$$\begin{aligned} & \Psi_j \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; g_1, g_2, h_1, h_2, 0, 0, N_1, N_2 \right) = \\ &= \operatorname{sgn} \delta \left(\frac{-N_1 N_2}{|\delta|} \right) (\gamma\tau + \delta)^5 \Psi_j(\tau; \alpha g_1, \alpha g_2, h_1, h_2, 0, 0, N_1, N_2) = \\ &= \operatorname{sgn} \delta \left(\frac{-\Delta}{|\delta|} \right) (\gamma\tau + \delta)^5 \Psi_j(\tau; g_1, g_2, h_1, h_2, 0, 0, N_1, N_2) \end{aligned}$$

for $j = 1$, by (2.1) and (2.16), and for $j = 2$, by (2.2), (2.19) and (2.20). Thus the functions $\Psi_1(\tau)$ and $\Psi_2(\tau)$ with $c_1 = c_2 = 0$ satisfy condition 2 of the definition.

III. From (13) it follows that, for $r = 1$, $s = 2$ and $r = 2$, $s = 1$,

$$\begin{aligned} & \vartheta'''_{g_r h_r}(\tau; 0, 2N_r) \vartheta'_{g_s h_s}(\tau; 0, 2N_s) = \pi^4 \sum_{m_r, m_s=-\infty}^{\infty} (-1)^{h_r m_r + h_s m_s} \times \\ & \quad \times (4N_r m_r + g_r)^3 (4N_s m_s + g_s) e(\Lambda\tau), \\ & \vartheta^{(4)}_{g_r h_r}(\tau; 0, 2N_r) \vartheta_{g_s h_s}(\tau; 0, 2N_s) = \pi^4 \sum_{m_r, m_s=-\infty}^{\infty} (-1)^{h_r m_r + h_s m_s} \times \\ & \quad \times (4N_r m_r + g_r)^4 e(\Lambda\tau) \end{aligned} \quad (2.21)$$

and also

$$\begin{aligned} & \vartheta''_{g_1 h_1}(\tau; 0, 2N_1) \vartheta''_{g_2 h_2}(\tau; 0, 2N_2) = \pi^4 \sum_{m_1, m_2=-\infty}^{\infty} (-1)^{h_1 m_1 + h_2 m_2} \times \\ & \quad \times (4N_1 m_1 + g_1)^2 (4N_2 m_2 + g_2)^2 e(\Lambda\tau), \end{aligned}$$

where

$$\Lambda = \sum_{k=1}^2 \frac{1}{4N_k} (2N_k m_k + g_k/2)^2 = \sum_{k=1}^2 (N_k m_k^2 + m_k g_k/2) + \frac{1}{4} \sum_{k=1}^2 g_k^2 / 4N_k,$$

by (2.14) and (2.15), is an integer. Thus, the functions $\Psi_1(\tau)$ and $\Psi_2(\tau)$ with $c_1 = c_2 = 0$ satisfy condition 3 of the definition.

IV. By Lemma 5 the functions $\Psi_1(\tau)$ and $\Psi_2(\tau)$ with $c_1 = c_2 = 0$ also satisfy condition 4 of the definition. \square

3.

Lemma 6. *For a given N let*

$$\begin{aligned} \Phi_1(\tau) &= \Phi_1(\tau; g_1, \dots, g_4, h_1, \dots, h_4, c_1, \dots, c_4, N_1, \dots, N_4) = \\ &= \frac{1}{N_1} \vartheta''_{g_1 h_1}(\tau; c_1, 2N_1) \vartheta'_{g_2 h_2}(\tau; c_2, 2N_2) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; c_k, 2N_k) - \\ &\quad - \frac{1}{N_2} \vartheta''_{g_2 h_2}(\tau; c_2, 2N_2) \vartheta'_{g_1 h_1}(\tau; c_1, 2N_1) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; c_k, 2N_k) \end{aligned} \quad (3.1)$$

and

$$\begin{aligned} \Phi_2(\tau) &= \Phi_2(\tau; g_1, \dots, g_4, h_1, \dots, h_4, c_1, \dots, c_4, N_1, \dots, N_4) = \\ &= \frac{1}{N_1^2} \vartheta^{(4)}_{g_1 h_1}(\tau; c_1, 2N_1) \vartheta_{g_2 h_2}(\tau; c_2, 2N_2) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; c_k, 2N_k) + \\ &\quad + \frac{1}{N_2^2} \vartheta^{(4)}_{g_2 h_2}(\tau; c_2, 2N_2) \vartheta_{g_1 h_1}(\tau; c_1, 2N_1) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; c_k, 2N_k) - \\ &\quad - \frac{6}{N_1 N_2} \vartheta''_{g_1 h_1}(\tau; c_1, 2N_1) \vartheta''_{g_2 h_2}(\tau; c_2, 2N_2) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; c_k, 2N_k), \end{aligned} \quad (3.2)$$

where

$$2|g_k, N_k|N \quad (k = 1, 2, 3, 4), \quad 4|N \sum_{k=1}^4 \frac{h_k}{N_k}. \quad (3.3)$$

For all substitutions from Γ in the neighborhood of each rational point $\tau = -\frac{\delta}{\gamma}$ ($\gamma \neq 0$, $(\gamma, \delta) = 1$), we then have

$$\begin{aligned} (\gamma\tau + \delta)^6 \Phi_j(\tau; g_1, \dots, g_4, h_1, \dots, h_4, 0, \dots, 0, N_1, \dots, N_4) &= \\ &= \sum_{n=0}^{\infty} D_n^{(j)} e\left(\frac{n}{4N} \frac{\alpha\tau + \beta}{\gamma\tau + \delta}\right) \quad (j = 1, 2). \end{aligned} \quad (3.4)$$

Proof. From Lemma 4 for $n = 3$, $n = 1$, and $n = 0$, as in the proof of Lemma 5, it follows that

$$\frac{1}{N_1} (\gamma\tau + \delta)^6 \vartheta''_{g_1 h_1}(\tau; 0, 2N_1) \vartheta'_{g_2 h_2}(\tau; 0, 2N_2) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) =$$

$$\begin{aligned}
&= e\left(\frac{3}{2} \operatorname{sgn} \gamma\right) \left(4\gamma^2 \left(\prod_{k=1}^4 N_k\right)^{1/2}\right)^{-1} \sum_{\substack{H_k \bmod 2N_k \\ (k=1,2,3,4)}} \prod_{k=1}^4 \varphi_{g'_k g_k h_k}(0, H_k; 2N_k) \times \\
&\quad \times \left\{ \left(\frac{1}{N_1} \vartheta'''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - \right. \right. \\
&\quad \left. \left. - 12\gamma\pi i(\gamma\tau + \delta) \vartheta'_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right) \times \right. \\
&\quad \left. \times \vartheta'_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \prod_{k=3}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) \right\} \quad (3.5)
\end{aligned}$$

and

$$\begin{aligned}
&\frac{1}{N_2} (\gamma\tau + \delta)^6 \vartheta''_{g'_2 h'_2}(\tau; 0, 2N_2) \vartheta'_{g'_1 h'_1}(\tau; 0, 2N_1) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\
&= e\left(\frac{3}{2} \operatorname{sgn} \gamma\right) \left(4\gamma^2 \left(\prod_{k=1}^4 N_k\right)^{1/2}\right)^{-1} \sum_{\substack{H_k \bmod 2N_k \\ (k=1,2,3,4)}} \prod_{k=1}^4 \varphi_{g'_k g_k h_k}(0, H_k; 2N_k) \times \\
&\quad \times \left\{ \left(\frac{1}{N_2} \vartheta'''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - \right. \right. \\
&\quad \left. \left. - 12\gamma\pi i(\gamma\tau + \delta) \vartheta'_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \right) \times \right. \\
&\quad \left. \times \vartheta'_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \prod_{k=3}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) \right\}. \quad (3.6)
\end{aligned}$$

As in the proof of Lemma 5, from Lemma 4 it follows that

1) for $n = 4$ and $n = 0$

$$\begin{aligned}
&\frac{1}{N_1^2} (\gamma\tau + \delta)^6 \vartheta^{(4)}_{g'_1 h'_1}(\tau; 0, 2N_1) \prod_{k=2}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\
&= e\left(\frac{3}{2} \operatorname{sgn} \gamma\right) \left(4\gamma^2 \left(\prod_{k=1}^4 N_k\right)^{1/2}\right)^{-1} \sum_{\substack{H_k \bmod 2N_k \\ (k=1,2,3,4)}} \prod_{k=1}^4 \varphi_{g'_k g_k h_k}(0, H_k; 2N_k) \times \\
&\quad \times \left\{ \left(\frac{1}{N_1^2} \vartheta^{(4)}_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - \right. \right. \\
&\quad \left. \left. - \frac{24\gamma\pi i}{N_1} (\gamma\tau + \delta) \vartheta''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - \right. \right. \\
&\quad \left. \left. - 48\gamma^2\pi^2 (\gamma\tau + \delta)^2 \vartheta'_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right) \times \right. \\
&\quad \left. \times \vartheta'_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \prod_{k=3}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) \right\}.
\end{aligned}$$

$$\times \prod_{k=2}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) \Big\} \quad (3.7)$$

and

$$\begin{aligned} & \frac{1}{N_2^2} (\gamma\tau + \delta)^6 \vartheta_{g_2 h_2}^{(4)}(\tau; 0, 2N_2) \vartheta_{g_1 h_1}(\tau; 0, 2N_1) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\ & = e \left(\frac{3}{2} \operatorname{sgn} \gamma \right) \left(4\gamma^2 \left(\prod_{k=1}^4 N_k \right)^{1/2} \right)^{-1} \sum_{\substack{H_k \bmod 2N_k \\ (k=1,2,3,4)}} \prod_{k=1}^4 \varphi_{g'_k g_k h_k}(0, H_k; 2N_k) \times \\ & \quad \times \left\{ \left(\frac{1}{N_2^2} \vartheta_{g'_2 h'_2}^{(4)} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) - \right. \right. \\ & \quad \left. \left. - \frac{24\gamma\pi i}{N_2} (\gamma\tau + \delta) \vartheta''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \right) \times \right. \\ & \quad \times \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \prod_{k=3}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) - \\ & \quad \left. \left. - 48\gamma^2 \pi^2 (\gamma\tau + \delta)^2 \prod_{k=1}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) \right\}, \quad (3.8) \end{aligned}$$

2) for $n = 2$ and $n = 0$

$$\begin{aligned} & \frac{6}{N_1 N_2} (\gamma\tau + \delta)^6 \vartheta''_{g_1 h_1}(\tau; 0, 2N_1) \vartheta''_{g_2 h_2}(\tau; 0, 2N_2) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\ & = e \left(\frac{3}{2} \operatorname{sgn} \gamma \right) \left(4\gamma^2 \left(\prod_{k=1}^4 N_k \right)^{1/2} \right)^{-1} \sum_{\substack{H_k \bmod 2N_k \\ (k=1,2,3,4)}} \varphi_{g'_k g_k h_k}(0, H_k; 2N_k) \times \\ & \quad \times \left\{ \left(\frac{6}{N_1 N_2} \vartheta''_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) - \right. \right. \\ & \quad \left. \left. - \frac{24\gamma\pi i}{N_2} (\gamma\tau + \delta) \vartheta_{g'_1 h'_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right) \times \right. \\ & \quad \times \vartheta''_{g'_2 h'_2} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_2, 2N_2 \right) \prod_{k=3}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) - \\ & \quad \left. \left. - \left(\frac{24\gamma\pi i}{N_1} (\gamma\tau + \delta) \vartheta''_{g_1 h_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) + \right. \right. \right. \\ & \quad \left. \left. \left. + 96\gamma^2 \pi^2 (\gamma\tau + \delta)^2 \vartheta_{g_1 h_1} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_1, 2N_1 \right) \right) \times \right. \end{aligned}$$

$$\times \prod_{k=2}^4 \vartheta_{g'_k h'_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; H_k, 2N_k \right) \}. \quad (3.9)$$

Subtracting (3.6) from (3.5), and (3.9) from the sum of (3.7) and (3.8), according to (3.1) and (3.2) respectively, we obtain

$$\begin{aligned} & (\gamma\tau + \delta)^6 \Phi_j(\tau; g_1, \dots, g_4, h_1, \dots, h_4, 0, \dots, N_1, \dots, N_4) = \\ & = e \left(\frac{3}{2} \operatorname{sgn} \gamma \right) \left(4\gamma^2 \left(\prod_{k=1}^4 N_k \right)^{1/2} \right)^{-1} \sum_{\substack{H_k \bmod 2N_k \\ (k=1,2,3,4)}} \varphi_{g'_k g_k h_k}(0, H_k; 2N_k) \times \\ & \quad \times \Phi_j \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; g_1, \dots, g_4, h_1, \dots, h_4, H_1, \dots, H_4, N_1, \dots, N_4 \right) \\ & \quad (j = 1, 2). \end{aligned} \quad (3.10)$$

Further, reasoning as in Lemma 5, from (3.10) we obtain (3.4). \square

Theorem 2. *For a given N the functions $\Phi_1(\tau)$ and $\Phi_2(\tau)$ with $c_1 = c_2 = c_3 = c_4 = 0$ are entire modular forms of weight 12 and character $\chi(\delta) = (\frac{\Delta}{|\delta|})$ (Δ is the determinant of an arbitrary positive quadratic form in 6 variables) for the congruence group $\Gamma_0(4N)$ if the following conditions hold:*

$$1) \quad 2|g_k, N_k|N \quad (k = 1, 2, 3, 4), \quad (3.11)$$

$$2) \quad 4|N \sum_{k=1}^4 \left(\frac{h_k^2}{N_k}, 4 \right) \sum_{k=1}^4 \frac{g_k^2}{4N_k}, \quad (3.12)$$

$$3) \quad \text{for all } \alpha \text{ and } \delta \text{ with } \alpha\delta \equiv 1 \pmod{4N}$$

$$\begin{aligned} & \left(\frac{\prod_{k=1}^4 N_k}{|\delta|} \right) \Phi_j(\tau; \alpha g_1, \dots, \alpha g_4, h_1, \dots, h_4, 0, \dots, 0, N_1, \dots, N_4) = \\ & = \left(\frac{\Delta}{|\delta|} \right) \Phi_j(\tau; g_1, \dots, g_4, h_1, \dots, h_4, 0, \dots, 0, N_1, \dots, N_4) \\ & \quad (j = 1, 2). \end{aligned} \quad (3.13)$$

Proof.

I. As in the case of Theorem 1, the functions $\Phi_1(\tau)$ and $\Phi_2(\tau)$ with $c_1 = c_2 = c_3 = c_4 = 0$ satisfy condition 1 and, by Lemma 6, also condition 4 of the definition.

II. From (3.12), since $\delta^2 \equiv 1 \pmod{4}$, it follows that

$$4|N\delta^2 \sum_{k=1}^4 \frac{h_k^2}{N_k}, 4| \sum_{k=1}^4 \frac{g_k^2}{4N_k} \delta^{2\varphi(2N_k)-2}. \quad (3.14)$$

By Lemma 3, for $n = 3$, $n = 1$ and $n = 0$, according to (3.11) and (3.14), for each substitution from $\Gamma_0(4N)$, we have

$$\begin{aligned} & \vartheta'''_{g_r h_r} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_r \right) \vartheta'_{g_s h_s} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_s \right) \times \\ & \quad \times \prod_{k=3}^4 \vartheta_{g_k h_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_k \right) = \\ & = (\gamma\tau + \delta)^6 \left(\frac{\prod_{k=1}^4 N_k}{|\delta|} \right) \vartheta'''_{\alpha g_r, h_r} (\tau; 0, 2N_r) \vartheta'_{\alpha g_s, h_s} (\tau; 0, 2N_s) \times \\ & \quad \times \prod_{k=3}^4 \vartheta_{\alpha g_k, h_k} (\tau; 0, 2N_k) \end{aligned} \tag{3.15}$$

for $r = 1$, $s = 2$ and $r = 2$, $s = 1$.

Analogously, by Lemma 3, we have:

1) for $n = 4$ and $n = 0$

$$\begin{aligned} & \vartheta^{(4)}_{g_r h_r} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_r \right) \vartheta_{g_s h_s} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_s \right) \times \\ & \quad \times \prod_{k=3}^4 \vartheta_{g_k h_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_k \right) = \\ & = (\gamma\tau + \delta)^6 \left(\frac{\prod_{k=1}^4 N_k}{|\delta|} \right) \vartheta^{(4)}_{\alpha g_r, h_r} (\tau; 0, 2N_r) \vartheta_{\alpha g_s, h_s} (\tau; 0, 2N_s) \times \\ & \quad \times \prod_{k=3}^4 \vartheta_{\alpha g_k, h_k} (\tau; 0, 2N_k) \end{aligned} \tag{3.16}$$

for $r = 1$, $s = 2$ and $r = 2$, $s = 1$,

2) for $n = 2$ and $n = 0$

$$\begin{aligned} & \prod_{k=1}^2 \vartheta''_{g_k h_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_k \right) \prod_{k=3}^4 \vartheta_{g_k h_k} \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; 0, 2N_k \right) = \\ & = (\gamma\tau + \delta)^6 \left(\frac{\prod_{k=1}^4 N_k}{|\delta|} \right) \prod_{k=1}^2 \vartheta''_{\alpha g_k, h_k} (\tau; 0, 2N_k) \prod_{k=3}^4 \vartheta_{\alpha g_k, h_k} (\tau; 0, 2N_k). \end{aligned}$$

Hence, according to (3.13), for all α and δ with $\alpha\delta \equiv 1 \pmod{4N}$, we have

$$\begin{aligned} & \Phi_j \left(\frac{\alpha\tau + \beta}{\gamma\tau + \delta}; g_1, \dots, g_4, h_1, \dots, h_4, 0, \dots, N_1, \dots, N_4 \right) = \\ & = \left(\frac{\prod_{k=1}^4 N_k}{|\delta|} \right) (\gamma\tau + \delta)^6 \times \end{aligned}$$

$$\begin{aligned} & \times \Phi_j(\tau; \alpha g_1, \dots, \alpha g_4, h_1, \dots, h_4, 0, \dots, 0, N_1, \dots, N_4) = \\ & = \left(\frac{\Delta}{|\delta|} \right) (\gamma\tau + \delta)^6 \Phi_j(\tau; g_1, \dots, g_4, h_1, \dots, h_4, 0, \dots, N_1, \dots, N_4) \end{aligned}$$

for $j = 1$, by (3.1) and (3.15), and for $j = 2$, by (3.2), (3.16), and (3.17). Thus the functions $\Phi_1(\tau)$ and $\Phi_2(\tau)$ satisfy condition 2 of the definition.

III. From (13) it follows that, for $r = 1, s = 2$ and $r = 2, s = 1$,

$$\begin{aligned} & \vartheta'''_{g_r h_r}(\tau; 0, 2N_r) \vartheta'_{g_s h_s}(\tau; 0, 2N_s) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\ & = \pi^4 \sum_{m_r, m_s, m_3, m_4 = -\infty}^{\infty} (-1)^{m_r + m_s + m_3 + m_4} (4N_r m_r + g_r)^3 (4N_s m_s + g_s) e(\Lambda\tau), \\ & \vartheta^{(4)}_{g_r h_r}(\tau; 0, 2N_r) \vartheta_{g_s h_s}(\tau; 0, 2N_s) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\ & = \pi^4 \sum_{m_r, m_s, m_3, m_4 = -\infty}^{\infty} (-1)^{m_r + m_s + m_3 + m_4} (4N_r m_r + g_r)^4 e(\Lambda\tau), \end{aligned}$$

and also

$$\begin{aligned} & \vartheta''_{g_1 h_1}(\tau; 0, 2N_1) \vartheta''_{g_2 h_2}(\tau; 0, 2N_2) \prod_{k=3}^4 \vartheta_{g_k h_k}(\tau; 0, 2N_k) = \\ & = \pi^4 \sum_{m_1, \dots, m_4 = -\infty}^{\infty} (-1)^{\sum_{k=1}^4 h_k m_k} (4N_1 m_1 + g_1)^2 (4N_2 m_2 + g_2)^2 e(\Lambda\tau), \end{aligned}$$

where

$$\Lambda = \sum_{k=1}^4 \frac{1}{4N_k} (2N_k m_k + \frac{g_k}{2})^2 = \sum_{k=1}^4 (N_k m_k^2 + \frac{1}{2} m_k g_k) + \frac{1}{4} \sum_{k=1}^4 \frac{g_k^2}{4N_k},$$

by (3.11) and (3.12), is an integer. Thus the functions $\Phi_1(\tau)$ and $\Phi_2(\tau)$ with $c_1 = c_2 = c_3 = c_4 = 0$ satisfy condition 3 of the definition. \square

REFERENCES

1. É. Goursat, Cours d'analyse mathématique. T. I. *Gauthier-Villars, Paris*, 1902.
2. H.D. Kloosterman, The behaviour of general theta functions under the modular group and the characters of binary modular congruence groups. I. *Ann. Math.* **47**(1946), No.3, 317-375.

3. G.A.Lomadze, On the representation of numbers by sums of generalized polygonal numbers. I. (Russian) *Trudy Tbiliss. Mat. Inst. Razmadze* **22**(1956), 77-102.
4. ——, On the representation of numbers by certain quadratic forms in six variables. I. (Russian) *Trudy Tbilis. Univ. Mat. Mech. Astron.* **117**(1966), 7-43.

(Received 17.11.1992)

Author's address:

Faculty of Mechanics and Mathematics
I. Javakhishvili Tbilisi State University
2 University St., Tbilisi 380043
Republic of Georgia