

**EXISTENCE OF CONJUGATE POINTS FOR
SECOND-ORDER LINEAR DIFFERENTIAL EQUATIONS**

A. LOMTATIDZE

ABSTRACT. The sufficient conditions are established under which the second-order linear differential equation is conjugate.

1. Consider the differential equation

$$u'' = p(t)u, \tag{1}$$

where $-\infty < a < b < +\infty$ and the function $p :]a, b[\rightarrow \mathbb{R}$ is Lebesgue integrable on each compact subset of $]a, b[$.

Definition 1. A function $p :]a, b[\rightarrow \mathbb{R}$ belongs to the class $\mathcal{O}(]a, b[)$ if the condition

$$\int_a^b (s-a)(b-s)|p(s)|ds < +\infty \tag{2}$$

is satisfied and the solution of equation (1), satisfying the initial conditions

$$u(a+) = 0, \quad u'(a+) = 1, \tag{3}$$

has at least one zero on $]a, b[$.

It is known [1] that if condition (2) is satisfied then any solution u of (1) has the finite left- and right-hand side limits $u(a+)$ and $u(b-)$ and problem (1),(3) is uniquely solvable.

Definition 2. A function $p :]a, b[\rightarrow \mathbb{R}$ belongs to the class $\mathcal{O}'(]a, b[)$ if

$$\int_a^b (s-a)|p(s)|ds < +\infty \tag{4}$$

and the derivative of the solution of problem (1),(3) has at least one zero on $]a, b[$.

1991 *Mathematics Subject Classification.* 34C10.

Key words and phrases. Second-order linear differential equation, conjugate point.

While investigating two-point singular boundary-value problems there arises a question as to an effective description of the classes $\mathbb{O}(]a, b[)$ and $\mathbb{O}'(]a, b[)$ (see, for example, [2]). Earlier attempts in this direction were undertaken in [3-7]. The statements given below complete the results of these papers.

2. On the Class $\mathbb{O}(]a, b[)$. In this section the function $p :]a, b[\rightarrow \mathbb{R}$ is assumed to satisfy (2).

Theorem 1. *Let there exist a point $t_0 \in]a, b[$ and absolutely continuous functions $f_1 : [a, t_0] \rightarrow [0, +\infty[$ and $f_2 : [t_0, b] \rightarrow [0, +\infty[$ such that $f_1(a) = f_2(b) = 0$, $f_1(t) > 0$ for $a < t \leq t_0$, $f_2(t) > 0$ for $t_0 \leq t < b$,*

$$g_1(t_0) = \int_a^{t_0} \frac{[f_1'(s)]^2}{f_1(s)} ds < +\infty, \quad g_2(t_0) = \int_{t_0}^b \frac{[f_2'(s)]^2}{f_2(s)} ds < +\infty \quad (5)$$

and

$$\begin{aligned} f_2(t_0) \int_a^{t_0} f_1(s)p(s)ds + f_1(t_0) \int_{t_0}^b f_2(s)p(s)ds &\leq \\ &\leq -\frac{1}{4}(g_1(t_0)f_2(t_0) + g_2(t_0)f_1(t_0)). \end{aligned} \quad (6)$$

Then $p \in \mathbb{O}(]a, b[)$.

Corollary 1. *Let there exist $t_0 \in]a, b[$ and $\alpha \in]1, +\infty[$ such that*

$$\begin{aligned} (b-t_0)^\alpha \int_a^{t_0} (s-a)^\alpha p(s)ds + (t_0-a)^\alpha \int_{t_0}^b (b-s)^\alpha p(s)ds &\leq \\ &\leq -\frac{\alpha^2}{4(\alpha-1)}[(t_0-a)(b-t_0)]^{\alpha-1}. \end{aligned}$$

Then $p \in \mathbb{O}(]a, b[)$.

When $t_0 = \frac{a+b}{2}$ and $\alpha = 2$ this proposition implies the result of R. Putnam ([4], p. 177).

Corollary 2. *Let one of the following three conditions hold:*

$$\begin{aligned} \int_a^b [(s-a)(b-s)]^{\frac{3}{2}} p(s)ds &\leq -\frac{9\pi(b-a)^2}{32}, \\ \int_a^b [(s-a)(b-s)]^n p(s)ds &\leq -\frac{n!^2}{2(n-1)(2n-1)!} (b-a)^{2n-1}, \\ n &\geq 2, \quad n \in \mathbb{N}, \\ \int_a^b \sin^2\left(\frac{\pi(s-a)}{b-a}\right) p(s)ds &\leq -\frac{\pi^2}{2(b-a)}. \end{aligned}$$

Then $p \in \mathbb{O}(]a, b[)$.

Corollary 3. *Let the inequality*

$$(b-t_0) \int_a^{t_0} (s-a)^{\frac{3}{2}} (b-s)^{\frac{1}{2}} p(s) ds + (t_0-a) \int_{t_0}^b (s-a)^{\frac{1}{2}} (b-s)^{\frac{3}{2}} p(s) ds \leq \\ \leq -\frac{5}{4} [(t_0-a)(b-t_0)]^{\frac{1}{2}} (b-a)$$

hold for some $t_0 \in]a, b[$. Then $p \in \mathbb{O}(]a, b[)$.

Theorem 2. *Let the inequality*

$$(b-t) \int_a^t (s-a)^{\frac{3}{2}} (b-s)^{\frac{1}{2}} [p(s)]_- ds + \\ + (t-a) \int_t^b (s-a)^{\frac{1}{2}} (b-s)^{\frac{3}{2}} [p(s)]_- ds < [(t-a)(b-t)]^{\frac{1}{2}} (b-a) \\ \text{for } a < t < b \quad (7)$$

hold, where $[p(t)]_- = \frac{|p(t)| - p(t)}{2}$. Then $p \notin \mathbb{O}(]a, b[)$.

3. On the Class $\mathbb{O}'(]a, b[)$. In this section the function $p :]a, b[\rightarrow \mathbb{R}$ is assumed to satisfy (4).

Theorem 3. *Let there exist $t_0 \in]a, b[$ and absolutely continuous functions $f_1 : [a, t_0] \rightarrow [0, +\infty[$ and $f_2 : [t_0, b] \rightarrow [0, +\infty[$ such that $f_1(a) = 0$, $f_1(t) > 0$ for $a < t < t_0$, $f_2(t) > 0$ for $t_0 < t < b$ and conditions (5) and (6) are satisfied. Then $p \in \mathbb{O}'(]a, b[)$.*

Corollary 4. *Let there exist $t_0 \in]a, b[$ and $\alpha \in]1, +\infty[$ such that*

$$\int_a^{t_0} (s-a)^\alpha p(s) ds + (t_0-a)^\alpha \int_{t_0}^b p(s) ds \leq -\frac{\alpha^2}{4(\alpha-1)} (t_0-a)^{\alpha-1}.$$

Then $p \in \mathbb{O}'(]a, b[)$.

Corollary 5. *Let the inequality*

$$\int_a^{t_0} (s-a)^{\frac{3}{2}} p(s) ds + (t_0-a) \int_{t_0}^b (s-a)^{\frac{1}{2}} p(s) ds \leq \\ \leq -\frac{5}{4} (t_0-a)^{\frac{1}{2}} + \frac{t_0-a}{8(b-a)^{\frac{1}{2}}}$$

hold for some $t_0 \in]a, b[$. Then $p \in \mathbb{O}'(]a, b[)$.

4. Proof of the Main results.

Proof of Theorem 1 (Theorem 3). Admit on the contrary that $p \notin \mathbb{O}(]a, b[)$ ($p \notin \mathbb{O}'(]a, b[)$). Then equation (1) has the solution u satisfying

$$\begin{aligned} & u(a+) = 0, \quad u'(a+) = 1, \quad u(t) > 0 \text{ for } a < t < b. \\ & \left(u(a+) = 1, \quad u'(b-) = 0, \quad u(t) > 0 \text{ for } a \leq t \leq b \right). \end{aligned}$$

Denote

$$\rho(t) = \frac{u'(t)}{u(t)} \quad \text{for } a < t < b.$$

It is clear that

$$\rho'(t) = p(t) - \rho^2(t) \quad \text{for } a < t < b. \quad (8)$$

Multiplying both sides of this equality by f_1 and integrating from $a + \varepsilon$ to t_0 where $\varepsilon \in]0, t_0 - a[$, we have

$$\begin{aligned} & - \int_{a+\varepsilon}^{t_0} f_1(s)p(s)ds + f_1(t_0)\rho(t_0) - f_1(a+\varepsilon)\rho(a+\varepsilon) = \\ & = \int_{a+\varepsilon}^{t_0} \left[f_1'(s)\rho(s) - f_1(s)\rho^2(s) \right] ds < \frac{1}{4} \int_{a+\varepsilon}^{t_0} \frac{[f_1'(s)]^2}{f_1(s)} ds < \frac{1}{4}g_1(t_0). \end{aligned}$$

From (5) it easily follows that

$$\lim_{t \rightarrow a+} f_1'(t) = 0.$$

Therefore

$$\lim_{t \rightarrow a+} f_1(t)\rho(t) = 0.$$

In view of this the last inequality can be rewritten as

$$- \int_a^{t_0} f_1(s)p(s)ds + f_1(t_0)\rho(t_0) < \frac{1}{4}g_1(t_0). \quad (9)$$

Now multiplying both sides of (8) by f_2 and integrating from t_0 to $b - \varepsilon$ where $\varepsilon \in]0, b - t_0[$, we have

$$\begin{aligned} & - \int_{t_0}^{b-\varepsilon} f_2(s)p(s)ds - f_2(t_0)\rho(t_0) + f_2(b-\varepsilon)\rho(b-\varepsilon) = \\ & = \int_{t_0}^{b-\varepsilon} \left[f_2'(s)\rho(s) - f_2(s)\rho^2(s) \right] ds < \frac{1}{4} \int_{t_0}^{b-\varepsilon} \frac{[f_2'(s)]^2}{f_2(s)} ds < \frac{1}{4}g_2(t_0). \end{aligned}$$

Taking into account

$$\lim_{t \rightarrow b-} f_2'(t) = 0 \quad \text{and} \quad \lim_{t \rightarrow b-} (b-t)u'(t) = 0,$$

from the last inequality we obtain

$$-\int_{t_0}^b f_2(s)p(s)ds - f_2(t_0)\rho(t_0) < \frac{1}{4}g_2(t_0). \quad (10)$$

From (9) and (10) we have

$$\begin{aligned} & -\left[f_2(t_0) \int_a^{t_0} f_1(s)p(s)ds + f_1(t_0) \int_{t_0}^b f_2(s)p(s)ds \right] < \\ & < \frac{1}{4} \left[f_2(t_0)g_1(t_0) + f_1(t_0)g_2(t_0) \right], \end{aligned}$$

which contradicts (6). \square

Proof of Theorem 2. Admit on the contrary that $p \in \mathbb{O}(]a, b[)$. Then equation (1) has the solution u satisfying

$$u(a+) = u(b_1) = 0, \quad u(t) > 0 \text{ for } a < t < b_1 \leq b.$$

According to the Green formula

$$\begin{aligned} & u(t) = \\ & = -\frac{1}{b_1 - a} \left[(b_1 - t) \int_a^t (s - a)p(s)u(s)ds + (t - a) \int_t^{b_1} (b_1 - s)p(s)u(s)ds \right] \\ & \quad \text{for } a \leq t \leq b_1. \end{aligned}$$

Hence we easily obtain

$$\begin{aligned} u(t) & \leq \frac{1}{b_1 - a} \left[(b_1 - t) \int_a^t (s - a)[p(s)]_- u(s)ds + \right. \\ & \left. + (t - a) \int_t^{b_1} (b_1 - s)[p(s)]_- u(s)ds \right] \text{ for } a \leq t \leq b_1, \end{aligned}$$

i.e.,

$$\begin{aligned} & v(t) \leq \\ & \leq \frac{1}{[(t - a)(b - t)]^{\frac{1}{2}}} \left[\frac{b_1 - t}{b_1 - a} \int_a^t (s - a)^{\frac{3}{2}} (b - s)^{\frac{1}{2}} [p(s)]_- v(s)ds + \right. \\ & \quad \left. + \frac{t - a}{b_1 - a} \int_t^b (s - a)^{\frac{1}{2}} (b_1 - s)(b - s)^{\frac{1}{2}} [p(s)]_- v(s)ds \right] < \\ & < \lambda \frac{1}{[(t - a)(b - t)]^{\frac{1}{2}}} \left[\frac{b - t}{b - a} \int_a^t (s - a)^{\frac{3}{2}} (b - s)^{\frac{1}{2}} [p(s)]_- ds + \right. \\ & \quad \left. + \frac{t - a}{b - a} \int_t^b (s - a)^{\frac{1}{2}} (b - s)^{\frac{3}{2}} [p(s)]_- ds \right] \text{ for } a < t < b_1, \end{aligned}$$

where

$$v(t) = \frac{u(t)}{[(t-a)(b-t)]^{\frac{1}{2}}} \quad \text{for } a < t < b_1$$

and

$$\lambda = \sup\{v(t) \mid t \in]a, b_1[\}.$$

Taking into account (7), we obtain the contradiction $\lambda < \lambda$. \square

Corollaries 1-5 are obtained from Theorems 1 and 3 by an appropriate choice of functions f_1 and f_2 .

REFERENCES

1. I. T. Kiguradze and A. G. Lomtadze, On certain boundary-value problems for second-order linear ordinary differential equations with singularities. *J. Math. Anal. Appl.* **101**(1984), No. 2, 325-347.
2. A. G. Lomtadze, On positive solutions of boundary-value problems for second-order ordinary differential equations with singularities. (Russian) *Differentsial'nye Uravneniya* **23**(1987), 1685-1692.
3. F. Hartman, Ordinary differential equations. *Wiley, New York/London/Sydney*, 1964.
4. M. A. Krasnosel'ski, Vector fields on the plane. (Russian) *Moscow, Nauka*, 1963.
5. N. I. Korshikova, On zeros of solutions of high-order linear equations. (Russian) *Differential equations and their applications. (Russian)* 143-148, *Moscow University Press, Moscow*, 1984.
6. A. G. Lomtadze, On oscillatory properties of solutions of second-order linear differential equations. (Russian) *Sem. I. Vekua Inst. Appl. Math. Tbiliss. St. Univ. Reports.* **19**(1985), 39-53.
7. O. Došlý, The multiplicity criteria for zero points of second-order differential equations. *Math. Slovaca* **42**(1992), No. 2, 181-193.

(Received 01.12.1993)

Author's address:

N.Muskhelishvili Institute of Computational Mathematics
 Georgian Academy of Sciences
 8, Akuri St., Tbilisi 380093
 Republic of Georgia