

Research Article

Weighted Iterated Radial Composition Operators between Some Spaces of Holomorphic Functions on the Unit Ball

Stevo Stević

Mathematical Institute of the Serbian Academy of Science and Arts, Knez Mihailova 36/III, 11000 Beograd, Serbia

Correspondence should be addressed to Stevo Stević, sstevic@ptt.rs

Received 26 September 2010; Accepted 26 October 2010

Academic Editor: Chaitan Gupta

Copyright © 2010 Stevo Stević. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The boundedness and compactness of weighted iterated radial composition operators from the mixed-norm space to the weighted-type space and the little weighted-type space on the unit ball are characterized here. We also calculate the Hilbert-Schmidt norm of the operator on the weighted Bergman-Hilbert space as well as on the Hardy H^2 space.

1. Introduction

Let $z = (z_1, \dots, z_n)$ and $w = (w_1, \dots, w_n)$ be points in \mathbb{C}^n , $\langle z, w \rangle = \sum_{k=1}^n z_k \bar{w}_k$, and $|z| = \sqrt{\langle z, z \rangle}$. Let $\mathbb{B} = \{z \in \mathbb{C}^n : |z| < 1\}$ be the open unit ball in \mathbb{C}^n , $\partial\mathbb{B}$ its boundary, and $H(\mathbb{B})$ the class of all holomorphic functions on \mathbb{B} .

For an $f \in H(\mathbb{B})$ with the Taylor expansion $f(z) = \sum_{|\beta| \geq 0} a_\beta z^\beta$, let

$$\Re f(z) = \sum_{|\beta| \geq 0} |\beta| a_\beta z^\beta \quad (1.1)$$

be the radial derivative of f , where $\beta = (\beta_1, \beta_2, \dots, \beta_n)$ is a multi-index, $|\beta| = \beta_1 + \dots + \beta_n$ and $z^\beta = z_1^{\beta_1} \dots z_n^{\beta_n}$ [1]. It is easy to see that

$$\Re f(z) = \langle \nabla f(z), \bar{z} \rangle, \quad (1.2)$$

where ∇f is the complex gradient of function f .

The iterated radial derivative operator $\mathfrak{R}^m f$ is defined inductively by

$$\mathfrak{R}^m f = \mathfrak{R}(\mathfrak{R}^{m-1} f), \quad m \in \mathbb{N} \setminus \{1\}. \quad (1.3)$$

A positive continuous function ν on the interval $[0, 1)$ is called normal [2] if there are $\delta \in [0, 1)$ and τ and t , $0 < \tau < t$ such that

$$\begin{aligned} \frac{\nu(r)}{(1-r)^\tau} \text{ is decreasing on } [\delta, 1), \quad \lim_{r \rightarrow 1} \frac{\nu(r)}{(1-r)^\tau} = 0, \\ \frac{\nu(r)}{(1-r)^t} \text{ is increasing on } [\delta, 1), \quad \lim_{r \rightarrow 1} \frac{\nu(r)}{(1-r)^t} = \infty. \end{aligned} \quad (1.4)$$

If we say that a function $\nu : \mathbb{B} \rightarrow [0, \infty)$ is normal, we also assume that it is radial, that is, $\nu(z) = \nu(|z|)$, $z \in \mathbb{B}$.

Strictly positive continuous functions on \mathbb{B} are called *weights*.

The weighted-type space $H_\mu^\infty(\mathbb{B}) = H_\mu^\infty$ consists of all $f \in H(\mathbb{B})$ such that

$$\|f\|_{H_\mu^\infty} := \sup_{z \in \mathbb{B}} \mu(z) |f(z)| < \infty, \quad (1.5)$$

where μ is a weight (see, e.g., [3, 4] as well as [5] for a related class of spaces).

The little weighted-type space $H_{\mu,0}^\infty(\mathbb{B}) = H_{\mu,0}^\infty$ is a subspace of H_μ^∞ consisting of all $f \in H(\mathbb{B})$ such that

$$\lim_{|z| \rightarrow 1} \mu(z) |f(z)| = 0. \quad (1.6)$$

For $0 < p, q < \infty$, and ϕ normal, the mixed-norm space $H(p, q, \phi)(\mathbb{B}) = H(p, q, \phi)$ consists of all functions $f \in H(\mathbb{B})$ such that

$$\|f\|_{H(p,q,\phi)} = \left(\int_0^1 M_q^p(f, r) \frac{\phi^p(r)}{1-r} dr \right)^{1/p} < \infty, \quad (1.7)$$

where

$$M_q(f, r) = \left(\int_{\partial \mathbb{B}} |f(r\zeta)|^q d\sigma(\zeta) \right)^{1/q}, \quad (1.8)$$

and $d\sigma$ is the normalized surface measure on $\partial \mathbb{B}$. For $p = q$, $\phi(r) = (1-r^2)^{(\alpha+1)/p}$, and $\alpha > -1$, the space is equivalent with the weighted Bergman space $A_\alpha^p(\mathbb{B}) = A_\alpha^p$, which is defined as the class of all $f \in H(\mathbb{B})$ such that

$$\|f\|_{A_\alpha^p}^p := \int_{\mathbb{B}} |f(z)|^p (1-|z|^2)^\alpha dV(z) < \infty, \quad (1.9)$$

where $dV(z)$ is the Lebesgue volume measure on \mathbb{B} . Some facts on mixed-norm spaces in various domains in \mathbb{C}^n can be found, for example, in [6–8] (see also the references therein).

For $0 < p < \infty$ the Hardy space $H^p(\mathbb{B}) = H^p$ consists of all $f \in H(\mathbb{B})$ such that

$$\|f\|_{H^p} := \sup_{0 < r < 1} \left(\int_{\partial\mathbb{B}} |f(r\zeta)|^p d\sigma(\zeta) \right)^{1/p} < \infty. \tag{1.10}$$

For $p = 2$ the Hardy and the weighted Bergman space are Hilbert.

Let φ be a holomorphic self-map of \mathbb{B} , $u \in H(\mathbb{B})$, and $m \in \mathbb{N}_0$. For $f \in H(\mathbb{B})$, the *weighted iterated radial composition operator* is defined by

$$\mathfrak{R}_{u,\varphi}^m(f)(z) = u(z)\mathfrak{R}^m f(\varphi(z)), \quad z \in \mathbb{B}. \tag{1.11}$$

Note that the operator is the composition of the multiplication, composition and the iterated radial operator, that is

$$\mathfrak{R}_{u,\varphi}^m = M_u \circ C_\varphi \circ \mathfrak{R}^m. \tag{1.12}$$

This is one of the product operators suggested by this author to be investigated at numerous talks (e.g., in [9]). Note that for $m = 0$ the operator $\mathfrak{R}_{u,\varphi}^m$ becomes the weighted composition operator (see, e.g., [4, 8, 10]). It is of interest to provide function-theoretic characterizations for when φ and u induce bounded or compact weighted iterated radial composition operators on spaces of holomorphic functions. Studying products of some concrete linear operators on spaces of analytic functions attracted recently some attention see, for example, [11–32] as well as the related references therein. Some operators on mixed-norm spaces have been studied, for example, in [8, 10, 11, 16, 25, 26, 29, 33] (see also the references therein).

Here we study the boundedness and compactness of weighted iterated radial composition operators from mixed-norm spaces to weighted-type spaces on the unit ball for the case $m \in \mathbb{N}$. We also calculate the Hilbert-Schmidt norm of the operator on the weighted Bergman-Hilbert space $A_\alpha^2(\mathbb{B})$ as well as on the Hardy $H^2(\mathbb{B})$ space.

In this paper, constants are denoted by C , they are positive and may differ from one occurrence to the other. The notation $a \leq b$ means that there is a positive constant C such that $a \leq Cb$. If both $a \leq b$ and $b \leq a$ hold, then one says that $a \asymp b$.

2. Auxiliary Results

In this section we quote several lemmas which are used in the proofs of the main results.

The next characterization of compactness is proved in a standard way, hence we omit its proof (see, e.g., [34]).

Lemma 2.1. *Assume $p, q > 0$, φ is a holomorphic self-map of \mathbb{B} , $u \in H(\mathbb{B})$, ϕ is normal and μ is a weight. Then the operator $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is compact if and only if for every bounded sequence $(f_k)_{k \in \mathbb{N}} \subset H(p, q, \phi)$ converging to 0 uniformly on compacts of \mathbb{B} as $k \rightarrow \infty$, one has*

$$\lim_{k \rightarrow \infty} \left\| \mathfrak{R}_{u,\varphi}^m f_k \right\|_{H_\mu^\infty} = 0. \tag{2.1}$$

The following lemma is a slight modification of Lemma 2.5 in [8] and is proved similar to Lemma 1 in [35].

Lemma 2.2. *Assume μ is a normal weight. Then a closed set K in $H_{\mu,0}^\infty$ is compact if and only if it is bounded and*

$$\lim_{|z| \rightarrow 1} \sup_{f \in K} \mu(z) |f(z)| = 0. \quad (2.2)$$

The following lemma is folklore and in the next form it can be found in [36].

Lemma 2.3. *Assume that $0 < p, q < \infty$, ϕ is normal, and $m \in \mathbb{N}$. Then for every $f \in H(\mathbb{B})$ the following asymptotic relationship holds:*

$$\int_0^1 M_q^p(f, r) \frac{\phi^p(r)}{1-r} dr \asymp |f(0)|^p + \int_0^1 M_q^p(\mathfrak{R}^m f, r) (1-r)^{mp} \frac{\phi^p(r)}{1-r} dr. \quad (2.3)$$

Lemma 2.4. *Assume that $m \in \mathbb{N}$, $0 < p, q < \infty$, ϕ is normal and $f \in H(p, q, \phi)$. Then, there is a positive constant C independent of f such that*

$$|\mathfrak{R}^m f(z)| \leq C \|f\|_{H(p,q,\phi)} \frac{|z|}{\phi(|z|) (1-|z|^2)^{n/q+m}}. \quad (2.4)$$

Proof. Let $g = \mathfrak{R}^{m-1} f$ and $z \in \mathbb{B}$. By the definition of the radial derivative, the Cauchy-Schwarz inequality and the Cauchy inequality, we have that

$$|\mathfrak{R}g(z)| \leq |z| |\nabla g(z)| \leq C |z| \frac{\sup_{B(z, 1-|z|/4)} |g(w)|}{1-|z|}. \quad (2.5)$$

From (2.3) with $m \rightarrow m-1$ we easily obtain the following inequality (see, e.g., [8, Lemma 2.1]):

$$|g(z)| \leq C \frac{\|f\|_{H(p,q,\phi)}}{\phi(|z|) (1-|z|^2)^{n/q+m-1}}. \quad (2.6)$$

From (2.5) and (2.6) and the asymptotic relations

$$1-|w| \asymp 1-|z|, \quad \phi(|z|) \asymp \phi(|w|), \quad \text{for } w \in B\left(z, \frac{1-|z|}{2}\right), \quad (2.7)$$

inequality (2.4) follows. \square

Lemma 2.5. *Let*

$$f_{a,s}(z) = \frac{1}{(1 - \langle z, a \rangle)^s}, \quad z \in \mathbb{B}. \tag{2.8}$$

Then,

$$\mathfrak{R}^m f_{a,s}(z) = s \frac{P_m(\langle z, a \rangle)}{(1 - \langle z, a \rangle)^{s+m}}, \tag{2.9}$$

where

$$P_m(w) = s^{m-1}w^m + p_{m-1}^{(m)}(s)w^{m-1} + \dots + p_2^{(m)}(s)w^2 + w, \tag{2.10}$$

and where $p_j^{(m)}(s)$, $j = 2, \dots, m - 1$ *are nonnegative polynomials for* $s > 0$.

Proof. We prove the lemma by induction. For $m = 1$,

$$\mathfrak{R} f_{a,s}(z) = s \frac{\langle z, a \rangle}{(1 - \langle z, a \rangle)^{s+1}}, \quad z \in \mathbb{B}, \tag{2.11}$$

which is formula (2.9) with $P_1(w) = w$.

Assume (2.9) is true for every $m \in \{1, \dots, l\}$. Taking the radial derivative operator on equality (2.9) with $m = l$, we obtain

$$\begin{aligned} \mathfrak{R}^{l+1} f_{a,s}(z) &= s \mathfrak{R} \frac{s^{l-1} \langle z, a \rangle^l + p_{l-1}^{(l)}(s) \langle z, a \rangle^{l-1} + \dots + p_2^{(l)}(s) \langle z, a \rangle^2 + \langle z, a \rangle}{(1 - \langle z, a \rangle)^{s+l}} \\ &= s \frac{(s+l) \left(s^{l-1} \langle z, a \rangle^{l+1} + p_{l-1}^{(l)}(s) \langle z, a \rangle^l + \dots + p_2^{(l)}(s) \langle z, a \rangle^3 + \langle z, a \rangle^2 \right)}{(1 - \langle z, a \rangle)^{s+l+1}} \\ &\quad + s \frac{\left(l s^{l-1} \langle z, a \rangle^l + (l-1) p_{l-1}^{(l)}(s) \langle z, a \rangle^{l-1} + \dots + 2 p_2^{(l)}(s) \langle z, a \rangle^2 + \langle z, a \rangle \right) (1 - \langle z, a \rangle)}{(1 - \langle z, a \rangle)^{s+l+1}}, \\ &= s \frac{s^l \langle z, a \rangle^{l+1} + \dots + \left[(s+l-2) p_2^{(l)}(s) + 3 p_3^{(l)}(s) \right] \langle z, a \rangle^3}{(1 - \langle z, a \rangle)^{s+l+1}} \\ &\quad + \frac{\left[(s+l-1) + 2 p_2^{(l)}(s) \right] \langle z, a \rangle^2 + \langle z, a \rangle}{(1 - \langle z, a \rangle)^{s+l+1}} \end{aligned} \tag{2.12}$$

from which the inductive proof easily follows. □

3. Boundedness and Compactness of $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ (or $H_{\mu,0}^\infty$)

This section characterizes the boundedness and compactness of $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ (or $H_{\mu,0}^\infty$).

Theorem 3.1. *Assume $m \in \mathbb{N}$, $0 < p, q < \infty$, ϕ is normal, μ is a weight, φ is a holomorphic self-map of \mathbb{B} , and $u \in H(\mathbb{B})$. Then $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded if and only if*

$$M_1 := \sup_{z \in \mathbb{B}} \frac{\mu(z)|u(z)||\varphi(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{n/q+m}} < \infty. \quad (3.1)$$

Moreover, if $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded, then the following asymptotic relationship holds

$$\left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty} \asymp M_1. \quad (3.2)$$

Proof. Assume (3.1) holds. Then by Lemma 2.4 for each $f \in H(p, q, \phi)$, we have that

$$\mu(z) \left| \mathfrak{R}_{u,\varphi}^m f(z) \right| \leq C \|f\|_{H(p,q,\phi)} \frac{\mu(z)|u(z)||\varphi(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{n/q+m}}. \quad (3.3)$$

Taking the supremum over the unit ball in (3.3) and using (3.1) the boundedness of operator $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ follows and

$$\left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty} \leq CM_1. \quad (3.4)$$

Now assume that operator $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded. By using the test functions

$$f_j(z) = z_j \in H(p, q, \phi), \quad j = 1, \dots, n, \quad (3.5)$$

we obtain

$$\mathfrak{R}_{u,\varphi}^m f_j \in H_\mu^\infty, \quad j = 1, \dots, n, \quad (3.6)$$

that is, for each $j = 1, \dots, n$, holds

$$\left\| \mathfrak{R}_{u,\varphi}^m f_j \right\|_{H_\mu^\infty} = \sup_{z \in \mathbb{B}} \mu(z)|u(z)||\varphi_j(z)| \leq \|z_j\|_{H(p,q,\phi)} \left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty} < \infty, \quad (3.7)$$

which implies that

$$\begin{aligned} \sup_{z \in \mathbb{B}} \mu(z)|u(z)||\varphi(z)| &\leq \sum_{j=1}^n \sup_{z \in \mathbb{B}} \mu(z)|u(z)||\varphi_j(z)| \\ &\leq \left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty} \sum_{j=1}^n \|z_j\|_{H(p,q,\phi)} < \infty. \end{aligned} \quad (3.8)$$

Let

$$f_w(z) = \frac{(1 - |w|^2)^{t+1}}{\phi(w)(1 - \langle z, w \rangle)^{n/q+t+1}}. \quad (3.9)$$

It is known that $L_1 := \sup_{w \in \mathbb{B}} \|f_w\|_{H(p,q,\phi)} < \infty$ (see [8, Theorem 3.3]). From this, using the boundedness of $\mathfrak{R}_{u,\varphi}^m : H(p,q,\phi) \rightarrow H_\mu^\infty$ and by Lemma 2.5, we have that for each $a \in \mathbb{B}$

$$\begin{aligned} L_1 \left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty} &\geq \left\| \mathfrak{R}_{u,\varphi}^m(f_{\varphi(a)}) \right\|_{H_\mu^\infty} = \sup_{z \in \mathbb{B}} \mu(z)|u(z)||\mathfrak{R}^m f_{\varphi(a)}(\varphi(z))| \\ &\geq \left(\frac{n}{q} + t + 1 \right) \frac{\mu(a)|u(a)|P_m(|\varphi(a)|^2)}{\phi(|\varphi(a)|)(1 - |\varphi(a)|^2)^{n/q+m}} \\ &\geq \left(\frac{n}{q} + t + 1 \right) \frac{\mu(a)|u(a)||\varphi(a)|^2}{\phi(|\varphi(a)|)(1 - |\varphi(a)|^2)^{n/q+m}}. \end{aligned} \quad (3.10)$$

From (3.10), we have that

$$\begin{aligned} \infty &> \sup_{|\varphi(z)| > 1/2} \frac{\mu(z)|u(z)||\varphi(z)|^2}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{n/q+m}} \\ &\geq \sup_{|\varphi(z)| > 1/2} \frac{\mu(z)|u(z)||\varphi(z)|}{2\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{n/q+m}}. \end{aligned} \quad (3.11)$$

On the other hand, from (3.8) and since ϕ is normal, we obtain

$$\sup_{|\varphi(z)| \leq 1/2} \frac{\mu(\varphi(z))|u(z)||\varphi(z)|^2}{\phi(\varphi(z))(1 - |\varphi(z)|^2)^{n/q+m}} \leq C \sup_{z \in \mathbb{B}} \mu(\varphi(z))|u(z)||\varphi(z)| < \infty. \quad (3.12)$$

From (3.8), (3.10), (3.11), and (3.12) condition (3.1) follows, and moreover

$$M_1 \leq C \left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty}. \quad (3.13)$$

From (3.4) and (3.13) asymptotic relationship (3.2) follows, finishing the proof of the theorem. \square

Theorem 3.2. *Assume $m \in \mathbb{N}$, $0 < p, q < \infty$, ϕ is normal, μ is a weight, φ is a holomorphic self-map of \mathbb{B} and $u \in H(\mathbb{B})$. Then $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is compact if and only if $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded and*

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(z)|u(z)||\varphi(z)|}{\phi(|\varphi(z)|)(1-|\varphi(z)|^2)^{n/q+m}} = 0. \quad (3.14)$$

Proof. Suppose that $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is compact. Then it is clear that $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded. If $\|\varphi\|_\infty < 1$, then (3.14) is vacuously satisfied. Hence assume that $\|\varphi\|_\infty = 1$. Let $(z_k)_{k \in \mathbb{N}}$ be a sequence in \mathbb{B} such that $|\varphi(z_k)| \rightarrow 1$ as $k \rightarrow \infty$, and $f_k(z) = f_{\varphi(z_k)}(z)$, $k \in \mathbb{N}$, where f_w is defined in (3.9). Then $\sup_{k \in \mathbb{N}} \|f_k\|_{H(p,q,\phi)} < \infty$, $f_k \rightarrow 0$ uniformly on compacts of \mathbb{B} as $k \rightarrow \infty$ since

$$\lim_{k \rightarrow \infty} \frac{(1-|\varphi(z_k)|^2)^{t+1}}{\phi(|\varphi(z_k)|)} = 0, \quad (3.15)$$

so that

$$\lim_{k \rightarrow \infty} \left\| \mathfrak{R}_{u,\varphi}^m f_k \right\|_{H_\mu^\infty} = 0. \quad (3.16)$$

On the other hand, by (3.10), we have

$$\frac{\mu(z_k)|u(z_k)||\varphi(z_k)|^2}{\phi(|\varphi(z_k)|)(1-|\varphi(z_k)|^2)^{n/q+m}} \leq C \left\| \mathfrak{R}_{u,\varphi}^m f_k \right\|_{H_\mu^\infty}. \quad (3.17)$$

From (3.16) and (3.17), equality (3.14) easily follows.

Conversely, assume that $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded and (3.14) holds. From the proof of Theorem 3.1 we know that (3.1) holds. On the other hand, from (3.14), we have that, for every $\varepsilon > 0$, there is a $\delta \in (0, 1)$, such that

$$\frac{\mu(z)|u(z)||\varphi(z)|}{\phi(|\varphi(z)|)(1-|\varphi(z)|^2)^{n/q+m}} < \varepsilon \quad (3.18)$$

whenever $\delta < |\varphi(z)| < 1$.

Assume $(f_k)_{k \in \mathbb{N}}$ is a sequence in $H(p, q, \phi)$ such that $\sup_{k \in \mathbb{N}} \|f_k\|_{H(p, q, \phi)} \leq L$ and f_k converges to 0 uniformly on compact subsets of \mathbb{B} as $k \rightarrow \infty$. Let $K = \{z \in \mathbb{B} : |\varphi(z)| \leq \delta\}$. Then from (3.18), and by Lemma 2.4, it follows that

$$\begin{aligned}
 \left\| \mathfrak{R}_{u, \varphi}^m f_k \right\|_{H_\mu^\infty} &= \sup_{z \in \mathbb{B}} \mu(z) |u(z)| \left| \mathfrak{R}^m f_k(\varphi(z)) \right| \\
 &\leq \sup_{z \in K} \mu(z) |u(z)| \left| \mathfrak{R}^m f_k(\varphi(z)) \right| + \sup_{z \in \mathbb{B} \setminus K} \mu(z) |u(z)| \left| \mathfrak{R}^m f_k(\varphi(z)) \right| \\
 &\leq \sup_{z \in K} \mu(z) |u(z)| |\varphi(z)| \left| \nabla \mathfrak{R}^{m-1} f_k(\varphi(z)) \right| \\
 &\quad + C \|f_k\|_{H(p, q, \phi)} \sup_{z \in \mathbb{B} \setminus K} \frac{\mu(z) |u(z)| |\varphi(z)|}{\phi(|\varphi(z)|) (1 - |\varphi(z)|^2)^{n/q+m}} \\
 &\leq K_2 C \sup_{|\zeta| \leq \delta} \left| \nabla \mathfrak{R}^{m-1} f_k(\zeta) \right| + C \varepsilon \|f_k\|_{H(p, q, \phi)},
 \end{aligned} \tag{3.19}$$

where $K_2 := \sup_{z \in \mathbb{B}} \mu(z) |u(z)| |\varphi(z)|$ (see (3.8)). Therefore

$$\left\| \mathfrak{R}_{u, \varphi}^m f_k \right\|_{H_\mu^\infty} \leq K_2 \sup_{|\zeta| \leq \delta} \left| \nabla \mathfrak{R}^{m-1} f_k(\zeta) \right| + C L \varepsilon. \tag{3.20}$$

Since $(f_k)_{k \in \mathbb{N}}$ converges to zero on compact subsets of \mathbb{B} as $k \rightarrow \infty$, by Cauchy's estimates it follows that the sequence $(|\nabla \mathfrak{R}^{m-1} f_k|)_{k \in \mathbb{N}}$ also converges to zero on compact subsets of \mathbb{B} as $k \rightarrow \infty$, in particular

$$\lim_{k \rightarrow \infty} \sup_{|\zeta| \leq \delta} \left| \nabla \mathfrak{R}^{m-1} f_k(\zeta) \right| = 0. \tag{3.21}$$

Using these facts and letting $k \rightarrow \infty$ in (3.20), we obtain that

$$\limsup_{k \rightarrow \infty} \left\| \mathfrak{R}_{u, \varphi}^m f_k \right\|_{H_\mu^\infty} \leq C L \varepsilon. \tag{3.22}$$

Since ε is an arbitrary positive number it follows that the last limit is equal to zero. Applying Lemma 2.1, the implication follows. \square

Theorem 3.3. *Assume $m \in \mathbb{N}$, $0 < p, q < \infty$, ϕ, μ are normal, φ is a holomorphic self-map of \mathbb{B} and $u \in H(\mathbb{B})$. Then $\mathfrak{R}_{u, \varphi}^m : H(p, q, \phi) \rightarrow H_{\mu, 0}^\infty$ is bounded if and only if $\mathfrak{R}_{u, \varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded and*

$$\lim_{|z| \rightarrow 1} \mu(z) |u(z)| |\varphi(z)| = 0. \tag{3.23}$$

Proof. First assume that $\mathfrak{R}_{u, \varphi}^m : H(p, q, \phi) \rightarrow H_{\mu, 0}^\infty$ is bounded. Then, it is clear that $\mathfrak{R}_{u, \varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded, and as in the proof of Theorem 3.1, by taking the test functions $f_j(z) = z_j$, $j = 1, \dots, n$, we obtain (3.23).

Conversely, assume that the operator $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$ is bounded and condition (3.23) holds. Then, for each polynomial p , we have

$$\mu(z) \left| \left(\mathfrak{R}_{u,\varphi}^m p \right) (z) \right| \leq \mu(z) |u(z)| \left| \left(\mathfrak{R}^m p \right) (\varphi(z)) \right| \leq \mu(z) |u(z)| |\varphi(z)| \left\| \nabla \mathfrak{R}^{m-1} p \right\|_\infty, \quad (3.24)$$

from which along with condition (3.23) it follows that $\mathfrak{R}_{u,\varphi}^m p \in H_{\mu,0}^\infty$. Since the set of all polynomials is dense in $H(p, q, \phi)$, we see that for every $f \in H(p, q, \phi)$ there is a sequence of polynomials $(p_k)_{k \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow \infty} \|f - p_k\|_{H(p,q,\phi)} = 0. \quad (3.25)$$

From this and by the boundedness of the operator $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_\mu^\infty$, we have that

$$\left\| \mathfrak{R}_{u,\varphi}^m f - \mathfrak{R}_{u,\varphi}^m p_k \right\|_{H_\mu^\infty} \leq \left\| \mathfrak{R}_{u,\varphi}^m \right\|_{H(p,q,\phi) \rightarrow H_\mu^\infty} \|f - p_k\|_{H(p,q,\phi)} \rightarrow 0 \quad (3.26)$$

as $k \rightarrow \infty$. Hence $\mathfrak{R}_{u,\varphi}^m(H(p, q, \phi)) \subseteq H_{\mu,0}^\infty$, and consequently $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_{\mu,0}^\infty$ is bounded. \square

Theorem 3.4. *Assume $m \in \mathbb{N}$, $0 < p, q < \infty$, ϕ, μ are normal, φ is a holomorphic self-map of \mathbb{B} and $u \in H(\mathbb{B})$. Then $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_{\mu,0}^\infty$ is compact if and only if*

$$\lim_{|z| \rightarrow 1} \frac{\mu(z) |u(z)| |\varphi(z)|}{\phi(|\varphi(z)|) \left(1 - |\varphi(z)|^2\right)^{n/q+m}} = 0. \quad (3.27)$$

Proof. From (3.27), we see that (3.1) hold. This fact along with (3.3) implies that the set $\mathfrak{R}_{u,\varphi}^m(\{f \mid \|f\|_{H(p,q,\phi)} \leq 1\})$ is bounded in H_μ^∞ , moreover in $H_{\mu,0}^\infty$. By taking the supremum in (3.3) over the unit ball in $H(p, q, \phi)$, using (3.27) and applying Lemma 2.2 we obtain that the operator $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_{\mu,0}^\infty$ is compact.

If $\mathfrak{R}_{u,\varphi}^m : H(p, q, \phi) \rightarrow H_{\mu,0}^\infty$ is compact, then by Theorem 3.2, we have that condition (3.14) holds, which implies that for every $\varepsilon > 0$ there is an $r \in (0, 1)$ such that

$$\frac{\mu(z) |u(z)| |\varphi(z)|}{\phi(|\varphi(z)|) \left(1 - |\varphi(z)|^2\right)^{n/q+m}} < \varepsilon, \quad (3.28)$$

for $r < |\varphi(z)| < 1$.

As in Theorem 3.3, we have that (3.23) holds. Thus there is a $\sigma \in (0, 1)$ such that

$$\mu(z) |u(z)| |\varphi(z)| < \varepsilon \left(1 - r^2\right)^{n/q+m} \inf_{|\varphi(z)| \leq r} \phi(|\varphi(z)|), \quad (3.29)$$

for $\sigma < |z| < 1$.

If $|\varphi(z)| \leq r$ and $\sigma < |z| < 1$, then from (3.29), we obtain

$$\frac{\mu(z)|u(z)||\varphi(z)|}{(1 - |\varphi(z)|^2)^{n/q+m} \phi(|\varphi(z)|)} \leq \frac{\mu(z)|u(z)||\varphi(z)|}{(1 - r^2)^{n/q+m} \inf_{|\varphi(z)| \leq r} \phi(|\varphi(z)|)} < \varepsilon. \quad (3.30)$$

Using (3.30) and the fact that from (3.28), we have

$$\frac{\mu(z)|u(z)||\varphi(z)|}{\phi(|\varphi(z)|)(1 - |\varphi(z)|^2)^{n/q+m}} < \varepsilon, \quad (3.31)$$

for $\sigma < |z| < 1$ and $r < |\varphi(z)| < 1$, we get (3.27). □

4. Hilbert-Schmidt Norm of $\mathfrak{R}_{u,\varphi}^m : A_\alpha^2 \rightarrow A_\alpha^2$ and $\mathfrak{R}_{u,\varphi}^m : H^2 \rightarrow H^2$

In this section we calculate Hilbert-Schmidt norm of the operators $\mathfrak{R}_{u,\varphi}^m : A_\alpha^2 \rightarrow A_\alpha^2$ and $\mathfrak{R}_{u,\varphi}^m : H^2 \rightarrow H^2$. For some related results see [37, 38].

If \mathcal{H} is a separable Hilbert space, then the Hilbert-Schmidt norm $\|T\|_{\text{HS}}$ of an operator $T : \mathcal{H} \rightarrow \mathcal{H}$ is defined by

$$\|T\|_{\text{HS}} = \left(\sum_{n=1}^{\infty} \|Te_n\|^2 \right)^{1/2}, \quad (4.1)$$

where $\{e_n\}_{n \in \mathbb{N}}$ is an orthonormal basis on \mathcal{H} . The right-hand side in (4.1) does not depend on the choice of basis. Hence, it is larger than the operator norm $\|T\|_{\text{op}}$ of T .

Let $\langle \cdot, \cdot \rangle_\alpha$, $\alpha \geq -1$, be the usual scalar product on A_α^2 , where we regard that $A_{-1}^2 = H_2$. Since for each multi-index $\beta = (\beta_1, \dots, \beta_n) \in \mathbb{N}_0^n$

$$\int_{\mathbb{B}} |z^\beta|^2 dv_\alpha(z) = \frac{\beta! \Gamma(n + \alpha + 1)}{\Gamma(n + |\beta| + \alpha + 1)}, \quad (4.2)$$

where $\beta! = \beta_1! \cdots \beta_n!$, and

$$\int_{\mathbb{B}} z^\beta z^\gamma dv_\alpha(z) = 0, \quad \beta \neq \gamma \quad (4.3)$$

(see, e.g., [1]), we have that the vectors

$$e_\beta(z) = \sqrt{\frac{\Gamma(n + |\beta| + \alpha + 1)}{\beta! \Gamma(n + \alpha + 1)}} z^\beta, \quad \beta \in \mathbb{N}_0^n, \quad (4.4)$$

form an orthonormal basis in A_α^2 .

Theorem 4.1. Let $m \in \mathbb{N}_0$. Then Hilbert-Schmidt norm of the operator $\mathfrak{R}_{u,\varphi}^m$ on A_α^2 , $\alpha > -1$ is

$$\|\mathfrak{R}_{u,\varphi}^m\|_{HS} = \left(\int_{\mathbb{B}} |u(z)|^2 \left(\mathfrak{R}^m \frac{1}{\left(1 - \sum_{j=1}^n \omega_j\right)^{n+\alpha+1}} \right) \Big|_{\omega_j=|\varphi_j(z)|^2} dv_\alpha(z) \right)^{1/2}. \tag{4.5}$$

Proof. By using the definition of the Hilbert-Schmidt norm and the monotone convergence theorem, we have

$$\begin{aligned} \|\mathfrak{R}_{u,\varphi}^m\|_{HS}^2 &= \sum_{\beta} \|\mathfrak{R}_{u,\varphi}^m(e_\beta)\|_{2,\alpha}^2 = \sum_{\beta} \frac{\Gamma(n + |\beta| + \alpha + 1)}{\beta! \Gamma(n + \alpha + 1)} \|\mathfrak{R}_{u,\varphi}^m(z^\beta)\|_{2,\alpha}^2 \\ &= \sum_{\beta} \frac{\Gamma(n + |\beta| + \alpha + 1)}{\beta! \Gamma(n + \alpha + 1)} |\beta|^m \int_{\mathbb{B}} |u(z)|^2 \prod_{j=1}^n |\varphi_j(z)|^{2\beta_j} dv_\alpha(z) \\ &= \int_{\mathbb{B}} |u(z)|^2 \sum_{\beta} |\beta|^m \frac{\Gamma(n + |\beta| + \alpha + 1)}{\beta! \Gamma(n + \alpha + 1)} \prod_{j=1}^n |\varphi_j(z)|^{2\beta_j} dv_\alpha(z). \end{aligned} \tag{4.6}$$

We also have that

$$\begin{aligned} \left(1 - \sum_{j=1}^n \omega_j\right)^{-(n+\alpha+1)} &= \sum_{k=0}^{\infty} \binom{n}{\sum_{j=1}^n \omega_j}^k \frac{\Gamma(n + \alpha + k + 1)}{k! \Gamma(n + \alpha + 1)} \\ &= \sum_{k=0}^{\infty} \sum_{|l|=k} \frac{k!}{l!} \prod_{j=1}^n \omega_j^{l_j} \frac{\Gamma(n + \alpha + k + 1)}{k! \Gamma(n + \alpha + 1)} \\ &= \sum_l \frac{\Gamma(n + \alpha + |l| + 1)}{l! \Gamma(n + \alpha + 1)} \prod_{j=1}^n \omega_j^{l_j}, \end{aligned} \tag{4.7}$$

from which by taking the radial derivatives it follows that

$$\mathfrak{R}^m \frac{1}{\left(1 - \sum_{j=1}^n \omega_j\right)^{n+\alpha+1}} = \sum_l |l|^m \frac{\Gamma(n + \alpha + |l| + 1)}{l! \Gamma(n + \alpha + 1)} \prod_{j=1}^n \omega_j^{l_j}. \tag{4.8}$$

From (4.6) and (4.8) the result easily follows. □

Similar to Theorem 4.1 the following result regarding the case of the Hardy space is proved. We omit the proof.

Theorem 4.2. Let $m \in \mathbb{N}_0$. Then, Hilbert-Schmidt norm of the operator $\mathfrak{R}_{u,\varphi}^m$ on H^2 , is

$$\|\mathfrak{R}_{u,\varphi}^m\|_{HS} = \sup_{0 < r < 1} \left(\int_S |u(r\zeta)|^2 \left(\mathfrak{R}^m \frac{1}{(1 - \sum_{j=1}^n \omega_j)^n} \right) \Big|_{\omega_j = |\varphi_j(r\zeta)|^2} d\sigma(\zeta) \right)^{1/2}. \quad (4.9)$$

References

- [1] W. Rudin, *Function Theory in the Unit Ball of \mathbb{C}^n* , vol. 241 of *Grundlehren der Mathematischen Wissenschaften*, Springer, New York, NY, USA, 1980.
- [2] A. L. Shields and D. L. Williams, "Bonded projections, duality, and multipliers in spaces of analytic functions," *Transactions of the American Mathematical Society*, vol. 162, pp. 287–302, 1971.
- [3] K. D. Bierstedt and W. H. Summers, "Biduals of weighted Banach spaces of analytic functions," *Journal of the Australian Mathematical Society. Series A*, vol. 54, no. 1, pp. 70–79, 1993.
- [4] X. Fu and X. Zhu, "Weighted composition operators on some weighted spaces in the unit ball," *Abstract and Applied Analysis*, vol. 2008, Article ID 605807, 8 pages, 2008.
- [5] K. L. Avetisyan, "Hardy-Bloch type spaces and lacunary series on the polydisk," *Glasgow Mathematical Journal*, vol. 49, no. 2, pp. 345–356, 2007.
- [6] K. L. Avetisyan, "Fractional integro-differentiation in harmonic mixed norm spaces on a half-space," *Commentationes Mathematicae*, vol. 42, no. 4, pp. 691–709, 2001.
- [7] K. L. Avetisyan, "Continuous inclusions and Bergman type operators in n -harmonic mixed norm spaces on the polydisc," *Journal of Mathematical Analysis and Applications*, vol. 291, no. 2, pp. 727–740, 2004.
- [8] S. Stević, "Weighted composition operators between mixed norm spaces and H_α^∞ spaces in the unit ball," *Journal of Inequalities and Applications*, vol. 2007, Article ID 28629, 9 pages, 2007.
- [9] S. Stević, "On some new integral-type operators on spaces of holomorphic functions on the unit ball," in *Conference: Spaces of Analytic and Smooth Functions III*, Bedlewo, Poland, September 2009.
- [10] A. K. Sharma and S. D. Sharma, "Weighted composition operators between Bergman-type spaces," *Communications of the Korean Mathematical Society*, vol. 21, no. 3, pp. 465–474, 2006.
- [11] S. Li, S. Stević, and S. Li, "Composition followed by differentiation from mixed-norm spaces to α -Bloch spaces," *Sbornik: Mathematics*, vol. 199, no. 12, pp. 1847–1857, 2008.
- [12] S. Li and S. Stević, "Generalized composition operators on Zygmund spaces and Bloch type spaces," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1282–1295, 2008.
- [13] S. Li and S. Stević, "Products of Volterra type operator and composition operator from H^∞ and Bloch spaces to Zygmund spaces," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 40–52, 2008.
- [14] Y. Liu and Y. Yu, "On a Li-Stević integral-type operator from Bloch-type spaces into logarithmic Bloch spaces," *Integral Transforms and Special Functions*, vol. 21, no. 1-2, pp. 93–103, 2010.
- [15] A. K. Sharma, "Products of multiplication, composition and differentiation between weighted Bergman-Nevanlinna and Bloch-type spaces," *Turkish Journal of Mathematics*, vol. 34, pp. 1–17, 2010.
- [16] S. Stević, "Generalized composition operators from logarithmic Bloch spaces to mixed-norm spaces," *Utilitas Mathematica*, vol. 77, pp. 167–172, 2008.
- [17] S. Stević, "Norms of some operators from Bergman spaces to weighted and Bloch-type spaces," *Utilitas Mathematica*, vol. 76, pp. 59–64, 2008.
- [18] S. Stević, "On a new operator from H^∞ to the Bloch-type space on the unit ball," *Utilitas Mathematica*, vol. 77, pp. 257–263, 2008.
- [19] S. Stević, "On a new operator from the logarithmic Bloch space to the Bloch-type space on the unit ball," *Applied Mathematics and Computation*, vol. 206, no. 1, pp. 313–320, 2008.
- [20] S. Stević, "Compactness of the Hardy-Littlewood operator on spaces of harmonic functions," *Siberian Mathematical Journal*, vol. 50, no. 1, pp. 167–180, 2009.
- [21] S. Stević, "Integral-type operators from a mixed norm space to a Bloch-type space on the unit ball," *Siberian Mathematical Journal*, vol. 50, no. 6, pp. 1098–1105, 2009.
- [22] S. Stević, "On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball," *Journal of Mathematical Analysis and Applications*, vol. 354, no. 2, pp. 426–434, 2009.

- [23] S. Stević, "On an integral operator from the Zygmund space to the Bloch-type space on the unit ball," *Glasgow Mathematical Journal*, vol. 51, no. 2, pp. 275–287, 2009.
- [24] S. Stević, "Products of composition and differentiation operators on the weighted Bergman space," *Bulletin of the Belgian Mathematical Society. Simon Stevin*, vol. 16, no. 4, pp. 623–635, 2009.
- [25] S. Stević, "Products of integral-type operators and composition operators from a mixed norm space to Bloch-type spaces," *Siberian Mathematical Journal*, vol. 50, no. 4, pp. 726–736, 2009.
- [26] S. Stević, "On an integral-type operator from logarithmic Bloch-type and mixed-norm spaces to Bloch-type spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 12, pp. 6323–6342, 2009.
- [27] S. Stević, "Boundedness and compactness of an integral operator between H^∞ and a mixed norm space on the polydisk," *Siberian Mathematical Journal*, vol. 50, no. 3, pp. 495–497, 2009.
- [28] S. Stević, "On an integral operator between Bloch-type spaces on the unit ball," *Bulletin des Sciences Mathematiques*, vol. 134, no. 4, pp. 329–339, 2010.
- [29] S. Stević, "On operator P_ϕ^S from the logarithmic Bloch-type space to the mixed-norm space on the unit ball," *Applied Mathematics and Computation*, vol. 215, no. 12, pp. 4248–4255, 2010.
- [30] S.-I. Ueki, "Weighted composition operators on some function spaces of entire functions," *Bulletin of the Belgian Mathematical Society. Simon Stevin*, vol. 17, no. 2, pp. 343–353, 2010.
- [31] W. Yang, "On an integral-type operator between Bloch-type spaces," *Applied Mathematics and Computation*, vol. 215, no. 3, pp. 954–960, 2009.
- [32] X. Zhu, "Generalized weighted composition operators from Bloch type spaces to weighted Bergman spaces," *Indian Journal of Mathematics*, vol. 49, no. 2, pp. 139–150, 2007.
- [33] S. Stević and S.-I. Ueki, "Integral-type operators acting between weighted-type spaces on the unit ball," *Applied Mathematics and Computation*, vol. 215, no. 7, pp. 2464–2471, 2009.
- [34] H. J. Schwartz, *Composition operators on H^p* , Ph.D. thesis, University of Toledo, Ann Arbor, Mich, USA, 1969.
- [35] K. Madigan and A. Matheson, "Compact composition operators on the Bloch space," *Transactions of the American Mathematical Society*, vol. 347, no. 7, pp. 2679–2687, 1995.
- [36] Z. Hu, "Extended Cesàro operators on mixed norm spaces," *Proceedings of the American Mathematical Society*, vol. 131, no. 7, pp. 2171–2179, 2003.
- [37] S. Stević, "Continuity with respect to symbols of composition operators on the weighted Bergman space," *Taiwanese Journal of Mathematics*, vol. 11, no. 4, pp. 1177–1188, 2007.
- [38] S.-I. Ueki, "Hilbert-Schmidt weighted composition operator on the Fock space," *International Journal of Mathematical Analysis*, vol. 1, no. 13-16, pp. 769–774, 2007.