

## Research Article

# Positive Periodic Solutions of Nicholson-Type Delay Systems with Nonlinear Density-Dependent Mortality Terms

Wei Chen<sup>1</sup> and Lijuan Wang<sup>2</sup>

<sup>1</sup> School of Mathematics and Information, Shanghai Lixin University of Commerce,  
Shanghai 201620, China

<sup>2</sup> College of Mathematics, Physics and Information Engineering, Jiaxing University, Zhejiang,  
Jiaxing 314001, China

Correspondence should be addressed to Lijuan Wang, wanglijuan1976@yahoo.com.cn

Received 3 October 2012; Revised 9 November 2012; Accepted 9 November 2012

Academic Editor: Allan Peterson

Copyright © 2012 W. Chen and L. Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the periodic solutions for a class of Nicholson-type delay systems with nonlinear density-dependent mortality terms. By using coincidence degree theory, some criteria are obtained to guarantee the existence of positive periodic solutions of the model. Moreover, an example and a numerical simulation are given to illustrate our main results.

## 1. Introduction

In the last twenty years, the delay differential equations have been widely studied both in a theoretical context and in that of related applications [1–4]. As a famous and common delay dynamic system, Nicholson's blowflies model and its modifications have made remarkable progress that has been collected in [5] and the references cited there in. Recently, to describe the dynamics for the models of marine protected areas and B-cell chronic lymphocytic leukemia dynamics which belong to the Nicholson-type delay differential systems, Berezhansky et al. [6], Wang et al. [7], and Liu [8] studied the problems on the permanence, stability, and periodic solution of the following Nicholson-type delay systems:

$$N_1'(t) = -\alpha_1(t)N_1(t) + \beta_1(t)N_2(t) + \sum_{j=1}^m c_{1j}(t)N_1(t - \tau_{1j}(t))e^{-\gamma_{1j}(t)N_1(t - \tau_{1j}(t))},$$

$$N_2'(t) = -\alpha_2(t)N_2(t) + \beta_2(t)N_1(t) + \sum_{j=1}^m c_{2j}(t)N_2(t - \tau_{2j}(t))e^{-\gamma_{2j}(t)N_2(t - \tau_{2j}(t))}, \quad (1.1)$$

where  $\alpha_i, \beta_i, c_{ij}, \gamma_{ij}, \tau_{ij} \in C(\mathbb{R}, (0 + \infty))$ , and  $i = 1, 2, j = 1, 2, \dots, m$ .

In [5], Berezansky et al. also pointed out that a new study indicates that a linear model of density-dependent mortality will be most accurate for populations at low densities and marine ecologists are currently in the process of constructing new fishery models with nonlinear density-dependent mortality rates. Consequently, Berezansky et al. [5] presented an open problem: to reveal the dynamics of the following Nicholson's blowflies model with a nonlinear density-dependent mortality term:

$$N'(t) = -D(N(t)) + PN(t - \tau)e^{-aN(t - \tau)}, \quad (1.2)$$

where  $P$  is a positive constant and function  $D$  might have one of the following forms:  $D(N) = aN/(N + b)$  or  $D(N) = a - be^{-N}$  with positive constants  $a, b > 0$ .

Most recently, based upon the ideas in [5–8], Liu and Gong [9] established the results on the permanence for the Nicholson-type delay system with nonlinear density-dependent mortality terms. Consequently, the problem on periodic solutions of Nicholson-type system with  $D(N) = a - be^{-N}$  has been studied extensively in [10–13]. However, to the best of our knowledge, there exist few results on the existence of the positive periodic solutions of Nicholson-type delay system with  $D(N) = aN/(N + b)$ . Motivated by this, the main purpose of this paper is to give the conditions to guarantee the existence of positive periodic solutions of the following Nicholson-type delay system with nonlinear density-dependent mortality terms:

$$\begin{aligned} N_1'(t) &= -D_{11}(t, N_1(t)) + D_{12}(t, N_2(t)) + \sum_{j=1}^l c_{1j}(t)N_1(t - \tau_{1j}(t))e^{-\gamma_{1j}(t)N_1(t - \tau_{1j}(t))}, \\ N_2'(t) &= -D_{22}(t, N_2(t)) + D_{21}(t, N_1(t)) + \sum_{j=1}^l c_{2j}(t)N_2(t - \tau_{2j}(t))e^{-\gamma_{2j}(t)N_2(t - \tau_{2j}(t))}, \end{aligned} \quad (1.3)$$

under the admissible initial conditions

$$x_{t_0} = \varphi, \quad \varphi \in C_+ = C([-r_1, 0], \mathbb{R}_+^1) \times C([-r_2, 0], \mathbb{R}_+^1), \quad \varphi_i(0) > 0, \quad (1.4)$$

where  $D_{ij}(t, N) = a_{ij}(t)N/(b_{ij}(t) + N)$ ,  $a_{ij}, b_{ij}, c_{ik}, \gamma_{ik} : \mathbb{R} \rightarrow (0, +\infty)$  and  $\tau_{ik} : \mathbb{R} \rightarrow [0, +\infty)$  are all bounded continuous functions, and  $r_i = \max_{1 \leq k \leq l} \{\sup_{t \in \mathbb{R}^1} \tau_{ik}(t)\}$ ,  $i, j = 1, 2, k = 1, \dots, l$ .

For convenience, we introduce some notations. Throughout this paper, given a bounded continuous function  $g$  defined on  $\mathbb{R}^1$ , let  $g^+$  and  $g^-$  be defined as

$$g^- = \inf_{t \in \mathbb{R}^1} g(t), \quad g^+ = \sup_{t \in \mathbb{R}^1} g(t). \quad (1.5)$$

We also assume that  $a_{ij}, b_{ij}, c_{ik}, \gamma_{ik} : R \rightarrow (0, +\infty)$  and  $\tau_{ik} : R \rightarrow [0, +\infty)$  are all  $\omega$ -periodic functions,  $r_i = \max_{1 \leq k \leq l} \{\tau_{ik}^+\}$ , and  $i, j = 1, 2, k = 1, \dots, l$ .

Set

$$\begin{aligned}
 A_i &= 2 \int_0^\omega \frac{a_{ii}(t)}{b_{ii}(t)} dt, & B_i &= \sum_{j=1}^l \int_0^\omega c_{ij}(t) dt, & \gamma_i^+ &= \max_{1 \leq j \leq l} \{\gamma_{ij}^+\}, & \gamma_i^- &= \min_{1 \leq j \leq l} \{\gamma_{ij}^-\}, \\
 D_1 &= \int_0^\omega a_{12}(t) dt, & D_2 &= \int_0^\omega a_{21}(t) dt, & C_i &= \int_0^\omega a_{ii}(t) dt, & i &= 1, 2.
 \end{aligned}
 \tag{1.6}$$

Let  $R^n(R_+^n)$  be the set of all (nonnegative) real vectors; we will use  $x = (x_1, x_2, \dots, x_n)^T \in R^n$  to denote a column vector, in which the symbol  $(^T)$  denotes the transpose of a vector. We let  $|x|$  denote the absolute-value vector given by  $|x| = (|x_1|, |x_2|, \dots, |x_n|)^T$  and define  $\|x\| = \max_{1 \leq i \leq n} |x_i|$ . For matrix  $A = (a_{ij})_{n \times n}$ ,  $A^T$  denotes the transpose of  $A$ . A matrix or vector  $A \geq 0$  means that all entries of  $A$  are greater than or equal to zero.  $A > 0$  can be defined similarly. For matrices or vectors  $A$  and  $B$ ,  $A \geq B$  (resp.  $A > B$ ) means that  $A - B \geq 0$  (resp.  $A - B > 0$ ). We also define the derivative and integral of vector function  $x(t) = (x_1(t), x_2(t))^T$  as  $x' = (x'_1(t), x'_2(t))^T$  and  $\int_0^\omega x(t) dt = (\int_0^\omega x_1(t) dt, \int_0^\omega x_2(t) dt)^T$ .

The organization of this paper is as follows. In the next section, some sufficient conditions for the existence of the positive periodic solutions of model (1.3) are given by using the method of coincidence degree. In Section 3, an example and numerical simulation are given to illustrate our results obtained in the previous section.

## 2. Existence of Positive Periodic Solutions

In order to study the existence of positive periodic solutions, we first introduce the continuation theorem as follows.

**Lemma 2.1** (continuation theorem [14]). *Let  $X$  and  $Z$  be two Banach spaces. Suppose that  $L : D(L) \subset X \rightarrow Z$  is a Fredholm operator with index zero and  $\widetilde{N} : X \rightarrow Z$  is  $L$ -compact on  $\overline{\Omega}$ , where  $\Omega$  is an open subset of  $X$ . Moreover, assume that all the following conditions are satisfied:*

- (1)  $Lx \neq \lambda \widetilde{N}x$ , for all  $x \in \partial\Omega \cap D(L)$ ,  $\lambda \in (0, 1)$ ;
- (2)  $\widetilde{N}x \notin \text{Im } L$ , for all  $x \in \partial\Omega \cap \text{Ker } L$ ;
- (3) the Brouwer degree

$$\text{deg}\{Q\widetilde{N}, \Omega \cap \text{Ker } L, 0\} \neq 0.
 \tag{2.1}$$

Then equation  $Lx = \widetilde{N}x$  has at least one solution in  $\text{dom } L \cap \overline{\Omega}$ .

Our main result is given in the following theorem.

**Theorem 2.2.** *Suppose*

$$C_i > 2D_i, \quad \ln \frac{2B_i}{A_i} > A_i, \quad i = 1, 2, \quad (2.2)$$

$$\sum_{j=1}^l \frac{c_{1j}^+}{a_{11}^- \gamma_{1j}^- e} + \frac{a_{12}^+}{a_{11}^-} < 1, \quad \sum_{j=1}^l \frac{c_{2j}^+}{a_{22}^- \gamma_{2j}^- e} + \frac{a_{21}^+}{a_{22}^-} < 1. \quad (2.3)$$

Then (1.3) has a positive  $\omega$ -periodic solution.

*Proof.* Set  $N(t) = (N_1(t), N_2(t))^T$  and  $N_i(t) = e^{x_i(t)}$  ( $i = 1, 2$ ). Then (1.3) can be rewritten as

$$\begin{aligned} x_1'(t) &= -\frac{a_{11}(t)}{b_{11}(t) + e^{x_1(t)}} + \frac{a_{12}(t)e^{x_2(t)-x_1(t)}}{b_{12}(t) + e^{x_2(t)}} \\ &\quad + \sum_{j=1}^l c_{1j}(t)e^{x_1(t-\tau_{1j}(t))-x_1(t)-\gamma_{1j}(t)e^{x_1(t-\tau_{1j}(t))}} := \Delta_1(x, t), \\ x_2'(t) &= -\frac{a_{22}(t)}{b_{22}(t) + e^{x_2(t)}} + \frac{a_{21}(t)e^{x_1(t)-x_2(t)}}{b_{21}(t) + e^{x_1(t)}} \\ &\quad + \sum_{j=1}^l c_{2j}(t)e^{x_2(t-\tau_{2j}(t))-x_2(t)-\gamma_{2j}(t)e^{x_2(t-\tau_{2j}(t))}} := \Delta_2(x, t). \end{aligned} \quad (2.4)$$

As usual, let  $X = Z = \{x = (x_1(t), x_2(t))^T \in C(\mathbb{R}, \mathbb{R}^2) : x(t + \omega) = x(t) \text{ for all } t \in \mathbb{R}\}$  be Banach spaces equipped with the supremum norm  $\|\cdot\|$ . For any  $x \in X$ , because of periodicity, it is easy to see that  $\Delta(x, \cdot) = (\Delta_1(x, \cdot), \Delta_2(x, \cdot))^T \in C(\mathbb{R}, \mathbb{R}^2)$  is  $\omega$ -periodic. Let

$$\begin{aligned} L : D(L) &= \left\{ x \in X : x \in C^1(\mathbb{R}, \mathbb{R}^2) \right\} \ni x \mapsto x' = (x_1', x_2')^T \in Z, \\ P : X \ni x &\mapsto \left( \frac{1}{\omega} \int_0^\omega x_1(s) ds, \frac{1}{\omega} \int_0^\omega x_2(s) ds \right)^T \in X, \\ Q : Z \ni z &\mapsto \left( \frac{1}{\omega} \int_0^\omega z_1(s) ds, \frac{1}{\omega} \int_0^\omega z_2(s) ds \right)^T \in Z, \\ \widetilde{N} : X \ni x &\mapsto \Delta(x, \cdot) \in Z. \end{aligned} \quad (2.5)$$

It is easy to see that

$$\begin{aligned} \text{Im } L &= \left\{ x \mid x \in Z, \int_0^\omega x(s) ds = (0, 0)^T \right\}, \quad \text{Ker } L = \mathbb{R}^2, \\ \text{Im } P &= \text{Ker } L, \quad \text{Ker } Q = \text{Im } L. \end{aligned} \quad (2.6)$$

Thus, the operator  $L$  is a Fredholm operator with index zero. Furthermore, denoting by  $L_P^{-1} : \text{Im } L \rightarrow D(L) \cap \text{Ker } P$  the inverse of  $L|_{D(L) \cap \text{Ker } P}$ , we have

$$\begin{aligned} L_P^{-1}y(t) &= -\frac{1}{\omega} \int_0^\omega \int_0^t y(s) ds dt + \int_0^t y(s) ds \\ &= \left( -\frac{1}{\omega} \int_0^\omega \int_0^t y_1(s) ds dt + \int_0^t y_1(s) ds, -\frac{1}{\omega} \int_0^\omega \int_0^t y_2(s) ds dt + \int_0^t y_2(s) ds \right)^T. \end{aligned} \tag{2.7}$$

It follows that

$$Q\widetilde{N}x = \frac{1}{\omega} \int_0^\omega \widetilde{N}x(t) dt = \left( \frac{1}{\omega} \int_0^\omega \Delta_1(x(t), t) dt, \frac{1}{\omega} \int_0^\omega \Delta_2(x(t), t) dt \right)^T, \tag{2.8}$$

$$\begin{aligned} L_P^{-1}(I - Q)\widetilde{N}x &= \int_0^t \widetilde{N}x(s) ds - \frac{t}{\omega} \int_0^\omega \widetilde{N}x(s) ds - \frac{1}{\omega} \int_0^\omega \int_0^t \widetilde{N}x(s) ds dt \\ &\quad + \frac{1}{\omega} \int_0^\omega \int_0^t Q\widetilde{N}x(s) ds dt. \end{aligned} \tag{2.9}$$

Obviously,  $Q\widetilde{N}$  and  $L_P^{-1}(I - Q)\widetilde{N}$  are continuous. It is not difficult to show that  $L_P^{-1}(I - Q)\widetilde{N}(\overline{\Omega})$  is compact for any open bounded set  $\Omega \subset X$  by using the Arzela-Ascoli theorem. Moreover,  $Q\widetilde{N}(\overline{\Omega})$  is clearly bounded. Thus  $\widetilde{N}$  is  $L$ -compact on  $\overline{\Omega}$  with any open bounded set  $\Omega \subset X$ .

Considering the operator equation  $Lx = \lambda\widetilde{N}x, \lambda \in (0, 1)$ , we have

$$x'(t) = (x'_1(t), x'_2(t))^T = \lambda\Delta(x, t) = (\lambda\Delta_1(x, t), \lambda\Delta_2(x, t))^T. \tag{2.10}$$

Suppose that  $x = (x_1(t), x_2(t))^T \in X$  is a solution of (2.10) for some  $\lambda \in (0, 1)$ .

Firstly, we claim that there exists a positive number  $H$  such that  $\|x\| < H$ . Integrating the first equation of (2.10) and in view of  $x \in X$ , it results that

$$0 = \int_0^\omega x'_1(t) dt = \lambda \int_0^\omega \Delta_1(x, t) dt, \tag{2.11}$$

which together with (2.4) implies that

$$\begin{aligned} &\int_0^\omega \left| \sum_{j=1}^l c_{1j}(t) e^{x_1(t-\tau_{1j}(t)) - x_1(t) - \gamma_j(t) e^{x_1(t-\tau_{1j}(t))}} + \frac{a_{12}(t) e^{x_2(t) - x_1(t)}}{b_{12}(t) + e^{x_2(t)}} \right| dt \\ &= \int_0^\omega \frac{a_{11}(t)}{b_{11}(t) + e^{x_1(t)}} dt \\ &< \int_0^\omega \frac{a_{11}(t)}{b_{11}(t)} dt. \end{aligned} \tag{2.12}$$

Similarly, we have

$$\begin{aligned}
 & \int_0^\omega \left| \sum_{j=1}^l c_{2j}(t) e^{x_2(t-\tau_{2j}(t))-x_2(t)-\gamma_{2j}(t)} e^{x_2(t-\tau_{2j}(t))} + \frac{a_{21}(t) e^{x_1(t)-x_2(t)}}{b_{21}(t) + e^{x_1(t)}} \right| dt \\
 &= \int_0^\omega \frac{a_{22}(t)}{b_{22}(t) + e^{x_2(t)}} dt \\
 &< \int_0^\omega \frac{a_{22}(t)}{b_{22}(t)} dt.
 \end{aligned} \tag{2.13}$$

It follows from (2.12) and (2.13) that

$$\begin{aligned}
 \int_0^\omega |x_1'(t)| dt &\leq \lambda \int_0^\omega \left| \sum_{j=1}^l c_{1j}(t) e^{x_1(t-\tau_{1j}(t))-x_1(t)-\gamma_{1j}(t)} e^{x_1(t-\tau_{1j}(t))} + \frac{a_{12}(t) e^{x_2(t)-x_1(t)}}{b_{12}(t) + e^{x_2(t)}} \right| dt \\
 &\quad + \lambda \int_0^\omega \left| \frac{a_{11}(t)}{b_{11}(t) + e^{x_1(t)}} \right| dt \\
 &< 2 \int_0^\omega \frac{a_{11}(t)}{b_{11}(t)} dt = A_1,
 \end{aligned} \tag{2.14}$$

$$\begin{aligned}
 \int_0^\omega |x_2'(t)| dt &\leq \lambda \int_0^\omega \left| \sum_{j=1}^l c_{2j}(t) e^{x_2(t-\tau_{2j}(t))-x_2(t)-\gamma_{2j}(t)} e^{x_2(t-\tau_{2j}(t))} + \frac{a_{21}(t) e^{x_1(t)-x_2(t)}}{b_{21}(t) + e^{x_1(t)}} \right| dt \\
 &\quad + \lambda \int_0^\omega \left| \frac{a_{22}(t)}{b_{22}(t) + e^{x_2(t)}} \right| dt \\
 &< 2 \int_0^\omega \frac{a_{22}(t)}{b_{22}(t)} dt = A_2.
 \end{aligned} \tag{2.15}$$

Since  $x \in X$ , there exist  $\xi_1, \xi_2, \eta_1, \eta_2 \in [0, \omega]$  such that

$$x_i(\xi_i) = \min_{t \in [0, \omega]} x_i(t), \quad x_i(\eta_i) = \max_{t \in [0, \omega]} x_i(t), \quad x_i'(\xi_i) = x_i'(\eta_i) = 0, \quad i = 1, 2. \tag{2.16}$$

It follows from (2.12) and (2.14) that

$$\begin{aligned}
 \frac{A_1}{2} &= \int_0^\omega \frac{a_{11}(t)}{b_{11}(t)} dt > \int_0^\omega \frac{a_{11}(t)}{b_{11}(t) + e^{x_1(t)}} dt \\
 &= \int_0^\omega \sum_{j=1}^l c_{1j}(t) e^{x_1(t-\tau_{1j}(t))-x_1(t)-\gamma_{1j}(t)} e^{x_1(t-\tau_{1j}(t))} dt + \int_0^\omega \frac{a_{12}(t) e^{x_2(t)-x_1(t)}}{b_{12}(t) + e^{x_2(t)}} dt
 \end{aligned}$$

$$\begin{aligned}
 &> e^{x_1(\xi_1)-x_1(\eta_1)-\gamma_1^+ e^{x_1(\eta_1)}} \sum_{j=1}^l \int_0^\omega c_{1j}(t) dt \\
 &= B_1 e^{x_1(\xi_1)-x_1(\eta_1)-\gamma_1^+ e^{x_1(\eta_1)}},
 \end{aligned} \tag{2.17}$$

which implies that

$$x_1(\xi_1) < \ln \frac{A_1}{2B_1} + x_1(\eta_1) + \gamma_1^+ e^{x_1(\eta_1)}. \tag{2.18}$$

Using (2.14) yields

$$x_1(t) \leq x_1(\xi_1) + \int_0^\omega |x_1'(t)| dt < \ln \frac{A_1}{2B_1} + x_1(\eta_1) + \gamma_1^+ e^{x_1(\eta_1)} + A_1, \quad t \in [0, \omega]. \tag{2.19}$$

In particular,

$$x_1(\eta_1) < x_1(\xi_1) + \int_0^\omega |x_1'(t)| dt < \ln \frac{A_1}{2B_1} + x_1(\eta_1) + \gamma_1^+ e^{x_1(\eta_1)} + A_1. \tag{2.20}$$

It follows that

$$x_1(\eta_1) > \ln \left( \frac{1}{\gamma_1^+} \left( \ln \frac{2B_1}{A_1} - A_1 \right) \right). \tag{2.21}$$

Again from (2.14), we have

$$x_1(t) \geq x_1(\eta_1) - \int_0^\omega |x_1'(t)| dt > \ln \left( \frac{1}{\gamma_1^+} \left( \ln \frac{2B_1}{A_1} - A_1 \right) \right) - A_1 := H_{11}, \quad t \in [0, \omega]. \tag{2.22}$$

Similarly, we can obtain

$$x_2(t) \geq x_2(\eta_2) - \int_0^\omega |x_2'(t)| dt > \ln \left( \frac{1}{\gamma_2^+} \left( \ln \frac{2B_2}{A_2} - A_2 \right) \right) - A_2 := H_{21}, \quad t \in [0, \omega]. \tag{2.23}$$

Since  $x_1'(\xi_1) = 0$ , from (2.10), we have

$$\begin{aligned}
 \frac{a_{11}(\xi_1)}{b_{11}(\xi_1) + e^{x_1(\xi_1)}} &= \sum_{j=1}^l c_{1j}(\xi_1) e^{x_1(\xi_1-\tau_{1j}(\xi_1))-x_1(\xi_1)-\gamma_j(\xi_1) e^{x_1(\xi_1-\tau_{1j}(\xi_1))}} \\
 &+ \frac{a_{12}(\xi_1) e^{x_2(\xi_1)-x_1(\xi_1)}}{b_{12}(\xi_1) + e^{x_2(\xi_1)}}.
 \end{aligned} \tag{2.24}$$

Hence, from (2.24) and the fact that  $\sup_{u \geq 0} ue^{-u} = 1/e$ , we have

$$\begin{aligned} \frac{e^{x_1(\xi_1)}}{b_{11}^+ + e^{x_1(\xi_1)}} &\leq \frac{e^{x_1(\xi_1)}}{b_{11}(\xi_1) + e^{x_1(\xi_1)}} \\ &= \sum_{j=1}^l \frac{c_{1j}(\xi_1)}{a_{11}(\xi_1)\gamma_{1j}(\xi_1)} \gamma_{1j}(\xi_1) e^{x_1(\xi_1 - \tau_{1j}(\xi_1))} e^{-\gamma_{1j}(\xi_1) e^{x_1(\xi_1 - \tau_{1j}(\xi_1))}} \\ &\quad + \frac{a_{12}(\xi_1)}{a_{11}(\xi_1)(1 + b_{12}(\xi_1)e^{-x_2(\xi_1)})} \\ &< \sum_{j=1}^l \frac{c_{1j}^+}{a_{11}^- \gamma_{1j}^- e} + \frac{a_{12}^+}{a_{11}^-}. \end{aligned} \quad (2.25)$$

Noting that  $u/(b_{11}^+ + u)$  is strictly monotone increasing on  $[0, +\infty)$  and

$$\sup_{u \geq 0} \frac{u}{b_{11}^+ + u} = 1 > \sum_{j=1}^l \frac{c_{1j}^+}{a_{11}^- \gamma_{1j}^- e} + \frac{a_{12}^+}{a_{11}^-}, \quad (2.26)$$

it is clear that there exists a constant  $k_1 > 0$  such that

$$\frac{u}{b_{11}^+ + u} > \sum_{j=1}^l \frac{c_{1j}^+}{a_{11}^- \gamma_{1j}^- e} + \frac{a_{12}^+}{a_{11}^-} \quad \forall u \in [k_1, +\infty). \quad (2.27)$$

In view of (2.25) and (2.27), we get

$$e^{x_1(\xi_1)} \leq k_1, \quad x_1(\xi_1) \leq \ln k_1. \quad (2.28)$$

In the same way, there exists a constant  $k_2 > 0$  such that

$$x_2(\xi_2) \leq \ln k_2. \quad (2.29)$$

Again from (2.14), (2.15), (2.28), and (2.29), we get

$$\begin{aligned} x_1(t) &\leq x_1(\xi_1) + \int_0^\omega |x_1'(t)| dt < \ln k_1 + A_1, \\ x_2(t) &\leq x_2(\xi_2) + \int_0^\omega |x_2'(t)| dt < \ln k_2 + A_2. \end{aligned} \quad (2.30)$$

Then, we can choose two sufficiently large positive constants  $H_{12} > \ln k_1 + A_1$  and  $H_{22} > \ln k_2 + A_2$  such that

$$x_1(t) < H_{12}, \quad x_2(t) < H_{22}, \quad \ln b_{11}^+ < H_{12}, \quad \ln b_{22}^+ < H_{22}. \quad (2.31)$$



Let  $H > \max\{|H_{11}|, |H_{21}|, H_{12}, H_{22}\}$  be a fix constant such that

$$e^H > \frac{1}{\gamma_1^-} \left( H - \ln \frac{C_1 - 2D_1}{2B_1} \right), \quad e^H > \frac{1}{\gamma_2^-} \left( H - \ln \frac{C_2 - 2D_2}{2B_2} \right). \quad (2.32)$$

Then (2.22), (2.23), and (2.31) imply that  $\|x\| < H$ , if  $x \in X$  is solution of (2.10). So we can define an open bounded set as  $\Omega = \{x \in X : \|x\| < H\}$  such that there is no  $\lambda \in (0, 1)$  and  $x \in \partial\Omega$  such that  $Lx = \lambda \widetilde{N}x$ . That is to say  $Lx \neq \lambda \widetilde{N}x$  for all  $x \in \partial\Omega \cap D(L), \lambda \in (0, 1)$ .

Secondly, we prove that  $\widetilde{N}x \notin \text{Im} L$  for all  $x \in \partial\Omega \cap \text{Ker} L$ . That is  $((Q\widetilde{N}(x))_1, (Q\widetilde{N}(x))_2)^T \neq (0, 0)^T$  for all  $x \in \partial\Omega \cap \text{Ker} L$ .

If  $x(t) = (x_1(t), x_2(t))^T \in \partial\Omega \cap \text{Ker} L$ , then  $x(t)$  is a constant vector in  $R^2$ , and there exists some  $i \in \{1, 2\}$  such that  $|x_i| = H$ . Assume  $|x_1| = H$ , so that  $x_1 = \pm H$ . Then, we claim

$$\left( Q\widetilde{N}(x) \right)_1 > 0 \quad \text{for } x_1 = -H, \quad \left( Q\widetilde{N}(x) \right)_1 < 0 \quad \text{for } x_1 = H. \quad (2.33)$$

If  $(Q\widetilde{N}(x))_1 \leq 0$  for  $x_1 = -H$ , it follows from (2.2) and (2.8) that

$$\int_0^\omega \Delta_1(x, t) dt \leq 0, \quad \text{for } x_1 = -H. \quad (2.34)$$

Hence,

$$\begin{aligned} \frac{A_1}{2} &= \int_0^\omega \frac{a_{11}(t)}{b_{11}(t)} dt \\ &> \int_0^\omega \frac{a_{11}(t)}{b_{11}(t) + e^{-H}} dt \\ &\geq \int_0^\omega \left[ \sum_{j=1}^l c_{1j}(t) e^{-\gamma_{1j}(t)e^{-H}} + \frac{a_{12}(t)e^{x_2+H}}{b_{12}(t) + e^{x_2}} \right] dt \\ &> \int_0^\omega \sum_{j=1}^l c_{1j}(t) e^{-\gamma_{1j}^+ e^{-H}} dt \\ &\geq e^{-\gamma_1^+ e^{-H}} \sum_{j=1}^l \int_0^\omega c_{1j}(t) dt \\ &= B_1 e^{-\gamma_1^+ e^{-H}}, \end{aligned} \quad (2.35)$$

which implies

$$-H > \ln \left( \frac{1}{\gamma_1^+} \ln \frac{2B_1}{A_1} \right) > \ln \left( \frac{1}{\gamma_1^+} \left( \ln \frac{2B_1}{A_1} - A_1 \right) \right) - A_1 = H_{11}. \quad (2.36)$$

This is a contradiction and implies that  $(Q\widetilde{N}(x))_1 > 0$  for  $x_1 = -H$ .

If  $(Q\widetilde{N}(x))_1 \geq 0$  for  $x_1 = H$ , it follows from (2.2) and (2.8) that

$$\begin{aligned}
 & \int_0^\omega \Delta_1(x, t) dt \geq 0, \quad \text{for } x_1 = H, \\
 \frac{C_1}{2} e^{-H} &= \int_0^\omega \frac{a_{11}(t)}{2e^H} dt \\
 &< \int_0^\omega \frac{a_{11}(t)}{b_{11}(t) + e^H} dt \\
 &\leq \int_0^\omega \sum_{j=1}^l c_{1j}(t) e^{-\gamma_j(t)e^H} dt + \int_0^\omega \frac{a_{12}(t) e^{x_2-H}}{b_{12}(t) + e^{x_2}} dt \\
 &\leq \int_0^\omega \sum_{j=1}^l c_{1j}(t) e^{-\gamma_j^- e^H} dt + \int_0^\omega \frac{a_{12}(t)}{b_{12}(t) e^{H-x_2} + e^H} dt \\
 &< e^{-\gamma_1^- e^H} \sum_{j=1}^l \int_0^\omega c_{1j}(t) dt + \int_0^\omega \frac{a_{12}(t)}{e^H} dt \\
 &= B_1 e^{-\gamma_1^- e^H} + D_1 e^{-H}.
 \end{aligned} \tag{2.37}$$

Consequently,

$$e^H < \frac{1}{\gamma_1^-} \left( H - \ln \frac{C_1 - 2D_1}{2B_1} \right), \tag{2.38}$$

a contradiction to the choice of  $H$ . Thus,  $(Q\widetilde{N}(x))_1 < 0$  for  $x_1 = H$ .

Similarly, if  $|x_2| = H$ , we obtain

$$(Q\widetilde{N}(x))_2 > 0 \quad \text{for } x_2 = -H, \quad (Q\widetilde{N}(x))_2 < 0 \quad \text{for } x_2 = H. \tag{2.39}$$

Consequently, (2.33) and (2.39) imply that  $\widetilde{N}x \notin \text{Im } L$  for all  $x \in \partial\Omega \cap \text{Ker } L$ .


Furthermore, let  $0 \leq \mu \leq 1$  and define continuous function  $H(x, \mu)$  by setting

$$H(x, \mu) = -(1 - \mu)x + \mu Q\widetilde{N}x. \tag{2.40}$$

For all  $x(t) = (x_1(t), x_2(t))^T \in \partial\Omega \cap \text{Ker } L$ , then there exists some  $i \in \{1, 2\}$  such that  $|x_i| = H$ . There are two cases:  $x_1 = \pm H$  or  $x_2 = \pm H$ . When  $x_1 = H$  or  $x_2 = H$ , from (2.33) and (2.39), it is obvious that  $(H(x, \mu))_1 < 0$  or  $(H(x, \mu))_2 < 0$ . Similarly, if  $x_1 = -H$  or  $x_2 = -H$ , it results that  $(H(x, \mu))_1 > 0$  or  $(H(x, \mu))_2 > 0$ . Hence  $H(x, \mu) \neq (0, 0)^T$  for all  $x \in \partial\Omega \cap \text{ker } L$ .

Finally, using the homotopy invariance theorem, we obtain

$$\deg\{Q\widetilde{N}, \Omega \cap \text{ker } L, (0, 0)^T\} = \deg\{-x, \Omega \cap \text{ker } L, (0, 0)^T\} \neq 0. \tag{2.41}$$



**Figure 1:** Numerical solution  $N(t) = (N_1(t), N_2(t))^T$  of systems (3.1) for initial value  $\varphi(t) \equiv (1, 2)^T$ .

It then follows from the continuation theorem that  $Lx = \widetilde{N}x$  has a solution

$$x^*(t) = (x_1^*(t), x_2^*(t))^T \in \text{Dom } L \cap \overline{\Omega}, \tag{2.42}$$

which is an  $\omega$ -periodic solution to (2.4). Therefore  $N^*(t) = (N_1^*(t), N_2^*(t))^T = (e^{x_1^*(t)}, e^{x_2^*(t)})^T$  is a positive  $\omega$ -periodic solution of (1.3) and the proof is complete.  $\square$

### 3. An Example

In this section, we give an example to demonstrate the results obtained in the previous section.

*Example 3.1.* Consider the following Nicholson-type delay system with nonlinear density-dependent mortality terms:

$$\begin{aligned} N_1'(t) &= -\frac{(5 + \sin t)N_1(t)}{5 + \sin t + N_1(t)} + \frac{(2 + \cos t)N_2(t)}{2 + \cos t + N_2(t)} \\ &\quad + e^{4\pi} \left(1 + \frac{\cos t}{4}\right) N_1(t - |2 + \cos t|) e^{-e^{4\pi + |\sin t|} N_1(t - |2 + \cos t|)} \\ &\quad + e^{4\pi} \left(1 + \frac{\sin t}{4}\right) N_1(t - |2 + \sin t|) e^{-e^{4\pi + |\cos t|} N_1(t - |2 + \sin t|)}, \\ N_2'(t) &= -\frac{(5 + \cos t)N_2(t)}{5 + \cos t + N_2(t)} + \frac{(2 + \sin t)N_1(t)}{2 + \sin t + N_1(t)} \end{aligned}$$

$$\begin{aligned}
& + e^{4\pi} \left( 1 + \frac{\sin t}{4} \right) N_2(t - |2 + \sin t|) e^{-e^{4\pi} |\cos t|} N_2(t - |2 + \sin t|) \\
& + e^{4\pi} \left( 1 + \frac{\cos t}{4} \right) N_2(t - |2 + \cos t|) e^{-e^{4\pi} |\sin t|} N_2(t - |2 + \cos t|).
\end{aligned} \tag{3.1}$$

Obviously,  $A_i = 4\pi$ ,  $B_i = 4\pi e^{4\pi}$ ,  $C_i = 10\pi$ ,  $D_i = 4\pi$  ( $i = 1, 2$ ),  $c_{ij}^+ = (5/4)e^{4\pi}$ ,  $\gamma_{ij}^- = e^{4\pi}$  ( $i, j = 1, 2$ ),  $a_{12}^+ = a_{21}^+ = 3$ ,  $a_{11}^- = a_{22}^- = 4$ , then

$$\begin{aligned}
\ln \frac{2B_i}{A_i} &= \ln 2 + 4\pi > 4\pi = A_i, & C_i &= 10\pi > 8\pi = 2D_i, & i &= 1, 2, \\
\sum_{j=1}^l \frac{c_{1j}^+}{a_{11}^- \gamma_{1j}^-} e + \frac{a_{12}^+}{a_{11}^-} &= \frac{5}{8e} + \frac{3}{4} \approx 0.9799 < 1, \\
\sum_{j=1}^l \frac{c_{2j}^+}{a_{22}^- \gamma_{2j}^-} e + \frac{a_{21}^+}{a_{22}^-} &= \frac{5}{8e} + \frac{3}{4} \approx 0.9799 < 1,
\end{aligned} \tag{3.2}$$

which means the conditions in Theorem 2.2 hold. Hence, the model (3.1) has a positive  $2\pi$ -periodic solution in  $\bar{\Omega}$ , where  $\Omega = \{x \in X : \|x\| < 10000\}$ . The fact is verified by the numerical simulation in Figure 1.

*Remark 3.2.* Equation (3.1) is a form of Nicholson's blowflies delayed systems with nonlinear density-dependent mortality terms, but as far as we know there are no that results can be applicable to (3.1) to obtain the existence of positive  $2\pi$ -periodic solutions. This implies the results of this paper are essentially new.

## Acknowledgments

The authors would like to express the sincere appreciation to the reviewers for their helpful comments in improving the presentation and quality of the paper. In particular, the authors expresses the sincere gratitude to Prof. Bingwen Liu for the helpful discussion when this work is carried out. This work was supported by National Natural Science Foundation of China (Grant nos. 11201184 and 11101283), Innovation Program of Shanghai Municipal Education Commission (Grant no. 13YZ127), the Natural Scientific Research Fund of Zhejiang Provincial of China (Grant nos. Y6110436, LY12A01018), and the Natural Scientific Research Fund of Zhejiang Provincial Education Department of China (Grant no. Z201122436).

## References

- [1] M. de la Sen and N. Luo, "On the uniform exponential stability of a wide class of linear time-delay systems," *Journal of Mathematical Analysis and Applications*, vol. 289, no. 2, pp. 456–476, 2004.
- [2] M. de la Sen, "Stability of impulsive time-varying systems and compactness of the operators mapping the input space into the state and output spaces," *Journal of Mathematical Analysis and Applications*, vol. 321, no. 2, pp. 621–650, 2006.

- [3] H. Lu and W. Wang, "Dynamics of a nonautonomous Leslie-Gower type food chain model with delays," *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 380279, 19 pages, 2011.
- [4] X. Fan, Z. Wang, and F. Jiang, "Dynamics of mutualism-competition-predator system with Beddington-DeAngelis functional responses and impulsive perturbations," *Abstract and Applied Analysis*, vol. 2012, Article ID 963486, 33 pages, 2012.
- [5] L. Berezensky, E. Braverman, and L. Idels, "Nicholson's blowflies differential equations revisited: main results and open problems," *Applied Mathematical Modelling*, vol. 34, no. 6, pp. 1405–1417, 2010.
- [6] L. Berezensky, L. Idels, and L. Troib, "Global dynamics of Nicholson-type delay systems with applications," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 1, pp. 436–445, 2011.
- [7] W. Wang, L. Wang, and W. Chen, "Existence and exponential stability of positive almost periodic solution for Nicholson-type delay systems," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 4, pp. 1938–1949, 2011.
- [8] B. Liu, "The existence and uniqueness of positive periodic solutions of Nicholson-type delay systems," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 6, pp. 3145–3151, 2011.
- [9] B. Liu and S. Gong, "Permanence for Nicholson-type delay systems with nonlinear density-dependent mortality terms," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 4, pp. 1931–1937, 2011.
- [10] W. Wang, "Positive periodic solutions of delayed Nicholson's blowflies models with a nonlinear density-dependent mortality term," *Applied Mathematical Modelling*, vol. 36, no. 10, pp. 4708–4713, 2012.
- [11] X. Hou and L. Duan, "New results on periodic solutions of delayed Nicholson's blowflies models," *Electronic Journal of Qualitative Theory of Differential Equations*, no. 24, pp. 1–11, 2012.
- [12] X. Hou, L. Duan, and Z. Huang, "Permanence and periodic solutions for a class of delay Nicholson's blowflies models," *Applied Mathematical Modelling*, vol. 37, pp. 1537–1544, 2013.
- [13] Z. Chen, "Periodic solutions for Nicholson-type delay system with nonlinear density-dependent mortality terms," *Electronic Journal of Qualitative Theory of Differential Equations*, vol. 56, pp. 1–9, 2012.
- [14] R. E. Gaines and J. L. Mawhin, *Coincidence Degree, and Nonlinear Differential Equations*, vol. 568 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1977.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

