

Research Article

The Univalence Conditions of Some Integral Operators

Laura Stanciu

Department of Mathematics, University of Pitești, Argeș, Romania

Correspondence should be addressed to Laura Stanciu, laura_stanciu_30@yahoo.com

Received 26 January 2012; Revised 4 March 2012; Accepted 8 March 2012

Academic Editor: Ondřej Došlý

Copyright © 2012 Laura Stanciu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce new integral operators of analytic functions f and g defined in the open unit disk \mathbb{U} . For these operators, we discuss some univalence conditions.

1. Introduction and Preliminaries

Let \mathcal{A} denote the class of all functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1.1)$$

which are analytic in the open unit disk

$$\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\} \quad (1.2)$$

and satisfy the following usual normalization condition:

$$f(0) = f'(0) - 1 = 0. \quad (1.3)$$

Also, let \mathcal{S} denote the subclass of \mathcal{A} consisting of functions f , which are univalent in \mathbb{U} (see, for details [1]; see also [2, 3]).

In [4, 5], Pescar gave the following univalence conditions for the functions $f \in \mathcal{A}$.

Theorem 1.1 (see [4]). Let α be a complex number, $\operatorname{Re} \alpha > 0$, and c a complex number, $|c| \leq 1$, $c \neq -1$, and $f(z) = z + \dots$ a regular function in \mathbb{U} . If

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zf''(z)}{\alpha f'(z)} \right| \leq 1, \quad (1.4)$$

for all $z \in \mathbb{U}$, then the function

$$F_\alpha(z) = \left(\alpha \int_0^z t^{\alpha-1} f'(t) dt \right)^{1/\alpha} = z + \dots \quad (1.5)$$

is regular and univalent in \mathbb{U} .

Theorem 1.2 (see [5]). Let α be a complex number, $\operatorname{Re} \alpha > 0$, and c a complex number, $|c| \leq 1$, $c \neq -1$, and $f \in \mathcal{A}$. If

$$\frac{1 - |z|^{2\operatorname{Re} \alpha}}{\operatorname{Re} \alpha} \left| \frac{zf''(z)}{f'(z)} \right| \leq 1 - |c| \quad (1.6)$$

for all $z \in \mathbb{U}$, then for any complex number β , $\operatorname{Re} \beta \geq \operatorname{Re} \alpha$, the function

$$F_\beta(z) = \left(\beta \int_0^z t^{\beta-1} f'(t) dt \right)^{1/\beta} \quad (1.7)$$

is in the class \mathcal{S} .

On the other hand, for the functions $f \in \mathcal{A}$, Ozaki and Nunokawa [6] proved another univalence condition asserted by Theorem 1.3.

Theorem 1.3 (see [6]). Let $f \in \mathcal{A}$ satisfy the condition

$$\left| \frac{z^2 f'(z)}{[f(z)]^2} - 1 \right| < 1 \quad (z \in \mathbb{U}). \quad (1.8)$$

Then f is univalent in \mathbb{U} .

In the paper [7], Pescar determined some univalence conditions for the following integral operators.

Theorem 1.4 (see [7]). Let the function g satisfy (1.8), M a positive real number fixed, and c a complex number. If $\alpha \in [(2M + 1)/(2M + 2), (2M + 1)/2M]$,

$$\begin{aligned} |c| &\leq 1 - \left| \frac{\alpha - 1}{\alpha} \right| (2M + 1), \quad c \neq -1, \\ |g(z)| &\leq M \end{aligned} \quad (1.9)$$

for all $z \in \mathbb{U}$, then the function

$$G_\alpha(z) = \left(\alpha \int_0^z [g(t)]^{\alpha-1} dt \right)^{1/\alpha} \quad (1.10)$$

is in the class \mathcal{S} .

Theorem 1.5 (see [7]). Let $g \in \mathcal{A}$, α a real number, $\alpha \geq 1$, and c a complex number, $|c| \leq 1/\alpha$, $c \neq -1$. If

$$\left| \frac{g''(z)}{g'(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad (1.11)$$

then the function

$$H_\alpha(z) = \left(\alpha \int_0^z [tg'(t)]^{\alpha-1} dt \right)^{1/\alpha} \quad (1.12)$$

is in the class \mathcal{S} .

Theorem 1.6 (see [7]). Let $g \in \mathcal{A}$ satisfies (1.8), α a complex number, $M > 1$ fixed, $\operatorname{Re} \alpha > 0$, and c a complex number, $|c| < 1$. If $|g(z)| \leq M$ for all $z \in \mathbb{U}$, then for any complex number β

$$\operatorname{Re} \beta \geq \operatorname{Re} \alpha \geq \frac{2M + 1}{|\alpha|(1 - |c|)} \quad (1.13)$$

the function

$$H_\beta(z) = \left(\beta \int_0^z t^{\beta-1} \left(\frac{g(t)}{t} \right)^{1/\alpha} dt \right)^{1/\beta} \quad (1.14)$$

is in the class \mathcal{S} .

In this paper, we introduce the following integral operators as follows:

$$F_1(f, g)(z) = \left(\alpha \int_0^z (f(t)e^{g(t)})^{\alpha-1} dt \right)^{1/\alpha} \quad (f, g \in \mathcal{A}; \alpha \in \mathbb{C}), \quad (1.15)$$

$$G_1(f, g)(z) = \left(\alpha \int_0^z (tf'(t)e^{g(t)})^{\alpha-1} dt \right)^{1/\alpha} \quad (f, g \in \mathcal{A}; \alpha \in \mathbb{C}), \quad (1.16)$$

$$H_1(f, g)(z) = \left(\beta \int_0^z t^{\beta-1} \left(\frac{f(t)}{t} e^{g(t)} \right)^{1/\alpha} dt \right)^{1/\beta} \quad (f, g \in \mathcal{A}; \alpha, \beta \in \mathbb{C} - \{0\}). \quad (1.17)$$

Remark 1.7. For $e^{g(z)} = 1$ and $f(z) = g(z)$, the integral operators (1.15), (1.16), and (1.17) would reduce to the integral operators (1.10), (1.12), and (1.14).

In this paper, we generalize the integral operators given by Pescar [7], and we study the univalence conditions for the integral operators defined by (1.15), (1.16), and (1.17).

For this purpose, we need the following result.

Lemma 1.8 (General Schwarz Lemma [8]). *Let the function f be regular in the disk $\mathbb{U}_R = \{z \in \mathbb{C} : |z| < R\}$, with $|f(z)| < M$ for fixed M . If f has one zero with multiplicity order bigger than m for $z = 0$, then*

$$|f(z)| \leq \frac{M}{R^m} |z|^m \quad (z \in \mathbb{U}_R). \quad (1.18)$$

The equality can hold only if

$$f(z) = e^{i\theta} \frac{M}{R^m} z^m, \quad (1.19)$$

where θ is constant.

2. Main Results

Theorem 2.1. *Let $f, g \in \mathcal{A}$, where g satisfies the condition (1.8), M_1 and M_2 are real positive numbers, and α a complex number, $\operatorname{Re} \alpha > 0$. If*

$$\left| \frac{f'(z)}{f(z)} \right| \leq M_1 \quad (z \in \mathbb{U}), \quad |g(z)| \leq M_2 \quad (z \in \mathbb{U}), \quad (2.1)$$

$$|c| \leq 1 - \left| \frac{\alpha - 1}{\alpha} \right| (M_1 + 2M_2^2 + 1), \quad c \in \mathbb{C}, \quad c \neq -1, \quad (2.2)$$

then the integral operator $F_1(f, g)(z)$ defined by (1.15) is in the class \mathcal{S} .

Proof. From (1.15), we have

$$F_1(f, g)(z) = \left(\alpha \int_0^z t^{\alpha-1} \left(\frac{f(t)}{t} e^{g(t)} \right)^{\alpha-1} dt \right)^{1/\alpha}. \quad (2.3)$$

Let us consider the function

$$h(z) = \int_0^z \left(\frac{f(t)}{t} e^{g(t)} \right)^{\alpha-1} dt. \quad (2.4)$$

The function h is regular in \mathbb{U} . From (2.4), we get

$$h'(z) = \left(\frac{f(z)}{z} e^{g(z)} \right)^{\alpha-1},$$

$$h''(z) = (\alpha - 1) \left(\frac{f(z)}{z} e^{g(z)} \right)^{\alpha-2} \left(\frac{zf'(z) - f(z)}{z^2} e^{g(z)} + \frac{f(z)}{z} g'(z) e^{g(z)} \right). \tag{2.5}$$

Also, a simple computation yields

$$\frac{zh''(z)}{h'(z)} = (\alpha - 1) \left[\left(\frac{zf'(z)}{f(z)} - 1 \right) + zg'(z) \right], \tag{2.6}$$

which readily shows that

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zh''(z)}{\alpha h'(z)} \right| = \left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{\alpha - 1}{\alpha} \left(\left(\frac{zf'(z)}{f(z)} - 1 \right) + zg'(z) \right) \right|$$

$$\leq |c| + \left| \frac{\alpha - 1}{\alpha} \right| \left(\left(\left| \frac{zf'(z)}{f(z)} \right| + 1 \right) + \left| \frac{z^2 g'(z)}{[g(z)]^2} \right| \left| \frac{[g(z)]^2}{z} \right| \right) \quad (z \in \mathbb{U}). \tag{2.7}$$

From the hypothesis of Theorem 2.1, we have

$$\left| \frac{f'(z)}{f(z)} \right| \leq M_1 \quad (z \in \mathbb{U}), \quad |g(z)| \leq M_2 \quad (z \in \mathbb{U}), \tag{2.8}$$

then by *General Schwarz Lemma* for the function g , we obtain

$$|g(z)| \leq M_2 |z| \quad (z \in \mathbb{U}). \tag{2.9}$$

Using the inequality (2.7), we have

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zh''(z)}{\alpha h'(z)} \right| \leq |c| + \left| \frac{\alpha - 1}{\alpha} \right| \left(M_1 + 1 + \left(\left| \frac{z^2 g'(z)}{[g(z)]^2} - 1 \right| + 1 \right) M_2^2 \right). \tag{2.10}$$

From (2.10) and since g satisfies the condition (1.8), we have

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zh''(z)}{\alpha h'(z)} \right| \leq |c| + \left| \frac{\alpha - 1}{\alpha} \right| (M_1 + 2M_2^2 + 1), \tag{2.11}$$

from which, by (2.2), we get

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zh''(z)}{\alpha h'(z)} \right| \leq 1 \quad (z \in \mathbb{U}). \tag{2.12}$$

Applying Theorem 1.1, we conclude that the integral operator $F_1(f, g)(z)$ defined by (1.15) is in the class \mathcal{S} . \square

Setting $M_1 = 1$ and $M_2 = 1$ in Theorem 2.1, we immediately arrive at the following application of Theorem 2.1.

Corollary 2.2. *Let $f, g \in \mathcal{A}$, where g satisfies the condition (1.8) and α a complex number, $\operatorname{Re} \alpha > 0$. If*

$$\begin{aligned} \left| \frac{f'(z)}{f(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad |g(z)| \leq 1 \quad (z \in \mathbb{U}), \\ |c| \leq 1 - \left| \frac{4(\alpha - 1)}{\alpha} \right|, \quad c \in \mathbb{C}, \quad c \neq -1, \end{aligned} \quad (2.13)$$

then the integral operator $F_1(f, g)(z)$ defined by (1.15) is in the class \mathcal{S} .

Theorem 2.3. *Let $f, g \in \mathcal{A}$, where g satisfies the inequality $|g(z)| \leq M$, $M \geq 1$. Also, let α be a real number, $\alpha \geq 1$, and c a complex number with*

$$|c| \leq 1 - \left| \frac{\alpha - 1}{\alpha} \right| (M + 1), \quad c \neq -1. \quad (2.14)$$

If

$$\left| \frac{f''(z)}{f'(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad \left| \frac{g'(z)}{g(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad (2.15)$$

then the integral operator $G_1(f, g)(z)$ defined by (1.16) is in the class \mathcal{S} .

Proof. We observe that

$$G_1(f, g)(z) = \left(\alpha \int_0^z t^{\alpha-1} (f'(t)e^{g(t)})^{\alpha-1} dt \right)^{1/\alpha}. \quad (2.16)$$

Let us consider the function

$$h(z) = \int_0^z (f'(t)e^{g(t)})^{\alpha-1} dt. \quad (2.17)$$

The function h is regular in \mathbb{U} . From (2.17), we have

$$\begin{aligned} h'(z) &= (f'(z)e^{g(z)})^{\alpha-1}, \\ \frac{zh''(z)}{h'(z)} &= (\alpha - 1) \left(\frac{zf''(z)}{f'(z)} + zg'(z) \right), \end{aligned} \quad (2.18)$$

which readily shows that

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zh''(z)}{\alpha h'(z)} \right| \leq |c| + \left| \frac{\alpha - 1}{\alpha} \right| \left(\left| \frac{zf''(z)}{f'(z)} \right| + \left| \frac{zg'(z)}{g(z)} \right| |g(z)| \right) \quad (z \in \mathbb{U}). \quad (2.19)$$

From (2.19) and the conditions of Theorem 2.3, we get

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zh''(z)}{\alpha h'(z)} \right| \leq |c| + \left| \frac{\alpha - 1}{\alpha} \right| (1 + M) \leq 1 \quad (z \in \mathbb{U}). \quad (2.20)$$

Applying Theorem 1.1, we conclude that the integral operator $G_1(f, g)(z)$ defined by (1.16) is in the class \mathcal{S} . \square

Setting $M = 1$ in Theorem 2.3, we obtain the following consequence of Theorem 2.3.

Corollary 2.4. *Let $f, g \in \mathcal{A}$, where g satisfies the condition $|g'(z)/g(z)| \leq 1$, α a real number, $\alpha \geq 1$, and c a complex number with $|c| \leq 1 - |2(\alpha - 1)/\alpha|$, $c \neq -1$. If*

$$\left| \frac{f''(z)}{f'(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad |g(z)| \leq 1 \quad (z \in \mathbb{U}), \quad (2.21)$$

then the integral operator $G_1(f, g)(z)$ defined by (1.16) is in the class \mathcal{S} .

Theorem 2.5. *Let $f, g \in \mathcal{A}$, where g satisfies the condition (1.8), α a complex number, $\operatorname{Re} \alpha > 0$, M_1 and M_2 are real positive numbers, and c a complex number, $|c| < 1$. If*

$$\left| \frac{f'(z)}{f(z)} \right| \leq M_1 \quad (z \in \mathbb{U}), \quad |g(z)| \leq M_2 \quad (z \in \mathbb{U}), \quad (2.22)$$

then for any complex number β ,

$$\operatorname{Re} \beta \geq \operatorname{Re} \alpha \geq \frac{M_1 + 2M_2^2 + 1}{|\alpha|(1 - |c|)}, \quad (2.23)$$

the integral operator $H_1(f, g)(z)$ defined by (1.17) is in the class \mathcal{S} .

Proof. Let us consider the function

$$h(z) = \int_0^z \left(\frac{f(t)}{t} e^{g(t)} \right)^{1/\alpha} dt. \quad (2.24)$$

The function h is regular in \mathbb{U} . From (2.24), we have

$$\begin{aligned} h'(z) &= \left(\frac{f(z)}{z} e^{g(z)} \right)^{1/\alpha}, \\ \frac{zh''(z)}{h'(z)} &= \frac{1}{\alpha} \left(\left(\frac{zf'(z)}{f(z)} - 1 \right) + zg'(z) \right), \end{aligned} \quad (2.25)$$

which readily shows that

$$\begin{aligned} \frac{1 - |z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \frac{1 - |z|^{2\operatorname{Re}\alpha}}{|\alpha| \operatorname{Re}\alpha} \left(\left| \frac{zf'(z)}{f(z)} \right| + 1 + \left| \frac{z^2 g'(z)}{[g(z)]^2} \right| \left| \frac{[g(z)]^2}{z} \right| \right) \\ &\leq \frac{1 - |z|^{2\operatorname{Re}\alpha}}{|\alpha| \operatorname{Re}\alpha} \left(\left| \frac{zf'(z)}{f(z)} \right| + 1 + \left(\left| \frac{z^2 g'(z)}{[g(z)]^2} - 1 \right| + 1 \right) \left| \frac{[g(z)]^2}{z} \right| \right). \end{aligned} \quad (2.26)$$

By the *General Schwarz Lemma* for the function g , we obtain

$$|g(z)| \leq M_2 |z| \quad (z \in \mathbb{U}), \quad (2.27)$$

and using the inequality (2.26), we have

$$\frac{1 - |z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \left| \frac{zh''(z)}{h'(z)} \right| \leq \frac{1 - |z|^{2\operatorname{Re}\alpha}}{|\alpha| \operatorname{Re}\alpha} \left(M_1 + 1 + \left(\left| \frac{z^2 g'(z)}{[g(z)]^2} - 1 \right| + 1 \right) M_2^2 \right). \quad (2.28)$$

From (2.28) and since g satisfies the condition (1.8), we get

$$\begin{aligned} \frac{1 - |z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \frac{1 - |z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \frac{(M_1 + 2M_2^2 + 1)}{|\alpha|} \\ &\leq \frac{M_1 + 2M_2^2 + 1}{|\alpha| \operatorname{Re}\alpha}. \end{aligned} \quad (2.29)$$

From (2.23), we have

$$\frac{M_1 + 2M_2^2 + 1}{|\alpha| \operatorname{Re}\alpha} \leq 1 - |c|, \quad (2.30)$$

and using (2.29), we obtain

$$\frac{1 - |z|^{2\operatorname{Re}\alpha}}{\operatorname{Re}\alpha} \left| \frac{zh''(z)}{h'(z)} \right| \leq 1 - |c| \quad (z \in \mathbb{U}). \quad (2.31)$$

Applying Theorem 1.2, we conclude that the integral operator $H_1(f, g)(z)$ defined by (1.17) is in the class \mathcal{S} . \square

Setting $M_1 = 1$ and $M_2 = 1$ in Theorem 2.5, we obtain the following corollary.

Corollary 2.6. *Let $f, g \in \mathcal{A}$, where g satisfies the condition (1.8), α a complex number, $\operatorname{Re} \alpha > 0$, and c a complex number, $|c| < 1$. If*

$$\left| \frac{f'(z)}{f(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad |g(z)| \leq 1 \quad (z \in \mathbb{U}), \quad (2.32)$$

then for any complex number β ,

$$\operatorname{Re} \beta \geq \operatorname{Re} \alpha \geq \frac{4}{|\alpha|(1-|c|)}, \quad (2.33)$$

the integral operator $H_1(f, g)(z)$ defined by (1.17) is in the class \mathcal{S} .

Acknowledgments

This paper was partially supported by the strategic project POSDRU 107/1.5/S/77265, inside POSDRU Romania 2007–2013 cofinanced by the European Social Fund-Investing in People.

References

- [1] D. Blezu, "On univalence criteria," *General Mathematics*, vol. 14, pp. 77–84, 2006.
- [2] D. Blezu and R. N. Pascu, "Univalence criteria for integral operators," *Glasnik Matematički*, vol. 36(56), no. 2, pp. 241–245, 2001.
- [3] V. Pescar, "Univalence of certain integral operators," *Acta Universitatis Apulensis*, no. 12, pp. 43–48, 2006.
- [4] V. Pescar, "A new generalization of Ahlfors's and Becker's criterion of univalence," *Bulletin of Malaysian Mathematical Society*, vol. 19, no. 2, pp. 53–54, 1996.
- [5] V. Pescar, "Univalence criteria of certain integral operators," *Acta Ciencia Indica, Mathematics*, vol. 29, no. 1, pp. 135–138, 2003.
- [6] S. Ozaki and M. Nunokawa, "The Schwarzian derivative and univalent functions," *Proceedings of the American Mathematical Society*, vol. 33, pp. 392–394, 1972.
- [7] V. Pescar, "On the univalence of some integral operators," *General Mathematics*, vol. 14, no. 2, pp. 77–84, 2006.
- [8] Z. Nehari, *Conformal Mapping*, Dover, New York, NY, USA, 1975.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

