

Research Article

On a Conjecture for a Higher-Order Rational Difference Equation

Maoxin Liao,^{1,2} Xianhua Tang,¹ and Changjin Xu^{1,3}

¹ School of Mathematical Sciences and Computing Technology, Central South University, Changsha, Hunan 410083, China

² School of Mathematics and Physics, University of South China, Hengyang, Hunan 421001, China

³ College of Science, Hunan Institute of Engineering, Xiangtan, Hunan 411104, China

Correspondence should be addressed to Maoxin Liao, maoxinliao@163.com

Received 30 December 2008; Revised 11 March 2009; Accepted 14 March 2009

Recommended by Jianshe Yu

This paper studies the global asymptotic stability for positive solutions to the higher order rational difference equation $x_n = (\prod_{j=1}^m(x_{n-k_j} + 1) + \prod_{j=1}^m(x_{n-k_j} - 1)) / (\prod_{j=1}^m(x_{n-k_j} + 1) - \prod_{j=1}^m(x_{n-k_j} - 1))$, $n = 0, 1, 2, \dots$, where m is odd and $x_{-k_m}, x_{-k_m+1}, \dots, x_{-1} \in (0, \infty)$. Our main result generalizes several others in the recent literature and confirms a conjecture by Berenhaut et al., 2007.

Copyright © 2009 Maoxin Liao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 2007, Berenhaut et al. [1] proved that every solution of the following rational difference equation

$$x_n = \frac{x_{n-k} + x_{n-m}}{1 + x_{n-k}x_{n-m}}, \quad n = 0, 1, 2, \dots \quad (1.1)$$

converges to its unique equilibrium 1, where $x_{-m}, x_{-m+1}, \dots, x_{-1} \in (0, \infty)$ and $1 \leq k < m$. Based on this fact, they put forward the following two conjectures.

Conjecture 1.1. Suppose that $1 \leq k < l < m$ and that $\{x_n\}$ satisfies

$$x_n = \frac{x_{n-k} + x_{n-l} + x_{n-m} + x_{n-k}x_{n-l}x_{n-m}}{1 + x_{n-k}x_{n-l} + x_{n-l}x_{n-m} + x_{n-m}x_{n-k}}, \quad n = 0, 1, 2, \dots \quad (1.2)$$

with $x_{-m}, x_{-m+1}, \dots, x_{-1} \in (0, \infty)$. Then, the sequence $\{x_n\}$ converges to the unique equilibrium 1.

Conjecture 1.2. Suppose that m is odd and $1 \leq k_1 < k_2 < \dots < k_m$, and define $S = \{1, 2, \dots, m\}$. If $\{x_n\}$ satisfies

$$x_n = \frac{f_1(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_m})}{f_2(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_m})}, \quad n = 0, 1, 2, \dots \quad (1.3)$$

with $x_{-k_m}, x_{-k_m+1}, \dots, x_{-1} \in (0, \infty)$, where

$$\begin{aligned} f_1(y_1, y_2, \dots, y_m) &= \sum_{j \in \{1, 3, \dots, m\}} \sum_{\{t_1, t_2, \dots, t_j\} \subset S; t_1 < t_2 < \dots < t_j} y_{t_1} y_{t_2} \cdots y_{t_j}, \\ f_2(y_1, y_2, \dots, y_m) &= 1 + \sum_{j \in \{2, 4, \dots, m-1\}} \sum_{\{t_1, t_2, \dots, t_j\} \subset S; t_1 < t_2 < \dots < t_j} y_{t_1} y_{t_2} \cdots y_{t_j}. \end{aligned} \quad (1.4)$$

Then the sequence $\{x_n\}$ converges to the unique equilibrium 1.

Motivated by [2], Berenhaut et al. started with the investigation of the following difference equation $y_n = A + (y_{n-k}/y_{n-m})^p$ for $p > 0$ (see, [3, 4]). Among others, in [3] they used a transformation method, which has turned out to be very useful in studying (1.1) and (1.2) as well as in confirming Conjecture 1.1; see [5].

Some particular cases of (1.2) had been studied previously by Li in [6, 7], by using semicycle analysis similar to that in [8]. The problem concerning periodicity of semicycles of difference equations was solved in very general settings by Berg and Stević in [9], partially motivated also by [10].

In the meantime, it turned out that the method used in [11] by Çinar et al. can be used in confirming Conjecture 1.2 (see also [12]). More precisely [11, 12] use Corollary 3 from [13] in solving similar problems. For example, Çinar et al. has shown, in an elegant way, that the main result in [14] is a consequence of Corollary 3 in [13]. With some calculations it can be also shown that Conjecture 1.2 can be confirmed in this way (see [15]).

Some other related results can be found in [16–24].

In this paper, we will prove that Conjecture 1.2 is correct by using a new method. Obviously, our results generalize the corresponding works in [1, 5–7] and other literature.

2. Preliminaries and Notations

Observe that

$$\begin{aligned} f_1(y_1, y_2, \dots, y_m) &= \frac{1}{2} \left[\prod_{j=1}^m (y_j + 1) + \prod_{j=1}^m (y_j - 1) \right], \\ f_2(y_1, y_2, \dots, y_m) &= \frac{1}{2} \left[\prod_{j=1}^m (y_j + 1) - \prod_{j=1}^m (y_j - 1) \right]. \end{aligned} \quad (2.1)$$

Define function G as follows:

$$G(y_1, y_2, \dots, y_m) = \frac{\prod_{j=1}^m (y_j + 1) + \prod_{j=1}^m (y_j - 1)}{\prod_{j=1}^m (y_j + 1) - \prod_{j=1}^m (y_j - 1)}, \quad y_1, y_2, \dots, y_m > 0. \quad (2.2)$$

Then we can rewrite (1.3) as

$$x_n = \frac{\prod_{j=1}^m (x_{n-k_j} + 1) + \prod_{j=1}^m (x_{n-k_j} - 1)}{\prod_{j=1}^m (x_{n-k_j} + 1) - \prod_{j=1}^m (x_{n-k_j} - 1)}, \quad n = 0, 1, 2, \dots, \quad (2.3)$$

or

$$x_n = G(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_m}), \quad n = 0, 1, 2, \dots, \quad (2.4)$$

where m is an odd integer and $x_{-k_m}, x_{-k_m+1}, \dots, x_{-1} \in (0, \infty)$.

The following lemma can be obtained by simple calculations.

Lemma 2.1. *Let G be defined by (2.2). Then*

$$\frac{\partial G}{\partial y_i} = \frac{4 \prod_{j=1, j \neq i}^m (y_j^2 - 1)}{\left[\prod_{j=1}^m (y_j + 1) - \prod_{j=1}^m (y_j - 1) \right]^2} \begin{cases} > 0, & \prod_{j=1, j \neq i}^m (y_j - 1) > 0, \\ < 0, & \prod_{j=1, j \neq i}^m (y_j - 1) < 0, \end{cases} \quad (2.5)$$

$i = 1, 2, \dots, m$.

Lemma 2.2. *Assume that $0 < \alpha < 1 < \beta < +\infty$. If $\alpha \leq y_1, y_2, \dots, y_m \leq \beta$, then*

$$\min\{A_1, A_3, \dots, A_m\} \leq G(y_1, y_2, \dots, y_m) \leq \max\{B_1, B_3, \dots, B_m\}, \quad (2.6)$$

where

$$A_i = \frac{(\alpha + 1)^i (\beta + 1)^{m-i} + (\alpha - 1)^i (\beta - 1)^{m-i}}{(\alpha + 1)^i (\beta + 1)^{m-i} - (\alpha - 1)^i (\beta - 1)^{m-i}}, \quad (2.7)$$

$$B_i = \frac{(\alpha + 1)^{m-i} (\beta + 1)^i + (\alpha - 1)^{m-i} (\beta - 1)^i}{(\alpha + 1)^{m-i} (\beta + 1)^i - (\alpha - 1)^{m-i} (\beta - 1)^i}$$

$i = 1, 3, \dots, m$.

Proof. Since $G(y_1, y_2, \dots, y_m)$ is symmetric in y_1, y_2, \dots, y_m , we can assume, without loss of generality, that $\alpha \leq y_1 \leq y_2 \leq \dots \leq y_m \leq \beta$. Then there are $m + 1$ possible cases:

- (1) $\alpha \leq 1 \leq y_1 \leq y_2 \leq \dots \leq y_m \leq \beta$;
- (2) $\alpha \leq y_1 \leq 1 \leq y_2 \leq \dots \leq y_m \leq \beta$;
- (3) $\alpha \leq y_1 \leq y_2 \leq 1 \leq \dots \leq y_m \leq \beta$;
- (4) $\alpha \leq y_1 \leq y_2 \leq y_3 \leq 1 \leq \dots \leq y_m \leq \beta$;
- \vdots
- (m+1) $\alpha \leq y_1 \leq y_2 \leq \dots \leq y_m \leq 1 \leq \beta$.

And, for the above cases (1)–(m+1), by the monotonicity of $G(y_1, y_2, \dots, y_m)$, in turn, we may get

- (1) $1 \leq G(y_1, y_2, \dots, y_m) \leq B_m$;
- (2) $A_1 \leq G(y_1, y_2, \dots, y_m) \leq 1$;
- (3) $1 \leq G(y_1, y_2, \dots, y_m) \leq B_{m-2}$;
- (4) $A_3 \leq G(y_1, y_2, \dots, y_m) \leq 1$;
- \vdots
- (m+1) $A_m \leq G(y_1, y_2, \dots, y_m) \leq 1$.

From the above inequalities, it follows that (2.6) holds. The proof is complete. \square

Lemma 2.3. Assume that $0 < \alpha < 1 < \beta < +\infty$. Then

$$A_i = \frac{(\alpha + 1)^i (\beta + 1)^{m-i} + (\alpha - 1)^i (\beta - 1)^{m-i}}{(\alpha + 1)^i (\beta + 1)^{m-i} - (\alpha - 1)^i (\beta - 1)^{m-i}} \geq \alpha, \quad (2.8)$$

$$B_i = \frac{(\alpha + 1)^{m-i} (\beta + 1)^i + (\alpha - 1)^{m-i} (\beta - 1)^i}{(\alpha + 1)^{m-i} (\beta + 1)^i - (\alpha - 1)^{m-i} (\beta - 1)^i} \leq \beta, \quad (2.9)$$

$i = 1, 3, \dots, m$.

Proof. For $i = 1, 3, \dots, m$, it is easy to see that

$$(\alpha - 1)^{i-1} (\beta - 1)^{m-i} \leq (\alpha + 1)^{i-1} (\beta + 1)^{m-i}, \quad (2.10)$$

which yields

$$(\alpha + 1)(\alpha - 1)^i (\beta - 1)^{m-i} \geq (\alpha - 1)(\alpha + 1)^i (\beta + 1)^{m-i}, \quad (2.11)$$

and so

$$\alpha \left[(\alpha + 1)^i (\beta + 1)^{m-i} - (\alpha - 1)^i (\beta - 1)^{m-i} \right] \leq (\alpha + 1)^i (\beta + 1)^{m-i} + (\alpha - 1)^i (\beta - 1)^{m-i}. \quad (2.12)$$

It follows that (2.8) holds. Similarly, for $i = 1, 3, \dots, m$, it is easy to see that

$$(\alpha - 1)^{m-i}(\beta - 1)^{i-1} \leq (\alpha + 1)^{m-i}(\beta + 1)^{i-1}, \tag{2.13}$$

which yields

$$(\beta + 1)(\alpha - 1)^{m-i}(\beta - 1)^i \leq (\beta - 1)(\alpha + 1)^{m-i}(\beta + 1)^i. \tag{2.14}$$

It follows that (2.9) holds. The proof is complete. □

Lemma 2.4. *Let*

$$\begin{aligned} \alpha_{j+1} &= \min\{A_{1j}, A_{3j}, \dots, A_{mj}\}, \\ \beta_{j+1} &= \max\{B_{1j}, B_{3j}, \dots, B_{mj}\}, \end{aligned} \tag{2.15}$$

where

$$\begin{aligned} A_{ij} &= \frac{(\alpha_j + 1)^i(\beta_j + 1)^{m-i} + (\alpha_j - 1)^i(\beta_j - 1)^{m-i}}{(\alpha_j + 1)^i(\beta_j + 1)^{m-i} - (\alpha_j - 1)^i(\beta_j - 1)^{m-i}}, \\ B_{ij} &= \frac{(\alpha_j + 1)^{m-i}(\beta_j + 1)^i + (\alpha_j - 1)^{m-i}(\beta_j - 1)^i}{(\alpha_j + 1)^{m-i}(\beta_j + 1)^i - (\alpha_j - 1)^{m-i}(\beta_j - 1)^i} \end{aligned} \tag{2.16}$$

$i = 1, 3, \dots, m; j = 0, 1, 2, \dots$. Assume that $0 < \alpha_0 < 1 < \beta_0 < +\infty$. Then

$$\lim_{j \rightarrow \infty} \alpha_j = \lim_{j \rightarrow \infty} \beta_j = 1. \tag{2.17}$$

Proof. By induction, we easily show that

$$0 < \alpha_j < 1 < \beta_j < +\infty, \quad j = 0, 1, 2, \dots \tag{2.18}$$

It follows from Lemma 2.3 that

$$\begin{aligned} A_{ij} &= \frac{(\alpha_j + 1)^i(\beta_j + 1)^{m-i} + (\alpha_j - 1)^i(\beta_j - 1)^{m-i}}{(\alpha_j + 1)^i(\beta_j + 1)^{m-i} - (\alpha_j - 1)^i(\beta_j - 1)^{m-i}} \geq \alpha_j, \\ B_{ij} &= \frac{(\alpha_j + 1)^{m-i}(\beta_j + 1)^i + (\alpha_j - 1)^{m-i}(\beta_j - 1)^i}{(\alpha_j + 1)^{m-i}(\beta_j + 1)^i - (\alpha_j - 1)^{m-i}(\beta_j - 1)^i} \leq \beta_j, \end{aligned} \tag{2.19}$$

$i = 1, 3, \dots, m; j = 0, 1, 2, \dots$. Hence, by (2.15) and (2.18), we have

$$\alpha_j \leq \alpha_{j+1} < 1 < \beta_{j+1} \leq \beta_j, \quad j = 0, 1, 2, \dots \tag{2.20}$$

Equation (2.20) implies that the limits $\lim_{j \rightarrow \infty} \alpha_j$ and $\lim_{j \rightarrow \infty} \beta_j$ exist, and

$$\alpha^* = \lim_{j \rightarrow \infty} \alpha_j \in [\alpha_0, 1], \quad \beta^* = \lim_{j \rightarrow \infty} \beta_j \in [1, \beta_0]. \quad (2.21)$$

It follows from (2.16) that

$$\begin{aligned} A_i^* &:= \lim_{j \rightarrow \infty} A_{ij} = \frac{(\alpha^* + 1)^i (\beta^* + 1)^{m-i} + (\alpha^* - 1)^i (\beta^* - 1)^{m-i}}{(\alpha^* + 1)^i (\beta^* + 1)^{m-i} - (\alpha^* - 1)^i (\beta^* - 1)^{m-i}}, \\ B_i^* &:= \lim_{j \rightarrow \infty} B_{ij} = \frac{(\alpha^* + 1)^{m-i} (\beta^* + 1)^i + (\alpha^* - 1)^{m-i} (\beta^* - 1)^i}{(\alpha^* + 1)^{m-i} (\beta^* + 1)^i - (\alpha^* - 1)^{m-i} (\beta^* - 1)^i} \end{aligned} \quad (2.22)$$

$i = 1, 3, \dots, m$. Let $j \rightarrow \infty$ in (2.15), we have

$$\begin{aligned} \alpha^* &= \min\{A_1^*, A_3^*, \dots, A_m^*\}, \\ \beta^* &= \max\{B_1^*, B_3^*, \dots, B_m^*\}. \end{aligned} \quad (2.23)$$

It follows that there exist $i, j \in \{1, 3, \dots, m\}$ such that

$$\begin{aligned} \alpha^* &= \frac{(\alpha^* + 1)^i (\beta^* + 1)^{m-i} + (\alpha^* - 1)^i (\beta^* - 1)^{m-i}}{(\alpha^* + 1)^i (\beta^* + 1)^{m-i} - (\alpha^* - 1)^i (\beta^* - 1)^{m-i}}, \\ \beta^* &= \frac{(\alpha^* + 1)^{m-j} (\beta^* + 1)^j + (\alpha^* - 1)^{m-j} (\beta^* - 1)^j}{(\alpha^* + 1)^{m-j} (\beta^* + 1)^j - (\alpha^* - 1)^{m-j} (\beta^* - 1)^j}. \end{aligned} \quad (2.24)$$

From (2.24), we have

$$\begin{aligned} (\alpha^* - 1) \left[(\alpha^* + 1)^{i-1} (\beta^* + 1)^{m-i} - (\alpha^* - 1)^{i-1} (\beta^* - 1)^{m-i} \right] &= 0, \\ (\beta^* - 1) \left[(\alpha^* + 1)^{m-j} (\beta^* + 1)^{j-1} - (\alpha^* - 1)^{m-j} (\beta^* - 1)^{j-1} \right] &= 0. \end{aligned} \quad (2.25)$$

Since

$$\begin{aligned} (\alpha^* + 1)^{i-1} (\beta^* + 1)^{m-i} - (\alpha^* - 1)^{i-1} (\beta^* - 1)^{m-i} &> 0, \\ (\alpha^* + 1)^{m-j} (\beta^* + 1)^{j-1} - (\alpha^* - 1)^{m-j} (\beta^* - 1)^{j-1} &> 0, \end{aligned} \quad (2.26)$$

it follows from (2.25) and (2.18) that $\alpha^* = \beta^* = 1$. The proof is complete. \square

3. Proof of Conjecture 1.2

Theorem 3.1. *Suppose that $0 < \alpha < 1 < \beta < +\infty$ and that*

$$x_{-k_m}, x_{-k_m+1}, \dots, x_{-1} \in [\alpha, \beta]. \quad (3.1)$$

Then the solution $\{x_n\}$ of (1.3) satisfies

$$x_n \in [\alpha, \beta], \quad \text{for } n = 0, 1, 2, \dots \quad (3.2)$$

Theorem 3.1 is a direct corollary of Lemmas 2.2 and 2.3.

Proof of Conjecture 1.2. Let $\{x_n\}$ be a solution of (1.3) with $x_{-k_m}, x_{-k_m+1}, \dots, x_{-1} \in (0, \infty)$. We need to prove that

$$\lim_{n \rightarrow \infty} x_n = 1. \quad (3.3)$$

Choose $\alpha_0 \in (0, 1)$ and $\beta_0 \in (1, +\infty)$ such that

$$x_{-k_m}, x_{-k_m+1}, \dots, x_{-1} \in [\alpha_0, \beta_0]. \quad (3.4)$$

In view of Theorem 3.1, we have

$$x_n \in [\alpha_0, \beta_0], \quad n = -k_m, -k_m + 1, -k_m + 2, \dots \quad (3.5)$$

Let $\alpha_j, \beta_j, A_{ij}$, and B_{ij} be defined as in Lemma 2.4. Then by (3.5) and Lemma 2.2, we have

$$\begin{aligned} \min\{A_{10}, A_{30}, \dots, A_{m0}\} &\leq G(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_m}) \\ &\leq \max\{B_{10}, B_{30}, \dots, B_{m0}\}, \quad n = 0, 1, 2, \dots \end{aligned} \quad (3.6)$$

That is

$$x_n \in [\alpha_1, \beta_1], \quad n = 0, 1, 2, \dots \quad (3.7)$$

By (3.7) and Lemma 2.2, we obtain

$$\begin{aligned} \min\{A_{11}, A_{31}, \dots, A_{m1}\} &\leq G(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_m}) \\ &\leq \max\{B_{11}, B_{31}, \dots, B_{m1}\}, \quad n = k_m, k_m + 1, k_m + 2, \dots \end{aligned} \quad (3.8)$$

That is

$$x_n \in [\alpha_2, \beta_2], \quad n = k_m, k_m + 1, k_m + 2, \dots \quad (3.9)$$

Repeating the above procedure, in general, we can obtain

$$x_n \in [\alpha_{j+1}, \beta_{j+1}], \quad n = jk_m, jk_m + 1, jk_m + 2, \dots, \quad j = 0, 1, 2, \dots \quad (3.10)$$

By Lemma 2.4, we have

$$\lim_{n \rightarrow \infty} x_n = \lim_{j \rightarrow \infty} \alpha_{j+1} = \lim_{j \rightarrow \infty} \beta_{j+1} = 1, \quad (3.11)$$

which implies that (3.3) holds. The proof of Conjecture 1.2 is complete. \square

Acknowledgments

The authors are grateful to the referees for their careful reading of the manuscript and many valuable comments and suggestions that greatly improved the presentation of this work. This work is supported partly by NNSF of China (Grant: 10771215, 10771094), Project of Hunan Provincial Youth Key Teacher and Project of Hunan Provincial Education Department (Grant: 07C639).

References

- [1] K. S. Berenhaut, J. D. Foley, and S. Stević, "The global attractivity of the rational difference equation $y_n = (y_{n-k} + y_{n-m}) / (1 + y_{n-k}y_{n-m})$," *Applied Mathematics Letters*, vol. 20, no. 1, pp. 54–58, 2007.
- [2] S. Stević, "On the recursive sequence $x_{n+1} = \alpha + x_{n-1}^p / x_n^p$," *Journal of Applied Mathematics & Computing*, vol. 18, no. 1-2, pp. 229–234, 2005.
- [3] K. S. Berenhaut, J. D. Foley, and S. Stević, "The global attractivity of the rational difference equation $y_n = 1 + (y_{n-k}) / (y_{n-m})$," *Proceedings of the American Mathematical Society*, vol. 135, no. 4, pp. 1133–1140, 2007.
- [4] K. S. Berenhaut, J. D. Foley, and S. Stević, "The global attractivity of the rational difference equation $y_n = A + (y_{n-k} / y_{n-m})^p$," *Proceedings of the American Mathematical Society*, vol. 136, no. 1, pp. 103–110, 2008.
- [5] K. S. Berenhaut and S. Stević, "The global attractivity of a higher order rational difference equation," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 940–944, 2007.
- [6] X. Li, "Qualitative properties for a fourth-order rational difference equation," *Journal of Mathematical Analysis and Applications*, vol. 311, no. 1, pp. 103–111, 2005.
- [7] X. Li, "Global behavior for a fourth-order rational difference equation," *Journal of Mathematical Analysis and Applications*, vol. 312, no. 2, pp. 555–563, 2005.
- [8] A. M. Amleh, N. Kruse, and G. Ladas, "On a class of difference equations with strong negative feedback," *Journal of Difference Equations and Applications*, vol. 5, no. 6, pp. 497–515, 1999.
- [9] L. Berg and S. Stević, "Linear difference equations mod 2 with applications to nonlinear difference equations," *Journal of Difference Equations and Applications*, vol. 14, no. 7, pp. 693–704, 2008.
- [10] L. Berg and S. Stević, "Periodicity of some classes of holomorphic difference equations," *Journal of Difference Equations and Applications*, vol. 12, no. 8, pp. 827–835, 2006.
- [11] C. Çinar, S. Stević, and I. Yalçinkaya, "A note on global asymptotic stability of a family of rational equations," *Rostocker Mathematisches Kolloquium*, no. 59, pp. 41–49, 2005.
- [12] S. Stević, "Global stability and asymptotics of some classes of rational difference equations," *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 60–68, 2006.
- [13] N. Kruse and T. Nesemann, "Global asymptotic stability in some discrete dynamical systems," *Journal of Mathematical Analysis and Applications*, vol. 235, no. 1, pp. 151–158, 1999.
- [14] X. Li and D. Zhu, "Global asymptotic stability in a rational equation," *Journal of Difference Equations and Applications*, vol. 9, no. 9, pp. 833–839, 2003.

- [15] M. Aloqeily, "Global stability of a rational symmetric difference equation," preprint, 2008.
- [16] L. Gutnik and S. Stević, "On the behaviour of the solutions of a second-order difference equation," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 27562, 14 pages, 2007.
- [17] G. Ladas, "A problem from the Putnam Exam," *Journal of Difference Equations and Applications*, vol. 4, no. 5, pp. 497–499, 1998.
- [18] "Putnam Exam," *The American Mathematical Monthly*, pp. 734–736, 1965.
- [19] S. Stević, "Asymptotics of some classes of higher-order difference equations," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 56813, 20 pages, 2007.
- [20] S. Stević, "Existence of nontrivial solutions of a rational difference equation," *Applied Mathematics Letters*, vol. 20, no. 1, pp. 28–31, 2007.
- [21] S. Stević, "Nontrivial solutions of a higher-order rational difference equation," *Matematičeskie Zametki*, vol. 84, no. 5, pp. 772–780, 2008.
- [22] T. Sun and H. Xi, "Global asymptotic stability of a higher order rational difference equation," *Journal of Mathematical Analysis and Applications*, vol. 330, no. 1, pp. 462–466, 2007.
- [23] X. Yang, F. Sun, and Y. Y. Tang, "A new part-metric-related inequality chain and an application," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 193872, 7 pages, 2008.
- [24] X. Yang, Y. Y. Tang, and J. Cao, "Global asymptotic stability of a family of difference equations," *Computers & Mathematics with Applications*, vol. 56, no. 10, pp. 2643–2649, 2008.