

## Research Article

# Periodic Problem with a Potential Landesman Lazer Condition

**Petr Tomiczek**

*Department of Mathematics, University of West Bohemia, Univerzitní 22, 306 14 Plzeň, Czech Republic*

Correspondence should be addressed to Petr Tomiczek, tomiczek@kma.zcu.cz

Received 6 January 2010; Revised 30 June 2010; Accepted 22 September 2010

Academic Editor: Pavel Drábek

Copyright © 2010 Petr Tomiczek. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We prove the existence of a solution to the periodic nonlinear second-order ordinary differential equation with damping  $u''(x) + r(x)u'(x) + g(x, u(x)) = f(x)$ ,  $u(0) = u(T)$ ,  $u'(0) = u'(T)$ . We suppose that  $\int_0^T r(x)dx = 0$ , the nonlinearity  $g$  satisfies the potential Landesman Lazer condition and prove that a critical point of a corresponding energy functional is a solution to this problem.

## 1. Introduction

Let us consider the nonlinear problem

$$\begin{aligned}u''(x) + r(x)u'(x) + g(x, u(x)) &= f(x), \quad x \in [0, T], \\u(0) &= u(T), \quad u'(0) = u'(T),\end{aligned}\tag{1.1}$$

where  $r \in L^1(0, T)$ , the nonlinearity  $g : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  is a Caratheodory function and  $f \in L^1(0, T)$ .

To state an existence result to (1.1) Amster [1] assumes that  $r$  is a nondecreasing function (see also [2]). He supposes that the nonlinearity  $g$  satisfies the growth condition  $(g(x, s) - g(x, t))/(s - t) \leq c_1$ ,  $c_1 < \lambda_1$  for  $x \in [0, T]$ ,  $s, t \in \mathbb{R}$ ,  $s \neq t$ , where  $\lambda_1$  is the first eigenvalue of the problem  $-u'' = \lambda u$ ,  $u(0) = u(T) = 0$  and there exist  $a^-, a^+$  such that  $g|_{[0, T] \times I_{a^+}} \geq \int_0^T p_1(x)f(x)dx / \|p_1\|_1 \geq g|_{[0, T] \times I_{a^-}}$ . An interval  $I_a$  is centered in  $a$  with the radius  $\delta_1|a| + \delta_2$  where  $\delta_1 = \sqrt{\lambda_1}c_1T / (\lambda_1 - c_1) < 1$ ,  $0 < \delta_2$  and  $p_1$  is a solution to the problem  $p_1' - rp_1 = k_1$ ,  $k_1 \in \mathbb{R}$  with  $p_1(0) = p_1(T) = 1$ .

In [3, 4] authors studied (1.1) with a constant friction term  $r(x) = c$  and results with repulsive singularities were obtained in [5, 6].

In this paper we present new assumptions, we suppose that the friction term  $r$  has zero mean value

$$\int_0^T r(x) dx = 0, \quad (1.2)$$

the nonlinearity  $g$  is bounded by a  $L^1$  function and satisfies the following potential Landesman-Lazer condition (see also [7, 8])

$$\int_0^T [R(x)^2 G_-(x)] dx < \int_0^T [R(x)^2 f(x)] dx < \int_0^T [R(x)^2 G_+(x)] dx, \quad (1.3)$$

where  $G(x, s) = \int_0^s g(x, t) dt$ ,  $G_+(x) = \liminf_{s \rightarrow +\infty} G(x, s)/s$ ,  $G_-(x) = \limsup_{s \rightarrow -\infty} (G(x, s)/s)$  and  $R(x) = e^{\int_0^x (1/2)r(\xi) d\xi}$ .

To obtain our result we use variational approach even if the linearization of the periodic problem (1.1) is a non-self-adjoint operator.

## 2. Preliminaries

*Notation.* We will use the classical space  $C^k(0, T)$  of functions whose  $k$ th derivative is continuous and the space  $L^p(0, T)$  of measurable real-valued functions whose  $p$ th power of the absolute value is Lebesgue integrable. We denote  $H$  the Sobolev space of absolutely continuous functions  $u : (0, T) \rightarrow \mathbb{R}$  such that  $u' \in L^2(0, T)$  and  $u(0) = u(T)$  with the norm  $\|u\| = (\int_0^T u^2(x) + u'^2(x) dx)^{1/2}$ . By a solution to (1.1) we mean a function  $u \in C^1(0, T)$  such that  $u'$  is absolutely continuous,  $u$  satisfies the boundary conditions and (1.1) is satisfied a.e. in  $(0, T)$ .

We denote  $R(x) = e^{\int_0^x (1/2)r(\xi) d\xi}$  and we study (1.1) by using variational methods. We investigate the functional  $J : H \rightarrow \mathbb{R}$ , which is defined by

$$J(u) = \frac{1}{2} \int_0^T [R^2(u')^2] dx - \int_0^T [R^2 G(x, u) - R^2 f u] dx, \quad (2.1)$$

where

$$G(x, s) = \int_0^s g(x, t) dt. \quad (2.2)$$

We say that  $u$  is a critical point of  $J$ , if

$$\langle J'(u), v \rangle = 0 \quad \forall v \in H. \quad (2.3)$$

We see that every critical point  $u \in H$  of the functional  $J$  satisfies

$$\int_0^T [R^2 u' v'] dx - \int_0^T [R^2 (g(x, u) - f) v] dx = 0 \quad (2.4)$$

for all  $v \in H$ .

Now we prove that any critical point of the functional  $J$  is a solution to (1.1) mentioned above.

**Lemma 2.1.** *Let the condition (1.2) be satisfied. Then any critical point of the functional  $J$  is a solution to (1.1).*

*Proof.* Setting  $v = 1$  in (2.4) we obtain

$$\int_0^T [R^2 (g(x, u) - f)] dx = 0. \quad (2.5)$$

We denote

$$\Phi(x) = \int_0^x [R(t)^2 (g(t, u(t)) - f(t))] dt \quad (2.6)$$

then previous equality (2.5) implies  $\Phi(0) = \Phi(T) = 0$  and by parts in (2.4) we have

$$\int_0^T [(R^2 u' + \Phi) v'] dx = 0 \quad (2.7)$$

for all  $v \in H$ . Hence there exists a constant  $c_u$  such that

$$R^2 u' + \Phi = c_u \quad (2.8)$$

on  $[0, T]$ . The condition (1.2) implies  $R(0) = R(T) = 1$  and from (2.8) we get  $u'(0) = R^2(0)u'(0) = -\Phi(0) + c_u = -\Phi(T) + c_u = u'(T)$ . Using  $(R^2)' = R^2 r$  and differentiating equality (2.8) with respect to  $x$  we obtain

$$R^2 (u'' + ru' + g(x, u) - f) = 0. \quad (2.9)$$

Thus  $u$  is a solution to (1.1).  $\square$

We say that  $J$  satisfies the *Palais-Smale condition* (PS) if every sequence  $(u_n)$  for which  $J(u_n)$  is bounded in  $H$  and  $J'(u_n) \rightarrow 0$  (as  $n \rightarrow \infty$ ) possesses a convergent subsequence.

To prove the existence of a critical point of the functional  $J$  we use the Saddle Point Theorem which is proved in Rabinowitz [9] (see also [10]).

**Theorem 2.2** (Saddle Point Theorem). *Let  $H = \widehat{H} \oplus \widetilde{H}$ ,  $\dim \widehat{H} < \infty$  and  $\dim \widetilde{H} = \infty$ . Let  $J : H \rightarrow \mathbb{R}$  be a functional such that  $J \in C^1(H, \mathbb{R})$  and*

- (a) *there exists a bounded neighborhood  $D$  of 0 in  $\widehat{H}$  and a constant  $\alpha$  such that  $J/\partial D \leq \alpha$ ,*
- (b) *there is a constant  $\beta > \alpha$  such that  $J/\widetilde{H} \geq \beta$ ,*
- (c)  *$J$  satisfies the Palais-Smale condition (PS).*

*Then, the functional  $J$  has a critical point in  $H$ .*

### 3. Main Result

We define

$$G_+(x) = \liminf_{s \rightarrow +\infty} \frac{G(x, s)}{s}, \quad G_-(x) = \limsup_{s \rightarrow -\infty} \frac{G(x, s)}{s}. \quad (3.1)$$

Assume that the following potential Landesman-Lazer type condition holds:

$$\int_0^T [R(x)^2 G_-(x)] dx < \int_0^T [R(x)^2 f(x)] dx < \int_0^T [R(x)^2 G_+(x)] dx. \quad (3.2)$$

We also suppose that there exists a function  $q(x) \in L^1(0, T)$  such that

$$|g(x, s)| \leq q(x), \quad x \in [0, T], \quad s \in \mathbb{R}. \quad (3.3)$$

**Theorem 3.1.** *Under the assumptions (1.2), (3.2), (3.3), problem (1.1) has at least one solution.*

*Proof.* We verify that the functional  $J$  satisfies assumptions of the Saddle Point Theorem 2.2 on  $H$ , then  $J$  has a critical point  $u$  and due to Lemma 2.1  $u$  is the solution to (1.1).

It is easy to see that  $J \in C^1(H, \mathbb{R})$ . Let  $\widetilde{H} = \{u \in H : \int_0^T u(x) dx = 0\}$  then  $H = \mathbb{R} \oplus \widetilde{H}$  and  $\dim(\widetilde{H}) = \infty$ .

In order to check assumption (a), we prove

$$\lim_{|s| \rightarrow \infty} J(s) = -\infty \quad (3.4)$$

by contradiction. Then, assume on the contrary there is a sequence of numbers  $(s_n) \subset \mathbb{R}$  such that  $|s_n| \rightarrow \infty$  and a constant  $c_1$  satisfying

$$\liminf_{n \rightarrow \infty} J(s_n) \geq c_1. \quad (3.5)$$

From the definition of  $J$  and from (3.5) it follows

$$\liminf_{n \rightarrow \infty} \int_0^T \frac{R^2(-G(x, s_n) + f s_n)}{|s_n|} dx \geq 0. \quad (3.6)$$

We note that from (3.2) it follows there exist constants  $s_+, s_-$  and functions  $A_+(x), A_-(x) \in L^1(0, T)$  such that  $A_+(x) \leq G(x, s), G(x, s) \leq A_-(x)$  for a.e.  $x \in (0, T)$  and for all  $s \geq s_+, s \leq s_-$ , respectively. We suppose that for this moment  $s_n \rightarrow +\infty$ . Using (3.6) and Fatou's lemma we obtain

$$\int_0^T [R(x)^2 f(x)] dx \geq \int_0^T [R(x)^2 G_+(x)] dx, \quad (3.7)$$

a contradiction to (3.2). We proceed for the case  $s_n \rightarrow -\infty$ . Then assumption (a) of Theorem 2.2 is verified.

(b) Now we prove that  $J$  is bounded from below on  $\widetilde{H}$ . For  $u \in \widetilde{H}$ , we have

$$\int_0^T (u')^2 dx = \|u\|^2 \quad (3.8)$$

and assumption (3.3) implies

$$|G(x, s)| \leq q(x)|s|, \quad x \in [0, T], \quad s \in \mathbb{R}. \quad (3.9)$$

Hence and due to compact imbedding  $H \subset C(0, T)$  ( $\|u\|_{C(0, T)} \leq c_2 \|u\|$ ) we obtain

$$\begin{aligned} J(u) &= \frac{1}{2} \int_0^T [R^2(u')^2] dx - \int_0^T [R^2 G(x, u) - R^2 f u] dx \\ &\geq \frac{1}{2} \min_{x \in [0, T]} R(x)^2 \int_0^T (u')^2 dx - \max_{x \in [0, T]} R(x)^2 \int_0^T (|q| + |f|) |u| dx \\ &\geq \frac{1}{2} \min_{x \in [0, T]} R(x)^2 \|u\|^2 - \max_{x \in [0, T]} R(x)^2 (\|q\|_1 + \|f\|_1) c_2 \|u\|. \end{aligned} \quad (3.10)$$

Since the function  $R$  is strictly positive equality (3.10) implies that the functional  $J$  is bounded from below.

Using (3.4), (3.10) we see that there exists a bounded neighborhood  $D$  of 0 in  $\mathbb{R} = \widehat{H}$ , a constant  $\alpha$  such that  $J/\partial D \leq \alpha$ , and there is a constant  $\beta > \alpha$  such that  $J/\widetilde{H} \geq \beta$ .

In order to check assumption (c), we show that  $J$  satisfies the Palais-Smale condition. First, we suppose that the sequence  $(u_n)$  is unbounded and there exists a constant  $c_3$  such that

$$\left| \frac{1}{2} \int_0^T [R^2(u'_n)^2] dx - \int_0^T [R^2(G(x, u_n) - f u_n)] dx \right| \leq c_3, \quad (3.11)$$

$$\lim_{n \rightarrow \infty} \|J'(u_n)\| = 0. \quad (3.12)$$

Let  $(w_k)$  be an arbitrary sequence bounded in  $H$ . It follows from (3.12) and the Schwarz inequality that

$$\begin{aligned} & \left| \lim_{\substack{n \rightarrow \infty \\ k \rightarrow \infty}} \int_0^T [R^2 u'_n w'_k] dx - \int_0^T [R^2 (g(x, u_n) w_k - f w_k)] dx \right| \\ &= \left| \lim_{\substack{n \rightarrow \infty \\ k \rightarrow \infty}} J'(u_n) w_k \right| \leq \lim_{\substack{n \rightarrow \infty \\ k \rightarrow \infty}} \|J'(u_n)\| \cdot \|w_k\| = 0. \end{aligned} \quad (3.13)$$

From (3.3) we obtain

$$\lim_{\substack{n \rightarrow \infty \\ k \rightarrow \infty}} \int_0^T \left[ \frac{R^2 g(x, u_n)}{\|u_n\|} w_k - \frac{R^2 f}{\|u_n\|} w_k \right] dx = 0. \quad (3.14)$$

Put  $v_n = u_n / \|u_n\|$  and  $w_k = v_n$  then (3.13), (3.14) imply

$$\lim_{n \rightarrow \infty} \int_0^T [R^2 (v'_n)^2] dx = 0. \quad (3.15)$$

Due to compact imbedding  $H \subset C(0, T)$  and (3.15) we have  $|v_n| \rightarrow d$  in  $C(0, T)$ ,  $d > 0$ . Suppose that  $v_n \rightarrow d$  and set  $w_k = v_n - d$  in (3.13), we get

$$\lim_{n \rightarrow \infty} \int_0^T [R^2 u'_n v'_n] dx - \int_0^T [R^2 (g(x, u_n) - f)(v_n - d)] dx = 0. \quad (3.16)$$

Because the nonlinearity  $g$  is bounded (assumption (3.3)) and  $v_n \rightarrow d$  the second integral in previous equality (3.16) converges to zero. Therefore

$$\lim_{n \rightarrow \infty} \int_0^T [R^2 u'_n v'_n] dx = 0. \quad (3.17)$$

Now we divide (3.11) by  $\|u_n\|$ . We get

$$\lim_{n \rightarrow \infty} \left\{ \frac{1}{2} \int_0^T [R^2 u'_n v'_n] dx - \int_0^T \frac{R^2 (G(x, u_n) - f u_n)}{\|u_n\|} dx \right\} = 0. \quad (3.18)$$

Equalities (3.17), (3.18) imply

$$\lim_{n \rightarrow \infty} \int_0^T R^2 \left( -\frac{G(x, u_n)}{u_n} + f \right) v_n dx = 0. \quad (3.19)$$

Because  $v_n \rightarrow d > 0$ ,  $\lim_{n \rightarrow \infty} u_n(x) = +\infty$ . Using Fatou's lemma and (3.19) we conclude

$$\int_0^T [R(x)^2 f(x)] dx \geq \int_0^T [R(x)^2 G_+(x)] dx, \quad (3.20)$$

a contradiction to (3.2). We proceed for the case  $v_n \rightarrow -d$  similarly. This implies that the sequence  $(u_n)$  is bounded. Then there exists  $u_0 \in H$  such that  $u_n \rightharpoonup u_0$  in  $H$ ,  $u_n \rightarrow u_0$  in  $L^2(0, T)$ ,  $C(0, T)$  (taking a subsequence if it is necessary). It follows from equality (3.13) that

$$\lim_{\substack{n \rightarrow \infty \\ m \rightarrow \infty \\ k \rightarrow \infty}} \left\{ \int_0^T [R^2(u_n - u_m)' w_k'] dx - \int_0^T [R^2(g(x, u_n) - g(x, u_m))] w_k dx \right\} = 0. \quad (3.21)$$

The strong convergence  $u_n \rightarrow u_0$  in  $C(0, T)$  and the assumption (3.3) imply

$$\lim_{\substack{n \rightarrow \infty \\ m \rightarrow \infty}} \int_0^T [R^2(g(x, u_n) - g(x, u_m))(u_n - u_m)] dx = 0. \quad (3.22)$$

If we set  $w_k = u_n$ ,  $w_k = u_m$  in (3.21) and subtract these equalities, then using (3.22) we have

$$\lim_{\substack{n \rightarrow \infty \\ m \rightarrow \infty}} \int_0^T [R^2(u_n' - u_m')^2] dx = 0. \quad (3.23)$$

Hence we obtain the strong convergence  $u_n \rightarrow u_0$  in  $H$ . This shows that  $J$  satisfies the Palais-Smale condition and the proof of Theorem 3.1 is complete.  $\square$

## Acknowledgment

This work was supported by Research Plan MSM 4977751301.

## References

- [1] P. Amster, "Nonlinearities in a second order ODE," in *Proceedings of the USA-Chile Workshop on Nonlinear Analysis*, vol. 6 of *Electron. J. Differ. Equ. Conf.*, pp. 13–21, Southwest Texas State Univ., San Marcos, Tex, USA, 2001.
- [2] P. Amster and M. C. Mariani, "A second order ODE with a nonlinear final condition," *Electronic Journal of Differential Equations*, vol. 2001, no. 75, p. 19, 2001.
- [3] H. Chen and Y. Li, "Rate of decay of stable periodic solutions of Duffing equations," *Journal of Differential Equations*, vol. 236, no. 2, pp. 493–503, 2007.
- [4] J. Mawhin and J. R. Ward, "Nonuniform nonresonance conditions at the two first eigenvalues for periodic solutions of forced Liénard and Duffing equations," *The Rocky Mountain Journal of Mathematics*, vol. 12, no. 4, pp. 643–654, 1982.
- [5] P. J. Torres, "Bounded solutions in singular equations of repulsive type," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 32, no. 1, pp. 117–125, 1998.
- [6] X. Li and Z. Zhang, "Periodic solutions for damped differential equations with a weak repulsive singularity," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 6, pp. 2395–2399, 2009.
- [7] A. C. Lazer, "On Schauder's fixed point theorem and forced second-order nonlinear oscillations," *Journal of Mathematical Analysis and Applications*, vol. 21, pp. 421–425, 1968.

- [8] P. Tomiczek, "Potential Landesman-Lazer type conditions and the Fučík spectrum," *Electronic Journal of Differential Equations*, vol. 2005, no. 94, pp. 1–12, 2005.
- [9] P. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Regional Conference Series in Mathematics, no. 65, American Mathematical Society, Providence, RI, USA, 1986.
- [10] P. H. Rabinowitz, "Some minimax theorems and applications to nonlinear partial differential equations," in *Nonlinear Analysis*, pp. 161–177, Academic Press, New York, NY, USA, 1978.