

Research Article

Positive Solution of a Nonlinear Fractional Differential Equation Involving Caputo Derivative

Changyou Wang,^{1,2,3} Haiqiang Zhang,² and Shu Wang³

¹ College of Mathematics and Physics, Chongqing University of Posts and Telecommunications, Chongqing 400065, China

² Key Laboratory of Industrial Internet of Things & Networked Control of Ministry of Education, Chongqing University of Posts and Telecommunications, Chongqing 400065, China

³ College of Applied Sciences, Beijing University of Technology, Beijing 100124, China

Correspondence should be addressed to Changyou Wang, wangcy@cqupt.edu.cn

Received 13 August 2012; Accepted 5 September 2012

Academic Editor: Seenith Sivasundaram

Copyright © 2012 Changyou Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with a nonlinear fractional differential equation involving Caputo derivative. By constructing the upper and lower control functions of the nonlinear term without any monotone requirement and applying the method of upper and lower solutions and the Schauder fixed point theorem, the existence and uniqueness of positive solution for the initial value problem are investigated. Moreover, the existence of maximal and minimal solutions is also obtained.

1. Introduction

Fractional differential equation can be extensively applied to various disciplines such as physics, mechanics, chemistry, and engineering, see [1–3]. Hence, in recent years, fractional differential equations have been of great interest and there have been many results on existence and uniqueness of the solution of FDE, see [4–8]. Especially, Diethelm and Ford [9] have gained existence, uniqueness, and structural stability of solution of the type of fractional differential equation

$$D^q(y - T_{m-1}[y])(x) = f(x, y(x)), \quad y^{(k)}(0) = y_0^{(k)}, \quad k = 0, 1, \dots, m-1, \quad (1.1)$$

where $q > 0$ is a real number, D^q denotes the Riemann-Liouville differential operator of order q , and $T_{m-1}[y]$ is the Taylor polynomial of order $(m-1)$ for the function $y(x)$ at

$x_0 = 0$. Recently, Daftardar-Gejji and Jafari [10] have discussed the existence, uniqueness, and stability of solution of the system of nonlinear fractional differential equation

$$D_*^\alpha y(t) = f(t, y(t)), \quad 0 < t < 1, \quad y^{(k)}(0) = c_k, \quad 0 \leq k \leq m, \quad 0 \leq c_k, \quad (1.2)$$

where $m < \alpha \leq m + 1$ and D_*^α denotes Caputo fractional derivative (see Definition 2.3). Delbosco and Rodino [11] have proved existence and uniqueness theorems for the nonlinear fractional equation

$$D^\delta u = f(t, u), \quad 0 < t < 1, \quad u(0) = 0, \quad (1.3)$$

where $0 < \delta < 1$, D^δ is the Riemann-Liouville fractional derivative. Zhang [12] used the method of the upper and lower solution and cone fixed point theorem to obtain the existence and uniqueness of positive solution to (1.3). Yao [13] considered the existence of positive solution to (1.3) controlled by the power function employing Krasnosel'skii fixed point theorem of cone expansion-compression type. The existence of the local and global solution for (1.3) was obtained by Lakshmikantham and Vatsala [14] utilizing classical differential equation theorem.

More recently, Zhang [15] shows the existence of positive solutions to the singular boundary value problem for fractional differential equation

$$\begin{aligned} D_{0+}^\alpha u(t) + g(t)f(u, u', \dots, u^{(n-2)}) &= 0, \quad 0 < t < 1, \\ u(0) = u'(0) = \dots = u^{(n-2)}(0) &= u^{(n-2)}(1) = 0, \end{aligned} \quad (1.4)$$

where D_{0+}^α is the Riemann-Liouville fractional derivative of order $n - 1 < \alpha \leq 2$, $n \geq 2$.

However, in the previous works, the nonlinear term has to satisfy the monotone or others control conditions. In fact, the fractional differential equations with nonmonotone function can respond better to impersonal law, so it is very important to weaken monotone condition. Considering this, in this paper, we mainly investigate the fractional differential Equation (1.2) without any monotone requirement on nonlinear term by constructing upper and lower control function and exploiting the method of upper and lower solutions and the Schauder fixed point theorem. The existence and uniqueness of positive solution for (1.2) are obtained. Some properties concerning the maximal and minimal solutions are also given. This work is motivated by the above references and my previous work [16, 17]. Other related results on the fractional differential equations can be found in [18–24].

This paper is organized as follow. In Section 2, we recall briefly some notions of the fractional calculus and the theory of the operators for integration and differentiation of fractional order. Section 3 is devoted to the study of the existence and uniqueness of positive solution for (1.2) utilizing the upper and lower solution method and the Schauder fixed point theorem. The existence of maximal and minimal solutions for (1.2) is given in Section 4.

2. Preliminaries and Notations

First, we give some basic definitions and theorems which are basically used throughout this paper. $C[0, 1]$ denotes the space of continuous functions defined on $[0, 1]$ and $C^n[0, 1]$

denotes the class of all real valued functions defined on $[0, 1]$ which have continuous n th order derivative.

Definition 2.1. Let $f(x) \in C[0, 1]$ and $\alpha > 0$, then the expression

$$I_0^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \quad 0 < x < 1 \quad (2.1)$$

is called the (left-sided) Riemann-Liouville integral of order α .

Definition 2.2. Let $n-1 < \alpha \leq n$, $n \in \mathbb{N}$, then the expression

$$D_0^\alpha f(x) = \frac{d^n}{dx^n} [I_0^{n-\alpha} f(x)], \quad 0 < x < 1 \quad (2.2)$$

is called the (left-sided) Riemann-Liouville derivative of $f(x)$ of order α whenever the expression on the right-hand side is defined.

Definition 2.3. Let $f(x) \in C^n[0, 1]$ and $n-1 < \alpha \leq n$, $n \in \mathbb{N}$, then the expression

$$D_{*0}^\alpha f(x) = I_0^{n-\alpha} f^{(n)}(x) \quad (2.3)$$

is called the (left-sided) Caputo derivative of $f(x)$ of order α .

In further discussion we will denote D_0^α , I_0^α , and D_{*0}^α as D^α , I^α , and D_*^α , respectively.

Lemma 2.4 (see [25, 26]). Let $f(x) \in C^n[0, 1]$ and $n-1 < \alpha \leq n$, $n \in \mathbb{N}$, then we one has

$$I^\mu I^\nu f(x) = I^{\mu+\nu} f(x), \quad \mu, \nu \geq 0, \quad (2.4)$$

$$D^\alpha I^\alpha f(x) = f(x), \quad (2.5)$$

$$I^\alpha D_*^\alpha f(x) = f(x) - \sum_{k=0}^{n-1} \frac{x^k}{k!} f^{(k)}(0+). \quad (2.6)$$

Lemma 2.5 (see, [10]). If the function $f(t, y(t))$ is $C^1[0, 1]$, then the initial value problem (1.2) is equivalent to the Volterra integral equations

$$y(t) = \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, y(t)), \quad 0 < t < 1, \quad m < \alpha \leq m+1. \quad (2.7)$$

Proof. Suppose $y(t)$ satisfies the initial value problem (1.2), then applying I^α to both sides of (1.2) and using Lemma 2.4 (2.7) follows. Conversely, suppose $y(t)$ satisfies (2.7). Then observe that $D^{(m+1)}y(t)$ exists and is integrable, because

$$\begin{aligned} y^{(m+1)}(t) &= D^{m+1} \left(\sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, y(t)) \right) \\ &= D^{m+1} I^\alpha f(t, y(t)) = DD^m I^m I^{\alpha-m} f(t, y(t)) \\ &= DI^{\alpha-m} f(t, y(t)) = D^{m+1-\alpha} f(t, y(t)), \end{aligned} \quad (2.8)$$

which exists and is integrable as $f(t, y(t))$ is $C^1[0, 1]$. Thus $I^{m+1-\alpha}y^{(m+1)}(t) = D_*^\alpha y(t)$ exists.

Applying D_*^α on both sides of (2.7), one has

$$\begin{aligned} D_*^\alpha y(t) &= I^{m+1-\alpha} D^{m+1} \left(\sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, y(t)) \right) \\ &= I^{m+1-\alpha} D^{m+1-\alpha} f(t, y(t)) = f(t, y(t)), \end{aligned} \quad (2.9)$$

as $f(t, y(t))$ is continuous and $0 < m + 1 - \alpha < 1$. Hence $y(t)$ satisfies (1.2). Moreover, from (2.4), $y^{(k)}(0) = c_k$, $0 \leq k \leq m$ hold. \square

Let $X = C[0, 1]$ be the Banach space endowed with the infinity norm and K a nonempty closed subset of X defined as $K = \{y(t) \in X \mid 0 < y(t) \leq l, 0 < t \leq 1, y^{(k)}(0) = c_k, 1 \leq k \leq m, 0 \leq c_k\}$. The positive solution which we consider in this paper is a function such that $y(t) \in K$.

According to Lemma 2.5, (1.2) is equivalent to the fractional integral Equation (2.7). The integral equation (2.7) is also equivalent to fixed point equation $Ty(t) = y(t)$, $y(t) \in C[0, 1]$, where operator $T : K \rightarrow K$ is defined as

$$Ty(t) = \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, y(t)), \quad (2.10)$$

then we have the following lemma.

Lemma 2.6. *Let $f : [0, 1] \times [0, l] \rightarrow R^+$ a given continuous function. Then the operator $T : K \rightarrow K$ is completely continuous.*

Proof. Let $M \subset K$ be bounded, that is, there exists a positive constant l^* such that $\|y\|_\infty \leq l^*$ for any $y(t) \in M$. Since $f(t, y(t))$ is a given continuous function, we have

$$\max_{0 \leq t \leq 1} f(t, y(t)) \leq \max_{(t, y) \in D} f(t, y), \quad \text{for any } y(t) \in M, \quad (2.11)$$

where $D = \{(t, y) \mid 0 \leq t \leq 1, 0 \leq y \leq l^*\}$.

Let $L = \max_{(t,y) \in D} f(t, y)$, then for any $y(t) \in M$, we have

$$\begin{aligned}
|Ty(t)| &= \left| \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, y(t)) \right| \\
&= \left| \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, y(s)) ds \right| \\
&\leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} |f(s, y(s))| ds \\
&\leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{L}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} ds \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{L}{\Gamma(1+\alpha)} t^\alpha.
\end{aligned} \tag{2.12}$$

Thus,

$$\|Ty\| \leq \sum_{k=0}^m \frac{c_k}{k!} + \frac{L}{\Gamma(1+\alpha)}. \tag{2.13}$$

Hence $T : K \rightarrow K$ is uniformly bounded.

Now, we prove that $T : K \rightarrow K$ is continuous. Since $f(t, y(t))$ is continuous function in a compact set $[0, 1] \times [0, l]$, then it is uniformly continuous there. Thus given $\varepsilon > 0$, we can find $\mu > 0$ such that $\|f(t, y) - f(t, z)\| < \varepsilon^*$ whenever $\|y - z\| < \mu$, where $\varepsilon^* = \varepsilon \Gamma(\alpha + 1)$. Then

$$\begin{aligned}
|Ty(t) - Tz(t)| &\leq \left| \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} [f(s, y(s)) - f(s, z(s))] ds \right| \\
&\leq \varepsilon^* \left| \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} ds \right| \\
&= \varepsilon^* \frac{t^\alpha}{\Gamma(\alpha + 1)} < \varepsilon,
\end{aligned} \tag{2.14}$$

proving the continuity of the operators $T : K \rightarrow K$.

Now, we will prove that the operator $T : K \rightarrow K$ is equicontinuous. For each $\mathbf{y}(t) \in M$, any $\varepsilon > 0$, $t_1, t_2 \in [0, 1]$ and $t_1 < t_2$. Let $\delta = \min\{(\varepsilon/2)(\sum_{k=1}^m (c_k/(k-1)!)^{-1}, (\varepsilon \Gamma(1+\alpha)/4L)^{1/\alpha}\}$, then when $|t_2 - t_1| < \delta$, we have

$$\begin{aligned}
|T\mathbf{y}(t_1) - T\mathbf{y}(t_2)| &= \left| \sum_{k=0}^m \frac{t_1^k}{k!} c_k + I^\alpha f(t_1, \mathbf{y}(t_1)) - \sum_{k=0}^m \frac{t_2^k}{k!} c_k - I^\alpha f(t_2, \mathbf{y}(t_2)) \right| \\
&\leq \left| \sum_{k=0}^m \frac{t_1^k}{k!} c_k - \sum_{k=0}^m \frac{t_2^k}{k!} c_k \right| \\
&\quad + \left| \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t_1 - s)^{\alpha-1} f(s, \mathbf{u}(s)) ds - \frac{1}{\Gamma(\alpha)} \int_0^{t_2} (t_2 - s)^{\alpha-1} f(s, \mathbf{u}(s)) ds \right| \\
&\leq (t_2 - t_1) \sum_{k=1}^m \frac{c_k}{(k-1)!} + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} |(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}| |f(s, \mathbf{u}(s))| ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} |f(s, \mathbf{u}(s))| ds \\
&\leq (t_2 - t_1) \sum_{k=1}^m \frac{c_k}{(k-1)!} + \frac{L}{\Gamma(\alpha)} \int_0^{t_1} ((t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}) ds \\
&\quad + \frac{L}{\Gamma(\alpha)} \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} ds \\
&= (t_2 - t_1) \sum_{k=1}^m \frac{c_k}{(k-1)!} \\
&\quad + \frac{L}{\Gamma(\alpha)} \left(\int_0^{t_1} (t_1 - s)^{\alpha-1} ds - \int_0^{t_1} (t_2 - s)^{\alpha-1} ds + \int_{t_1}^{t_2} (t_2 - s)^{\alpha-1} ds \right) \\
&= (t_2 - t_1) \sum_{k=1}^m \frac{c_k}{(k-1)!} + \frac{L}{\Gamma(1+\alpha)} [t_1^\alpha + (t_2 - t_1)^\alpha - t_2^\alpha + (t_2 - t_1)^\alpha] \\
&\leq (t_2 - t_1) \sum_{k=1}^m \frac{c_k}{(k-1)!} + \frac{2L}{\Gamma(1+\alpha)} (t_2 - t_1)^\alpha < \delta \sum_{k=1}^m \frac{c_k}{(k-1)!} + \frac{2L}{\Gamma(1+\alpha)} \delta^\alpha = \varepsilon.
\end{aligned} \tag{2.15}$$

The Arzela-Ascoli Theorem implies that T is completely continuous. The proof is therefore completed. \square

Lemma 2.7. *If the operator $A : X \rightarrow X$ is the contraction mapping, where X is the Banach space, then A has a unique fixed point in X .*

Let $f : [0, 1] \times [0, l] \rightarrow \mathbb{R}^+$ be a given function. Take $a, b \in \mathbb{R}^+$, and $a < b < l$. For any $\mathbf{y} \in [a, b]$ one defines the upper-control function $H(t, \mathbf{y}) = \sup_{a \leq \eta \leq \mathbf{y}} f(t, \eta)$, and lower-control function $h(t, \mathbf{y}) = \inf_{\mathbf{y} \leq \eta \leq b} f(t, \eta)$, obviously $H(t, \mathbf{y}), h(t, \mathbf{y})$ is monotonous nondecreasing on \mathbf{y} and $h(t, \mathbf{y}) \leq f(t, \mathbf{y}) \leq H(t, \mathbf{y})$.

Definition 2.8. Let $\tilde{y}(t), \hat{y}(t) \in K$, $b \geq \tilde{y}(t) \geq \hat{y}(t) \geq a$, and satisfy

$$\begin{aligned} D_*^\alpha \tilde{y}(t) &\geq f(t, \tilde{y}(t)), \quad \tilde{y}^{(k)}(0) \geq c_k, \quad m < \alpha \leq m+1, \quad 1 \leq k \leq m, \quad c_k \geq 0, \\ &\left(\text{or } \tilde{y}(t) \geq \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, \tilde{y}(t)), \quad m < \alpha \leq m+1 \right), \\ D_*^\alpha \hat{y}(t) &\leq f(t, \hat{y}(t)), \quad \hat{y}^{(k)}(0) \leq c_k, \quad m < \alpha \leq m+1, \quad 1 \leq k \leq m, \quad c_k \geq 0, \\ &\left(\text{or } \hat{y}(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, \hat{y}(t)), \quad m < \alpha \leq m+1 \right), \end{aligned} \quad (2.16)$$

then the functions $\tilde{y}(t), \hat{y}(t)$ are called a pair of order upper and lower solutions for (1.2).

3. Existence and Uniqueness of Positive Solution

Now, we give and prove the main results of this paper.

Theorem 3.1. *Assume $f : [0, 1] \times [0, l] \rightarrow [0, +\infty)$ is continuous, and $\tilde{y}(t), \hat{y}(t)$ are a pair of order upper and lower solutions of (1.2), then the boundary value problem (1.2) exists one solution $y(t) \in C[0, 1]$; moreover,*

$$\tilde{y}(t) \geq y(t) \geq \hat{y}(t), \quad t \in [0, 1]. \quad (3.1)$$

Proof. Let

$$S = \{z(t) \mid z(t) \in K, \hat{y}(t) \leq z(t) \leq \tilde{y}(t), t \in [0, 1]\}, \quad (3.2)$$

endowed with the norm $\|z\| = \max_{t \in [0, 1]} z(t)$, then we have $\|z\| \leq b$. Hence S is a convex, bounded, and closed subset of the Banach space X . According to Lemma 2.6, the operator $T : K \rightarrow K$ is completely continuous. Then we need only to prove $T : S \rightarrow S$.

For any $z(t) \in S$, we have $\tilde{y}(t) \geq z(t) \geq \hat{y}(t)$, then

$$\begin{aligned} Tz(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, z(t)) \\ &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, z(s)) ds \\ &\leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} H(s, z(s)) ds \\ &\leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} H(s, \tilde{y}(s)) ds \\ &\leq \tilde{y}(t), \end{aligned}$$

$$\begin{aligned}
Tz(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, z(t)) \\
&= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, z(s)) ds \\
&\geq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} h(s, z(s)) ds \\
&\geq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} h(s, \hat{y}(s)) ds \\
&\geq \hat{y}(t).
\end{aligned} \tag{3.3}$$

Hence $\tilde{y}(t) \geq Tz(t) \geq \hat{y}(t)$, $1 > t > 0$, that is, $T : S \rightarrow S$. According to Schauder fixed point theorem, the operator T exists at least one fixed point $y(t) \in S$, $0 < t < 1$. Therefore the boundary value problem (1.2) exists at least one solution $y(t) \in C[0, 1]$, and $\tilde{y}(t) \geq y(t) \geq \hat{y}(t)$, $t \in [0, 1]$. \square

Corollary 3.2. *Assume $f : [0, 1] \times [0, l] \rightarrow [0, +\infty)$ is continuous, and there exist $p_2 > p_1 \geq 0$, such that*

$$p_1 \leq f(t, s) \leq p_2, \quad (t, s) \in [0, 1] \times [0, l], \tag{3.4}$$

then the boundary value problem (1.2) exists at least one positive solution $y(t) \in C[0, 1]$, moreover

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{p_1 t^\alpha}{\Gamma(\alpha+1)} \leq y(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{p_2 t^\alpha}{\Gamma(\alpha+1)}. \tag{3.5}$$

Proof. By assumption (3.4) and the definition of control function, we have

$$p_1 \leq h(t, s) \leq H(t, s) \leq p_2, \quad (t, s) \in [0, 1] \times [a, b]. \tag{3.6}$$

Now, we consider the equation

$$D_*^\alpha w(t) = p_2, \quad w^{(k)}(0) = c_k, \quad m < \alpha \leq m+1, \quad 1 \leq k \leq m, \quad c_k \geq 0. \tag{3.7}$$

Obviously, (3.7) has a positive solution

$$\begin{aligned}
 w(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} p_2 ds = \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{p_2 t^\alpha}{\Gamma(\alpha+1)}, \quad t \in [0, 1], \\
 w(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{p_2 t^\alpha}{\Gamma(\alpha+1)} \\
 &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} p_2 ds \\
 &\geq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} H(t, w(t)) ds,
 \end{aligned} \tag{3.8}$$

namely, $w(t)$ is a upper solution of (1.2). In the similar way, we obtain $v(t) = \sum_{k=0}^m ((t^k/k!)c_k) + p_1 t^\alpha/\Gamma(\alpha+1)$ is the lower solution of (1.2). An application of Theorem 3.1 now yields that the boundary value problem (1.2) exists at least one positive solution $y(t) \in C[0, 1]$, moreover

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{p_1 t^\alpha}{\Gamma(\alpha+1)} \leq y(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{p_2 t^\alpha}{\Gamma(\alpha+1)}. \tag{3.9}$$

□

Corollary 3.3. Assume $f : [0, 1] \times [0, +\infty) \rightarrow [c, +\infty)$ is continuous, where $c > 0$, moreover

$$c < \lim_{y \rightarrow +\infty} f(t, y) < +\infty, \quad t \in [0, 1], \tag{3.10}$$

then the boundary value problem (1.2) has at least one positive solution $u(t) \in C[0, 1]$.

Proof. By assumption (3.9), there are positive constants N, R , such that $f(t, y) \leq N$ whenever $u > R$. Let $M = \max_{0 \leq t \leq 1, 0 \leq y \leq R} f(t, y)$, then $f(t, y) \leq N + M, 0 \leq y < +\infty$. By the definition of control function, one has $H(t, y) \leq N + M, 0 \leq t \leq 1, 0 \leq y < +\infty$.

Now, we consider the equation

$$D_*^\alpha w(t) = N + M, \quad w^{(k)}(0) = c_k, \quad m < \alpha \leq m + 1, \quad 1 \leq k \leq m, \quad c_k \geq 0. \tag{3.11}$$

Obviously, (3.11) has a positive solution

$$\begin{aligned}
 w(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (N+M) ds = \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{(N+M)t^\alpha}{\Gamma(\alpha+1)}, \quad t \in [0, 1], \\
 w(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{(N+M)t^\alpha}{\Gamma(\alpha+1)} \\
 &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (N+M) ds \\
 &\geq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} H(t, w(t)) ds,
 \end{aligned} \tag{3.12}$$

namely, $w(t)$ is the upper solution of (1.2). In the similar way, we obtain $v(t) = \sum_{k=0}^m (t^k/k!)c_k + ct^\alpha/\Gamma(\alpha+1)$ is the lower solution of (1.2). Therefore, the boundary value problem of (1.2) has at least one positive solution $y(t) \in C[0, 1]$, what is more, we have

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{ct^\alpha}{\Gamma(\alpha+1)} \leq y(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{(N+M)t^\alpha}{\Gamma(\alpha+1)}. \tag{3.13}$$

□

Corollary 3.4. Assume $f : [0, 1] \times [0, +\infty) \rightarrow [c, +\infty)$ is continuous, where $c > 0$, moreover

$$c < \lim_{y \rightarrow +\infty} \max_{0 \leq t \leq 1} \frac{f(t, y)}{y} = M < \Gamma(\alpha+1), \tag{3.14}$$

then the boundary value problem (1.2) exists at least at one positive solution $u(t) \in C[0, 1]$.

Proof. According to $c < \lim_{y \rightarrow +\infty} \max_{0 \leq t \leq 1} (f(t, y)/y) = M < +\infty$, there exists $D > 0$, such that for any $y(t) \in X$, we have

$$f(t, y(t)) \leq My(t) + D. \tag{3.15}$$

By the definition of control function, we have

$$H(t, y(t)) \leq My(t) + D. \tag{3.16}$$

We now consider the equation

$$D_*^\alpha w(t) = My(t) + D, \quad w^{(k)}(0) = c_k, \quad m < \alpha \leq m+1, \quad 1 \leq k \leq m, \quad c_k \geq 0. \tag{3.17}$$

According to Lemma 2.5, (3.17) is equivalent to the integral equation

$$\begin{aligned} y(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha (My(t) + D) \\ &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (My(s) + D) ds, \quad m < \alpha \leq m+1. \end{aligned} \quad (3.18)$$

Let $A : K \rightarrow K$ be an operator as follows:

$$\begin{aligned} A(y)(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha (My(t) + D) \\ &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (My(s) + D) ds, \quad m < \alpha \leq m+1. \end{aligned} \quad (3.19)$$

by Lemma 2.6, the operator A is completely continuous.

Let

$$B_R = \left\{ u(t) \in K \mid \left\| u - \frac{Dt^\alpha}{\Gamma(\alpha+1)} \right\| \leq R < +\infty \right\}, \quad (3.20)$$

where $R > 0$ and satisfies that $\sum_{k=0}^m (c_k/k!) + (M/\Gamma(\alpha+1))[(D/\Gamma(\alpha+1)) + R] - R \leq 0$, then B_R is convex, bounded, and closed subset of the Banach space $C[0, 1]$. For any $y(t) \in B_R$, we have

$$\|u\| \leq \frac{D}{\Gamma(\alpha+1)} + R, \quad (3.21)$$

then

$$\begin{aligned} \left\| Ay(t) - \frac{Dt^\alpha}{\Gamma(\alpha+1)} \right\| &= \left\| \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (My(s) + D) ds - \frac{Dt^\alpha}{\Gamma(\alpha+1)} \right\| \\ &= \left\| \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{M}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds \right\| \\ &\leq \sum_{k=0}^m \frac{t^k}{k!} c_k + M \|y(t)\| \frac{t^\alpha}{\Gamma(\alpha+1)} \\ &\leq \sum_{k=0}^m \frac{c_k}{k!} + \frac{M}{\Gamma(\alpha+1)} \left(R + \frac{D}{\Gamma(\alpha+1)} \right) \leq R, \end{aligned} \quad (3.22)$$

thus

$$\left\| Ay(t) - \frac{Dt^\alpha}{\Gamma(\alpha+1)} \right\| \leq R. \quad (3.23)$$

Hence, the Schauder fixed theorem assures that the operator A has at least one fixed point and then (3.17) has at least one positive solution $y^*(t)$, therefore we have

$$y^*(t) = \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} (My^*(s) + D) ds, \quad m < \alpha \leq m+1. \quad (3.24)$$

Combining condition (3.16), we have

$$y^*(t) \geq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} H(s, y^*(s)) ds, \quad m < \alpha \leq m+1. \quad (3.25)$$

Obviously, $y^*(t)$ is the upper solution of initial value problem (1.2), and $v(t) = \sum_{k=0}^m (c_k/k!) + ct^\alpha/\Gamma(\alpha+1)$ is the lower solution. By Theorem 3.1, system (1.2) has at least one positive solution $u(t) \in C[0, 1]$. \square

Corollary 3.5. *Assume $f : [0, 1] \times [0, +\infty) \rightarrow [c, +\infty)$ is continuous and there exists $d > 0$, $e > 0$, such that*

$$\max\{f(t, l) : (t, l) \in [0, 1] \times [0, d]\} \leq e, \quad (3.26)$$

then the boundary value problem (1.2) has at least one positive solution $y(t) \in C[0, 1]$, moreover

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{c t^\alpha}{\Gamma(\alpha+1)} \leq y(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{e t^\alpha}{\Gamma(\alpha+1)}. \quad (3.27)$$

Proof. By the definition of control function, we have

$$c \leq h(t, l) \leq H(t, l) \leq e, \quad (t, l) \in [0, 1] \times [0, d]. \quad (3.28)$$

By Corollary 3.2, the boundary value problem (1.2) has at least one positive solution $y(t) \in C[0, 1]$, moreover

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{ct^\alpha}{\Gamma(\alpha+1)} \leq y(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{e t^\alpha}{\Gamma(\alpha+1)}. \quad (3.29) \quad \square$$

Theorem 3.6. *Let the conditions in Theorem 3.1 hold. Moreover for any $y_1(t), y_2(t) \in X$, $0 < t < 1$, there exists $l > 0$, such that*

$$|f(t, y_1) - f(t, y_2)| \leq l|y_1 - y_2|, \quad (3.30)$$

then when $l/\Gamma(\alpha+1) < 1$, the boundary value problem (1.2) has a unique positive solution $y(t) \in S$.

Proof. According to Theorem 3.1, if the conditions in Theorem 3.1 hold, then the boundary value problem (1.2) has at least one positive solution in S . Hence we need only to prove that the operator T defined in (2.10) is the contraction mapping in X . In fact, for any $y_1(t), y_2(t) \in X$, by assumption (3.30), we have

$$\begin{aligned} |Ty_1(t) - Ty_2(t)| &= \left| \sum_{k=0}^m \frac{t^k}{k!} c_k + I^\alpha f(t, y_1(t)) - \sum_{k=0}^m \frac{t^k}{k!} c_k - I^\alpha f(t, y_2(t)) \right| \\ &= \left| \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, y_1(s)) ds - \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, y_2(s)) ds \right| \\ &\leq \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} |f(s, y_1(s)) - f(s, y_2(s))| ds \\ &\leq \frac{lt^\alpha}{\Gamma(\alpha+1)} |y_1 - y_2| < \frac{l}{\Gamma(\alpha+1)} |y_1 - y_2|. \end{aligned} \tag{3.31}$$

Thus, when $l/\Gamma(\alpha+1) < 1$, the operator T is the contraction mapping. Then by Lemma 2.7, the boundary value problem (1.2) has a unique positive solution $y(t) \in S$. \square

4. Maximal and Minimal Solutions Theorem

In this section, we consider the existence of maximal and minimal solutions for (1.2).

Definition 4.1. Let $m(t)$ be a solution of (1.2) in $[0, 1]$, then $m(t)$ is said to be a maximal solution of (1.2), if for every solution $y(t)$ of (1.2) existing on $[0, 1]$ the inequality $y(t) \leq m(t)$, $t \in [0, 1]$ holds. A minimal solution may be defined similarly by reversing the last inequality.

Theorem 4.2. Let $f : [0, 1] \times [0, +\infty) \rightarrow [0, +\infty)$ be a given continuous and monotone nondecreasing with respect to the second variable. Assume that there exist two positive constants λ, μ ($\mu > \lambda$) such that

$$\lambda \leq f(t, y) \leq \mu. \tag{4.1}$$

Then there exist maximal solution $\varphi(t)$ and minimal solution $\eta(t)$ of (1.2) on $[0, 1]$, moreover

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{\lambda t^\alpha}{\Gamma(\alpha+1)} \leq \eta(t) \leq \varphi(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{\mu t^\alpha}{\Gamma(\alpha+1)}, \quad 0 \leq t \leq 1. \tag{4.2}$$

Proof. It is easy to know that $\sum_{k=0}^m ((t^k/k!)c_k) + ((\mu t^\alpha)/\Gamma(\alpha+1))$ and $\sum_{k=0}^m ((t^k/k!)c_k) + ((\lambda t^\alpha)/\Gamma(\alpha+1))$ are the upper and lower solutions of (1.2), respectively. Then by using $\bar{y}^{(0)} = \sum_{k=0}^m ((t^k/k!)c_k) + \mu t^\alpha/\Gamma(\alpha+1)$, $\underline{y}^{(0)} = \sum_{k=0}^m ((t^k/k!)c_k) + \lambda t^\alpha/\Gamma(\alpha+1)$ as a pair of

coupled initial iterations we construct two sequences $\{\bar{y}^{(m)}\}, \{\underline{y}^{(m)}\}$ from the following linear iteration process:

$$\begin{aligned}\bar{y}^{(m)}(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, \bar{y}^{(m-1)}(t)) ds, \\ \underline{y}^{(m)}(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, \underline{y}^{(m-1)}(t)) ds.\end{aligned}\tag{4.3}$$

It is easy to show from the monotone property of $f(t, y)$ and condition (4.1) that the sequences $\{\bar{y}^{(m)}\}, \{\underline{y}^{(m)}\}$ possess the following monotone property:

$$\begin{aligned}\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{\lambda t^\alpha}{\Gamma(\alpha+1)} &= \underline{y}^{(0)} \leq \underline{y}^{(m)} \leq \underline{y}^{(m+1)} \leq \bar{y}^{(m+1)} \leq \bar{y}^{(m)} \leq \bar{y}^{(0)} \\ &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{\mu t^\alpha}{\Gamma(\alpha+1)} \quad (m = 1, 2, \dots).\end{aligned}\tag{4.4}$$

The above property implies that

$$\lim_{m \rightarrow \infty} \bar{y}^{(m)}(t) = \varphi(t), \quad \lim_{m \rightarrow \infty} \underline{y}^{(m)}(t) = \eta(t)\tag{4.5}$$

exist and satisfy the relation

$$\sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{\lambda t^\alpha}{\Gamma(\alpha+1)} \leq \eta(t) \leq \varphi(t) \leq \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{\mu t^\alpha}{\Gamma(\alpha+1)}, \quad 0 \leq t \leq 1.\tag{4.6}$$

Letting $m \rightarrow \infty$ in (4.3) shows that $\varphi(t)$ and $\eta(t)$ satisfy the equations

$$\begin{aligned}\varphi(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, \varphi(t)) ds, \\ \eta(t) &= \sum_{k=0}^m \frac{t^k}{k!} c_k + \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, \eta(t)) ds.\end{aligned}\tag{4.7}$$

It is easy to verify that the limits $\varphi(t)$ and $\eta(t)$ are maximal and minimal solutions of (1.2) in $S^* = \{\varphi(t) \mid \varphi(t) \in K, \sum_{k=0}^m ((t^k/k!)c_k) + \lambda t^\alpha/\Gamma(\alpha+1) \leq \varphi(t) \leq \sum_{k=0}^m ((t^k/k!)c_k) + \mu t^\alpha/\Gamma(\alpha+1), st \in [0, 1], \|\varphi(t)\| = \max_{0 \leq t \leq 1} \varphi(t)\}$, respectively, furthermore, if $\varphi(t) = \eta(t) (\equiv \zeta(t))$ then $\zeta(t)$ is the unique solution in S^* , and hence the proof is completed. \square

Acknowledgments

The authors are grateful to the referee for the comments. This work is supported by the Science and Technology Project of Chongqing Municipal Education Committee (Grants nos.

KJ110501 and KJ120520) of China, Natural Science Foundation Project of CQ CSTC (Grant no. cstc2012jjA20016) of China, and the NSFC (Grant no. 11101298) of China.

References

- [1] F. Mainardi, "The fundamental solutions for the fractional diffusion-wave equation," *Applied Mathematics Letters*, vol. 9, no. 6, pp. 23–28, 1996.
- [2] E. Buckwar and Y. Luchko, "Invariance of a partial differential equation of fractional order under the Lie group of scaling transformations," *Journal of Mathematical Analysis and Applications*, vol. 227, no. 1, pp. 81–97, 1998.
- [3] Z. Zheng-you, L. Gen-guo, and C. Chang-jun, "Quasi-static and dynamical analysis for viscoelastic Timoshenko beam with fractional derivative constitutive relation," *Applied Mathematics and Mechanics*, vol. 23, no. 1, pp. 1–12, 2002.
- [4] R. Gorenflo and S. Vessella, *Abel Integral Equations*, vol. 1461 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1991.
- [5] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, A Wiley-Interscience Publication, John Wiley & Sons, New York, NY, USA, 1993.
- [6] B. Ross and B. K. Sachdeva, "The solution of certain integral equations by means of operators of arbitrary order," *The American Mathematical Monthly*, vol. 97, no. 6, pp. 498–503, 1990.
- [7] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives*, Gordon and Breach Science, Yverdon, Switzerland, 1993.
- [8] H. M. Srivastava and R. G. Buschman, *Theory and Applications of Convolution Integral Equations*, vol. 79 of *Mathematics and Its Applications*, Kluwer Academic Publishers Group, Dordrecht, The Netherlands, 1992.
- [9] K. Diethelm and N. J. Ford, "Analysis of fractional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 265, no. 2, pp. 229–248, 2002.
- [10] V. Daftardar-Gejji and H. Jafari, "Analysis of a system of nonautonomous fractional differential equations involving Caputo derivatives," *Journal of Mathematical Analysis and Applications*, vol. 328, no. 2, pp. 1026–1033, 2007.
- [11] D. Delbosco and L. Rodino, "Existence and uniqueness for a nonlinear fractional differential equation," *Journal of Mathematical Analysis and Applications*, vol. 204, no. 2, pp. 609–625, 1996.
- [12] S. Zhang, "The existence of a positive solution for a nonlinear fractional differential equation," *Journal of Mathematical Analysis and Applications*, vol. 252, no. 2, pp. 804–812, 2000.
- [13] Q. L. Yao, "Existence of positive solution to a class of sublinear fractional differential equations," *Acta Mathematicae Applicatae Sinica*, vol. 28, no. 3, pp. 429–434, 2005 (Chinese).
- [14] V. Lakshmikantham and A. S. Vatsala, "Basic theory of fractional differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 8, pp. 2677–2682, 2008.
- [15] S. Zhang, "Positive solutions to singular boundary value problem for nonlinear fractional differential equation," *Computers & Mathematics with Applications*, vol. 59, no. 3, pp. 1300–1309, 2010.
- [16] C. Wang, "Existence and stability of periodic solutions for parabolic systems with time delays," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 2, pp. 1354–1361, 2008.
- [17] C. Wang, R. Wang, S. Wang, and C. Yang, "Positive solution of singular boundary value problem for a nonlinear fractional differential equation," *Boundary Value Problems*, vol. 2011, Article ID 297026, 12 pages, 2011.
- [18] V. Lakshmikantham, S. Leela, and J. Vasundhara Devi, *Theory of Fractional Dynamic Systems*, Cambridge Academic, Cambridge, UK, 2009.
- [19] Y.-K. Chang and J. J. Nieto, "Some new existence results for fractional differential inclusions with boundary conditions," *Mathematical and Computer Modelling*, vol. 49, no. 3-4, pp. 605–609, 2009.
- [20] N. Kosmatov, "Integral equations and initial value problems for nonlinear differential equations of fractional order," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 7, pp. 2521–2529, 2009.
- [21] G. Jumarie, "An approach via fractional analysis to non-linearity induced by coarse-graining in space," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 1, pp. 535–546, 2010.
- [22] R. P. Agarwal, V. Lakshmikantham, and J. J. Nieto, "On the concept of solution for fractional differential equations with uncertainty," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 6, pp. 2859–2862, 2010.

- [23] Z. Wei, Q. Li, and J. Che, "Initial value problems for fractional differential equations involving Riemann-Liouville sequential fractional derivative," *Journal of Mathematical Analysis and Applications*, vol. 367, no. 1, pp. 260–272, 2010.
- [24] K. Balachandran and J. J. Trujillo, "The nonlocal Cauchy problem for nonlinear fractional integrodifferential equations in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 12, pp. 4587–4593, 2010.
- [25] Y. Luchko and R. Gorenflo, "An operational method for solving fractional differential equations with the Caputo derivatives," *Acta Mathematica Vietnamica*, vol. 24, no. 2, pp. 207–233, 1999.
- [26] I. Podlubny, *Fractional Differential Equations*, vol. 198 of *Mathematics in Science and Engineering*, Academic Press, San Diego, Calif, USA, 1999.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

