

Research Article

Strong Convergence Theorem by Monotone Hybrid Algorithm for Equilibrium Problems, Hemirelatively Nonexpansive Mappings, and Maximal Monotone Operators

Yun Cheng and Ming Tian

College of Science, Civil Aviation University of China, Tianjin 300300, China

Correspondence should be addressed to Yun Cheng, mathscy@126.com

Received 17 June 2008; Accepted 11 November 2008

Recommended by Wataru Takahashi

We introduce a new hybrid iterative algorithm for finding a common element of the set of fixed points of hemirelatively nonexpansive mappings and the set of solutions of an equilibrium problem and for finding a common element of the set of zero points of maximal monotone operators and the set of solutions of an equilibrium problem in a Banach space. Using this theorem, we obtain three new results for finding a solution of an equilibrium problem, a fixed point of a hemirelatively nonexpansive mapping, and a zero point of maximal monotone operators in a Banach space.

Copyright © 2008 Y. Cheng and M. Tian. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let E be a Banach space, let C be a closed convex subset of E , and let f be a bifunction from $C \times C$ to \mathbb{R} , where \mathbb{R} is the set of real numbers. The equilibrium problem is to find

$$x^* \in C \quad \text{such that} \quad f(x^*, y) \geq 0 \quad \forall y \in C. \quad (1.1)$$

The set of such solutions x^* is denoted by $EP(f)$.

In 2006, Martinez-Yanes and Xu [1] obtained strong convergence theorems for finding a fixed point of a nonexpansive mapping by a new hybrid method in a Hilbert space. In particular, Takahashi and Zembayashi [2] established a strong convergence theorem for finding a common element of the set of solutions of an equilibrium problem and the set of fixed points of a nonexpansive mapping in a uniformly convex and uniformly smooth

Banach space. Very recently, Su et al. [3] proved the following theorem by a monotone hybrid method.

Theorem 1.1 (see Su et al. [3]). *Let E be a uniformly convex and uniformly smooth real Banach space, let C be a nonempty closed convex subset of E , and let $T : C \rightarrow C$ be a closed hemirelatively nonexpansive mapping such that $F(T) \neq \emptyset$. Assume that α_n is a sequence in $[0, 1]$ such that $\limsup_{n \rightarrow \infty} \alpha_n < 1$. Define a sequence x_n in C by the following:*

$$\begin{aligned} x_0 &\in C, \text{ chosen arbitrarily,} \\ y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JT x_n), \\ C_n &= \{z \in C_{n-1} \cap Q_{n-1} : \phi(z, y_n) \leq \phi(z, x_n)\}, \\ C_0 &= \{z \in C : \phi(z, y_0) \leq \phi(z, x_0)\}, \\ Q_n &= \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\ Q_0 &= C, \\ x_{n+1} &= \Pi_{C_n \cap Q_n}(x_0), \end{aligned} \tag{1.2}$$

where J is the duality mapping on E . Then, x_n converges strongly to $\Pi_{F(T)}x_0$, where $\Pi_{F(T)}$ is the generalized projection from C onto $F(T)$.

In this paper, motivated by Su et al. [3], we prove a strong convergence theorem for finding a common element of the set of solutions of an equilibrium problem and the set of fixed points of a hemirelatively nonexpansive mapping and for finding a common element of the set of zero points of maximal monotone operators and the set of solutions of an equilibrium problem in a Banach space by using the monotone hybrid method. Using this theorem, we obtain three new strong convergence results for finding a solution of an equilibrium problem, a fixed point of a hemirelatively nonexpansive mapping, and a zero point of maximal monotone operators in a Banach space.

2. Preliminaries

Let E be a real Banach space with dual E^* . We denote by J the normalized duality mapping from E to 2^{E^*} defined by

$$Jx = \{f \in E^* : \langle x, f \rangle = \|x\|^2 = \|f\|^2\}, \tag{2.1}$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. It is well known that if E^* is uniformly convex, then J is uniformly continuous on bounded subsets of E . In this case, J is single valued and also one to one.

Let E be a smooth, strictly convex, and reflexive Banach space and let C be a nonempty closed convex subset of E . Throughout this paper, we denote by ϕ the function defined by

$$\phi(y, x) = \|y\|^2 - 2\langle y, Jx \rangle + \|x\|^2. \tag{2.2}$$

Following Alber [4], the generalized projection $\Pi_C : E \rightarrow C$ from E onto C is defined by

$$\Pi_C(x) = \arg \min_{y \in C} \phi(y, x) \quad \forall x \in E. \quad (2.3)$$

The generalized projection Π_C from E onto C is well defined and single valued, and it satisfies

$$(\|x\| - \|y\|)^2 \leq \phi(y, x) \leq (\|x\| + \|y\|)^2 \quad \forall x, y \in E. \quad (2.4)$$

If E is a Hilbert space, then $\phi(y, x) = \|y - x\|^2$ and Π_C is the metric projection of E onto C .

If E is a reflexive strict convex and smooth Banach space, then for $x, y \in E$, $\phi(x, y) = 0$ if and only if $\|x\| = \|y\|$. It is sufficient to show that if $\phi(x, y) = 0$, then $x = y$. From (2.4), we have $\|x\| = \|y\|$. This implies $\langle x, Jy \rangle = \|x\|^2 = \|Jy\|^2$. From the definition of J , we have $Jx = Jy$, that is, $x = y$.

Let C be a closed convex subset of E and let T be a mapping from C into itself. We denote by $F(T)$ the set of fixed points of T . T is called hemirelatively nonexpansive if $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$.

A point p in C is said to be an asymptotic fixed point of T [5] if C contains a sequence x_n which converges weakly to p such that the strong $\lim_{n \rightarrow \infty} (Tx_n - x_n) = 0$. The set of asymptotic fixed points of T will be denoted by $\hat{F}(T)$. A hemirelatively nonexpansive mapping T from C into itself is called relatively nonexpansive [1, 5, 6] if $\hat{F}(T) = F(T)$.

We need the following lemmas for the proof of our main results.

Lemma 2.1 (see Alber [4]). *Let C be a nonempty closed convex subset of a smooth, strictly convex, and reflexive Banach space E . Then,*

$$\phi(x, \Pi_C y) + \phi(\Pi_C y, y) \leq \phi(x, y) \quad \forall x \in C, y \in E. \quad (2.5)$$

Lemma 2.2 (see Alber [4]). *Let C be a nonempty closed convex subset of a smooth, strictly convex, and reflexive Banach space, let $x \in E$, and let $z \in C$. Then,*

$$z = \Pi_C x \iff \langle y - z, Jx - Jz \rangle \leq 0 \quad \forall y \in C. \quad (2.6)$$

Lemma 2.3 (see Kamimura and Takahashi [7]). *Let E be a smooth and uniformly convex Banach space and let $\{x_n\}$ and $\{y_n\}$ be sequences in E such that either $\{x_n\}$ or $\{y_n\}$ is bounded. If $\lim_{n \rightarrow \infty} \phi(x_n, y_n) = 0$. Then $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.*

Lemma 2.4 (see Xu [8]). *Let E be a uniformly convex Banach space and let $r > 0$. Then, there exists a strictly increasing, continuous, and convex function $g : [0, 2r] \rightarrow \mathbb{R}$ such that $g(0) = 0$ and*

$$\|tx + (1-t)y\|^2 \leq t\|x\|^2 + (1-t)\|y\|^2 - t(1-t)g(\|x - y\|) \quad \forall x, y \in B_r, t \in [0, 1], \quad (2.7)$$

where $B_r = \{z \in E : \|z\| \leq r\}$.

Lemma 2.5 (see Kamimura and Takahashi [7]). *Let E be a smooth and uniformly convex Banach space and let $r > 0$. Then, there exists a strictly increasing, continuous, and convex function $g : [0, 2r] \rightarrow \mathbb{R}$ such that $g(0) = 0$ and*

$$g(\|x - y\|) \leq \phi(x, y) \quad \forall x, y \in B_r. \quad (2.8)$$

For solving the equilibrium problem, let us assume that a bifunction f satisfies the following conditions:

- (A1) $f(x, x) = 0$ for all $x \in C$;
- (A2) f is monotone, that is, $f(x, y) + f(y, x) \leq 0$ for all $x, y \in C$;
- (A3) for all $x, y, z \in C$, $\limsup_{t \rightarrow 0} f(tz + (1-t)x, y) \leq f(x, y)$;
- (A4) for all $x \in C$, $f(x, \cdot)$ is convex.

Lemma 2.6 (see Blum and Oettli [9]). *Let C be a closed convex subset of a smooth, strictly convex, and reflexive Banach space E , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), let $r > 0$, and let $x \in E$. Then, there exists $z \in C$ such that*

$$f(z, y) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0 \quad \forall y \in C. \quad (2.9)$$

Lemma 2.7 (see Takahashi and Zembayashi [10]). *Let C be a closed convex subset of a uniformly smooth, strictly convex, and reflexive Banach space E , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let $x \in E$, for $r > 0$. Define a mapping $T_r : E \rightarrow 2^C$ as follows:*

$$T_r(x) = \left\{ z \in C : f(z, y) + \frac{1}{r} \langle y - z, Jz - Jx \rangle \geq 0 \quad \forall y \in C \right\} \quad \forall x \in E. \quad (2.10)$$

Then, the following holds:

- (1) T_r is single valued;
- (2) T_r is a firmly nonexpansive-type mapping [11], that is, for all $x, y \in E$,

$$\langle T_r x - T_r y, JT_r x - JT_r y \rangle \leq \langle T_r x - T_r y, Jx - Jy \rangle; \quad (2.11)$$

- (3) $F(T_r) = \widehat{F}(T_r) = \text{Ep}(f)$;
- (4) $\text{Ep}(f)$ is closed and convex.

Lemma 2.8 (see Takahashi and Zembayashi [10]). *Let C be a closed convex subset of a smooth, strictly convex, and reflexive Banach space E and let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4). Then, for $r > 0$ and $x \in E$, and $q \in F(T_r)$,*

$$\phi(q, T_r x) + \phi(T_r x, x) \leq \phi(q, x). \quad (2.12)$$

Lemma 2.9 (see Su et al. [3]). *Let E be a strictly convex and smooth real Banach space, let C be a closed convex subset of E , and let T be a hemirelatively nonexpansive mapping from C into itself. Then, $F(T)$ is closed and convex.*

Recall that an operator T in a Banach space is called closed, if $x_n \rightarrow x$, $Tx_n \rightarrow y$, then $Tx = y$.

3. Strong convergence theorem

Theorem 3.1. *Let E be a uniformly convex and uniformly smooth real Banach space, let C be a nonempty closed convex subset of E , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let $T : C \rightarrow C$ be a closed hemirelatively nonexpansive mapping such that $F(T) \cap \text{EP}(f) \neq \emptyset$. Define a sequence $\{x_n\}$ in C by the following:*

$$\begin{aligned}
x_0 &\in C, \text{ chosen arbitrarily,} \\
y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JTz_n), \\
z_n &= J^{-1}(\beta_n Jx_n + (1 - \beta_n)JTx_n), \\
u_n &\in C \text{ such that } f(u_n, y) + \frac{1}{r_n} \langle y - u_n, Ju_n - Jy_n \rangle \geq 0, \quad \forall y \in C, \\
C_n &= \{z \in C_{n-1} \cap Q_{n-1} : \phi(z, u_n) \leq \phi(z, x_n)\}, \\
C_0 &= \{z \in C : \phi(z, u_0) \leq \phi(z, x_0)\}, \\
Q_n &= \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\
Q_0 &= C, \\
x_{n+1} &= \Pi_{C_n \cap Q_n}(x_0),
\end{aligned} \tag{3.1}$$

for every $n \in \mathbb{N} \cup \{0\}$, where J is the duality mapping on E , $\{\alpha_n\}$, $\{\beta_n\}$ are sequences in $[0, 1]$ such that $\liminf_{n \rightarrow \infty} (1 - \alpha_n)\beta_n(1 - \beta_n) > 0$ and $\{r_n\} \subset [a, \infty)$ for some $a > 0$. Then, $\{x_n\}$ converges strongly to $\Pi_{F(T) \cap \text{EP}(f)}x_0$, where $\Pi_{F(T) \cap \text{EP}(f)}$ is the generalized projection of E onto $F(T) \cap \text{EP}(f)$.

Proof. First, we can easily show that C_n and Q_n are closed and convex for each $n \geq 0$.

Next, we show that $F(T) \cap \text{EP}(f) \subset C_n$ for all $n \geq 0$. Let $u \in F(T) \cap \text{EP}(f)$. Putting $u_n = T_{r_n}y_n$ for all $n \in \mathbb{N}$, from Lemma 2.8, we have T_{r_n} relatively nonexpansive. Since T_{r_n} are relatively nonexpansive and T is hemirelatively nonexpansive, we have

$$\begin{aligned}
\phi(u, z_n) &= \phi(u, J^{-1}(\beta_n Jx_n + (1 - \beta_n)JTx_n)) \\
&= \|u\|^2 - 2\langle u, \beta_n Jx_n + (1 - \beta_n)JTx_n \rangle + \|\beta_n Jx_n + (1 - \beta_n)JTx_n\|^2 \\
&\leq \|u\|^2 - 2\beta_n \langle u, Jx_n \rangle - 2(1 - \beta_n) \langle u, JTx_n \rangle + \beta_n \|x_n\|^2 + (1 - \beta_n) \|Tx_n\|^2 \\
&= \beta_n \phi(u, x_n) + (1 - \beta_n) \phi(u, Tx_n) \\
&\leq \phi(u, x_n), \\
\phi(u, u_n) &= \phi(u, T_{r_n}y_n) \leq \phi(u, y_n) \leq \alpha_n \phi(u, x_n) + (1 - \alpha_n) \phi(u, z_n) \leq \phi(u, x_n).
\end{aligned} \tag{3.2}$$

Hence, we have

$$F(T) \cap EP(f) \subset C_n \quad \forall n \geq 0. \quad (3.3)$$

Next, we show that $F(T) \cap EP(f) \subset Q_n$ for all $n \geq 0$. We prove this by induction. For $n = 0$, we have

$$F(T) \cap EP(f) \subset Q_0 = C. \quad (3.4)$$

Suppose that $F(T) \cap EP(f) \subset Q_n$, by Lemma 2.2, we have

$$\langle x_{n+1} - z, Jx_0 - Jx_{n+1} \rangle \geq 0 \quad \forall z \in C_n \cap Q_n. \quad (3.5)$$

As $F(T) \cap EP(f) \subset C_n \cap Q_n$, by the induction assumptions, the last inequality holds, in particular, for all $z \in F(T) \cap EP(f)$. This, together with the definition of Q_{n+1} , implies that $F(T) \cap EP(f) \subset Q_{n+1}$. So, $\{x_n\}$ is well defined.

Since $x_{n+1} = \Pi_{C_n \cap Q_n} x_0$ and $C_n \cap Q_n \subset C_{n-1} \cap Q_{n-1}$ for all $n \geq 1$, we have

$$\phi(x_n, x_0) \leq \phi(x_{n+1}, x_0) \quad \forall n \geq 0. \quad (3.6)$$

Therefore, $\{\phi(x_n, x_0)\}$ is nondecreasing. In addition, from the definition of Q_n and Lemma 2.2, $x_n = \Pi_{Q_n} x_0$. Therefore, for each $u \in F(T) \cap EP(f)$, we have

$$\phi(x_n, x_0) = \phi(\Pi_{Q_n} x_0, x_0) \leq \phi(u, x_0) - \phi(u, x_n) \leq \phi(u, x_0). \quad (3.7)$$

Therefore, $\phi(x_n, x_0)$ and $\{x_n\}$ are bounded. This, together with (3.6), implies that the limit of $\{\phi(x_n, x_0)\}$ exists. From Lemma 2.1, we have, for any positive integer m ,

$$\phi(x_{n+m}, x_n) = \phi(x_{n+m}, \Pi_{Q_n} x_0) \leq \phi(x_{n+m}, x_0) - \phi(\Pi_{Q_n} x_0, x_0) = \phi(x_{n+m}, x_0) - \phi(x_n, x_0) \quad \forall n \geq 0. \quad (3.8)$$

Therefore,

$$\lim_{n \rightarrow \infty} \phi(x_{n+m}, x_n) = 0. \quad (3.9)$$

From (3.11), we can prove that $\{x_n\}$ is a Cauchy sequence. Therefore, there exists a point $\hat{x} \in C$ such that $\{x_n\}$ converges strongly to \hat{x} .

Since $x_{n+1} \in C_n$, we have

$$\phi(x_{n+1}, u_n) \leq \phi(x_{n+1}, x_n). \quad (3.10)$$

Therefore, we have

$$\phi(x_{n+1}, u_n) \longrightarrow 0. \quad (3.11)$$

From Lemma 2.3, we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - u_n\| = \lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (3.12)$$

So, we have

$$\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0. \quad (3.13)$$

Since J is uniformly norm-to-norm continuous on bounded sets, we have

$$\lim_{n \rightarrow \infty} \|Jx_n - Ju_n\| = 0. \quad (3.14)$$

Let $r = \sup_{n \in \mathbb{N}} \{\|x_n\|, \|Tx_n\|\}$. Since E is a uniformly smooth Banach space, we know that E^* is a uniformly convex Banach space. Therefore, from Lemma 2.4, there exists a continuous, strictly increasing, and convex function g with $g(0) = 0$, such that

$$\|\alpha x^* + (1 - \alpha)y^*\|^2 \leq \alpha \|x^*\|^2 + (1 - \alpha)\|y^*\|^2 - \alpha(1 - \alpha)g(\|x^* - y^*\|) \quad (3.15)$$

for $x^*, y^* \in B_r$, and $\alpha \in [0, 1]$. So, we have that for $u \in F(T) \cap EP(f)$,

$$\begin{aligned} \phi(u, z_n) &= \phi(u, J^{-1}(\beta_n Jx_n + (1 - \beta_n)JT x_n)) \\ &= \|u\|^2 - 2\langle u, \beta_n Jx_n + (1 - \beta_n)JT x_n \rangle + \|\beta_n Jx_n + (1 - \beta_n)JT x_n\|^2 \\ &\leq \phi(u, x_n) - \beta_n(1 - \beta_n)g(\|Jx_n - JT x_n\|), \\ \phi(u, u_n) &\leq \alpha_n \phi(u, x_n) + (1 - \alpha_n)\phi(u, z_n) \\ &\leq \phi(u, x_n) - (1 - \alpha_n)\beta_n(1 - \beta_n)g(\|Jx_n - JT x_n\|). \end{aligned} \quad (3.16)$$

Therefore, we have

$$(1 - \alpha_n)\beta_n(1 - \beta_n)g(\|Jx_n - JT x_n\|) \leq \phi(u, x_n) - \phi(u, u_n). \quad (3.17)$$

Since

$$\phi(u, x_n) - \phi(u, u_n) = \|x_n\|^2 - \|u_n\|^2 - 2\langle u, Jx_n - Ju_n \rangle \leq \|x_n - u_n\|(\|x_n\| + \|u_n\|) + 2\|u\|\|Jx_n - Ju_n\|, \quad (3.18)$$

we have

$$\lim_{n \rightarrow \infty} \phi(u, x_n) - \phi(u, u_n) = 0. \quad (3.19)$$

From $\liminf_{n \rightarrow \infty} (1 - \alpha_n)\beta_n(1 - \beta_n) > 0$, we have

$$\lim_{n \rightarrow \infty} g(\|Jx_n - JT x_n\|) = 0. \quad (3.20)$$

Therefore, from the property of g , we have

$$\lim_{n \rightarrow \infty} \|Jx_n - JT x_n\| = 0. \quad (3.21)$$

Since J^{-1} is uniformly norm-to-norm continuous on bounded sets, we have

$$\lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0. \quad (3.22)$$

Since T is a closed operator and $x_n \rightarrow \hat{x}$, then \hat{x} is a fixed point of T .

On the other hand,

$$\phi(u_n, y_n) = \phi(T_{r_n} y_n, y_n) \leq \phi(u, y_n) - \phi(u, T_{r_n} y_n) \leq \phi(u, x_n) - \phi(u, T_{r_n} y_n) = \phi(u, x_n) - \phi(u, u_n). \quad (3.23)$$

So, we have from (3.21) that

$$\lim_{n \rightarrow \infty} \phi(u_n, y_n) = 0. \quad (3.24)$$

From Lemma 2.3, we have that

$$\lim_{n \rightarrow \infty} \|u_n - y_n\| = 0. \quad (3.25)$$

From $x_n \rightarrow \hat{x}$ and $\|x_n - u_n\| \rightarrow 0$, we have $y_n \rightarrow \hat{x}$.

From (3.27), we have

$$\lim_{n \rightarrow \infty} \|J u_n - J y_n\| = 0. \quad (3.26)$$

From $r_n \geq a$, we have

$$\lim_{n \rightarrow \infty} \frac{\|J u_n - J y_n\|}{r_n} = 0. \quad (3.27)$$

By $u_n = T_{r_n} y_n$, we have

$$f(u_n, y) + \frac{1}{r_n} \langle y - u_n, J u_n - J y_n \rangle \geq 0 \quad \forall y \in C. \quad (3.28)$$

From (A2), we have that

$$\frac{1}{r_n} \langle y - u_n, J u_n - J y_n \rangle \geq -f(u_n, y) \geq f(y, u_n) \quad \forall y \in C. \quad (3.29)$$

From (3.29) and (A4), we have

$$f(\mathbf{y}, \hat{x}) \leq 0 \quad \forall \mathbf{y} \in C. \quad (3.30)$$

For t with $0 < t \leq 1$ and $\mathbf{y} \in C$, let $\mathbf{y}_t = t\mathbf{y} + (1-t)\hat{x}$. We have $f(\mathbf{y}_t, \hat{x}) \leq 0$. So, from (A1), we have

$$0 = f(\mathbf{y}_t, \mathbf{y}_t) \leq tf(\mathbf{y}_t, \mathbf{y}) + (1-t)f(\mathbf{y}_t, \hat{x}) \leq tf(\mathbf{y}_t, \mathbf{y}). \quad (3.31)$$

Dividing by t , we have

$$f(\mathbf{y}_t, \mathbf{y}) \geq 0 \quad \forall \mathbf{y} \in C. \quad (3.32)$$

Letting $t \rightarrow 0$, from (A3), we have

$$f(\hat{x}, \mathbf{y}) \geq 0 \quad \forall \mathbf{y} \in C. \quad (3.33)$$

Therefore, $\hat{x} \in \text{EP}(f)$. Finally, we prove that $\hat{x} = \Pi_{F(T) \cap \text{EP}(f)} x_0$. From Lemma 2.1, we have

$$\phi(\hat{x}, \Pi_{F(T) \cap \text{EP}(f)} x_0) + \phi(\Pi_{F(T) \cap \text{EP}(f)} x_0, x_0) \leq \phi(\hat{x}, x_0). \quad (3.34)$$

Since $x_{n+1} = \Pi_{C_n \cap Q_n} x_0$ and $\hat{x} \in F(T) \cap \text{EP}(f) \subset C_n \cap Q_n$, for all $n \geq 0$, we get from Lemma 2.1 that

$$\phi(\Pi_{F(T) \cap \text{EP}(f)} x_0, x_{n+1}) + \phi(x_{n+1}, x_0) \leq \phi(\Pi_{F(T) \cap \text{EP}(f)} x_0, x_0). \quad (3.35)$$

By the definition of $\phi(x, y)$, it follows that $\phi(\hat{x}, x_0) \leq \phi(\Pi_{F(T) \cap \text{EP}(f)} x_0, x_0)$ and $\phi(\hat{x}, x_0) \geq \phi(\Pi_{F(T) \cap \text{EP}(f)} x_0, x_0)$, whence $\phi(\hat{x}, x_0) = \phi(\Pi_{F(T) \cap \text{EP}(f)} x_0, x_0)$. Therefore, it follows from the uniqueness of $\Pi_{F(T) \cap \text{EP}(f)} x_0$ that $\hat{x} = \Pi_{F(T) \cap \text{EP}(f)} x_0$. This completes the proof. \square

Corollary 3.2. *Let E be a uniformly convex and uniformly smooth real Banach space, let C be a nonempty closed convex subset of E , and let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4). Define a sequence $\{x_n\}$ in C by the following:*

$$\begin{aligned} & x_0 \in C, \text{ chosen arbitrarily,} \\ & u_n \in C \text{ such that } f(u_n, \mathbf{y}) + \frac{1}{r_n} \langle \mathbf{y} - u_n, Ju_n - Jx_n \rangle \geq 0, \quad \forall \mathbf{y} \in C, \\ & C_n = \{z \in C_{n-1} \cap Q_{n-1} : \phi(z, u_n) \leq \phi(z, x_n)\}, \\ & C_0 = \{z \in C : \phi(z, u_0) \leq \phi(z, x_0)\}, \\ & Q_n = \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\ & Q_0 = C, \\ & x_{n+1} = \Pi_{C_n \cap Q_n}(x_0), \end{aligned} \quad (3.36)$$

for every $n \in \mathbb{N} \cup \{0\}$, where J is the duality mapping on E and $\{r_n\} \subset [a, \infty)$ for some $a > 0$. Then, $\{x_n\}$ converges strongly to $\Pi_{\text{EP}(f)}x_0$.

Proof. Putting $T = I$ in Theorem 3.1, we obtain Corollary 3.2. \square

Corollary 3.3. *Let E be a uniformly convex and uniformly smooth real Banach space, let C be a nonempty closed convex subset of E , and let $T : C \rightarrow C$ be a closed hemirelatively nonexpansive mapping. Define a sequence $\{x_n\}$ in C by the following:*

$$\begin{aligned}
x_0 &\in C, \text{ chosen arbitrarily,} \\
y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JTz_n), \\
z_n &= J^{-1}(\beta_n Jx_n + (1 - \beta_n)JTz_n), \\
u_n &= \Pi_C y_n, \\
C_n &= \{z \in C_{n-1} \cap Q_{n-1} : \Phi(z, u_n) \leq \Phi(z, x_n)\}, \\
C_0 &= \{z \in C : \Phi(z, u_0) \leq \Phi(z, x_0)\}, \\
Q_n &= \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\
Q_0 &= C, \\
x_{n+1} &= \Pi_{C_n \cap Q_n}(x_0),
\end{aligned} \tag{3.37}$$

for every $n \in \mathbb{N} \cup \{0\}$, where J is the duality mapping on E , $\{\alpha_n\}, \{\beta_n\}$ are sequences in $[0, 1]$ such that $\liminf_{n \rightarrow \infty} (1 - \alpha_n)\beta_n(1 - \beta_n) > 0$. Then, $\{x_n\}$ converges strongly to $\Pi_{F(T)}x_0$.

Proof. Putting $f(x, y) = 0$ for all $x, y \in C$ and $r_n = 1$ for all n in Theorem 3.1, we obtain Corollary 3.3. \square

Corollary 3.4. *Let E be a uniformly convex and uniformly smooth real Banach space, let C be a nonempty closed convex subset of E , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let $T : C \rightarrow C$ be a closed relatively nonexpansive mapping such that $F(T) \cap \text{EP}(f) \neq \emptyset$. Define a sequence $\{x_n\}$ in C by the following:*

$$\begin{aligned}
x_0 &\in C, \text{ chosen arbitrarily,} \\
y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JTz_n), \\
z_n &= J^{-1}(\beta_n Jx_n + (1 - \beta_n)JTz_n), \\
u_n &\in C \text{ such that } f(u_n, y) + \frac{1}{r_n} \langle y - u_n, Ju_n - Jy_n \rangle \geq 0, \quad \forall y \in C, \\
C_n &= \{z \in C_{n-1} \cap Q_{n-1} : \Phi(z, u_n) \leq \Phi(z, x_n)\}, \\
C_0 &= \{z \in C : \Phi(z, u_0) \leq \Phi(z, x_0)\}, \\
Q_n &= \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\
Q_0 &= C, \\
x_{n+1} &= \Pi_{C_n \cap Q_n}(x_0),
\end{aligned} \tag{3.38}$$

for every $n \in \mathbb{N} \cup \{0\}$, where J is the duality mapping on E , $\{\alpha_n\}$, $\{\beta_n\}$ are sequences in $[0, 1]$ such that $\liminf_{n \rightarrow \infty} (1 - \alpha_n)\beta_n(1 - \beta_n) > 0$ and $\{r_n\} \subset [a, \infty)$ for some $a > 0$. Then, $\{x_n\}$ converges strongly to $\Pi_{F(T) \cap \text{EP}(f)}x_0$.

Proof. Since every relatively nonexpansive mapping is a hemirelatively one, Corollary 3.4 is implied by Theorem 3.1. \square

Remark 3.5 (see Rockafellar [12]). Let E be a reflexive, strictly convex, and smooth Banach space and let A be a monotone operator from E to E^* . Then, A is maximal if and only if $R(J + rA) = E^*$ for all $r > 0$.

Let E be a reflexive, strictly convex, and smooth Banach space and let A be a maximal monotone operator from E to E^* . Using Remark 3.5 and strict convexity of E , we obtain that for every $r > 0$ and $x \in E$, there exists a unique $x_r \in D(A)$ such that $Jx \in Jx_r + rAx_r$. If $J_r x = x_r$, then we can define a single-valued mapping $J_r : E \rightarrow D(A)$ by $J_r = (J + rA)^{-1}J$, and such a J_r is called the resolvent of A . We know that $A^{-1}0 = F(J_r)$ for all $r > 0$ and J_r is relatively nonexpansive mapping (see [2] for more details). Using Theorem 3.1, we can consider the problem of strong convergence concerning maximal monotone operators in a Banach space.

Theorem 3.6. *Let E be a uniformly convex and uniformly smooth real Banach space, let C be a nonempty closed convex subset of E , let f be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let J_r be a resolvent of A and a closed mapping such that $A^{-1}0 \cap \text{EP}(f) \neq \emptyset$, where $r > 0$. Define a sequence $\{x_n\}$ in C by the following:*

$$\begin{aligned}
& x_0 \in C, \text{ chosen arbitrarily,} \\
& y_n = J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JJ_r x_n), \\
& z_n = J^{-1}(\beta_n Jx_n + (1 - \beta_n)JJ_r x_n), \\
& u_n \in C \text{ such that } f(u_n, y) + \frac{1}{r_n} \langle y - u_n, Ju_n - Jy_n \rangle \geq 0, \quad \forall y \in C, \\
& C_n = \{z \in C_{n-1} \cap Q_{n-1} : \phi(z, u_n) \leq \phi(z, x_n)\}, \\
& C_0 = \{z \in C : \phi(z, u_0) \leq \phi(z, x_0)\}, \\
& Q_n = \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx_0 - Jx_n \rangle \geq 0\}, \\
& Q_0 = C, \\
& x_{n+1} = \Pi_{C_n \cap Q_n}(x_0),
\end{aligned} \tag{3.39}$$

for every $n \in \mathbb{N} \cup \{0\}$, where J is the duality mapping on E , $\{\alpha_n\}$ is a sequences in $[0, 1]$ such that $\liminf_{n \rightarrow \infty} (1 - \alpha_n)\beta_n(1 - \beta_n) > 0$ and $\{r_n\} \subset [a, \infty)$ for some $a > 0$, Then, $\{x_n\}$ converges strongly to $\Pi_{A^{-1}0 \cap \text{EP}(f)}x_0$.

Proof. Since J_r is a closed relatively nonexpansive mapping and $A^{-1}0 = F(J_r)$, from Corollary 3.4, we obtain Theorem 3.6. \square

Acknowledgment

This work is supported by Tianjin Natural Science Foundation in China Grant no. 06YFJMJC12500.

References

- [1] C. Martinez-Yanes and H.-K. Xu, "Strong convergence of the CQ method for fixed point iteration processes," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 11, pp. 2400–2411, 2006.
- [2] W. Takahashi and K. Zembayashi, "Strong convergence theorem by a new hybrid method for equilibrium problems and relatively nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2008, Article ID 528476, 11 pages, 2008.
- [3] Y. Su, D. Wang, and M. Shang, "Strong convergence of monotone hybrid algorithm for hemi-relatively nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2008, Article ID 284613, 8 pages, 2008.
- [4] Y. I. Alber, "Metric and generalized projection operators in Banach spaces: properties and applications," in *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*, A. G. Kartsatos, Ed., vol. 178 of *Lecture Notes in Pure and Applied Mathematics*, pp. 15–50, Marcel Dekker, New York, NY, USA, 1996.
- [5] D. Butnariu, S. Reich, and A. J. Zaslavski, "Asymptotic behavior of relatively nonexpansive operators in Banach spaces," *Journal of Applied Analysis*, vol. 7, no. 2, pp. 151–174, 2001.
- [6] S.-Y. Matsushita and W. Takahashi, "A strong convergence theorem for relatively nonexpansive mappings in a Banach space," *Journal of Approximation Theory*, vol. 134, no. 2, pp. 257–266, 2005.
- [7] S. Kamimura and W. Takahashi, "Strong convergence of a proximal-type algorithm in a Banach space," *SIAM Journal on Optimization*, vol. 13, no. 3, pp. 938–945, 2002.
- [8] H. K. Xu, "Inequalities in Banach spaces with applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 16, no. 12, pp. 1127–1138, 1991.
- [9] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [10] W. Takahashi and K. Zembayashi, "Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 1, pp. 45–57, 2009.
- [11] F. Kohsaka and W. Takahashi, "Existence and approximation of fixed points of firmly nonexpansive type mappings in Banach spaces," to appear in *SIAM Journal on Optimization*.
- [12] R. T. Rockafellar, "On the maximality of sums of nonlinear monotone operators," *Transactions of the American Mathematical Society*, vol. 149, pp. 75–88, 1970.