

Research Article

Existence and Localization Results for $p(x)$ -Laplacian via Topological Methods

B. Cekic and R. A. Mashiyev

Department of Mathematics, Dicle University, 21280 Diyarbakir, Turkey

Correspondence should be addressed to B. Cekic, bilalcekic@gmail.com

Received 23 February 2010; Revised 16 April 2010; Accepted 20 June 2010

Academic Editor: J. Mawhin

Copyright © 2010 B. Cekic and R. A. Mashiyev. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We show the existence of a weak solution in $W_0^{1,p(x)}(\Omega)$ to a Dirichlet problem for $-\Delta_{p(x)}u = f(x, u)$ in Ω , and its localization. This approach is based on the nonlinear alternative of Leray-Schauder.

1. Introduction

In this work, we consider the boundary value problem

$$\begin{aligned} -\Delta_{p(x)}u &= f(x, u) \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{P}$$

where $\Omega \subset \mathbb{R}^N$, $N \geq 2$, is a nonempty bounded open set with smooth boundary $\partial\Omega$, $\Delta_{p(x)}u = \operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ is the so-called $p(x)$ -Laplacian operator, and (CAR): $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Caratheodory function which satisfies the growth condition

$$|f(x, s)| \leq a(x) + C|s|^{q(x)/q'(x)} \quad \text{for a.e. } x \in \Omega \text{ and all } s \in \mathbb{R}, \tag{1.1}$$

with $C = \operatorname{const.} > 0$, $1/q(x) + 1/q'(x) = 1$ for a.e. $x \in \Omega$, and $a \in L^{q'(x)}(\Omega)$, $a(x) \geq 0$ for a.e. $x \in \Omega$.

We recall in what follows some definitions and basic properties of variable exponent Lebesgue and Sobolev spaces $L^{p(x)}(\Omega)$, $W^{1,p(x)}(\Omega)$, and $W_0^{1,p(x)}(\Omega)$. In that context, we refer to [1, 2] for the fundamental properties of these spaces.

Set

$$L_+^\infty(\Omega) = \left\{ p : p \in L^\infty(\Omega), \operatorname{ess\,inf}_{x \in \Omega} p(x) > 1 \right\}. \quad (1.2)$$

For $p \in L_+^\infty(\Omega)$, let $p_1 := \operatorname{ess\,inf}_{x \in \Omega} p(x) \leq p(x) \leq p_2 := \operatorname{ess\,sup}_{x \in \Omega} p(x) < \infty$, for a.e. $x \in \Omega$.

Let us define by $\mathcal{M}(\Omega)$ the set of all measurable real functions defined on Ω . For any $p \in L_+^\infty(\Omega)$, we define the variable exponent Lebesgue space by

$$L^{p(x)}(\Omega) = \left\{ u \in \mathcal{M}(\Omega) : \rho_{p(x)}(u) = \int_\Omega |u(x)|^{p(x)} dx < \infty \right\}. \quad (1.3)$$

We define a norm, the so-called *Luxemburg norm*, on this space by the formula

$$\|u\|_{p(x)} = \inf \left\{ \delta > 0 : \rho_{p(x)}\left(\frac{u}{\delta}\right) \leq 1 \right\}, \quad (1.4)$$

and $(L^{p(x)}(\Omega), \|\cdot\|_{p(x)})$ becomes a Banach space.

The variable exponent Sobolev space $W^{1,p(x)}(\Omega)$ is

$$W^{1,p(x)}(\Omega) = \left\{ u \in L^{p(x)}(\Omega) : \frac{\partial u}{\partial x_i} \in L^{p(x)}(\Omega), i = 1, \dots, N \right\} \quad (1.5)$$

and we define on this space the norm

$$\|u\| = \|u\|_{p(x)} + \|\nabla u\|_{p(x)} \quad (1.6)$$

for all $u \in W^{1,p(x)}(\Omega)$. The space $W_0^{1,p(x)}(\Omega)$ is the closure of $C_0^\infty(\Omega)$ in $W^{1,p(x)}(\Omega)$.

Proposition 1.1 (see [1, 2]). *If $p \in L_+^\infty(\Omega)$, then the spaces $L^{p(x)}(\Omega)$, $W^{1,p(x)}(\Omega)$, and $W_0^{1,p(x)}(\Omega)$ are separable and reflexive Banach spaces.*

Proposition 1.2 (see [1, 2]). *If $u \in L^{p(x)}(\Omega)$ and $p_2 < \infty$, then we have*

- (i) $\|u\|_{p(x)} < 1 (= 1; > 1) \Leftrightarrow \rho_{p(x)}(u) < 1 (= 1; > 1)$,
- (ii) $\|u\|_{p(x)} > 1 \Rightarrow \|u\|_{p(x)}^{p_1} \leq \rho_{p(x)}(u) \leq \|u\|_{p(x)}^{p_2}$,
- (iii) $\|u\|_{p(x)} < 1 \Rightarrow \|u\|_{p(x)}^{p_2} \leq \rho_{p(x)}(u) \leq \|u\|_{p(x)}^{p_1}$,
- (iv) $\|u\|_{p(x)} = a > 0 \Leftrightarrow \rho_{p(x)}(u/a) = 1$.

Proposition 1.3 (see [3]). Assume that Ω is bounded and smooth. Denote by $C_+(\overline{\Omega}) = \{h \in C(\overline{\Omega}) : h_1 > 1\}$.

(i) Let $p, q \in C_+(\overline{\Omega})$. If

$$q(x) < p^*(x) = \begin{cases} \frac{Np(x)}{N-p(x)} & \text{if } p(x) < N, \\ +\infty & \text{if } p(x) \geq N, \end{cases} \quad (1.7)$$

then $(W_0^{1,p(x)}(\Omega), \|\cdot\|)$ is compactly imbedded in $L^{q(x)}(\Omega)$.

(ii) (Poincaré inequality, see [1, Theorem 2.7]). If $p \in C_+(\overline{\Omega})$, then there is a constant $C > 0$ such that

$$\|u\|_{p(x)} \leq C \|\nabla u\|_{p(x)}, \quad \forall u \in W_0^{1,p(x)}(\Omega). \quad (1.8)$$

Consequently, $\|u\|_{1,p(x)} = \|\nabla u\|_{p(x)}$ and $\|u\|$ are equivalent norms on $W_0^{1,p(x)}(\Omega)$. In what follows, $W_0^{1,p(x)}(\Omega)$, with $p \in C_+(\overline{\Omega})$, will be considered as endowed with the norm $\|u\|_{1,p(x)}$.

Lemma 1.4. Assume that $r \in L_+^\infty(\Omega)$ and $p \in C_+(\overline{\Omega})$. If $|u|^{r(x)} \in L^{p(x)}(\Omega)$, then we have

$$\min\left\{\|u\|_{r(x)p(x)}^{r_1}, \|u\|_{r(x)p(x)}^{r_2}\right\} \leq \left\| |u|^{r(x)} \right\|_{p(x)} \leq \max\left\{\|u\|_{r(x)p(x)}^{r_1}, \|u\|_{r(x)p(x)}^{r_2}\right\}. \quad (1.9)$$

Proof. By Proposition 1.2 (iv), we have

$$\begin{aligned} 1 &= \int_{\Omega} \left| \frac{|u|^{r(x)}}{\left\| |u|^{r(x)} \right\|_{p(x)}} \right|^{p(x)} dx \\ &= \int_{\Omega} \left| \frac{|u|}{\|u\|_{r(x)p(x)}} \right|^{r(x)p(x)} \frac{\|u\|_{r(x)p(x)}^{r(x)p(x)}}{\left\| |u|^{r(x)} \right\|_{p(x)}^{p(x)}} dx \\ &\leq \int_{\Omega} \left| \frac{|u|}{\|u\|_{r(x)p(x)}} \right|^{r(x)p(x)} \frac{\max\left\{\|u\|_{r(x)p(x)}^{r_1 p(x)}, \|u\|_{r(x)p(x)}^{r_2 p(x)}\right\}}{\left\| |u|^{r(x)} \right\|_{p(x)}^{p(x)}} dx. \end{aligned} \quad (1.10)$$

By the mean value theorem, there exists $\xi \in \overline{\Omega}$ such that

$$1 \leq \frac{\max\left\{\|u\|_{r(x)p(x)}^{r_1 p(\xi)}, \|u\|_{r(x)p(x)}^{r_2 p(\xi)}\right\}}{\left\| |u|^{r(x)} \right\|_{p(x)}^{p(\xi)}} \int_{\Omega} \left| \frac{|u|}{\|u\|_{r(x)p(x)}} \right|^{r(x)p(x)} dx \quad (1.11)$$

and we have

$$\left\| |u|^{r(x)} \right\|_{p(x)} \leq \max \left\{ \|u\|_{r(x)p(x)}^{r_1}, \|u\|_{r(x)p(x)}^{r_2} \right\}. \quad (1.12)$$

Similarly

$$1 \geq \frac{\min \left\{ \|u\|_{r(x)p(x)}^{r_1 p(\xi)}, \|u\|_{r(x)p(x)}^{r_2 p(\xi)} \right\}}{\left\| |u|^{r(x)} \right\|_{p(x)}^{p(\xi)}} dx, \quad (1.13)$$

$$\left\| |u|^{r(x)} \right\|_{p(x)} \geq \min \left\{ \|u\|_{r(x)p(x)}^{r_1}, \|u\|_{r(x)p(x)}^{r_2} \right\}.$$

□

Remark 1.5. If $r(x) = r = \text{const.}$, then

$$\left\| |u|^r \right\|_{p(x)} = \|u\|_{rp(x)}^r. \quad (1.14)$$

For simplicity of notation, we write

$$X = W_0^{1,p(x)}(\Omega), \quad X^* = \left(W_0^{1,p(x)}(\Omega) \right)^*, \quad Y = L^{q(x)}(\Omega), \quad Y^* = L^{q'(x)}(\Omega), \quad (1.15)$$

$$\|\cdot\|_X = \|\cdot\|_{1,p(x)}, \quad \|\cdot\|_Y = \|\cdot\|_{q(x)}.$$

In [4], a topological method, based on the fundamental properties of the Leray-Schauder degree, is used in proving the existence of a weak solution in X to the Dirichlet problem (P) that is an adaptation of that used by Dinca et al. for Dirichlet problems with classical p -Laplacian [5]. In this work, we use the nonlinear alternative of Leray-Schauder and give the existence of a solution and its localization. This method is used for finding solutions in Hölder spaces, while in [6], solutions are found in Sobolev spaces.

Let us recall some results borrowed from Dinca [4] about $p(x)$ -Laplacian and Nemytskii operator N_f . Firstly, since $q(x) < p(x) < p^*(x)$ for all $x \in \overline{\Omega}$, X is compactly embedded in Y . Denote by i the compact injection of X in Y and by $i^* : Y^* \rightarrow X^*$, $i^*v = v \circ i$ for all $v \in Y^*$, its adjoint.

Since the Caratheodory function f satisfies (CAR), the Nemytskii operator N_f generated by f , $(N_f u)(x) = f(x, u(x))$, is well defined from Y into Y^* , continuous, and bounded ([3, Proposition 2.2]). In order to prove that problem (P) has a weak solution in X it is sufficient to prove that the equation

$$-\Delta_{p(x)} u = (i^* N_f i) u \quad (1.16)$$

has a solution in X .

Indeed, if $u \in X$ satisfies (1.16) then, for all $v \in X$, one has

$$\langle -\Delta_{p(x)}u, v \rangle_{X, X^*} = \langle (i^* N_f i)u, v \rangle_{X, X^*} = \langle N_f(iu), iv \rangle_{Y, Y^*} \quad (1.17)$$

which rewrites as

$$\int_{\Omega} |\nabla u|^{p(x)} \nabla u \nabla v dx = \int_{\Omega} f v dx \quad (1.18)$$

and tells us that u is a weak solution in X to problem (P).

Since $-\Delta_{p(x)}$ is a homeomorphism of X onto X^* , (1.16) may be equivalently written as

$$u = (-\Delta_{p(x)})^{-1} (i^* N_f i)u. \quad (1.19)$$

Thus, proving that problem (P) has a weak solution in X reduces to proving that the compact operator

$$K = (-\Delta_{p(x)})^{-1} (i^* N_f i) : X \rightarrow X \quad (1.20)$$

has a fixed point.

Theorem 1.6 (Alternative of Leray-Schauder, [7]). *Let $B[0, R]$ denote the closed ball in a Banach space E , $\{u \in E : \|u\| \leq R\}$, and let $K : B[0, R] \rightarrow E$ be a compact operator. Then either*

- (i) *the equation $\lambda K u = u$ has a solution in $B[0, R]$ for $\lambda = 1$ or*
- (ii) *there exists an element $u \in E$ with $\|u\| = R$ satisfying $\lambda K u = u$ for some $\lambda \in (0, 1)$.*

2. Main Results

In this work, we present new existence and localization results for X -solutions to problem (P), under (CAR) condition on f . Our approach is based on regularity results for the solutions of Dirichlet problems and again on the nonlinear alternative of Leray-Schauder.

We start with an existence and localization principle for problem (P).

Theorem 2.1. *Assume that there is a constant $R > 0$, independent of $\lambda > 0$, with $\|u\|_X \neq R$ for any solution $u \in X$ to*

$$\begin{aligned} -\Delta_{p(x)}u &= \lambda f(x, u) \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega \end{aligned} \quad (P_\lambda)$$

and for each $\lambda \in (0, 1)$. Then the Dirichlet problem (P) has at least one solution $u \in X$ with $\|u\|_X \leq R$.

Proof. By [3, Theorem 3.1], $-\Delta_{p(x)}$ is a homeomorphism of X onto X^* . We will apply Theorem 2.1 to $E = X$ and to operator $K : X \rightarrow X$,

$$K u = (-\Delta_{p(x)})^{-1} (i^* N_f i)u, \quad (2.1)$$

where $i^*N_f i : X \rightarrow X^*$ is given by $(N_f u)(x) = f(x, u(x))$. Notice that, according to a well-known regularity result [4], the operator $(-\Delta_{p(x)})^{-1}$ from X to X is well defined, continuous, and order preserving. Consequently, K is a compact operator. On the other hand, it is clear that the fixed points of K are the solutions of problem (P). Now the conclusion follows from Theorem 1.6 since condition (ii) is excluded by hypothesis. \square

Theorem 2.2 immediately yields the following existence and localization result.

Theorem 2.2. *Let $\Omega \subset \mathbb{R}^N$, $N \geq 2$, be a smooth bounded domain and let $p, q \in C_+(\overline{\Omega})$ be such that $q(x) < p(x)$ for all $x \in \overline{\Omega}$. Assume that $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Caratheodory function which satisfies the growth condition (CAR).*

Suppose, in addition, that

$$C \|i^*\|_{Y^* \rightarrow X^*} \max \left\{ \|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1} \right\} < 1, \quad (2.2)$$

where C is the constant appearing in condition (CAR). Let $R \geq 1$ be a constant such that

$$R \geq \left(\frac{\|i^*\|_{Y^* \rightarrow X^*} \|a\|_{Y^*}}{1 - C \|i^*\|_{Y^* \rightarrow X^*} \max \left\{ \|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1} \right\}} \right)^{1/(p_1-1)}. \quad (2.3)$$

Then the Dirichlet problem (P) has at least a solution in X with $\|u\|_X \leq R$.

Proof. Let $u \in X$ be a solution of problem (P_λ) with $\|u\|_X = R \geq 1$, corresponding to some $\lambda \in (0, 1)$. Then by Propositions 1.2, 1.3, and Lemma 1.4, we obtain

$$\begin{aligned} \|u\|_X^{p_1} &\leq \int_{\Omega} |\nabla u|^{p(x)} dx = \lambda \langle (i^* N_f i) u, u \rangle_{X, X^*} = \lambda \langle N_f(iu), iu \rangle_{Y, Y^*} \\ &\leq \lambda \|i^*\|_{Y^* \rightarrow X^*} \|N_f(iu)\|_{Y^*} \|u\|_X \\ &\leq \lambda \|i^*\|_{Y^* \rightarrow X^*} \|u\|_X \left(\|a\|_{Y^*} + C \max \left\{ \|iu\|_Y^{q_1-1}, \|iu\|_Y^{q_2-1} \right\} \right) \\ &\leq \lambda \|i^*\|_{Y^* \rightarrow X^*} \|u\|_X \left(\|a\|_{Y^*} + C \|u\|_X^{q_2-1} \max \left\{ \|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1} \right\} \right) \\ &\leq \lambda \|i^*\|_{Y^* \rightarrow X^*} \|u\|_X \left(\|a\|_{Y^*} + C \|u\|_X^{p_1-1} \max \left\{ \|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1} \right\} \right). \end{aligned} \quad (2.4)$$

Therefore, we have

$$\|u\|_X^{p_1-1} \leq \frac{\lambda \|i^*\|_{Y^* \rightarrow X^*} \|a\|_{Y^*}}{1 - \lambda C \|i^*\|_{Y^* \rightarrow X^*} \max \left\{ \|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1} \right\}}. \quad (2.5)$$

Substituting $\|u\|_X = R$ in the above inequality, we obtain

$$R \leq \left(\frac{\lambda \|i^*\|_{Y^* \rightarrow X^*} \|a\|_{Y^*}}{1 - \lambda C \|i^*\|_{Y^* \rightarrow X^*} \max\{\|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1}\}} \right)^{1/(p_1-1)}, \quad (2.6)$$

which, taking into account (2.3) and $\lambda \in (0, 1)$, gives

$$\begin{aligned} R &\leq \lambda^{(1/p_1-1)} \left(\frac{\|i^*\|_{Y^* \rightarrow X^*} \|a\|_{Y^*}}{1 - C \lambda \|i^*\|_{Y^* \rightarrow X^*} \max\{\|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1}\}} \right)^{1/(p_1-1)} \\ &\leq \lambda^{(1/p_1-1)} \left(\frac{\|i^*\|_{Y^* \rightarrow X^*} \|a\|_{Y^*}}{1 - C \|i^*\|_{Y^* \rightarrow X^*} \max\{\|i\|_{X \rightarrow Y}^{q_1-1}, \|i\|_{X \rightarrow Y}^{q_2-1}\}} \right)^{1/(p_1-1)} \\ &\leq \lambda^{(1/p_1-1)} R < R, \end{aligned} \quad (2.7)$$

a contradiction. Theorem 2.1 applies. \square

Acknowledgment

The authors would like to thank the referees for their valuable and useful comments.

References

- [1] X. Fan and D. Zhao, "On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$," *Journal of Mathematical Analysis and Applications*, vol. 263, no. 2, pp. 424–446, 2001.
- [2] O. Kováčik and J. Rákosník, "On spaces $L^{p(x)}$ and $W^{k,p(x)}$," *Czechoslovak Mathematical Journal*, vol. 41(116), no. 4, pp. 592–618, 1991.
- [3] X.-L. Fan and Q.-H. Zhang, "Existence of solutions for $p(x)$ -Laplacian Dirichlet problem," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 52, no. 8, pp. 1843–1852, 2003.
- [4] G. Dinca, "A fixed point method for the $p(x)$ -Laplacian," *Comptes Rendus Mathématique*, vol. 347, no. 13-14, pp. 757–762, 2009.
- [5] G. Dinca, P. Jebelean, and J. Mawhin, "Variational and topological methods for Dirichlet problems with p -Laplacian," *Portugaliae Mathematica*, vol. 58, no. 3, pp. 339–378, 2001.
- [6] D. O'Regan and R. Precup, *Theorems of Leray-Schauder Type and Applications*, vol. 3 of *Series in Mathematical Analysis and Applications*, Gordon and Breach, Amsterdam, The Netherlands, 2001.
- [7] J. Dugundji and A. Granas, *Fixed Point Theory. I*, vol. 61 of *Monografie Matematyczne*, PWN-Polish Scientific, Warsaw, Poland, 1982.