

Research Article

Convergence Theorems of Modified Ishikawa Iterative Scheme for Two Nonexpansive Semigroups

Kriengsak Wattanawitton and Poom Kumam

Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), Bangmod, Thungkru, Bangkok 10140, Thailand

Correspondence should be addressed to Poom Kumam, poom.kum@kmutt.ac.th

Received 26 September 2009; Accepted 24 November 2009

Academic Editor: Tomonari Suzuki

Copyright © 2010 K. Wattanawitton and P. Kumam. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We prove convergence theorems of modified Ishikawa iterative sequence for two nonexpansive semigroups in Hilbert spaces by the two hybrid methods. Our results improve and extend the corresponding results announced by Saejung (2008) and some others.

1. Introduction

Let C be a subset of real Hilbert spaces H with the inner product $\langle \cdot, \cdot \rangle$ and the norm $\| \cdot \|$. $T : C \rightarrow C$ is called a nonexpansive mapping if

$$\|Tx - Ty\| \leq \|x - y\| \quad \forall x, y \in C. \quad (1.1)$$

We denote by $F(T)$ the set of fixed points of T , that is, $F(T) = \{x \in C : x = Tx\}$.

Let $\{T(t) : t \geq 0\}$ be a family of mappings from a subset C of H into itself. We call it a nonexpansive semigroup on C if the following conditions are satisfied:

- (i) $T(0)x = x$ for all $x \in C$;
- (ii) $T(s + t) = T(s)T(t)$ for all $s, t \geq 0$;
- (iii) for each $x \in C$ the mapping $t \mapsto T(t)x$ is continuous;
- (iv) $\|T(t)x - T(t)y\| \leq \|x - y\|$ for all $x, y \in C$ and $t \geq 0$.

The Mann's iterative algorithm was introduced by Mann [1] in 1953. This iterative process is now known as Mann's iterative process, which is defined as

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) T x_n, \quad n \geq 0, \quad (1.2)$$

where the initial guess x_0 is taken in C arbitrarily and the sequence $\{\alpha_n\}_{n=0}^{\infty}$ is in the interval $[0, 1]$.

In 1967, Halpern [2] first introduced the following iterative scheme:

$$\begin{aligned} x_0 &= u \in C \text{ chosen arbitrarily,} \\ x_{n+1} &= \alpha_n u + (1 - \alpha_n) T x_n, \end{aligned} \quad (1.3)$$

see also Browder [3]. He pointed out that the conditions $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$ are necessary in the sense that, if the iteration (1.3) converges to a fixed point of T , then these conditions must be satisfied.

On the other hand, in 2002, Suzuki [4] was the first to introduce the following implicit iteration process in Hilbert spaces:

$$x_n = \alpha_n u + (1 - \alpha_n) T(t_n)(x_n), \quad n \geq 1, \quad (1.4)$$

for the nonexpansive semigroup. In 2005, Xu [5] established a Banach space version of the sequence (1.4) of Suzuki [4].

In 2007, Chen and He [6] studied the viscosity approximation process for a nonexpansive semigroup and prove another strong convergence theorem for a nonexpansive semigroup in Banach spaces, which is defined by

$$x_{n+1} = \alpha_n f(x_n) + (1 - \alpha_n) T(t_n)x_n, \quad \forall n \in \mathbb{N}, \quad (1.5)$$

where $f : C \rightarrow C$ is a fixed contractive mapping.

Recently He and Chen [7] is proved a strong convergence theorem for nonexpansive semigroups in Hilbert spaces by hybrid method in the mathematical programming. Very recently, Saejung [8] proved a convergence theorem by the new iterative method introduced by Takahashi et al. [9] without Bochner integrals for a nonexpansive semigroup $\{T(t) : t \geq 0\}$ with $F := \bigcap_{t=0}^{\infty} F(T(t)) \neq \emptyset$ in Hilbert spaces:

$$\begin{aligned} x_0 &\in H \text{ taken arbitrary,} \\ C_1 &= C, \\ x_1 &= P_{C_1} x_0, \\ y_n &= \alpha_n x_n + (1 - \alpha_n) T(t_n)x_n, \\ C_{n+1} &= \{z \in C_n : \|y_n - z\| \leq \|x_n - z\|\}, \\ x_{n+1} &= P_{C_{n+1}}(x_0), \end{aligned} \quad (1.6)$$

where P_C denotes the metric projection from H onto a closed convex subset C of H .

In 1974, Ishikawa [10] introduced a new iterative scheme, which is defined recursively by

$$\begin{aligned}y_n &= \beta_n x_n + (1 - \beta_n) T x_n, \\x_{n+1} &= \alpha_n x_n + (1 - \alpha_n) T y_n,\end{aligned}\tag{1.7}$$

where the initial guess x_0 is taken in C arbitrarily and the sequences $\{\alpha_n\}$ and $\{\beta_n\}$ are in the interval $[0, 1]$.

In this paper, motivated by the iterative sequences (1.6) given by Saejung in [8] and Ishikawa [10], we introduce the modified Ishikawa iterative scheme for two nonexpansive semigroups in Hilbert spaces. Further, we obtain strong convergence theorems by using the hybrid methods. This result extends and improves the result of Saejung [8] and some others.

2. Preliminaries

This section collects some lemmas which will be used in the proofs for the main results in the next section.

It is known that every Hilbert space H satisfies the Opial's condition [11], that is,

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|, \quad \forall y \in X, y \neq x.\tag{2.1}$$

Recall that the metric (nearest point) projection P_C from a Hilbert space H to a closed convex subset C of H is defined as follows. Given $x \in H$, $P_C x$ is the only point in C with the property

$$\|x - P_C x\| = \inf\{\|x - y\| : y \in C\}.\tag{2.2}$$

$P_C x$ is characterized as follows.

Lemma 2.1. *Let H be a real Hilbert space, C a closed convex subset of H . Given $x \in H$ and $y \in C$. Then $y = P_C x$ if and only if there holds the inequality*

$$\langle x - y, y - z \rangle \geq 0, \quad \forall z \in C.\tag{2.3}$$

Lemma 2.2. *There holds the identity in a Hilbert space H*

$$\|\lambda x + (1 - \lambda)y\|^2 = \lambda\|x\|^2 + (1 - \lambda)\|y\|^2 - \lambda(1 - \lambda)\|x - y\|^2\tag{2.4}$$

for all $x, y \in H$ and $\lambda \in [0, 1]$.

Lemma 2.3 (see [12, Lemma 1]). *Let $\{t_n\}$ be a real sequence and let τ be a real number such that $\liminf_n t_n \leq \tau \leq \limsup_n t_n$. Suppose that either of the following holds:*

$$(i) \limsup_n (t_{n+1} - t_n) \leq 0 \text{ or}$$

$$(ii) \liminf_n (t_{n+1} - t_n) \geq 0,$$

then τ is a cluster point of $\{t_n\}$. Moreover, for $\varepsilon > 0$, $k, m \in \mathbb{N}$, there exists $m_0 \geq m$ such that $|t_j - \tau| < \varepsilon$ for every integer j with $m_0 \leq j \leq m_0 + k$.

3. Main Results

3.1. The Shrinking Projection Method

In this section, we prove strong convergence of an iterative sequence generated by the shrinking hybrid projection method in mathematical programming.

Theorem 3.1. *Let C be a closed convex subset of a real Hilbert space H . Let $\{T(t) : t \geq 0\}$ and $\{S(t) : t \geq 0\}$ be nonexpansive semigroups on C with a nonempty common fixed point set F , that is, $F := (\bigcap_{t=0}^{\infty} F(T(t))) \cap (\bigcap_{t=0}^{\infty} F(S(t))) \neq \emptyset$. Let $\{\alpha_n\} \subset [0, a] \subset [0, 1)$, $\{\beta_n\} \subset [b, c] \subset (0, 1)$ and $\{t_n\}$ be the sequences such that $\liminf_{n \rightarrow \infty} t_n = 0$, $\limsup_{n \rightarrow \infty} t_n > 0$, and $\lim_{n \rightarrow \infty} (t_{n+1} - t_n) = 0$. Suppose that $\{x_n\}$ is a sequence generated by the following iterative scheme:*

$$\begin{aligned} x_0 &\in H \text{ taken arbitrary,} \\ C_1 &= C, \\ x_1 &= P_{C_1}(x_0), \\ z_n &= \beta_n x_n + (1 - \beta_n) T(t_n) x_n, \\ y_n &= \alpha_n x_n + (1 - \alpha_n) S(t_n) z_n, \\ C_{n+1} &= \{u \in C_n : \|y_n - u\| \leq \|x_n - u\|\}, \\ x_{n+1} &= P_{C_{n+1}}(x_0), \end{aligned} \tag{3.1}$$

then $\{x_n\}$ converges strongly to $P_F(x_0)$.

Proof. We first show that C_{n+1} is closed and convex for each $n \geq 0$. From the definition of C_{n+1} it is obvious that C_{n+1} is closed for each $n \geq 0$. We show that C_{n+1} is convex for any $n \geq 0$. Since

$$\|y_n - u\| \leq \|x_n - u\| \iff 2\langle x_n - y_n, u \rangle \leq \|x_n\|^2 - \|y_n\|^2, \tag{3.2}$$

and hence C_{n+1} is convex. Next we show that $F \subset C_{n+1}$ for all $n \geq 0$. Let $p \in F$, then we have

$$\begin{aligned} \|z_n - p\| &= \|\beta_n x_n + (1 - \beta_n)T(t_n)x_n - p\| \\ &\leq \beta_n \|x_n - p\| + (1 - \beta_n)\|T(t_n)x_n - p\| \\ &\leq \beta_n \|x_n - p\| + (1 - \beta_n)\|x_n - p\| \\ &\leq \|x_n - p\|, \end{aligned} \quad (3.3)$$

$$\begin{aligned} \|y_n - p\| &= \|\alpha_n x_n + (1 - \alpha_n)S(t_n)z_n - p\| \\ &\leq \alpha_n \|x_n - p\| + (1 - \alpha_n)\|S(t_n)z_n - p\| \\ &\leq \alpha_n \|x_n - p\| + (1 - \alpha_n)\|z_n - p\|. \end{aligned} \quad (3.4)$$

Substituting (3.3) into (3.4), we have

$$\|y_n - p\| \leq \|x_n - p\|. \quad (3.5)$$

This means that $p \in C_{n+1}$ for all $n \geq 0$. Thus, $\{x_n\}$ is well defined. Since $x_n = P_{C_n}(x_0)$ and $x_{n+1} \in C_{n+1} \subset C_n$, we get

$$\langle x_0 - x_n, x_n - x_{n+1} \rangle \geq 0 \quad \forall n \in \mathbb{N}. \quad (3.6)$$

Consequently,

$$\begin{aligned} 0 &\leq \langle x_0 - x_n, x_n - x_{n+1} \rangle \\ &= \langle x_0 - x_n, x_n - x_0 + x_0 - x_{n+1} \rangle \\ &= -\langle x_n - x_0, x_n - x_0 \rangle + \langle x_0 - x_n, x_0 - x_{n+1} \rangle \\ &\leq -\|x_n - x_0\|^2 + \|x_0 - x_n\|\|x_0 - x_{n+1}\|, \end{aligned} \quad (3.7)$$

for $n \in \mathbb{N}$. This implies that

$$\|x_0 - x_n\| \leq \|x_0 - x_{n+1}\| \quad \forall n \in \mathbb{N}. \quad (3.8)$$

Therefore, $\{\|x_0 - x_n\|\}$ is nondecreasing. From $x_n = P_{C_n}(x_0)$, we also have $\langle x_0 - x_n, x_n - p \rangle \geq 0$, for all $p \in C_n$.

Since $F \subseteq C_n$, we get

$$\langle x_0 - x_n, x_n - p \rangle \geq 0 \quad \forall p \in F. \quad (3.9)$$

Thus, for $p \in F$, we obtain

$$\begin{aligned}
0 &\leq \langle x_0 - x_n, x_n - p \rangle \\
&= -\langle x_n - x_0, x_n - x_0 \rangle + \langle x_0 - x_n, x_0 - p \rangle \\
&\leq -\|x_n - x_0\|^2 + \|x_0 - x_n\| \|x_0 - p\|.
\end{aligned} \tag{3.10}$$

Thus, $\|x_n - x_0\| \leq \|x_0 - p\|$, for all $p \in F$ and $n \in \mathbb{N}$. Then $\lim_{n \rightarrow \infty} \|x_n - x_0\|$ exists and $\{x_n\}$ is bounded.

Next, we show that $\|x_{n+1} - x_n\| \rightarrow 0$ as $n \rightarrow \infty$. From (3.6) we have

$$\begin{aligned}
\|x_n - x_{n+1}\|^2 &= \|x_n - x_0 + x_0 - x_{n+1}\|^2 \\
&= \|x_n - x_0\|^2 + 2\langle x_n - x_0, x_0 - x_{n+1} \rangle + \|x_0 - x_{n+1}\|^2 \\
&= \|x_n - x_0\|^2 + 2\langle x_n - x_0, x_0 - x_n + x_n - x_{n+1} \rangle + \|x_0 - x_{n+1}\|^2 \\
&= \|x_n - x_0\|^2 - 2\langle x_0 - x_n, x_0 - x_n \rangle - 2\langle x_0 - x_n, x_n - x_{n+1} \rangle + \|x_0 - x_{n+1}\|^2 \\
&\leq \|x_n - x_0\|^2 - 2\|x_n - x_0\|^2 + \|x_0 - x_{n+1}\|^2 \\
&= -\|x_n - x_0\|^2 + \|x_0 - x_{n+1}\|^2.
\end{aligned} \tag{3.11}$$

Since $\lim_{n \rightarrow \infty} \|x_n - x_0\|$ exists, then

$$\lim_{n \rightarrow \infty} \|x_n - x_{n+1}\| = 0. \tag{3.12}$$

Further, as in the proof of [8, page 3], we have $\{x_n\}$ which is a Cauchy sequence. So, we have $x_n \rightarrow z$. By definition of y_n , we have

$$\|y_n - x_n\| = (1 - \alpha_n) \|S(t_n)z_n - x_n\|. \tag{3.13}$$

Since $x_{n+1} \in C_{n+1}$ and (3.12), we obtain

$$\begin{aligned}
\|S(t_n)z_n - x_n\| &= \frac{1}{1 - \alpha_n} \|y_n - x_n\| \\
&\leq \frac{1}{1 - \alpha_n} (\|y_n - x_{n+1}\| + \|x_{n+1} - x_n\|) \\
&\leq \frac{1}{1 - \alpha_n} (\|x_n - x_{n+1}\| + \|x_{n+1} - x_n\|) \\
&\leq \frac{2}{1 - \alpha_n} \|x_n - x_{n+1}\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.
\end{aligned} \tag{3.14}$$

We now show that $\|T(t_n)x_n - x_n\| \rightarrow 0$.

For $p \in F$, we have $\|x_n - p\| \leq \|x_n - S(t_n)z_n\| \leq \|S(t_n)z_n - p\|$. This implies that $0 \leq \|x_n - p\| - \|z_n - p\| \leq \|x_n - S(t_n)z_n\| \rightarrow 0$ and hence $\|x_n - p\|^2 - \|z_n - p\|^2 \rightarrow 0$. Moreover, since

$$\|z_n - p\|^2 = \beta_n \|x_n - p\|^2 + (1 - \beta_n) \|T(t_n)x_n - p\|^2 - \beta_n(1 - \beta_n) \|x_n - T(t_n)x_n\|^2, \quad (3.15)$$

we have

$$\begin{aligned} \beta_n \|x_n - T(t_n)x_n\|^2 &\leq \beta_n(1 - \beta_n) \|x_n - T(t_n)x_n\|^2 \\ &\leq \beta_n \|x_n - p\|^2 + (1 - \beta_n) \|T(t_n)x_n - p\|^2 - \|z_n - p\|^2 \\ &\leq \|x_n - p\|^2 - \|z_n - p\|^2 \rightarrow 0. \end{aligned} \quad (3.16)$$

And since $S(t_n)$ is a nonexpansive mapping, we obtain

$$\begin{aligned} \|x_n - S(t_n)x_n\| &\leq \|x_n - S(t_n)z_n\| + \|S(t_n)z_n - S(t_n)x_n\|, \\ &\leq \|x_n - S(t_n)z_n\| + \|z_n - x_n\|. \end{aligned} \quad (3.17)$$

Since $\|z_n - x_n\| = (1 - \beta_n) \|T(t_n)x_n - x_n\| \rightarrow 0$ and $\|x_n - S(t_n)z_n\| \rightarrow 0$, we obtain

$$\lim_{n \rightarrow \infty} \|x_n - S(t_n)x_n\| = 0. \quad (3.18)$$

As in the proof of [12, Theorem 4], by Lemma 2.3, we can choose a sequence $\{t_{n_k}\}$ of positive real numbers such that

$$t_{n_k} \rightarrow 0, \quad \frac{1}{t_{n_k}} \|x_{n_k} - T(t_{n_k})x_{n_k}\| \rightarrow 0, \quad \text{as } k \rightarrow \infty. \quad (3.19)$$

In similar way, we also have

$$t_{n_k} \rightarrow 0, \quad \frac{1}{t_{n_k}} \|x_{n_k} - S(t_{n_k})x_{n_k}\| \rightarrow 0, \quad \text{as } k \rightarrow \infty. \quad (3.20)$$

Next, we show that $z \in F$. To see this, we fix $t > 0$,

$$\begin{aligned}
\|x_{n_k} - T(t)z\| &\leq \sum_{j=0}^{\lfloor t/t_{n_k} \rfloor - 1} \|T(jt_{n_k})x_{n_k} - T((j+1)t_{n_k})x_{n_k}\| \\
&\quad + \left\| T\left(\left[\frac{t}{t_{n_k}}\right]t_{n_k}\right)x_{n_k} - T\left(\left[\frac{t}{t_{n_k}}\right]t_{n_k}\right)z \right\| + \left\| T\left(\left[\frac{t}{t_{n_k}}\right]t_{n_k}\right)z - T(t)z \right\| \\
&\leq \left[\frac{t}{t_{n_k}}\right] \|x_{n_k} - T(t_{n_k})x_{n_k}\| + \|x_{n_k} - z\| + \left\| T\left(t - \left[\frac{t}{t_{n_k}}\right]t_{n_k}\right)z - z \right\| \\
&\leq \frac{t}{t_{n_k}} \|x_{n_k} - T(t_{n_k})x_{n_k}\| + \|x_{n_k} - z\| + \sup\{\|T(s)z - z\| : 0 \leq s \leq t_{n_k}\}.
\end{aligned} \tag{3.21}$$

As $x_{n_k} \rightarrow z$ and (3.19), we obtain $x_{n_k} \rightarrow T(t)z$ and so $T(t)z = z$. Similarly, we have $S(t)z = z$. Thus $z \in F$.

Finally, we show that $z = P_F(x_0)$. Since $F \subset C_{n+1}$ and $x_{n+1} = P_{C_{n+1}}(x_0)$,

$$\|x_{n+1} - x_0\| \leq \|q - x_0\| \quad \forall n \in \mathbb{N}, q \in F. \tag{3.22}$$

But $x_n \rightarrow z$ as $n \rightarrow \infty$, we have

$$\|z - x_0\| \leq \|q - x_0\| \quad \forall q \in F. \tag{3.23}$$

Hence $z = P_F(x_0)$ as required. This completes the proof. \square

Corollary 3.2. *Let C be a closed convex subset of a real Hilbert space H . Let $\{T(t) : t \geq 0\}$ be nonexpansive semigroups on C with a nonempty common fixed point set F , that is, $F := \bigcap_{t=0}^{\infty} F(T(t)) \neq \emptyset$. Let $\{\alpha_n\} \subset [0, a] \subset [0, 1)$, $\{\beta_n\} \subset [b, c] \subset (0, 1)$ and $\{t_n\}$ be the sequences such that $\liminf_{n \rightarrow \infty} t_n = 0$, $\limsup_{n \rightarrow \infty} t_n > 0$, and $\lim_{n \rightarrow \infty} (t_{n+1} - t_n) = 0$. Suppose that $\{x_n\}$ is a sequence iteratively generated by the following iterative scheme:*

$$\begin{aligned}
x_0 &\in H \text{ taken arbitrary,} \\
C_1 &= C, \\
x_1 &= P_{C_1}(x_0), \\
y_n &= \alpha_n x_n + (1 - \alpha_n)T(t_n)z_n, \\
z_n &= \beta_n x_n + (1 - \beta_n)T(t_n)x_n, \\
C_{n+1} &= \{u \in C_n : \|y_n - u\| \leq \|x_n - u\|\}, \\
x_{n+1} &= P_{C_{n+1}}(x_0),
\end{aligned} \tag{3.24}$$

then $\{x_n\}$ converges strongly to $P_F(x_0)$.

Proof. Putting $S(t_n) = T(t_n)$, in Theorem 3.1, we obtain the conclusion immediately. \square

Corollary 3.3 (see [8, Theorem 2.1]). *Let C be a closed convex subset of a real Hilbert space H . Let $\{T(t) : t \geq 0\}$ be a nonexpansive semigroups on C with a nonempty common fixed point set F , that is, $F := \bigcap_{t=0}^{\infty} F(T(t)) \neq \emptyset$. Let $\{\alpha_n\} \subset [0, a] \subset [0, 1)$ and $\{t_n\}$ be the sequences such that $\liminf_{n \rightarrow \infty} t_n = 0$, $\limsup_{n \rightarrow \infty} t_n > 0$, and $\lim_{n \rightarrow \infty} (t_{n+1} - t_n) = 0$. Suppose that $\{x_n\}$ is a sequence iteratively generated by the following iterative scheme:*

$$\begin{aligned} x_0 &\in H \text{ taken arbitrary,} \\ C_1 &= C, \\ x_1 &= P_{C_1}(x_0), \\ z_n &= \alpha_n x_n + (1 - \alpha_n)T(t_n)x_n, \\ C_{n+1} &= \{u \in C_n : \|y_n - u\| \leq \|x_n - u\|\}, \\ x_{n+1} &= P_{C_{n+1}}(x_0), \end{aligned} \tag{3.25}$$

then $x_n \rightarrow P_F(x_0)$.

Proof. If $S(t_n) = T(t_n)$ for all $n \in \mathbb{N}$ and $T(t) = I$ for every $t > 0$ in Theorem 3.1 then (3.1) reduced to (3.25). By using Theorem 3.1, we get the following conclusion. \square

3.2. The CQ Hybrid Method

In this section, we consider the modified Ishikawa iterative scheme computing by the CQ hybrid method [13–15]. We use the same idea as Saejung's Theorem 2.2 in [8] and our Theorem 3.1 to obtain the following result and the proof is omitted.

Theorem 3.4. *Let C be a closed convex subset of a real Hilbert space H . Let $\{T(t) : t \geq 0\}$ and $\{S(t) : t \geq 0\}$ be nonexpansive semigroups on C with a nonempty common fixed point set F , that is, $F := (\bigcap_{t=0}^{\infty} F(T(t))) \cap (\bigcap_{t=0}^{\infty} F(S(t))) \neq \emptyset$. Let $\{\alpha_n\} \subset [0, a] \subset [0, 1)$, $\{\beta_n\} \subset [b, c] \subset (0, 1)$ and $\{t_n\}$ be the sequences such that $\liminf_{n \rightarrow \infty} t_n = 0$, $\limsup_{n \rightarrow \infty} t_n > 0$, and $\lim_{n \rightarrow \infty} (t_{n+1} - t_n) = 0$. Suppose that $\{x_n\}$ is a sequence generated by the following iterative scheme:*

$$\begin{aligned} x_0 &\in H \text{ taken arbitrary,} \\ y_n &= \alpha_n x_n + (1 - \alpha_n)S(t_n)z_n, \\ z_n &= \beta_n x_n + (1 - \beta_n)T(t_n)x_n, \\ C_n &= \{u \in C : \|y_n - u\| \leq \|x_n - u\|\}, \\ Q_n &= \{u \in C : \langle x_n - x_0, u - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n}(x_0), \end{aligned} \tag{3.26}$$

then $\{x_n\}$ converges strongly to $P_F(x_0)$.

Proof. First, we show that both C_n and Q_n are closed and convex, and $C_n \cap Q_n \neq \emptyset$ for all $n \in \mathbb{N} \cup \{0\}$. It follows easily from the definition that C_n and Q_n are just intersection of C and the half-spaces see also [9]. As in the proof of the preceding theorem, we have $F \subset C_n$ for all $n \in \mathbb{N} \cup \{0\}$. Clearly, $F \subset C = Q_0$. Suppose that $F \subset Q_k$ for some $k \in \mathbb{N} \cup \{0\}$,

we have $p \in C_k \cap Q_k$. In particular, $\langle x_{k+1} - x_0, p - x_{k+1} \rangle \geq 0$, that is, $p \in Q_{k+1}$. It follows from the induction that $F \subset Q_n$ for all $n \in \mathbb{N} \cup \{0\}$. This proves the claim.

Next, we show that $\|x_n - T(t_n)x_n\| \rightarrow 0$, and $\|x_n - S(t_n)x_n\| \rightarrow 0$.

We first claim that $\|x_{n+1} - x_n\| \rightarrow 0$. Indeed, as $x_{n+1} \in Q_n$ and $x_n = P_{Q_n}(x_0)$,

$$\|x_n - x_0\| \leq \|x_{n+1} - x_0\| \quad \forall n \in \mathbb{N}. \quad (3.27)$$

For fixed $z \in F$. It follows from $F \subset Q_n$ for all $n \in \mathbb{N}$ that

$$\|x_n - x_0\| \leq \|z - x_0\| \quad \forall n \in \mathbb{N}. \quad (3.28)$$

This implies that sequence $\{x_n\}$ is bounded and

$$\lim_{n \rightarrow \infty} \|x_n - x_0\| \text{ exists.} \quad (3.29)$$

Notice that

$$\langle x_{n+1} - x_n, x_n - x_0 \rangle \geq 0. \quad (3.30)$$

This implies that

$$\begin{aligned} \|x_{n+1} - x_n\|^2 &= \|x_{n+1} - x_0\|^2 - 2\langle x_{n+1} - x_n, x_n - x_0 \rangle - \|x_0 - x_n\|^2 \\ &\leq \|x_{n+1} - x_0\|^2 - \|x_n - x_0\|^2 \rightarrow 0. \end{aligned} \quad (3.31)$$

By using the same argument of Saejung [8, Theorem 2.2, page 6] and in the proof of Theorem 3.1, we have $\|T(t_n)x_n - x_n\| \rightarrow 0$ and $\|S(t_n)x_n - x_n\| \rightarrow 0$. And we can choose a subsequence $\{n_k\}$ of $\{n\}$ such that $x_{n_k} \rightarrow z \in C$, $t_{n_k} \rightarrow 0$, $(1/t_{n_k})\|x_{n_k} - T(t_{n_k})x_{n_k}\| \rightarrow 0$ and $(1/t_{n_k})\|x_{n_k} - S(t_{n_k})x_{n_k}\| \rightarrow 0$ as $k \rightarrow \infty$.

From (3.21), we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} \|x_{n_k} - T(t)z\| &\leq \limsup_{k \rightarrow \infty} \|x_{n_k} - z\|, \\ \limsup_{k \rightarrow \infty} \|x_{n_k} - S(t)z\| &\leq \limsup_{k \rightarrow \infty} \|x_{n_k} - z\|. \end{aligned} \quad (3.32)$$

By the Opial's condition of H , we have $z = T(t)z$ and $z = S(t)z$ for all $t > 0$, that is, $z \in F$.

We note that

$$\|x_0 - P_F(x_0)\| \leq \|x_0 - z\| \leq \liminf_{k \rightarrow \infty} \|x_0 - x_{n_k}\| \leq \limsup_{k \rightarrow \infty} \|x_0 - x_{n_k}\| \leq \|x_0 - P_F(x_0)\|. \quad (3.33)$$

This implies that

$$\lim_{k \rightarrow \infty} \|x_0 - x_{n_k}\| = \|x_0 - P_F(x_0)\| = \|x_0 - z\|. \quad (3.34)$$

Therefore,

$$x_{n_k} \longrightarrow P_F(x_0) = z, \quad \text{as } k \longrightarrow \infty. \quad (3.35)$$

Hence the whole sequence must converge to $P_F(x_0) = z$, as required. This completes the proof. \square

Corollary 3.5. *Let C be a closed convex subset of a real Hilbert space H . Let $\{T(t) : t \geq 0\}$ be nonexpansive semigroups on C with a nonempty common fixed point set F , that is, $F := \bigcap_{t=0}^{\infty} F(T(t)) \neq \emptyset$. Let $\{\alpha_n\} \subset [0, a] \subset [0, 1)$, $\{\beta_n\} \subset [b, c] \subset (0, 1)$ and $\{t_n\}$ be the sequences such that $\liminf_{n \rightarrow \infty} t_n = 0$, $\limsup_{n \rightarrow \infty} t_n > 0$, and $\lim_{n \rightarrow \infty} (t_{n+1} - t_n) = 0$. Suppose that $\{x_n\}$ is a sequence iteratively generated by the following iterative scheme:*

$$\begin{aligned} x_0 &\in H \text{ taken arbitrary,} \\ y_n &= \alpha_n x_n + (1 - \alpha_n) T(t_n) z_n, \\ z_n &= \beta_n x_n + (1 - \beta_n) T(t_n) x_n, \\ C_n &= \{u \in C : \|y_n - u\| \leq \|x_n - u\|\}, \\ Q_n &= \{u \in C : \langle x_n - x_0, u - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n}(x_0), \end{aligned} \quad (3.36)$$

then $\{x_n\}$ converges strongly to $P_F(x_0)$.

Proof. If $S(t_n) = T(t_n)$ for all $n \in \mathbb{N} \cup \{0\}$, in Theorem 3.4 then (3.26) reduced to (3.36). So, we obtain the result immediately. \square

We also deduce the following corollary.

Corollary 3.6 (see [8, Theorem 2.2]). *Let C be a closed convex subset of a real Hilbert space H . Let $\{T(t) : t \geq 0\}$ be a nonexpansive semigroups on C with a nonempty common fixed point set F , that is, $F := \bigcap_{t=0}^{\infty} F(T(t)) \neq \emptyset$. Let $\{\alpha_n\} \subset [0, a] \subset [0, 1)$ and $\{t_n\}$ be the sequences such that $\liminf_{n \rightarrow \infty} t_n = 0$, $\limsup_{n \rightarrow \infty} t_n > 0$ and $\lim_{n \rightarrow \infty} (t_{n+1} - t_n) = 0$. Suppose that $\{x_n\}$ is a sequence iteratively generated by the following iterative scheme:*

$$\begin{aligned} x_0 &\in H \text{ taken arbitrary,} \\ z_n &= \alpha_n x_n + (1 - \alpha_n) T(t_n) x_n, \\ C_n &= \{u \in C : \|y_n - u\| \leq \|x_n - u\|\}, \\ Q_n &= \{u \in C : \langle x_n - x_0, u - x_n \rangle \geq 0\}, \\ x_{n+1} &= P_{C_n \cap Q_n}(x_0), \end{aligned} \quad (3.37)$$

then $x_n \rightarrow P_F(x_0)$.

Acknowledgments

The authors would like to thank the editors and the anonymous referees for their valuable suggestions which help to improve this paper. This research was supported by the Computational Science and Engineering Research Cluster, King Mongkut's University of Technology Thonburi (KMUTT) (National Research Universities under CSEC Project no. E01008).

References

- [1] W. R. Mann, "Mean value methods in iteration," *Proceedings of the American Mathematical Society*, vol. 4, pp. 506–510, 1953.
- [2] B. Halpern, "Fixed points of nonexpanding maps," *Bulletin of the American Mathematical Society*, vol. 73, pp. 957–961, 1967.
- [3] F. E. Browder, "Fixed-point theorems for noncompact mappings in Hilbert space," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 53, pp. 1272–1276, 1965.
- [4] T. Suzuki, "On strong convergence to common fixed points of nonexpansive semigroups in Hilbert spaces," *Proceedings of the American Mathematical Society*, vol. 131, no. 7, pp. 2133–2136, 2002.
- [5] H.-K. Xu, "A strong convergence theorem for contraction semigroups in Banach spaces," *Bulletin of the Australian Mathematical Society*, vol. 72, no. 3, pp. 371–379, 2005.
- [6] R. Chen and H. He, "Viscosity approximation of common fixed points of nonexpansive semigroups in Banach space," *Applied Mathematics Letters*, vol. 20, no. 7, pp. 751–757, 2007.
- [7] H. He and R. Chen, "Strong convergence theorems of the CQ method for nonexpansive semigroups," *Fixed Point Theory and Applications*, vol. 2007, Article ID 59735, 8 pages, 2007.
- [8] S. Saejung, "Strong convergence theorems for nonexpansive semigroups without Bochner integrals," *Fixed Point Theory and Applications*, vol. 2008, Article ID 745010, 7 pages, 2008.
- [9] W. Takahashi, Y. Takeuchi, and R. Kubota, "Strong convergence theorems by hybrid methods for families of nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 276–286, 2008.
- [10] S. Ishikawa, "Fixed points by a new iteration method," *Proceedings of the American Mathematical Society*, vol. 44, pp. 147–150, 1974.
- [11] Z. Opial, "Weak convergence of the sequence of successive approximations for nonexpansive mappings," *Bulletin of the American Mathematical Society*, vol. 73, pp. 591–597, 1967.
- [12] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [13] Y. Haugazeau, *Sur les Inéquations variationnelles et la minimisation de fonctionnelles convexes*, Ph.D. thesis, Université Paris, Paris, France, 1968.
- [14] K. Nakajo and W. Takahashi, "Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups," *Journal of Mathematical Analysis and Applications*, vol. 279, no. 2, pp. 372–379, 2003.
- [15] K. Nakajo, K. Shimoji, and W. Takahashi, "Strong convergence theorems by the hybrid method for families of nonexpansive mappings in Hilbert spaces," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 339–360, 2006.