

Research Article

Fixed Points and Random Stability of a Generalized Apollonius Type Quadratic Functional Equation

Min June Kim,¹ Seung Won Shin,¹ Dohyeong Ki,¹
Jaewon Chang,¹ and Ji-Hye Kim²

¹ Mathematics Branch, Seoul Science High School, Seoul 110-530, Republic of Korea

² Department of Mathematics, Research Institute for Natural Sciences, Hanyang University,
Seoul 133-791, Republic of Korea

Correspondence should be addressed to Ji-Hye Kim, saharin@hanyang.ac.kr

Received 23 December 2010; Revised 28 February 2011; Accepted 28 February 2011

Academic Editor: Yeol J. Cho

Copyright © 2011 Min June Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Using the fixed-point method, we prove the generalized Hyers-Ulam stability of a generalized Apollonius type quadratic functional equation in random Banach spaces.

1. Introduction

The stability problem of functional equations was originated from a question of Ulam [1] concerning the stability of group homomorphisms. Hyers [2] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' theorem was generalized by Aoki [3] for additive mappings and by Th. M. Rassias [4] for linear mappings by considering an unbounded Cauchy difference. The paper of Th. M. Rassias [4] has provided a lot of influence in the development of what we call *generalized Hyers-Ulam stability* of functional equations. A generalization of the Th. M. Rassias theorem was obtained by Găvruta [5] by replacing the unbounded Cauchy difference by a general control function in the spirit of the Th. M. Rassias' approach.

On the other hand, in 1982–1998, J. M. Rassias generalized the Hyers' stability result by presenting a weaker condition controlled by a product of different powers of norms.

Theorem 1.1 (see [6–12]). *Assume that there exist constants $\Theta \geq 0$ and $p_1, p_2 \in \mathbb{R}$ such that $p = p_1 + p_2 \neq 1$, and $f : E \rightarrow E'$ is a mapping from a normed space E into a Banach space E' , such that the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon \|x\|^{p_1} \|y\|^{p_2}, \quad (1.1)$$

for all $x, y \in E$, then there exists a unique additive mapping $T : E \rightarrow E'$, such that

$$\|f(x) - L(x)\| \leq \frac{\Theta}{2 - 2^p} \|x\|^p, \quad (1.2)$$

for all $x \in E$.

The control function $\|x\|^p \cdot \|y\|^q + \|x\|^{p+q} + \|y\|^{p+q}$ was introduced by Ravi et al. [13] and was used in several papers (see [14–19]).

The functional equation $f(x + y) + f(x - y) = 2f(x) + 2f(y)$ is called a *quadratic functional equation*. In particular, every solution of the quadratic functional equation is said to be a *quadratic mapping*. The generalized Hyers-Ulam stability of the quadratic functional equation was proved by Skof [20] for mappings $f : X \rightarrow Y$, where X is a normed space and Y is a Banach space. Cholewa [21] noticed that the theorem of Skof is still true if the relevant domain X is replaced by an Abelian group. Czerwik [22] proved the generalized Hyers-Ulam stability of the quadratic functional equation. The stability problems of several functional equations have been extensively investigated by a number of authors, and there are many interesting results concerning this problem (see [23–44]).

In [45], Park and Th. M. Rassias defined and investigated the following generalized Apollonius type quadratic functional equation:

$$\begin{aligned} & Q\left(\left(\sum_{i=1}^n z_i\right) - \left(\sum_{i=1}^n x_i\right)\right) + Q\left(\left(\sum_{i=1}^n z_i\right) - \left(\sum_{i=1}^n y_i\right)\right) \\ &= \frac{1}{2}Q\left(\left(\sum_{i=1}^n x_i\right) - \left(\sum_{i=1}^n y_i\right)\right) + 2Q\left(\left(\sum_{i=1}^n z_i\right) - \frac{(\sum_{i=1}^n x_i) - (\sum_{i=1}^n y_i)}{2}\right) \end{aligned} \quad (1.3)$$

in Banach spaces.

2. Preliminaries

We define the notion of a random normed space, which goes back to Šerstnev et al. (see, e.g., [46, 47]).

In the sequel, we adopt the usual terminology, notations, and conventions of the theory of random normed spaces, as in [47, 48]. Throughout this paper, let Δ^+ be the space of distribution functions, that is,

$$\begin{aligned} \Delta^+ := \{F : \mathbb{R} \cup \{-\infty, \infty\} \rightarrow [0, 1] : F \text{ is leftcontinuous,} \\ \text{nondecreasing on } \mathbb{R}, F(0) = 0 \text{ and } F(+\infty) = 1\}, \end{aligned} \quad (2.1)$$

and the subset $D^+ \subseteq \Delta^+$ is the set $D^+ = \{F \in \Delta^+ : l^-F(+\infty) = 1\}$, where $l^-f(x)$ denotes the left limit of the function f at the point x . The space Δ^+ is partially ordered by the usual pointwise

ordering of functions, that is, $F \leq G$ if and only if $F(t) \leq G(t)$ for all $t \in \mathbb{R}$. The maximal element for Δ^+ in this order is the distribution function given by

$$\varepsilon_0(t) = \begin{cases} 0 & \text{if } t \leq 0, \\ 1 & \text{if } t > 0. \end{cases} \quad (2.2)$$

Definition 2.1 (see [48]). A function $T : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is a continuous triangular norm (briefly, a t -norm) if T satisfies the following conditions:

- (TN₁) T is commutative and associative;
- (TN₂) T is continuous;
- (TN₃) $T(a, 1) = a$ for all $a \in [0, 1]$;
- (TN₄) $T(a, b) \leq T(c, d)$ whenever $a \leq c$ and $b \leq d$ for all $a, b, c, d \in [0, 1]$.

Typical examples of continuous t -norms are $T_P(a, b) = ab$, $T_M(a, b) = \min(a, b)$ and $T_L(a, b) = \max(a + b - 1, 0)$ (the Łukasiewicz t -norm). Recall (see [46, 49]) that if T is a t -norm and $\{x_n\}$ is a given sequence of numbers in $[0, 1]$, $T_{i=1}^n x_i$ is defined recurrently by

$$T_{i=1}^n x_i = \begin{cases} x_1 & \text{if } n = 1, \\ T(T_{i=1}^{n-1} x_i, x_n) & \text{if } n \geq 2. \end{cases} \quad (2.3)$$

$T_{i=n}^\infty x_i$ is defined as $T_{i=1}^\infty x_{n+i}$.

Definition 2.2 (see [47]). A random normed space (briefly, RN-space) is a triple (X, Λ, T) , where X is a vector space, T is a continuous t -norm, and Λ is a mapping from X into D^+ , such that the following conditions hold:

- (RN₁) $\Lambda_x(t) = \varepsilon_0(t)$ for all $t > 0$ if and only if $x = 0$;
- (RN₂) $\Lambda_{\alpha x}(t) = \Lambda_x(t/|\alpha|)$ for all $x \in X$, $\alpha \neq 0$;
- (RN₃) $\Lambda_{x+y}(t+s) \geq T(\Lambda_x(t), \Lambda_y(s))$ for all $x, y \in X$ and all $t, s \geq 0$.

Every normed space $(X, \|\cdot\|)$ defines a random normed space (X, Λ, T_M) , where $\Lambda_u(t) = t/(t + \|u\|)$ for all $t > 0$, and T_M is the minimum t -norm. This space is called the induced random normed space.

Definition 2.3. Let (X, Λ, T) be an RN-space.

- (1) A sequence $\{x_n\}$ in X is said to be *convergent* to x in X if, for every $\varepsilon > 0$ and $\lambda > 0$, there exists a positive integer N , such that $\Lambda_{x_n-x}(\varepsilon) > 1 - \lambda$ whenever $n \geq N$.
- (2) A sequence $\{x_n\}$ in X is called *Cauchy* if, for every $\varepsilon > 0$ and $\lambda > 0$, there exists a positive integer N , such that $\Lambda_{x_n-x_m}(\varepsilon) > 1 - \lambda$ whenever $n \geq m \geq N$.
- (3) An RN-space (X, Λ, T) is said to be *complete* if every Cauchy sequence in X is convergent to a point in X . A complete RN-space is said to be a *random Banach space*.

Theorem 2.4 (see [48]). *If (X, Λ, T) is an RN-space and $\{x_n\}$ is a sequence, such that $x_n \rightarrow x$, then $\lim_{n \rightarrow \infty} \Lambda_{x_n}(t) = \Lambda_x(t)$ almost everywhere.*

Starting with the paper [50], the stability of some functional equations in the framework of fuzzy normed spaces or random normed spaces has been investigated in [51–57].

Let X be a set. A function $d: X \times X \rightarrow [0, \infty]$ is called a *generalized metric* on X if d satisfies

- (1) $d(x, y) = 0$ if and only if $x = y$;
- (2) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (3) $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in X$.

Let (X, d) be a generalized metric space. An operator $T: X \rightarrow X$ satisfies a Lipschitz condition with Lipschitz constant L if there exists a constant $L \geq 0$ such that $d(Tx, Ty) \leq Ld(x, y)$ for all $x, y \in X$. If the Lipschitz constant L is less than 1, then the operator T is called a strictly contractive operator. Note that the distinction between the generalized metric and the usual metric is that the range of the former is permitted to include the infinity. We recall the following theorem by Diaz and Margolis.

Theorem 2.5 (see [58, 59]). *Let (X, d) be a complete generalized metric space and let $J: X \rightarrow X$ be a strictly contractive mapping with Lipschitz constant $L < 1$, then for each given element $x \in X$, either*

$$d(J^n x, J^{n+1} x) = \infty, \quad (2.4)$$

for all nonnegative integers n , or there exists a positive integer n_0 , such that

- (1) $d(J^n x, J^{n+1} x) < \infty$, for all $n \geq n_0$;
- (2) the sequence $\{J^n x\}$ converges to a fixed-point y^* of J ;
- (3) y^* is the unique fixed-point of J in the set $Y = \{y \in X \mid d(J^{n_0} x, y) < \infty\}$;
- (4) $d(y, y^*) \leq (1/(1-L))d(y, Jy)$ for all $y \in Y$.

In 1996, Isac and Th. M Rassias [60] were the first to provide applications of stability theory of functional equations for the proof of new fixed-point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [61–67]).

In this paper, we prove the generalized Hyers-Ulam stability of the generalized Apollonius type quadratic functional equation (1.3) in random Banach space by using the fixed point method.

Throughout this paper, assume that X is a vector spaces and (Y, μ, T) is a complete RN-space.

3. Generalized Hyers-Ulam Stability of the Quadratic Functional Equation (1.3) in RN-Spaces

Let $x = \sum_i^n x_i$, $y = \sum_i^n y_i$, $z = \sum_i^n z_i$, and for a given mapping $Q: X \rightarrow Y$, consider the mapping $DQ: X^3 \rightarrow Y$, defined by

$$DQ(z, x, y) = Q(z - x) + Q((z - y)) - \frac{1}{2}Q(x - y) + 2Q\left(z - \frac{x + y}{2}\right), \quad (3.1)$$

for all $x, y, z \in X$.

Using the fixed-point method, we prove the generalized Hyers-Ulam stability of the quadratic functional equation $DQ(z, x, y) = 0$ in complete RN-spaces.

Theorem 3.1. *Let $\rho: X^3 \rightarrow D^+$ be a mapping ($\rho(z_1, \dots, z_n, x_1, \dots, x_n, y_1, \dots, y_n)$) is denoted by $\rho_{z,x,y}$) such that, for some $0 < \alpha < 4$,*

$$\rho_{2z,2x,2y}(t) \geq \rho_{z,x,y}(t), \quad (3.2)$$

for all $x, y, z \in X$ and all $t > 0$. Suppose that an even mapping $Q: X \rightarrow Y$ with $Q(0) = 0$ satisfies the inequality

$$\mu_{DQ(z,x,y)}(t) \geq \rho_{z,x,y}(t), \quad (3.3)$$

for all $x, y, z \in X$ and all $t > 0$, then there exists a unique quadratic mapping $R: X \rightarrow Y$, such that

$$\mu_{Q(x)-R(x)}\left(\frac{t}{4-\alpha}\right) \geq \rho_{x,x,-x}(t), \quad (3.4)$$

for all $x \in X$ and all $t > 0$.

Proof. Putting $z = x$ and $y = -x$ in (3.3), we get

$$\mu_{Q(2x)-4Q(x)}(t) \geq \rho_{x,x,-x}(t), \quad (3.5)$$

for all $x \in X$ and all $t > 0$. Therefore,

$$\mu_{Q(2x)/4-Q(x)}(t) \geq \rho_{x,x,-x}(4t), \quad (3.6)$$

for all $x \in X$ and all $t > 0$.

Let S be the set of all even mappings $h: X \rightarrow Y$ with $h(0) = 0$ and introduce a generalized metric on S as follows:

$$d(h, k) = \inf\{u \in \mathbb{R}^+ : \mu_{h(x)-k(x)}(ut) \geq \rho_{x,x,-x}(t), \forall x \in X, \forall t > 0\}, \quad (3.7)$$

where, as usual, $\inf \emptyset = +\infty$. It is easy to show that (S, d) is a generalized complete metric space (see [68, Lemma 2.1]).

Now, we define the mapping $J : S \rightarrow S$

$$Jh(x) := \frac{h(2x)}{4}, \quad (3.8)$$

for all $h \in S$ and $x \in X$. Let $f, g \in S$ such that $d(f, g) < \varepsilon$. Therefore,

$$\begin{aligned} \mu_{Jg(x)-Jf(x)}\left(\frac{\alpha\varepsilon}{4}t\right) &= \mu_{g(2x)/4-f(2x)/4}\left(\frac{\alpha\varepsilon}{4}t\right) = \mu_{g(2x)-f(2x)}(\alpha\varepsilon t) \\ &\geq \rho_{2x,2x,-2x}(\alpha t) \geq \rho_{x,x,-x}(t), \end{aligned} \quad (3.9)$$

that is, if $d(f, g) < \varepsilon$, we have $d(Jf, Jg) < (\alpha/4)\varepsilon$. Hence,

$$d(Jf, Jg) \leq \frac{\alpha}{4}d(f, g), \quad (3.10)$$

for all $f, g \in S$, that is, J is a strictly contractive self-mapping on S with the Lipschitz constant $L = \alpha/4 < 1$.

It follows from (3.6) that

$$\mu_{JQ(x)-Q(x)}\left(\frac{1}{4}t\right) \geq \rho_{x,x,-x}(t), \quad (3.11)$$

for all $x \in X$ and all $t > 0$, which means that $d(JQ, Q) \leq 1/4$.

By Theorem 2.5, there exists a unique mapping $R : X \rightarrow Y$, such that R is a fixed point of J , that is, $R(2x) = 4R(x)$ for all $x \in X$.

Also, $d(J^m Q, R) \rightarrow 0$ as $m \rightarrow \infty$, which implies the equality

$$\lim_{m \rightarrow \infty} \frac{Q(2^m x)}{2^{2m}} = R(x), \quad (3.12)$$

for all $x \in X$.

It follows from (3.2) and (3.3) that

$$\begin{aligned} \mu_{DQ(2^m z, 2^m x, 2^m y)/2^{2m}}(t) &\geq \rho_{2^m z, 2^m x, 2^m y}(2^{2m}t) = \rho_{2^m z, 2^m x, 2^m y}\left(\alpha^m \left(\frac{4}{\alpha}\right)^m t\right) \\ &\geq \rho_{z,x,y}\left(\left(\frac{4}{\alpha}\right)^m t\right), \end{aligned} \quad (3.13)$$

for all $x, y, z \in X$ and all $t > 0$. Letting $m \rightarrow \infty$ in (3.13), we find that $\mu_{DR(z,x,y)}(t) = 1$ for all $t > 0$, which implies $DR(z, x, y) = 0$. By [45, Lemma 2.1], the mapping, $R : X \rightarrow Y$ is quadratic.

Since R is the unique fixed point of J in the set $\Omega = \{g \in S : d(f, g) < \infty\}$, R is the unique mapping such that

$$\mu_{Q(x)-R(x)}(ut) \geq \rho_{x,x,-x}(t), \quad (3.14)$$

for all $x, y, z \in X$ and all $t > 0$. Using the fixed-point alternative, we obtain that

$$d(Q, R) \leq \frac{1}{1-L} d(Q, JQ) \leq \frac{1}{4(1-L)} = \frac{1}{4-\alpha}, \quad (3.15)$$

which implies the inequality

$$\mu_{Q(x)-R(x)}\left(\frac{1}{4-\alpha}t\right) \geq \rho_{x,x,-x}(t), \quad (3.16)$$

for all $x \in X$ and all $t > 0$. So

$$\mu_{Q(x)-R(x)}(t) \geq \rho_{x,x,-x}((4-\alpha)t), \quad (3.17)$$

for all $x \in X$ and all $t > 0$. This completes the proof. \square

Theorem 3.2. Let $\rho : X^3 \rightarrow D^+$ be a mapping ($\rho(z_1, \dots, z_n, x_1, \dots, x_n, y_1, \dots, y_n)$ is denoted by $\rho_{z,x,y}$) such that, for some $\alpha > 4$,

$$\rho_{z/2, x/2, y/2}(t) \geq \rho_{z,x,y}(\alpha t), \quad (3.18)$$

for all $x, y, z \in X$ and all $t > 0$. Suppose that an even mapping $Q : X \rightarrow Y$ satisfying $Q(0) = 0$ and (3.3), then there exists a unique quadratic mapping $R : X \rightarrow Y$, such that

$$\mu_{Q(x)-R(x)}(t) \geq \rho_{x,x,-x}((\alpha-4)t), \quad (3.19)$$

for all $x \in X$ and all $t > 0$.

Proof. Let (S, d) be the generalized metric space defined in the proof of Theorem 3.1.

We consider the mapping $J : S \rightarrow S$ defined by

$$Jh(x) := 4h\left(\frac{x}{2}\right), \quad (3.20)$$

for all $h \in S$ and $x \in X$. Let $f, g \in S$, such that $d(f, g) < \varepsilon$, then

$$\begin{aligned} \mu_{Jg(x)-Jf(x)}\left(\frac{4\varepsilon}{\alpha}t\right) &= \mu_{4g(x/2)-4f(x/2)}\left(\frac{4\varepsilon}{\alpha}t\right) = \mu_{g(x/2)-f(x/2)}\left(\frac{\varepsilon}{\alpha}t\right) \\ &\geq \rho_{x/2, x/2, -x/2}\left(\frac{t}{\alpha}\right) \geq \rho_{x,x,-x}(t), \end{aligned} \quad (3.21)$$

that is, if $d(f, g) < \varepsilon$, we have $d(Jf, Jg) < (4/\alpha)\varepsilon$. This means that

$$d(Jf, Jg) \leq \frac{4}{\alpha}d(f, g), \quad (3.22)$$

for all $f, g \in S$, that is, J is a strictly contractive self-mapping on S with the Lipschitz constant $L = 4/\alpha < 1$.

By Theorem 2.5, there exists a unique mapping $R: X \rightarrow Y$, such that R is a fixed point of J , that is, $R(x/2) = (1/4)R(x)$ for all $x \in X$.

Also, $d(J^m Q, R) \rightarrow 0$ as $m \rightarrow \infty$, which implies the equality

$$\lim_{m \rightarrow \infty} 2^{2m} Q\left(\frac{x}{2^m}\right) = R(x), \quad (3.23)$$

for all $x \in X$.

It follows from (3.5) that

$$\mu_{JQ(x)-Q(x)}\left(\frac{1}{\alpha}t\right) \geq \rho_{x/2, x/2, -x/2}\left(\frac{t}{\alpha}\right) \geq \rho_{x, x, -x}(t), \quad (3.24)$$

for all $x \in X$ and all $t > 0$, which implies that $d(JQ, Q) \leq 1/\alpha$.

Since R is the unique fixed point of J in the set $\Omega = \{g \in S : d(f, g) < \infty\}$, and R is the unique mapping, such that

$$\mu_{Q(x)-R(x)}(ut) \geq \rho_{x, x, -x}(t), \quad (3.25)$$

for all $x \in X$ and all $t > 0$. Using the fixed point alternative, we obtain that

$$d(Q, R) \leq \frac{1}{1-L} d(Q, JQ) \leq \frac{1}{\alpha(1-L)} = \frac{1}{\alpha(1-4/\alpha)}, \quad (3.26)$$

which implies the inequality

$$\mu_{Q(x)-R(x)}\left(\frac{1}{\alpha-4}t\right) \geq \rho_{x, x, -x}(t), \quad (3.27)$$

for all $x \in X$ and all $t > 0$. So

$$\mu_{Q(x)-R(x)}(t) \geq \rho_{x, x, -x}((\alpha-4)t), \quad (3.28)$$

for all $x \in X$ and all $t > 0$.

The rest of the proof is similar to the proof of Theorem 3.1. □

Acknowledgments

The fifth author was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (NRF-2009-0070788) and by Hi Seoul Science Fellowship from Seoul Scholarship Foundation. This paper was dedicated to Professor Sehie Park.

References

- [1] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience, New York, NY, USA, 1960.
- [2] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [3] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [4] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [5] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [6] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [7] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques*, vol. 108, no. 4, pp. 445–446, 1984.
- [8] J. M. Rassias, "Solution of a problem of Ulam," *Journal of Approximation Theory*, vol. 57, no. 3, pp. 268–273, 1989.
- [9] J. M. Rassias, "On the stability of the Euler-Lagrange functional equation," *Chinese Journal of Mathematics*, vol. 20, no. 2, pp. 185–190, 1992.
- [10] J. M. Rassias, "On the stability of the non-linear Euler-Lagrange functional equation in real normed linear spaces," *Journal of Mathematical and Physical Sciences*, vol. 28, no. 5, pp. 231–235, 1994.
- [11] J. M. Rassias, "On the stability of the general Euler-Lagrange functional equation," *Demonstratio Mathematica*, vol. 29, no. 4, pp. 755–766, 1996.
- [12] J. M. Rassias, "Solution of the Ulam stability problem for Euler-Lagrange quadratic mappings," *Journal of Mathematical Analysis and Applications*, vol. 220, no. 2, pp. 613–639, 1998.
- [13] K. Ravi, M. Arunkumar, and J. M. Rassias, "Ulam stability for the orthogonally general Euler-Lagrange type functional equation," *International Journal of Mathematics and Statistics*, vol. 3, no. A08, pp. 36–46, 2008.
- [14] H.-X. Cao, J.-R. Lv, and J. M. Rassias, "Superstability for generalized module left derivations and generalized module derivations on a Banach module. I," *Journal of Inequalities and Applications*, vol. 2009, Article ID 718020, 10 pages, 2009.
- [15] H.-X. Cao, J.-R. Lv, and J. M. Rassias, "Superstability for generalized module left derivations and generalized module derivations on a Banach module. II," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 10, no. 3, article 85, pp. 1–8, 2009.
- [16] M. E. Gordji, S. Zolfaghari, J. M. Rassias, and M. B. Savadkouhi, "Solution and stability of a mixed type cubic and quartic functional equation in quasi-Banach spaces," *Abstract and Applied Analysis*, vol. 2009, Article ID 417473, 14 pages, 2009.
- [17] K. Ravi, J. M. Rassias, M. Arunkumar, and R. Kodandan, "Stability of a generalized mixed type additive, quadratic, cubic and quartic functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 10, no. 4, article 114, pp. 1–29, 2009.
- [18] M. E. Gordji and H. Khodaei, "On the generalized Hyers-Ulam-Rassias stability of quadratic functional equations," *Abstract and Applied Analysis*, vol. 2009, Article ID 923476, 11 pages, 2009.
- [19] M. B. Savadkouhi, M. E. Gordji, J. M. Rassias, and N. Ghobadipour, "Approximate ternary Jordan derivations on Banach ternary algebras," *Journal of Mathematical Physics*, vol. 50, no. 4, Article ID 042303, 9 pages, 2009.
- [20] F. Skof, "Proprietà locali e approssimazione di operatori," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 53, pp. 113–129, 1983.
- [21] P. W. Cholewa, "Remarks on the stability of functional equations," *Aequationes Mathematicae*, vol. 27, no. 1-2, pp. 76–86, 1984.
- [22] St. Czerwik, "On the stability of the quadratic mapping in normed spaces," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 62, pp. 59–64, 1992.
- [23] J. Aczél and J. Dhombres, *Functional Equations in Several Variables*, vol. 31 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 1989.

- [24] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, USA, 2002.
- [25] M. E. Gordji, M. B. Ghaemi, and H. Majani, "Generalized Hyers-Ulam-Rassias theorem in Menger probabilistic normed spaces," *Discrete Dynamics in Nature and Society*, vol. 2010, Article ID 162371, 11 pages, 2010.
- [26] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, Progress in Nonlinear Differential Equations and Their Applications, 34, Birkhäuser, Boston, Mass, USA, 1998.
- [27] S.-M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, Fla, USA, 2001.
- [28] A. Najati, "Hyers-Ulam stability of an n -Apollonius type quadratic mapping," *Bulletin of the Belgian Mathematical Society. Simon Stevin*, vol. 14, no. 4, pp. 755–774, 2007.
- [29] A. Najati, "Fuzzy stability of a generalized quadratic functional equation," *Communications of the Korean Mathematical Society*, vol. 25, no. 3, pp. 405–417, 2010.
- [30] A. Najati and F. Moradlou, "Hyers-Ulam-Rassias stability of the Apollonius type quadratic mapping in non-Archimedean spaces," *Tamsui Oxford Journal of Mathematical Sciences*, vol. 24, no. 4, pp. 367–380, 2008.
- [31] A. Najati and C. Park, "The Pexiderized Apollonius-Jensen type additive mapping and isomorphisms between C^* -algebras," *Journal of Difference Equations and Applications*, vol. 14, no. 5, pp. 459–479, 2008.
- [32] S.-Q. Oh and C.-G. Park, "Linear functional equations in a Hilbert module," *Taiwanese Journal of Mathematics*, vol. 7, no. 3, pp. 441–448, 2003.
- [33] C.-G. Park, "On the stability of the linear mapping in Banach modules," *Journal of Mathematical Analysis and Applications*, vol. 275, no. 2, pp. 711–720, 2002.
- [34] C.-G. Park, "Modified Trif's functional equations in Banach modules over a C^* -algebra and approximate algebra homomorphisms," *Journal of Mathematical Analysis and Applications*, vol. 278, no. 1, pp. 93–108, 2003.
- [35] C.-G. Park, "Lie $*$ -homomorphisms between Lie C^* -algebras and Lie $*$ -derivations on Lie C^* -algebras," *Journal of Mathematical Analysis and Applications*, vol. 293, no. 2, pp. 419–434, 2004.
- [36] C.-G. Park, "Homomorphisms between Lie JC^* -algebras and Cauchy-Rassias stability of Lie JC^* -algebra derivations," *Journal of Lie Theory*, vol. 15, no. 2, pp. 393–414, 2005.
- [37] C.-G. Park, "Approximate homomorphisms on JB^* -triples," *Journal of Mathematical Analysis and Applications*, vol. 306, no. 1, pp. 375–381, 2005.
- [38] C.-G. Park, "Homomorphisms between Poisson JC^* -algebras," *Bulletin of the Brazilian Mathematical Society*, vol. 36, no. 1, pp. 79–97, 2005.
- [39] C.-G. Park, "Isomorphisms between unital C^* -algebras," *Journal of Mathematical Analysis and Applications*, vol. 307, no. 2, pp. 753–762, 2005.
- [40] C. G. Park, J. C. Hou, and S. Q. Oh, "Homomorphisms between JC^* -algebras and Lie C^* -algebras," *Acta Mathematica Sinica*, vol. 21, no. 6, pp. 1391–1398, 2005.
- [41] C.-G. Park and W.-G. Park, "On the Jensen's equation in Banach modules," *Taiwanese Journal of Mathematics*, vol. 6, no. 4, pp. 523–531, 2002.
- [42] Th. M. Rassias, "The problem of S. M. Ulam for approximately multiplicative mappings," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 352–378, 2000.
- [43] Th. M. Rassias, "On the stability of functional equations and a problem of Ulam," *Acta Applicandae Mathematicae*, vol. 62, no. 1, pp. 23–130, 2000.
- [44] Th. M. Rassias and P. Šemrl, "On the behavior of mappings which do not satisfy Hyers-Ulam stability," *Proceedings of the American Mathematical Society*, vol. 114, no. 4, pp. 989–993, 1992.
- [45] C.-G. Park and Th. M. Rassias, "Hyers-Ulam stability of a generalized Apollonius type quadratic mapping," *Journal of Mathematical Analysis and Applications*, vol. 322, no. 1, pp. 371–381, 2006.
- [46] O. Hadžić and E. Pap, *Fixed Point Theory in PM Spaces*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2001.
- [47] A. N. Šerstnev, "On the notion of a random normed space," *Doklady Akademii Nauk SSSR*, vol. 149, pp. 280–283, 1963.
- [48] B. Schweizer and A. Sklar, *Probabilistic Metric Spaces*, North-Holland Series in Probability and Applied Mathematics, North-Holland, New York, NY, USA, 1983.

- [49] O. Hadžić, E. Pap, and M. Budinčević, "Countable extension of triangular norms and their applications to the fixed point theory in probabilistic metric spaces," *Kybernetika*, vol. 38, no. 3, pp. 363–382, 2002.
- [50] A. K. Mirmostafae, M. Mirzavaziri, and M. S. Moslehian, "Fuzzy stability of the Jensen functional equation," *Fuzzy Sets and Systems*, vol. 159, no. 6, pp. 730–738, 2008.
- [51] D. Miheţ, "The fixed point method for fuzzy stability of the Jensen functional equation," *Fuzzy Sets and Systems*, vol. 160, no. 11, pp. 1663–1667, 2009.
- [52] D. Miheţ, R. Saadati, and S. M. Vaezpour, "The stability of the quartic functional equation in random normed spaces," *Acta Applicandae Mathematicae*, vol. 110, no. 2, pp. 797–803, 2010.
- [53] D. Miheţ, R. Saadati, and S. M. Vaezpour, "The stability of an additive functional equation in Menger probabilistic φ -normed spaces," *Mathematica Slovaca*. In press.
- [54] A. K. Mirmostafae, "A fixed point approach to almost quartic mappings in quasi fuzzy normed spaces," *Fuzzy Sets and Systems*, vol. 160, no. 11, pp. 1653–1662, 2009.
- [55] A. K. Mirmostafae and M. S. Moslehian, "Fuzzy versions of Hyers-Ulam-Rassias theorem," *Fuzzy Sets and Systems*, vol. 159, no. 6, pp. 720–729, 2008.
- [56] A. K. Mirmostafae and M. S. Moslehian, "Fuzzy approximately cubic mappings," *Information Sciences*, vol. 178, no. 19, pp. 3791–3798, 2008.
- [57] A. K. Mirmostafae and M. S. Moslehian, "Stability of additive mappings in non-Archimedean fuzzy normed spaces," *Fuzzy Sets and Systems*, vol. 160, no. 11, pp. 1643–1652, 2009.
- [58] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, 2003.
- [59] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, pp. 305–309, 1968.
- [60] G. Isac and Th. M. Rassias, "Stability of ψ -additive mappings: applications to nonlinear analysis," *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 2, pp. 219–228, 1996.
- [61] L. Cădariu and V. Radu, "On the stability of the Cauchy functional equation: a fixed point approach," in *Iteration Theory (ECIT '02)*, vol. 346 of *Grazer Math. Ber.*, pp. 43–52, Karl-Franzens-Univ. Graz, Graz, Austria, 2004.
- [62] L. Cădariu and V. Radu, "Fixed point methods for the generalized stability of functional equations in a single variable," *Fixed Point Theory and Applications*, vol. 2008, Article ID 749392, 15 pages, 2008.
- [63] J. R. Lee, J.-H. Kim, and C. Park, "A fixed point approach to the stability of an additive-quadratic-cubic-quartic functional equation," *Fixed Point Theory and Applications*, vol. 2010, Article ID 185780, 16 pages, 2010.
- [64] M. Mirzavaziri and M. S. Moslehian, "A fixed point approach to stability of a quadratic equation," *Bulletin of the Brazilian Mathematical Society*, vol. 37, no. 3, pp. 361–376, 2006.
- [65] C. Park, "Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras," *Fixed Point Theory and Applications*, vol. 2007, Article ID 50175, 15 pages, 2007.
- [66] C. Park, "Generalized Hyers-Ulam stability of quadratic functional equations: a fixed point approach," *Fixed Point Theory and Applications*, vol. 2008, Article ID 493751, 9 pages, 2008.
- [67] V. Radu, "The fixed point alternative and the stability of functional equations," *Fixed Point Theory*, vol. 4, no. 1, pp. 91–96, 2003.
- [68] D. Miheţ and V. Radu, "On the stability of the additive Cauchy functional equation in random normed spaces," *Journal of Mathematical Analysis and Applications*, vol. 343, no. 1, pp. 567–572, 2008.