

## EXISTENCE OF GLOBAL SOLUTION FOR A DIFFERENTIAL SYSTEM WITH INITIAL DATA IN $L^p$

PETER BATES, FENGXIN CHEN, and PING WANG

(Received 15 October 1997)

**ABSTRACT.** In this paper, we study the system governing flows in the magnetic field within the earth. The system is similar to the magnetohydrodynamic (MHD) equations. By establishing a new priori estimates and following Calderón's procedure for the Navier Stokes equations [1], we obtained, for initial data in space  $L^p$ , the global in time existence and uniqueness of weak solution of the system subject to appropriate conditions.

**Keywords and phrases.** Fluid, magnetic field, global solution.

1991 Mathematics Subject Classification. 35K55, 35Q35, 76D99.

**1. Introduction.** We consider in this work the following differential system arising from geophysics (cf. Hide [7]), which governs the flow of an electrically-conducting fluid in the presence of a magnetic field, when referred to a frame which rotates with angular velocity  $\Omega$  relative to an inertial frame

$$\begin{aligned}\frac{\partial v}{\partial t} + (v \cdot \nabla)v &= \nu \Delta v - \frac{1}{\rho} \nabla p - 2\Omega \times v - \frac{1}{\rho\mu} (\nabla \times b) \times b + f(x), \\ \frac{\partial b}{\partial t} &= \lambda \Delta b - \nabla \times (v \times b) - \frac{1}{\mu} \nabla q + g(x), \\ \operatorname{div} v &= 0, \quad \operatorname{div} b = 0,\end{aligned}\tag{1.1}$$

where  $v$  is the Eulerian flow velocity,  $\rho$  is the density,  $b$  is the magnetic field,  $p$  is the pressure,  $\nu, \mu$  are, respectively, constants of kinematical viscosity, magnetic permeability,  $\lambda = \eta/\mu$  with electrical resistivity  $\eta$ , and  $f(x), g(x)$  are volume forces.

The initial conditions are as follows:

$$v(x, 0) = v_0, \quad b(x, 0) = b_0 \quad \text{for } x \in R^n.\tag{1.2}$$

The existence of solutions of system (1.1) and (1.2) in  $L^2$  has been proved in [9]. Some regularity properties and large time behaviors of the solutions for a similar system, the MHD equations, are obtained in Sermange [10] and Temam [12]. More recently, we obtained in [2] the local in time existence and uniqueness of weak solutions of the system in  $L^p$  with  $p > n$ .

Motivated by Calderón's work on the Navier Stokes equations [1], we consider in this paper the initial value problem for the above system in the infinite cylinder  $S = (0, \infty) \times R^n$  with initial data  $v_0, b_0 \in L^p$  with  $p \leq n$ .

This article is arranged in the following order: in Section 2, we introduce some notations and definitions. Applying Calderón's partition lemma, we introduce in Section 3

Leray’s approximating system for our problem. In Section 4, we state and briefly prove some lemmas similar to those for Navier-Stokes equations. Finally, in Section 5, by establishing a priori estimates for our system and adapting Calderón’s technique, we prove the global in time existence and uniqueness of weak solution of (1.1) and (1.2) for initial data in  $L^p$ .

**2. Notations and definition of weak solution.** In this section, we introduce some notations and the definition of a weak solution of the differential system (1.1) and (1.2).

Denote by  $L^{p,q}(S_T)$  the standard functional space consisting of Lebesgue measurable vector functions  $u = (u_1, u_2, \dots, u_n)$  with the following property:

$$\|u\|_{p,q} = \sum_{j=1}^n \left[ \int_0^T \left( \int_{R^n} |u_j(x,t)|^p dx \right)^{q/p} \right]^{1/q} < \infty, \tag{2.1}$$

where  $S_T = (0, T) \times R^n$ . Let  $u^* = \sup_t |u|$  and define  $\|u^*\|_p(T) = (\int (\sup_{0 < t < T} |u|)^p dx)^{1/p}$ . Let  $\mathcal{L}^{p,q}(S_T) = L^{p,q}(S_T) \times L^{p,q}(S_T)$  with the standard product norm  $\|(v, b)\|_{p,q} = \|v\|_{p,q} + \|b\|_{p,q}$  and  $\mathbb{L}^p(R^n) = L^p(R^n) \times \dots \times L^p(R^n)$  with the norm  $\|g\|_p = \sum_{i=1}^n \|g_i\|_p$  for  $g \in \mathbb{L}^p(R^n)$ .

Let  $\mathcal{S}(R^n)$  denote the space of rapidly decreasing functions on  $R^n$ ,  $\mathcal{S}'(R^n)$  the space of tempered distributions, and  $\mathcal{D}_T$  the space of functions  $\phi(x, t) = (\phi_1(x, t), \dots, \phi_n(x, t))$  with the properties:  $\phi_i \in \mathcal{S}(R^{n+1})$ ,  $\phi_i(x, t) = 0$  for  $t \geq T$ ;  $\text{div } \phi = \sum_{i=1}^n D_{x_i} \times \phi(x, t) = 0$  for all  $t$ .

**DEFINITION 2.1.** A function  $u = (v, b)$  is a weak solution of (1.1) and (1.2) with initial divergence free data  $(v_0, b_0) \in \mathbb{L}^p(R^n) \times \mathbb{L}^p(R^n)$  if the following conditions hold

- (1)  $u(x, t) \in \mathcal{L}^{p,q}(S_T)$  for some  $p, q$  with  $p, q \geq 2$ ;
- (2) for  $\phi, \psi \in \mathcal{D}_T$ ,

$$\begin{aligned} & \int_0^T \int_{R^n} \langle v, (v\Delta + D_t)\phi \rangle dx dt + \int_0^T \int_{R^n} \langle v, (\nabla\phi)v \rangle dx dt \\ & \quad + \int_0^T \int_{R^n} \langle v, 2\Omega \times \phi \rangle dx dt - \frac{1}{\rho\mu} \int_0^T \int_{R^n} \langle b, (\nabla\phi)b \rangle dx dt \\ & = - \int_{R^n} \langle v_0, \phi(x, 0) \rangle dx + \int_0^T \int_{R^n} \langle f(x, t), \phi \rangle dx dt; \\ & \int_0^T \int_{R^n} \langle b, (\lambda\Delta + D_t)\psi \rangle dx dt + \int_0^T \int_{R^n} \langle v, (\nabla\psi)b \rangle dx dt \\ & \quad - \int_0^T \int_{R^n} \langle b, (\nabla\psi)v \rangle dx dt \\ & = - \int_{R^n} \langle b_0, \psi(x, 0) \rangle dx + \int_0^T \int_{R^n} \langle g(x, t), \psi \rangle dx dt; \end{aligned} \tag{2.2}$$

- (3) for almost every  $t \in [0, T]$ ,  $\text{div } v(x, t) = \text{div } b(x, t) = 0$  in the distributional sense.

Following Fabes et al. [4], we can find a divergence free matrix fundamental solution  $E_{i,j}$  for the  $n$ -dimensional heat equation. We define matrices  $(E_{i,j}^k)$ ,  $k = 1, 2$  as follows:

$$E_{i,j}^k = \delta_{i,j} \Gamma_k(x, t) - R_i R_j \Gamma_k(x, t), \tag{2.3}$$

where

$$\Gamma_1 = \frac{e^{-|x|^2/4vt}}{(4\pi vt)^{n/2}}, \quad \Gamma_2 = \frac{e^{-|x|^2/4\lambda t}}{(4\pi \lambda t)^{n/2}}, \tag{2.4}$$

$R_j$  is the  $j$ th Riesz transform, namely,  $R_j$  is a singular integral operator on  $L^p(\mathbb{R}^n)$ ,  $1 < p < \infty$ , defined as

$$R_j(f) = \text{P.V.C}_j \int_{\mathbb{R}^n} (x_j - y_j) |x - y|^{-n-1} f(y) dy. \tag{2.5}$$

Now, we define an integral operator  $A(v, w)$  for  $v = (v_1, \dots, v_n)$ ,  $w = (w_1, \dots, w_n)$ . Denote

$$B_k(v, w)(x, t) = \int_0^t \int_{\mathbb{R}^n} \langle v(y, s), \nabla E^k(x - y, t - s) \rangle w(y, s) dy ds; \quad \text{for } k = 1, 2. \tag{2.6}$$

$$D(v)(x, t) = \int_0^t \int_{\mathbb{R}^n} \langle v(y, s), 2\Omega \times E^1(x - y, t - s) \rangle dy ds. \tag{2.7}$$

For  $u_1 = (v_1, b_1)$ ,  $u_2 = (v_2, b_2)$ , let

$$A(u_1, u_2) = \left( \begin{array}{c} B_1(v_1, v_2) - \frac{1}{\rho\mu} B_1(b_1, b_2) \\ \frac{1}{2} [B_2(v_1, b_1) - B_2(b_1, v_1) + B_2(v_2, b_2) - B_2(b_2, v_2)] \end{array} \right). \tag{2.8}$$

**3. Approximating system.** The following result was obtained in [2].

**THEOREM 3.1.** *Let  $v_0, b_0 \in L^r$ ,  $1 \leq r < \infty$ , be divergence free weakly.  $u(x, t) = (v(x, t), b(x, t)) \in \mathcal{L}^{p,q}(S_T)$ ,  $p, q \geq 2$ ,  $p < \infty$ , is a weak solution of (1.1) and (1.2) with initial value  $(v_0, b_0)$  if and only if  $u$  is a solution of the following integral equation:*

$$u + A(u, u) + D(u) = u^0 + f^0, \tag{3.1}$$

where

$$u^0 = \left( \begin{array}{c} \int_{\mathbb{R}^n} \Gamma_1(x - y, t) v_0(y) dy \\ \int_{\mathbb{R}^n} \Gamma_2(x - y, t) b_0(y) dy \end{array} \right), \tag{3.2}$$

$$f^0 = \left( \begin{array}{c} \int_0^t \int_{\mathbb{R}^n} E^1(x - y, t - s) f(y, s) dy ds \\ \int_0^t \int_{\mathbb{R}^n} E^2(x - y, t - s) g(y, s) dy ds \end{array} \right).$$

We need the following lemmas obtained by Calderón [1].

**LEMMA 3.1.** *Let  $f \in L^p(\mathbb{R}^n)$ ,  $2 < p < n$ , be a given vector function such that  $\text{div } f = 0$  in the distributional sense. Then, for each  $s > 0$ ,  $f$  can be expressed as  $g + h$ , where*

$$\|g\|_n \leq cs^{1-(p/n)} \|f\|_p^{p/n}, \quad \text{div } g = 0, \tag{3.3}$$

$$\|h\|_2 \leq cs^{1-(p/2)} \|f\|_p^{p/2}, \quad \text{div } h = 0,$$

where the constant  $c$  depends only on  $n$  and  $p$ .

**LEMMA 3.2.** *Let  $T(u, v) = B(u, v) + l(u) + F$  ( $B(u, v)$  is bilinear and  $l(u)$  is linear) satisfy*

$$\|T(u, v)\| \leq c_1 \|u\| \|v\| + c_2 \|u\| + \|F\| \tag{3.4}$$

*with the same norm in a Banach space. Then the quadratic operator  $T(u, v)$  maps the ball  $\{\|u\| \leq s_1\}$  into itself if  $s_1$  is the smallest root of*

$$c_1 s^2 + (c_2 - 1)s + \|F\| = 0, \tag{3.5}$$

*provided that  $c_1, c_2$ , and  $\|F\|$  satisfy*

$$(1 - c_2)^2 > 4c_1 \|F\|, \quad c_1 > 0, \quad 0 \leq c_2 < 1. \tag{3.6}$$

*If  $2s_1 c_1 + c_2 < 1$ ,  $T(u, v)$  is a contraction mapping in the ball of radius  $s_1$ . In particular,  $T(u, v)$  is a contraction mapping in the ball of radius  $s_1$  if*

$$2c_1 \|F\| ((1 - c_2)^2 - 4c_1 \|F\|)^{-1/2} + c_2 < 1. \tag{3.7}$$

Consider the following system in  $v_1, v_2, b_1, b_2, p_1, p_2, q_1$ , and  $q_2$

$$\begin{aligned} L_1 v_1 + (\nabla v_1) v_1 - \frac{1}{\rho\mu} (\nabla b_1) b_1 + \nabla p_1 &= 0, \\ L_1 v_2 + (\nabla v_2) v_2 + (\nabla v_2) v_1 + (\nabla v_1) v_2 - \frac{1}{\rho\mu} (\nabla b_2) b_2 + (\nabla b_2) b_1 + (\nabla b_1) b_2 + \nabla p_2 &= 0, \\ L_2 b_1 + (\nabla b_1) v_1 - (\nabla v_1) b_1 + \nabla q_1 &= 0, \\ L_2 b_2 + (\nabla b_2) v_2 + (\nabla b_2) v_1 + (\nabla v_1) v_2 - (\nabla v_2) b_2 - (\nabla v_2) b_1 - (\nabla v_1) b_2 + \nabla q_2 &= 0, \\ \operatorname{div} v_i &= 0, \quad \operatorname{div} b_i = 0, \quad i = 1, 2, \\ v_i(x, 0) &= h_i(x), \quad b_i(x, 0) = k_i(x), \quad i = 1, 2, \end{aligned} \tag{3.8}$$

where  $L_1 = \partial/\partial t - \nu\Delta$ ,  $L_2 = \partial/\partial t - \lambda\Delta$ . We have the following definition.

**DEFINITION 3.1.** The vector  $((v_1, v_2), (u_1, u_2))$  is said to be a weak solution of (3.8) if  $((v_1 + v_2), (b_1 + b_2))$  is a weak solution of (1.1) and (1.2) with initial data  $(h_1 + h_2, k_1 + k_2)$ .

It then follows from Theorem 3.1 that

**THEOREM 3.2.** *The vector functions  $((v_1, v_2), (b_1, b_2)) \in \mathcal{L}^{p,q}(S_T)^4$ ,  $2 \leq p, q \leq \infty$ , are weak solutions of (3.8) if and only if they are solutions of the following integral equations:*

$$\begin{aligned} v_1 + B_1(v_1, v_1) - \frac{1}{\rho\mu} B_1(b_1, b_1) &= v_1^0, \\ v_2 + B_1(v_2, v_2) + B_1(v_1, v_2) + B_1(v_2, v_1) - \frac{1}{\rho\mu} [B_1(b_2, b_2) + B_1(b_1, b_2) + B_1(b_2, b_1)] &= v_2^0, \\ b_1 + B_2(v_1, b_1) - B_2(b_1, v_1) &= b_1^0, \\ b_2 + B_2(v_2, b_2) + B_2(v_1, b_2) + B_2(v_2, b_1) - [B_2(b_2, v_2) + B_2(b_1, v_2) + B_2(b_2, v_1)] &= b_2^0, \end{aligned} \tag{3.9}$$

where

$$v_i^0 = \Gamma_1 * h_i, \quad b_i^0 = \Gamma_2 * k_i, \quad i = 1, 2. \quad (3.10)$$

Now, let us introduce Leray's approximating system. Let  $\alpha(x)$  be a  $C^\infty$  nonnegative, compact supported function on  $R^n$  with integral equal to 1,  $\alpha_\varepsilon(x) = \varepsilon^{-n}\alpha(\varepsilon^{-1}x)$ . Denote the modifying function of  $u(x, t)$  by  $u^\#(x, t)$ , i.e.,  $u^\# = \alpha_\varepsilon * u$ . For each  $\varepsilon$ , consider the following approximating system

$$L_1 v_1 + (\nabla u) v_1^\# - \frac{1}{\rho\mu} (\nabla b_1) b_1^\# + \nabla p_1 = 0 \quad (3.11)$$

$$L_1 v_2 + \nabla(v_1 + v_2) v_2^\# + (\nabla v_2) v_1^\# - \frac{1}{\rho\mu} ((\nabla b_2) b_2^\# + (\nabla b_2) b_1^\# + (\nabla b_1) b_2^\#) + \nabla p_2 = 0, \quad (3.12)$$

$$L_2 b_1 + (\nabla b_1) v_1^\# - (\nabla v_1) b_1^\# + \nabla q_1 = 0, \quad (3.13)$$

$$L_2 b_2 + (\nabla b_2) v_2^\# + (\nabla b_2) v_1^\# + (\nabla b_1) v_2^\# - (\nabla v_2) b_2^\# - (\nabla v_2) b_1^\# - (\nabla v_1) b_2^\# + \nabla q_2 = 0, \quad (3.14)$$

$$\operatorname{div} v_i = 0, \quad \operatorname{div} b_i = 0, \quad i = 1, 2, \quad (3.15)$$

$$v_i(x, 0) = v_i^{\prime\#}(x), \quad b_i(x, 0) = b_i^{\prime\#}(x), \quad i = 1, 2, \quad (3.16)$$

where  $v_i', b_i', i = 1, 2$ , are partitions of initial data  $v_0, b_0$ , respectively, in the sense of Lemma 3.1, i.e.,  $v_0 = v_1' + v_2', b_0 = b_1' + b_2'$ . From Lemma 3.1, we have

$$\begin{aligned} \|v_1^{\prime\#}\|_n &\leq \|v_1'\|_n \leq c s^{1-(p/n)} \|v_0\|_n, \\ \|v_2^{\prime\#}\|_2 &\leq \|v_2'\|_2 \leq c s^{1-(p/2)} \|v_0\|_2, \\ \|b_1^{\prime\#}\|_n &\leq \|b_1'\|_n \leq c s^{1-(p/n)} \|b_0\|_n, \\ \|b_2^{\prime\#}\|_2 &\leq \|b_2'\|_2 \leq c s^{1-(p/2)} \|b_0\|_2. \end{aligned} \quad (3.17)$$

**4. Some lemmas.** In this section, we present some lemmas without providing much of the details of their proofs for the arguments involved are similar to those used in [1].

First, we consider (3.11), (3.13), and (3.15) with corresponding data  $v_1(x, 0) = v_1^{\prime\#}$ ,  $b_1(x, 0) = b_1^{\prime\#}$ . The problem is equivalent to the following integral equations (cf. [2])

$$\begin{aligned} v_1 + B_1(v_1, v_1^\#) - \frac{1}{\rho\mu} B_1(b_1, b_2^\#) &= v_1^{0\#}, \\ b_1 + B_2(v_1, b_1^\#) - B_2(b_1, v_1^\#) &= b_1^{0\#}, \end{aligned} \quad (4.1)$$

where  $v_1^{0\#}, v_2^{0\#}$  are defined by (3.10) with  $h_i, k_i$  being replaced by  $v_1^{\prime\#}, b_1^{\prime\#}$ , respectively. Denote  $u_1 = (v_1, b_1)$ , the solution of (4.1). Define, for  $s > 0$ , and a function  $w, w^s = w$  if  $|w| < s, w = 0$  otherwise. We have the following lemma:

**LEMMA 4.1.** *The system (4.1), including the limit case, i.e., when  $u_1 = u_1^\#, (v_1^0, b_1^0) = (v_1^{0\#}, b_1^{0\#})$ , admits a unique solution  $u_1 = (v, b)$ , for all  $t$ , satisfying*

$$\|u_1^*\|_n(\infty) \leq c s^{1-(p/n)} \|u_1^0\|_p^{p/n}, \tag{4.2}$$

provided that  $s^{1-(p/n)} \|u_1^0\|_p^{p/n} < \varepsilon_0$ , where  $u_1^0 = (v_1^0, b_1^0)$ , and

$$\|u_1^*\|_p(\infty) \leq c \max\left(s^{1-(p/n)} \|u_1^0\|_p^{p/n}, \|(u_1^0)^s\|_p\right), \tag{4.3}$$

provided that  $\max(s^{1-(p/n)} \|u_1^0\|_p^{p/n}, \|(u_1^0)^s\|_p) < \varepsilon_0$ , where  $(u_1^0)^s = ((v_1^0)^s, (b_1^0)^s)$ ,  $\varepsilon_0$  is a fixed and a small constant and  $c$  depends only on  $\varepsilon_0$ .

**PROOF.** The proof is a direct extension of that of [1, Lem. III.1]. □

**LEMMA 4.2.** *Let  $u'_1 = (v'_1, b'_1)$  be chosen such that  $\|(v'_1, b'_1)\|_p$  is so small that the existence of solution  $u_1$  is assured by Lemma 4.1 and such that, for all  $t$ ,*

$$\|u_1^*\|_n < a_0 < c_0^{-1}, \tag{4.4}$$

where  $c_0$  is an independent constant. Suppose that  $u_2 = (v_2, b_2)$  is a solution of (3.12), (3.14), (3.15), and (3.16) and suppose that  $\nabla v_2, \nabla b_2, (\partial/\partial t)v_2, (\partial/\partial t)b_2 \in L^2(S_T)$ . Then  $u_2$  satisfies the following estimate:

$$|u_2(t)|_2^2 + 2(1 - c_0 a_0) \int_0^t \|\nabla u_2\|_2^2 dt \leq |u_2(0)|_2^2, \tag{4.5}$$

where

$$\begin{aligned} |u_2|_2^2 &= \left(\|v_2\|_2^2 + \frac{1}{\rho\mu} \|b_2\|_2^2\right), \\ \|\nabla u_2\|_2^2 &= \left(v\|\nabla v_2\|_2^2 + \frac{1}{\rho\mu} \|\nabla b_2\|_2^2\right). \end{aligned} \tag{4.6}$$

**PROOF.** Multiplying (3.12) and (3.14) by  $v_2, b_2$ , respectively and integrating over  $R^n$ , we get

$$\begin{aligned} \frac{1}{2} \frac{d\|v_2\|_2^2}{dt} + v\|\nabla v_2\|_2^2 + ((\nabla v_1)v_2^\#, v_2) \\ - \frac{1}{\rho\mu} \left[ ((\nabla b_1)b_2^\#, v_2) + ((\nabla b_2)b_1^\#, v_2) + ((\nabla b_2)b_2^\#, v_2) \right] = 0, \end{aligned} \tag{4.7}$$

$$\begin{aligned} \frac{1}{2} \frac{d\|b_2\|_2^2}{dt} + \lambda\|\nabla b_2\|_2^2 + ((\nabla b_1)v_2^\#, b_2) \\ - \left[ ((\nabla v_1)b_2^\#, b_2) + ((\nabla v_2)b_2^\#, b_2) - ((\nabla v_2)b_1^\#, b_2) \right] = 0. \end{aligned} \tag{4.8}$$

Note that, for functions  $a, b, c$ , and exponents  $r, n, 2$  such that  $(1/r) + (1/n) + (1/2) = 1$ , we have

$$|((\nabla a)b, c)| \leq \|\nabla a\|_2 \|b\|_r \|c\|_n. \tag{4.9}$$

Multiplying (4.8) by  $(1/\rho\mu)$  and adding the resulting equation to (4.7), we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |(v_2, b_2)|^2 + \|(\nabla v_2, \nabla b_2)\|_2^2 \\ \leq c_1 (\|v_1\|_n + \|b_1\|_n) (\|\nabla v_2\|_2^2 + \|\nabla b_2\|_2^2) \\ \leq c_2 (\|v_1\|_n + \|b_1\|_n) \|(v, b)\|_2^2. \end{aligned} \tag{4.10}$$

It is then standard to obtain (4.5). □

Now, let us consider the existence of a weak solution of (3.12), (3.14), (3.15), and (3.16). It is easy to see that the system is equivalent to the following

$$\begin{aligned} v_2 + \mathfrak{B}_1(u_1, u_2) &= v_2^{0\#}, \\ b_2 + \mathfrak{B}_2(u_1, u_2) &= b_2^{0\#}, \end{aligned} \tag{4.11}$$

where

$$\begin{aligned} \mathfrak{B}_1(u_1, u_2) &= B_1(v_2, v_2^\#) + B_1(v_1, v_2^\#) + B_1(v_2, v_1^\#) \\ &\quad - \frac{1}{\rho\mu} B_1(b_2, b_2^\#) + B_1(b_1, b_2^\#) + B_1(b_2, b_1^\#), \\ \mathfrak{B}_2(u_1, u_2) &= B_2(v_2, b_2^\#) + B_2(v_1, b_2^\#) + B_2(v_2, b_1^\#) \\ &\quad - B_2(b_2, v_2^\#) + B_2(b_1, v_2^\#) + B_2(b_2, v_1^\#). \end{aligned} \tag{4.12}$$

**LEMMA 4.3.** *If  $T$  is suitably small, then there exists a solution  $u_2$  of (4.11) such that*

$$\|u_2^*\|_2(T) < \infty. \tag{4.13}$$

**PROOF.** Applying the standard estimate on  $E^i$ , the definition of  $B_i$  (cf. (2.6)) and the Hardy-Littlewood-Sobolev potential inequality we can prove that

$$\|\mathfrak{B}_i(u_1, u_2)\|_2(T) \leq c \left( \varepsilon^{-n/2} \|u_2^*\|_2(T) + \|u_1^*\|_n(T) \right) \|u_2^\#\|_2(T), \quad i = 1, 2. \tag{4.14}$$

Taking  $\varepsilon^{-n/2} T^{1/2}$  and  $\|u_1^*\|_n$  small enough, we can apply Lemma 3.2 to obtain the existence of  $u_2$ . □

Using the arguments in the proofs of Lemmas [1, III.3, III.4], one can similarly prove the next two lemmas.

**LEMMA 4.4.** *Let  $u_1(x, t) = (v_1, b_1)$  be the solution obtained in Lemma 4.1 solving (4.1). Then, for all  $T > 0$  and all  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$ , we have*

$$\begin{aligned} \|D^\alpha u_1\|_n(S_T) &< \infty, \\ \|D_t D^\alpha u_1\|_n(S_T) &< \infty. \end{aligned} \tag{4.15}$$

**LEMMA 4.5.** *Consider the following integral equations of unknown  $u_2 = (v_2, b_2)$ :*

$$\begin{aligned} v_2 + \mathfrak{B}_1(u_1, u_2) &= F_1(x, t), \\ b_2 + \mathfrak{B}_2(u_1, u_2) &= F_2(x, t), \end{aligned} \tag{4.16}$$

where  $\mathfrak{B}_i$ ,  $i = 1, 2$  are defined in (4.12),  $u_1$  is the solution of (4.1), and  $F_1, F_2$  are functions satisfying

$$\|D^\alpha F_i\|_2(S_T) < \infty, \quad \|D_t D^\alpha F_i\|_2(S_T) < \infty. \tag{4.17}$$

If we denote  $T > 0$  the existence interval for  $t$  of solution of (4.16) by the standard fixed point argument, then

$$\|D^\alpha u_2\|_2(S_T) < \infty, \tag{4.18}$$

$$\|D_t D^\alpha u_2\|_2(S_T) < \infty. \tag{4.19}$$

Using the above estimates, we can prove the following theorem.

**LEMMA 4.6.** *The solution obtained in Lemma 4.3 can be extended to all time  $t > 0$  and it satisfies (4.5) for all  $t$ .*

**PROOF.** We only give a sketch of the proof here. The existence time  $T$  obtained in Lemma 4.3 by the standard fixed point argument depends only on the  $L^2$  norm of the initial data and  $\|u_1^*\|_n$ . Lemma 4.2 implies that  $\|u_2(t)\|_2$  is uniformly bounded by the corresponding norm of the initial data when  $u_2$  satisfies the regularity conditions of Lemma 4.2, which is guaranteed by Lemmas 4.4 and 4.5. Therefore, (4.5) holds for all  $t$  by moving from  $[0, T]$  to  $[T, 2T]$  to  $[2T, 3T]$  and so on. And then the interval of existence can be extended to  $(0, \infty)$ .  $\square$

**5. The global existence theorem.** In this section, we establish some a priori estimate for the solution of (4.11) and obtain, by following Calderón’s procedure [1], the global existence and uniqueness of solution of (1.1) and (1.2).

To adapt Leray’s argument [8] to prove Lemmas 5.2 and 5.3 that we state later, we need to establish the following a priori estimate.

**LEMMA 5.1.** *For a  $C^\infty$  function  $\beta(x)$  satisfying  $\beta(x) = 1$ , if  $|x| > N$ ;  $\beta(x) = 0$ , if  $|x| < N/2$ , and  $\|\nabla\beta\| \leq C/N$ , the solution  $u_2$  of (4.11) satisfies the following inequality*

$$\begin{aligned} & \frac{1}{2} \int_{R^n} \beta(x) \left[ |v_2|^2 + \frac{1}{\rho\mu} |b_2|^2 \right] dx + \int_0^t \int_{R^n} \beta(x) \left[ v |\nabla v_2|^2 + \frac{1}{\rho\mu} |\nabla b_2|^2 \right] dx dt \\ & \leq \frac{1}{2} \int_{R^n} \beta(x) \left[ |v_2(0)|^2 + \frac{1}{\rho\mu} |b_2(0)|^2 \right] dx + C \left( \frac{1+t}{N} \right) |u_2(0)|_2^2 dx dt \tag{5.1} \\ & \quad + \frac{C}{N} \|u_1\|_{n,\infty}(T) |u_2(0)|_2^2 + C \|\beta u_1\|_{n,\infty}(T) |u_2(0)|_2^2, \end{aligned}$$

where  $|u_2|_2$  is defined by (4.6).

**PROOF.** Multiplying equations (3.12) and (3.14) by  $\beta v_2$  and  $\beta b_2$ , respectively, and integrating over  $R^n$ , we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{R^n} \beta(x) |v_2|^2 dx + v (\nabla v_2, \nabla(\beta v_2)) + ((\nabla v_1) v_2^\#, \beta v_2) \\ & \quad + ((\nabla v_2) v_2^\#, \beta v_2) + ((\nabla v_2) v_1^\#, \beta v_2) \\ & \quad - \frac{1}{\rho\mu} \left[ ((\nabla b_1) b_2^\#, \beta v_2) + ((\nabla b_2) b_2^\#, \beta v_2) + ((\nabla b_2) b_1^\#, \beta v_2) \right] \tag{5.2} \\ & \quad - \int_{R^n} (\nabla \beta, v_2) p_2 dx = 0, \end{aligned}$$



$$\begin{aligned}
 \frac{1}{2} \frac{d}{dt} \int_{R^n} \beta(x) |b_2|^2 dx + \lambda (\nabla b_2, \nabla (\beta b_2)) + ((\nabla b_1) v_2^\#, \beta b_2) \\
 + ((\nabla b_2) v_2^\#, \beta b_2) + ((\nabla b_2) v_1^\#, \beta b_2) \\
 - \left[ ((\nabla v_1) b_2^\#, \beta b_2) + ((\nabla v_2) b_2^\#, \beta b_2) + ((\nabla v_2) b_1^\#, \beta b_2) \right] \\
 + \int_{R^n} (\nabla \beta, b_2) q_2 dx = 0.
 \end{aligned}
 \tag{5.3}$$

Let us now separately estimate terms on the left-hand sides of (5.2) and (5.3). First, we deal with the terms on the left side of (5.2). For the second term, we have

$$\begin{aligned}
 v (\nabla v_2, \nabla (\beta v_2)) &\geq v \int_{R^n} \beta |\nabla v_2|^2 dx - v (\nabla v_2, \nabla \beta v_2) \\
 &\geq v \int_{R^n} \beta |\nabla v_2|^2 dx - \frac{C}{N} \|\nabla v_2\|_2 \|v_2\|_2.
 \end{aligned}
 \tag{5.4}$$

For the third term, we apply Hölder’s inequality for exponents,  $r, 2, n$ , to get

$$\begin{aligned}
 ((\nabla v_1) v_2^\#, \beta v_2) &= -((\nabla (\beta v_2)) v_2^\#, v_1) = -((\nabla \beta v_2) v_2^\#, v_1) - ((\nabla v_2) v_2^\#, \beta v_1) \\
 &\leq \frac{C}{N} \|v_1\|_n \|v_2\|_r \|v_2^\#\|_2 + \|\beta v_1\|_n \|\nabla v_2\|_2 \|v_2^\#\|_r \\
 &\leq \frac{C}{N} \|v_1\|_n \|\nabla v_2\|_2^2 + \|\beta v_1\|_n \|\nabla v_2\|_2^2,
 \end{aligned}
 \tag{5.5}$$

where  $1/r = (1/2) - (1/n)$ . For the fourth term, integration by parts, Hölder’s inequality, and then Sobolov’s inequality yield

$$\begin{aligned}
 |((\nabla v_2) v_2^\#, \beta v_2)| &= \left| \frac{1}{2} ((v_2 \nabla \beta) v_2^\#, v_2) \right| \\
 &\leq \frac{C}{N} \|v_2\|_2 (\|\nabla v_2\|_2^2 + \|v_2\|_2^2).
 \end{aligned}
 \tag{5.6}$$

For the fifth term, we have

$$|((\nabla v_2) v_1^\#, \beta v_2)| \leq \|\beta v_1\|_n \|\nabla v_2\| \|v_2\|.
 \tag{5.7}$$

The estimates on the sixth and eighth terms can be obtained, respectively, as

$$\begin{aligned}
 \left| \frac{1}{\rho \mu} ((\nabla b_1) b_2^\#, \beta v_2) \right| &\leq \frac{C}{N} \|b_1\|_n \|\nabla v_2\|_2 \|b_2\|_2 + C \|\beta b_1\|_n \|\nabla v_2\|_2 \|\nabla b_2\|_2, \\
 \left| \frac{1}{\rho \mu} ((\nabla b_2) b_1^\#, \beta v_2) \right| &\leq C \|\beta b_1\|_n \|\nabla b_2\|_2 \|\nabla v_2\|_2.
 \end{aligned}
 \tag{5.8}$$

We do not need to estimate the seventh term because it will be canceled with part of the seventh term in (5.3).

Now, let us check terms on the left-hand side of (5.3). Similarly, for the second term, we have

$$\lambda (\nabla b_2, \nabla (\beta b_2)) \geq \lambda \int_{R^n} \beta |\nabla b_2|^2 dx - \frac{C}{N} \|\nabla b_2\|_2 \|b_2\|_2.
 \tag{5.9}$$

For the third term, we have

$$((\nabla b_1) v_2^\#, \beta b_2) \leq \frac{C}{N} \|b_1\|_n \|\nabla v_2\|_2 \|\nabla b_2\|_2 + \|\beta b_1\|_n \|\nabla v_2\|_2 \|b_2\|_2.
 \tag{5.10}$$

For the fourth term, we get

$$|((\nabla b_2)v_2^\#, \beta b_2)| \leq \frac{C}{N} \|b_2\|_2 (\|\nabla b_2\|_2^2 + \|b_2\|_2^2). \quad (5.11)$$

For the fifth and sixth terms, we have, respectively,

$$\begin{aligned} |((\nabla b_2)v_1^\#, \beta b_2)| &\leq \|\beta v_1\|_n \|\nabla b_2\| \|b_2\|, \\ |((\nabla v_1)b_2^\#, \beta b_2)| &\leq \frac{C}{N} \|v_1\|_n \|\nabla b_2\|_2 \|b_2\|_2 + C \|\beta v_1\|_n \|\nabla b_2\|_2 \|b_2\|_2. \end{aligned} \quad (5.12)$$

The seventh term

$$| -((\nabla v_2)b_2^\#, \beta b_2) - ((\nabla b_2)b_2^\#, \beta v_2) | \leq \frac{C}{N} \|v_2\|_2 (\|\nabla b_2\|_2^2 + \|b_2\|_2^2). \quad (5.13)$$

A multiple of the second term on the left-hand side of the above inequality cancels out the seventh term on the left-hand side of (5.2). The estimate on the eighth term can be obtained as

$$|((\nabla v_2)b_1^\#, \beta b_2)| \leq C \|\beta b_1\|_n \|\nabla v_2\|_2 \|\nabla b_2\|_2. \quad (5.14)$$

Now, multiplying (5.3) by  $1/\rho\mu$ , adding the resulting equation to (5.2), and applying the above estimates yield

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{R^n} \beta(x) (|v_2|^2 + \frac{1}{\rho\mu} |b_2|^2) + \int_{R^n} \beta(x) (v|\nabla v_2|^2 + \frac{1}{\rho\mu} |b_2|^2) dx \\ &\leq \frac{C}{N} \|u_1\|_n \|\nabla u_2\|_2 \|u_2\|_2 + C \|\beta u_1\|_n \|\nabla u_2\|_2 \|\nabla u_2\|_2 \\ &\quad + \int_{R^n} (\nabla \beta, v_2) p_2 dx + \int_{R^n} (\nabla \beta, v_2) q_2 dx. \end{aligned} \quad (5.15)$$

We use Riesz transformation to express  $p_2$  and  $q_2$  as

$$\begin{aligned} p_2 &= R_i R_j \left( v_{1i}(v_2)_j^\# + v_{2i}(v_2)_j^\# + v_{2i}(v_1)_j^\# \right. \\ &\quad \left. - \frac{1}{\rho\mu} b_{1i}(b_2)_j^\# + b_{2i}(b_2)_j^\# + b_{2i}(b_1)_j^\# \right), \\ q_2 &= R_i R_j \left( b_{1i}(v_2)_j^\# + b_{2i}(v_2)_j^\# + b_{2i}(v_1)_j^\# - v_{1i}(b_2)_j^\# - v_{2i}(b_2)_j^\# - v_{2i}(b_1)_j^\# \right). \end{aligned} \quad (5.16)$$

Since  $R_i$  is a continuous map from  $L^p$  to itself for  $p > 1$ , we have

$$\begin{aligned} &\left| \int_{R^n} (\nabla \beta, v_2) p_2 dx + \int_{R^n} (\nabla \beta, v_2) q_2 dx \right| \\ &\leq \frac{C}{N} \|u_1\|_n \|\nabla u_2\|_2 \|u_2\|_2 + C \|\beta u_1\|_n \|\nabla u_2\|_2 \|\nabla u_2\|_2. \end{aligned} \quad (5.17)$$

Plugging this inequality into (5.15) and integrating over  $[0, t]$ , we complete the proof of the lemma.  $\square$

Applying Lemma 5.1 and following the procedure adapted in [1], we can similarly prove the next two lemmas.

**LEMMA 5.2.** (1) Let  $u'_1(0) = (v'_1(0), b'_1(0))$  be the partition by Lemma 3.1 such that Lemma 4.2 holds. There is a  $T > 0$ , depending only on the norm of  $u'_1(0)$ ,

$$\|u'_1(0)\| = \|u'_1(0)\|_n + \|u'_1(0)\|_{n/\beta}, \quad 0 < \beta < 1, \tag{5.18}$$

such that  $u_1(x, t)$ , as a family depending on parameter  $\varepsilon$  and  $0 < t < T$ , is compact in  $L^n$ ;

(2) the size of  $T$  is determined by

$$T^{(1-\beta)/2} \left( s^{1-(\beta p/n)} \|u(0)\|_p^{\beta p/n} + s^{1-(\beta/n)} \|u(0)\|_p^{p/n} \right) < \varepsilon_0; \tag{5.19}$$

(3) the following inequalities hold

$$\begin{aligned} \|u_1^*\|_n &\leq c_1(\varepsilon_0) \left( s^{1-(\beta p/n)} \|u(0)\|_p^{\beta p/n} + s^{1-(p/n)} \|u(0)\|_p^{p/n} \right), \\ \|u_1^*\|_p &\leq c_2(\varepsilon_0) \left( s^{1-(\beta p/n)} \|u(0)\|_p^{\beta p/n} + \|u^s(0)\|_p \right). \end{aligned} \tag{5.20}$$

**LEMMA 5.3.** The solution  $u_2(x, t) = (v_2, b_2)$  of (3.12), (3.14), (3.15), and (3.16), as a family depending on the parameter  $\varepsilon$ , contains a subfamily that converges in  $L^2$  of any subset  $S_T$ , for  $n = 3, 4, T > 0$ .

We are now ready to state and prove the main result of this paper.

**THEOREM 5.1.** Assume that the initial data  $(v_0, b_0) \in L^p(R^n)$ ,  $2 < p < n$ ,  $n = 3, 4$ ,  $\text{div } v_0 = \text{div } b_0 = 0$ . Then there exists a weak solution  $u(x, t) = (v(x, t), b(x, t))$  of (1.1) and (1.2) for all time  $t$ , such that, for  $0 < t < T$ , where  $T$  can be arbitrarily large, we have

$$\|u\|_{p,2} < C, \tag{5.21}$$

where the constant  $C$  depends on  $T, \|u_0\|_p$ .

**PROOF.** From Lemmas 5.2 and 5.3, we have a sequence of solutions  $u_{1m}, u_{2m}$  of (3.11), (3.12), (3.13), (3.14), (3.15), and (3.16) such that  $u_{1m}, u_{2m}$  converge in  $L^n(S_T), L^2(S_T)$  to  $u_1, u_2$ , respectively. Sending  $m$  to  $\infty$ , we see that  $u_1 + u_2$  is a weak solution of (3.9) for some  $T > 0$ .

By Lemma 5.2, we have

$$\|u_{1m}^*\|_p \leq c_2(\varepsilon_0) \left( s^{1-(\beta p/n)} \|u(0)\|_p^{\beta p/n} + \|u^s(0)\|_p \right), \quad [0, T]. \tag{5.22}$$

Fatou's theorem implies that

$$\|u_1^*\|_{p,2} \leq T^{1/2} c_2(\varepsilon_0) \left( s^{1-(\beta p/n)} \|u(0)\|_p^{\beta p/n} + \|u^s(0)\|_p \right), \quad [0, T]. \tag{5.23}$$

Now, for  $u_{2m}$ , from a priori estimate for  $u_{2m}$ , we have

$$\begin{aligned} \|u_{2m}\|_{p,2} &\leq c_1 (\|\nabla u_{2m}\|_{2,2} + \|u_{2m}\|_2) \\ &\leq c_2 \|u_{2m}(0)\|_2 \leq c_2 s^{1-(p/2)} \|u(0)\|_p^{p/2}. \end{aligned} \tag{5.24}$$

Fatou's theorem implies that

$$\|u_2\|_{p,2} \leq c_2 s^{1-(p/2)} \|u(0)\|_p^{p/2}. \tag{5.25}$$

(5.23) and (5.25) implies (5.21).

Due to a priori estimates, we can extend the interval of existence of solution  $u$  from  $[0, T]$  to  $[T, T_1]$ , from  $[T, T_1]$  to  $[T_1, T_2]$ , and so on in such a way that, in each step,

we make sure that  $T_{k+1} - T_k > \delta_0$ —a fixed constant. Therefore, we obtain the weak solution  $u$  for all  $t$ .  $\square$

For  $n \geq 3$ , adapting Calderón's approach [1], one can also prove the global existence result for system (1.1) and (1.2) as long as the  $L^n$  norm of the initial data is suitably small.

#### REFERENCES

- [1] C. P. Calderón, *Existence of weak solutions for the Navier-Stokes equations with initial data in  $L^p$* , Trans. Amer. Math. Soc. **318** (1990), no. 1, 179–200. MR 90k:35199. Zbl 707.35118.
- [2] F. Chen, P. Wang, and C. Qu, *On the differential system governing flows in magnetic field with data in  $L^p$* , Internat. J. Math. Math. Sci. **21** (1998), no. 2, 299–30. Zbl 908.35058.
- [3] P. Constantin and C. Foias, *Navier-Stokes equations*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, 1988. MR 90b:35190. Zbl 687.35071.
- [4] E. B. Fabes, B. F. Jones, and N. M. Riviere, *The initial value problem for the Navier-Stokes equations with data in  $L^p$* , Arch. Rational Mech. Anal. **45** (1972), 222–240. MR 47 5463. Zbl 254.35097.
- [5] J. K. Hale, *Asymptotic behavior of dissipative systems*, Mathematical Surveys and Monographs, vol. 25, American Mathematical Society, Providence, RI, 1988. MR 89g:58059. Zbl 642.58013.
- [6] D. Henry, *Geometric theory of semilinear parabolic equations*, Lecture Notes in Mathematics, vol. 840, Springer-Verlag, Berlin, Heidelberg, New York, 1981. MR 83j:35084. Zbl 456.35001.
- [7] R. Hide, *On planetary atmospheres and interiors*, Mathematical Problems in the Geophysical Sciences I (Providence, RI) (W. H. Reid, ed.), Amer. Math. Soc., 1971, pp. 229–353.
- [8] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math. **63** (1934), 193–248.
- [9] C. Qu, S. Song, and P. Wang, *On the equations for the flow and the magnetic field within the earth*, J. Math. Anal. Appl. **187** (1994), no. 3, 1003–1018. MR 95k:86009. Zbl 812.76091.
- [10] M. Sermange and R. Temam, *Some mathematical questions related to the MHD equations*, Commun. Pure Appl. Math. **36** (1983), no. 5, 635–664. MR 85k:76042. Zbl 524.76099.
- [11] R. Temam, *Navier-Stokes equations and nonlinear functional analysis*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 41, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, 1983. MR 86f:35152. Zbl 522.35002.
- [12] ———, *Infinite-dimensional dynamical systems in mechanics and physics*, Applied Mathematical Sciences, vol. 68, Springer-Verlag, New York, Heidelberg, Berlin, 1988. MR 89m:58056. Zbl 662.35001.

BATES AND CHEN: DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UTAH 84602, USA

WANG: DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, SCHUYLKILL HVEN, PA 17972, USA

## Special Issue on Time-Dependent Billiards

### Call for Papers

This subject has been extensively studied in the past years for one-, two-, and three-dimensional space. Additionally, such dynamical systems can exhibit a very important and still unexplained phenomenon, called as the Fermi acceleration phenomenon. Basically, the phenomenon of Fermi acceleration (FA) is a process in which a classical particle can acquire unbounded energy from collisions with a heavy moving wall. This phenomenon was originally proposed by Enrico Fermi in 1949 as a possible explanation of the origin of the large energies of the cosmic particles. His original model was then modified and considered under different approaches and using many versions. Moreover, applications of FA have been of a large broad interest in many different fields of science including plasma physics, astrophysics, atomic physics, optics, and time-dependent billiard problems and they are useful for controlling chaos in Engineering and dynamical systems exhibiting chaos (both conservative and dissipative chaos).

We intend to publish in this special issue papers reporting research on time-dependent billiards. The topic includes both conservative and dissipative dynamics. Papers discussing dynamical properties, statistical and mathematical results, stability investigation of the phase space structure, the phenomenon of Fermi acceleration, conditions for having suppression of Fermi acceleration, and computational and numerical methods for exploring these structures and applications are welcome.

To be acceptable for publication in the special issue of Mathematical Problems in Engineering, papers must make significant, original, and correct contributions to one or more of the topics above mentioned. Mathematical papers regarding the topics above are also welcome.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	December 1, 2008
First Round of Reviews	March 1, 2009
Publication Date	June 1, 2009

### Guest Editors

**Edson Denis Leonel**, Departamento de Estatística, Matemática Aplicada e Computação, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, Avenida 24A, 1515 Bela Vista, 13506-700 Rio Claro, SP, Brazil ; [edleonel@rc.unesp.br](mailto:edleonel@rc.unesp.br)

**Alexander Loskutov**, Physics Faculty, Moscow State University, Vorob'evy Gory, Moscow 119992, Russia; [loskutov@chaos.phys.msu.ru](mailto:loskutov@chaos.phys.msu.ru)