

ASYMPTOTIC TRACTS OF HARMONIC FUNCTIONS III

KARL F. BARTH

Department of Mathematics
Syracuse University
Syracuse, NY 13244, USA
e-mail: kfbarth@mailbox.syr.edu

DAVID A. BRANNAN

The Open University
Department of Pure Mathematics
Milton Keynes MK7 6AA, United Kingdom
e-mail: D.A.Brannan@open.ac.uk

(Received June 30, 1995)

ABSTRACT. A tract (or asymptotic tract) of a real function u harmonic and nonconstant in the complex plane \mathcal{C} is one of the n_c components of the set $\{z : u(z) \neq c\}$, and the order of a tract is the number of non-homotopic curves from any given point to ∞ in the tract. The authors prove that if $u(z)$ is an entire harmonic polynomial of degree n , if the critical points of any of its analytic completions f lie on the level sets $\tau_j = \{z : u(z) = c_j\}$, where $1 \leq j \leq p$ and $p \leq n - 1$, and if the total order of all the critical points of f on τ_j is denoted by σ_j , then

$$\{n_c : c \in \mathbb{R}\} = \{n + 1\} \cup \{n + 1 + \sigma_j : 1 \leq j \leq p\}.$$

KEY WORDS AND PHRASES: Asymptotic tracts, harmonic functions.

1991 SUBJECT CLASSIFICATION CODES: Primary: 31A05; Secondary: 30C35.

1. INTRODUCTION

This paper continues a study, begun in [1] and [2], of the asymptotic tracts of functions harmonic in \mathcal{C} (entire harmonic functions).

Definition 1. An asymptotic tract (or tract) of a real function $u(z)$ harmonic and nonconstant in \mathcal{C} is a component of the set $\{z : u(z) \neq c\}$ for some real number c .

It was shown in [1] that each tract T is necessarily simply-connected and unbounded, and that u is necessarily unbounded in each tract T ; in addition, ∞ is an accessible boundary point (in \mathcal{C}) of each tract T . The local mapping properties of analytic functions show that the set $\{z : u(z) \neq c\}$ consists of a finite or countable number of curves which are locally analytic, except at the zeros of $f'(z)$ (where f' is any analytic completion of u)—where the set $\{z : u(z) = c\}$ branches. Observe that the angle between the ‘branches’ must be equal to $2\pi/n$ for some $n \geq 1$.

We continue the study of harmonic polynomials in the plane initiated in [3], where it was shown that, if $u(z)$ is a harmonic polynomial in \mathcal{C} of degree n , then the number, k , of tracts of u satisfies the sharp inequality

$$n + 1 \leq k \leq 2n \tag{1}$$

A special case of our results, putting Example 2 together with Theorem 1, shows that, given any pair of positive integers n and k that satisfy the inequality (1) there is a harmonic polynomial $u(z)$ of degree n with k tracts. This is stronger than [3, Theorem 3] where it was shown that there exists a harmonic polynomial of degree n that has $2n$ tracts for the case $c = 0$. We also discover a restriction, for each given harmonic polynomial $u(z)$ in \mathcal{C} , on the number of tracts of $u(z) - c$, as the constant c varies over \mathbb{R} .

Definition 2. An unbounded simply-connected domain T in \mathcal{C} is said to be branched of order n_T (possibly $n_T = +\infty$) if it has the following property: There exists a family \mathcal{T}_T of n_T non-homotopic (in T) and disjoint (except for the end-point z_T) Jordan curves in T connecting some fixed point in T , z_T say, to ∞ ; in addition, any Jordan curve in T joining z_T to ∞ is homotopic

(in T) to one of the elements of \mathcal{T}_T . If $n_T = 1$, we say that T is unbranched; if $n_T < +\infty$, we say that T is finitely branched; if $n_T = +\infty$, we say that T is infinitely-branched.

2. NUMBERS OF TRACTS

Let $u(z)$ be an entire harmonic polynomial of degree n . Then, if $z = re^{i\theta}$, we have that

$$u(z) = a_n r^n \cos(n\theta + \theta_n) + O(r^{n-1}), \text{ where } a_n \neq 0. \quad (2)$$

It follows that near ∞ there must be on $\{z : |z| = r\}$ at least n arcs (each of angular length about π/n) on which $u(z) > 0$, and at least n arcs (each of angular length about π/n) on which $u(z) < 0$. Since u is a polynomial of degree n and so can have at most $2n$ zeros on $\{z : |z| = r\}$, it follows that for sufficiently large r there are precisely n arcs of each type. Also, it is easy to prove that the boundaries separating the $2n$ regions comprising $\{z : |z| = r, u(z) \neq 0\}$ tend to radial lines of angular separation π/n as $r \rightarrow +\infty$.

We will denote by n_c the number of components of the set $\{z : u(z) - c \neq 0\}$. It will be useful to examine how n_c varies with c . For sufficiently large r , the set $\{z : |z| > R\} \cap \{z : u(z) \neq 0\}$ consists of precisely $2n$ unbounded disjoint domains. Then, for such an r , we define

$$M = 1 + \max\{u(z) : |z| \leq r\}. \quad (3)$$

It follows that the set $\{z : u(z) - M \neq 0\}$ has exactly n components in which $\{z : u(z) - M > 0\}$ and exactly one component in which $\{z : u(z) - M < 0\}$. Thus $n_M = n + 1$. Also, it follows from the Phragmen-Lindelöf Principle that $n_c = n + 1$ when $c > M$. We now look at how n_c varies as c decreases from the value M . The components (tracts) of $\{z : u(z) - c \neq 0\}$ vary continuously with c , in terms of kernel convergence. Hence, as c decreases, n_c is an integer and varies continuously with c (hence remains constant)—except at those values of c for which a critical point of the analytic completion of u lies on the set $\{z : u(z) = c\}$.

Now two tracts of $u(z) - c$ in which $u(z) - c$ has opposite signs can never lie in a single tract of $u(z) - c_1$, for $c_1 \neq c$, since u is unbounded in any tract; however their boundaries may meet in a point or in an arc. No two tracts of $u(z) - c$ can have the property that their boundaries meet in a set with more than one component: for, if they did, then there would be a bounded (non-empty) domain on whose boundary $u(z) = c$, and so we would have $u(z) \equiv c$ in \mathcal{C} .

Suppose that T_1 and T_2 are two tracts of $u(z) - c$ in which $u(z) - c > 0$; we will call such tracts upper tracts (for the value c). (Lower tracts are defined similarly.) It may be that $\partial T_1 \cap \partial T_2 = \emptyset$. However we cannot have a situation where $\partial T_1 \cap \partial T_2$ contains an arc in \mathcal{C} , by the Maximum Principle. It follows, then, that, if ∂T_1 meets ∂T_2 , the set $\partial T_1 \cap \partial T_2$ must be a singleton.

If T_1 and T_2 are both upper tracts or both lower tracts for which $\partial T_1 \cap \partial T_2 = \{z_0\}$, then there must exist an equal number of upper and lower tracts whose boundaries contain z_0 . Since z_0 must thus be a critical point of any analytic completion of u , there can be at most $(n - 1)$ such points z_0 (since u is a polynomial of degree n). Note also that, as c decreases, the upper tracts individually increase in size. Hence their total number must decrease as c decreases.

Our main result in this Section is the following.

Theorem 1. *Let $u(z)$ be an entire harmonic polynomial of degree n . Let the critical points of any of its analytic completions f lie on the level sets $\tau_j = \{z : u(z) = c_j\}$, where $1 \leq j \leq p$ and $p \leq n - 1$, and let the total order of all the critical points of f on τ_j be denoted by σ_j . (In particular, $\sum_{j=1}^p \sigma_j = n - 1$.) Then $\{n_c : c \in \mathbb{R}\} = \{n + 1\} \cup \{n + 1 + \sigma_j : 1 \leq j \leq p\}$.*

Proof. Let f be any analytic completion of u .

Case 1. All the critical points of f lie on different level sets for u .

Assume first that all the critical points of f are simple; then we may choose our notation so that they lie on the level sets $\tau_j = \{z : u(z) = c_j\}$, $1 \leq j \leq n - 1$, where $c_1 > c_2 > \dots > c_{n-1}$. Then, by the previous comments, for $c > c_1$ (for example, when $c = M$ (see (3))), we have $n_c = n + 1$ and there are n upper tracts of u and one lower tract. Next, $n_{c_1} = n + 2$ and there are, for the value $c = c_1$, n upper tracts and two lower tracts (the lower tract has 'split' in two). Finally, for $c_1 > c > c_2$, we have $n_c = n + 1$, and there are $(n - 1)$ upper tracts (two upper tracts have 'combined') and 2 lower tracts.

As c decreases further, a similar argument holds for each c_j in turn, $2 \leq j \leq n - 1$. For $c_{j-1} > c > c_j$, we have $n_c = n + 1$ and there are $(n + 1 - j)$ upper tracts and j lower tracts; when $c = c_j$, we have $n_{c_j} = n + 2$ and there are $(n + 1 - j)$ upper tracts and $(j + 1)$ lower tracts; and, for $c_j > c > c_{j+1}$ (with the convention that $c_n = -\infty$), we have $n_c = n + 1$ and there are $(n - j)$ upper tracts and $(j + 1)$ lower tracts.

Assume next that the critical points of f are not necessarily simple. First, suppose that the level set $\{z : u(z) = c_j\}$, for some particular value of j , contains a critical point of f (at z_j where f' has a zero of order b_j). Let I be an open interval of \mathbb{R} that contains c_j but contains no other c 's corresponding to critical points of f . Then, for a sufficiently small neighborhood \mathcal{U} of z_j there are $(2b_j + 2)$ tracts of $u(z) - c_j$ that meet \mathcal{U} , namely $(b_j + 1)$ upper tracts and $(b_j + 1)$ lower tracts. However, when $c > c_j$, $c \in I$ and $c - c_j$ is sufficiently small, there are only $(b_j + 2)$ tracts of $u(z) - c$ that meet \mathcal{U} , namely $(b_j + 1)$ upper tracts and 1 lower tract; similarly, when $c < c_j$, $c \in I$ and $c_j - c$ is sufficiently small, there are $(b_j + 2)$ tracts of $u(z) - c$ that meet \mathcal{U} , namely $(b_j + 1)$ lower tracts and 1 upper tract.

Now consider the level set $\{z : u(z) = c\}$ for an arbitrary c . Since, except for values of c corresponding to critical points of f (and even then locally only in small neighborhoods of the critical points themselves) the tracts vary continuously with c (in the sense of kernel convergence), it follows from the above argument that there is some number N such that, for $|c - c_j|$ sufficiently small and non-zero, we have $n_c = N + 1$ whereas $n_{c_j} = N + 1 + b_j$. But $n_M = n + 1$, so that we must have $N = n$. This completes the proof of Case 1 of the theorem.

Case 2. More than one critical point of f lies on a given level set for u .

Assume first that, for some c_j , the level set $\{z : u(z) = c_j\}$ contains just two branch points, z_1 and z_2 , of orders b_1 and b_2 respectively, and that z_1 and z_2 lie on different components, C_1 and C_2 respectively, of $\{z : u(z) = c_j\}$; thus $C_1 \cap C_2 = \emptyset$. It follows that there exists some Jordan curve from ∞ to ∞ that separates C_1 from C_2 ; this curve can be chosen to lie either in a single component of $\{z : u(z) > c_j\}$ or in a single component of $\{z : u(z) < c_j\}$. By considering the local behavior of u near z_1 and z_2 , and by using the fact that components of $\{z : u(z) - d \neq 0\}$ vary continuously with d (except when their boundaries coalesce), it follows that, when $|d - c_j|$ is sufficiently small, we have $n_d = n + 1$ and $n_{c_j} = (n + 1) + b_1 + b_2$. A similar argument works in the case of more than two branch points on a single level set of u , so long as each such branch point lies on a different component of that level set.

Assume next that, for some c_j , the level set $\{z : u(z) = c_j\}$ contains just two branch points, z_1 and z_2 , of orders b_1 and b_2 respectively (corresponding to zeros of f' of these orders), and that z_1 and z_2 lie on the same component, C , of $\{z : u(z) = c_j\}$. Then there is a Jordan subarc Γ of C joining z_1 to z_2 ; let z' be any interior point of this subarc. Since C cannot contain any closed Jordan curves, it follows that there are precisely two tracts, T_1 and T_2 , say, of $u(z) - c_j$ that have $\Gamma - \{z_1, z_2\}$ as part of their boundaries; we may assume that $u(z) > c_j$ in T_1 and so that $u(z) < c_j$ in T_2 . Similar considerations also show that there is a Jordan curve J_1 in $T_1 \cup \{z'\}$ that joins z' to ∞ inside T_1 , and a Jordan curve J_2 in $T_2 \cup \{z'\}$ that joins z' to ∞ inside T_2 .

We define $J' = J_1 \cup J_2$. Then J' plays the same role as J did earlier (when it separated C_1 from C_2), and a similar argument to the previous one shows that

$$n_d = \begin{cases} n + 1, & \text{if } d \neq c_j, \text{ and } |d - c_j| \text{ sufficiently small,} \\ n + 1 + (b_1 + b_2), & \text{if } d = c_j. \end{cases} \quad (4)$$

Again a similar argument can be used even when there are more than two branch points on the same component of the level set.

The result of this theorem is stronger than [3, Theorem 1], where it was shown that $\{n_c : c \in \mathbb{R}\}$ is a subset of $\{n + 1, n + 2, \dots, 2n\}$.

Notice that for the function $u_1(z) = \operatorname{Re}(z^n)$ we have $n_0 = 2n$ and $n_1 = n + 1$, and that in fact $\{n_c : c \in \mathbb{R}\} = \{n + 1, 2n\}$. The next two examples show that, while this particular function u_1 is extremal in a certain sense, the conclusion of Theorem 1 concerning the range of possible values of n_c (as c varies) is best-possible.

Example 1. *There exists a harmonic polynomial u of degree n for which $\{n_c : c \in \mathbb{R}\} = \{n + 1, n + 2\}$, and all the critical points of any analytic completion f of u are simple and lie on different level sets of u .*

Let $u(z) = \operatorname{Re}(z^n - Az)$, for a complex number A yet to be specified. The analytic completion $f(z) \equiv z^n - Az$ of u has critical points where $nz^{n-1} - A = 0$; that is, where

$$z = z_k = \left(\frac{A}{n}\right)^{\frac{1}{n-1}} \exp\left(\frac{2\pi i k}{n-1}\right), \quad 1 \leq k \leq n-1. \quad (5)$$

Now $u(z_k) = \operatorname{Re}(\frac{z_k A(1-n)}{n})$; and it follows that, if A is chosen with $|A| = 1$ and with $\arg A$ not a rational multiple of 2π , then all the values of $\{u(z_k)\}_1^{n-1}$ are distinct. Thus u has the desired properties.

Example 2. Let p be an integer, such that $1 \leq p \leq n-1$, and let b_1, b_2, \dots, b_p be any integers in $[1, n-2]$ for which $\sum_{j=1}^p b_j = n-1$. There exists a harmonic polynomial $u(z)$ of degree n with the properties that any analytic completion f of u has critical points of orders b_1, b_2, \dots, b_p , and that all these critical points lie on different level sets of u . Hence (from Theorem 1)

$$\{n_c : c \in \mathbb{R}\} = \{n+1\} \cup \{n+1+b_j : 1 \leq j \leq p\}.$$

First, let f_1 be the polynomial given by $f_1(0) = 0$ and

$$f_1'(z) = (z-1)^{b_1}(z-a_2)^{b_2}z^{(n-1)-(b_1+b_2)}, \quad (6)$$

where a_2 is chosen to be either $\frac{1}{2}$ or to be very close to $\frac{1}{2}$; in particular, we make our choice of a_2 to ensure that $u_1(1) \neq 0$, where $u_1(z) \equiv \operatorname{Ref}_1(z)$. It follows from Rolle's Theorem that the points 1 and a_2 lie on different level sets of u_1 .

Next, let f_2 be the polynomial given by $f_2(0) = 0$ and

$$f_2'(z) = (z-1)^{b_1}(z-a_2)^{b_2}(z-a_3)^{b_3}z^{(n-1)-(b_1+b_2+b_3)}, \quad (7)$$

where a_3 is positive but small. By continuity arguments (on f) we see that we may choose a_3 sufficiently near to 0 that 1 and a_2 lie on different level sets of $u_2(z) \equiv \operatorname{Ref}_2(z)$.

We have to check that a_3 can be chosen so that a_3 lies on a different level set of u_2 from those that contain either 1 or a_2 . But, if a_3 is sufficiently small, we have that

$$u_2(1) \approx \int_0^1 (t-1)^{b_1}(t-a_2)^{b_2}t^{(n-1)-(b_1+b_2)} dt, \quad (8)$$

$$u_2(a_2) \approx \int_0^{a_2} (t-1)^{b_1}(t-a_2)^{b_2}t^{(n-1)-(b_1+b_2)} dt, \quad (9)$$

and

$$u_2(a_3) \approx (-1)^{b_1+b_2+b_3}(a_2)^{b_2}(a_3)^{(n-1)-(b_1+b_2)} \int_0^1 (1-x)^{b_3}x^{(n-1)-(b_1+b_2+b_3)} dx;$$

hence, for all sufficiently small a_3 , the values of $u_2(1)$, $u_2(a_2)$ and $u_2(a_3)$ are all distinct.

A similar argument shows, after a further $(p-3)$ steps, that the polynomial $u(z) \equiv \operatorname{Ref}(z)$, where $f(0) = 0$ and

$$f'(z) = (z-1)^{b_1}(z-a_2)^{b_2}(z-a_3)^{b_3}(z-a_4)^{b_4} \dots (z-a_p)^{b_p}, \quad (10)$$

and the sequence $\{a_j\}_{j=3}^p$ decreases to 0 sufficiently quickly, has the desired properties.

Finally, as was mentioned in the Introduction, suppose n and k are positive integers such that $n+1 \leq k \leq 2n$. If $k = n+1$, we see from Example 1 that there exists a harmonic polynomial $u(z)$ such that $n_c = \{n+1, n+2\}$. If $n+1 < k \leq 2n$ and we set $b_1 = k - (n+1)$ and $b_2 = 2n - k$, Example 2 shows that there exists a harmonic polynomial $u(z)$ such that

$$\{n_c\} = \{n+1, n+1+b_1, n+1+b_2\} = \{n+1, k, 3n+1-k\}.$$

ACKNOWLEDGMENT. The first author gratefully acknowledges support from the Science and Engineering Research Council of the United Kingdom.

REFERENCES

1. K. F. Barth and D. A. Brannan, *Asymptotic tracts of harmonic functions I*, Acad. Sci. Fenn. Series A. I. Math. **11** (1986), 215–232.
2. ———, *Asymptotic tracts of harmonic functions II*, Proc. Edinburgh Math. Soc. **38** (1995), 35–52.
3. W. K. Hayman D. A. Brannan, W. H. J. Fuchs and Ü. Kuran, *A characterisation of harmonic polynomials in the plane*, Proc. London Math. Soc. (3) **32** (1976), 213–229.

Special Issue on Modeling Experimental Nonlinear Dynamics and Chaotic Scenarios

Call for Papers

Thinking about nonlinearity in engineering areas, up to the 70s, was focused on intentionally built nonlinear parts in order to improve the operational characteristics of a device or system. Keying, saturation, hysteretic phenomena, and dead zones were added to existing devices increasing their behavior diversity and precision. In this context, an intrinsic nonlinearity was treated just as a linear approximation, around equilibrium points.

Inspired on the rediscovering of the richness of nonlinear and chaotic phenomena, engineers started using analytical tools from “Qualitative Theory of Differential Equations,” allowing more precise analysis and synthesis, in order to produce new vital products and services. Bifurcation theory, dynamical systems and chaos started to be part of the mandatory set of tools for design engineers.

This proposed special edition of the *Mathematical Problems in Engineering* aims to provide a picture of the importance of the bifurcation theory, relating it with nonlinear and chaotic dynamics for natural and engineered systems. Ideas of how this dynamics can be captured through precisely tailored real and numerical experiments and understanding by the combination of specific tools that associate dynamical system theory and geometric tools in a very clever, sophisticated, and at the same time simple and unique analytical environment are the subject of this issue, allowing new methods to design high-precision devices and equipment.

Authors should follow the Mathematical Problems in Engineering manuscript format described at <http://www.hindawi.com/journals/mpe/>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2009
First Round of Reviews	May 1, 2009
Publication Date	August 1, 2009

Guest Editors

José Roberto Castilho Piqueira, Telecommunication and Control Engineering Department, Polytechnic School, The University of São Paulo, 05508-970 São Paulo, Brazil; piqueira@lac.usp.br

Elbert E. Neher Macau, Laboratório Associado de Matemática Aplicada e Computação (LAC), Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil ; elbert@lac.inpe.br

Celso Grebogi, Department of Physics, King's College, University of Aberdeen, Aberdeen AB24 3UE, UK; grebogi@abdn.ac.uk