

## Research Article

# Convergence and Divergence of the Solutions of a Neutral Difference Equation

**G. E. Chatzarakis and G. N. Miliaras**

*Department of Electrical Engineering Educators, School of Pedagogical and Technological Education (ASPETE), N. Heraklion, 14121 Athens, Greece*

Correspondence should be addressed to G. E. Chatzarakis, [geaxatz@otenet.gr](mailto:geaxatz@otenet.gr)

Received 9 June 2011; Accepted 7 August 2011

Academic Editor: Alexandre Carvalho

Copyright © 2011 G. E. Chatzarakis and G. N. Miliaras. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the asymptotic behavior of the solutions of a neutral type difference equation of the form  $\Delta[x(n) + cx(\tau(n))] + p(n)x(\sigma(n)) = 0$ , where  $\tau(n)$  is a general retarded argument,  $\sigma(n)$  is a general deviated argument (retarded or advanced),  $c \in \mathbb{R}$ ,  $(p(n))_{n \geq 0}$  is a sequence of positive real numbers such that  $p(n) \geq p$ ,  $p \in \mathbb{R}_+$ , and  $\Delta$  denotes the forward difference operator  $\Delta x(n) = x(n+1) - x(n)$ . Also, we examine the asymptotic behavior of the solutions in case they are continuous and differentiable with respect to  $c$ .

## 1. Introduction

Neutral type differential equations are differential equations in which the highest-order derivative of the unknown function appears in the equation both with and without delays (or delays advanced). See Driver [1], Bellman and Cooke [2], and Hale [3] for questions of existence, uniqueness, and continuous dependence.

It is to be noted that, in general, the theory of neutral differential equations presents extra complications, and basic results which are true for delay differential equations may not be true for neutral equations. For example, Snow [4] has shown that, even though the characteristic roots of a neutral differential equation may all have negative real parts, it is still possible for some solutions to be unbounded.

The discrete counterparts of neutral differential equations are called *neutral difference equations*, and it is a well-known fact that there is a similarity between the qualitative theories of neutral differential equations and neutral difference equations.

Besides its theoretical interest, strong interest in the study of the asymptotic and oscillatory behavior of solutions of neutral type equations (difference or differential) is motivated by the fact that they arise in many areas of applied mathematics, such as circuit

theory [5, 6], bifurcation analysis [7], population dynamics [8, 9], stability theory [10], and dynamical behavior of delayed network systems [11]. See, also, Driver [12], Hale [3], Brayton and Willoughby [13], and the references cited therein. This is the reason that during the last few decades these equations are in the main interest of the literature.

In the present paper, we are interested in the first-order neutral type difference equation of the form

$$\Delta[x(n) + cx(\tau(n))] + p(n)x(\sigma(n)) = 0, \quad n \geq 0 \quad (\text{E})$$

where  $(p(n))_{n \geq 0}$  is a sequence of positive real numbers such that  $p(n) \geq p$ ,  $p \in \mathbb{R}_+$ ,  $c \in \mathbb{R}$ ,  $(\tau(n))_{n \geq 0}$  is an increasing sequence of integers which satisfies

$$\tau(n) \leq n - 1 \quad \forall n \geq 0, \quad \lim_{n \rightarrow \infty} \tau(n) = +\infty, \quad (1.1)$$

and  $(\sigma(n))_{n \geq 0}$  is an increasing sequence of integers such that

$$\sigma(n) \leq n - 1 \quad \forall n \geq 0, \quad \lim_{n \rightarrow \infty} \sigma(n) = +\infty, \quad (1.2)$$

or

$$\sigma(n) \geq n + 1 \quad \forall n \geq 0. \quad (1.3)$$

Define

$$\begin{aligned} k_1 &= -\min_{n \geq 0} \tau(n), & k_2 &= -\min_{n \geq 0} \sigma(n), \\ k &= \max\{k_1, k_2\}. \end{aligned} \quad (1.4)$$

(Clearly,  $k$  is a positive integer.)

By a *solution* of the neutral type difference equation (E), we mean a sequence of real numbers  $(x(n))_{n \geq -k}$  which satisfies (E) for all  $n \geq 0$ . It is clear that, for each choice of real numbers  $c_{-k}, c_{-k+1}, \dots, c_{-1}, c_0$ , there exists a unique solution  $(x(n))_{n \geq -k}$  of (E) which satisfies the initial conditions  $x(-k) = c_{-k}$ ,  $x(-k+1) = c_{-k+1}, \dots, x(-1) = c_{-1}$ ,  $x(0) = c_0$ .

A solution  $(x(n))_{n \geq -k}$  of the neutral type difference equation (E) is called *oscillatory* if for every positive integer  $n$  there exist  $n_1, n_2 \geq n$  such that  $x(n_1)x(n_2) \leq 0$ . In other words, a solution  $(x(n))_{n \geq -k}$  is oscillatory if it is neither eventually positive nor eventually negative. Otherwise, the solution is said to be *nonoscillatory*.

In the special case where  $\tau(n) = n - a$  and  $\sigma(n) = n - b$ ,  $a, b \in \mathbb{N}$ , (E) takes the form

$$\Delta[x(n) + cx(n - a)] + p(n)x(n - b) = 0, \quad n \geq 0. \quad (\text{E}_1)$$

Equation (E) represents a discrete analogue of the neutral type differential equations

$$\frac{d}{dt}[x(t) + cx(\tau(t))] + p(t)x(\sigma(t)) = 0, \quad t \geq 0, \quad (\text{E}_2)$$

which, in the special case where  $\tau(t) = t - \tau_0$  and  $\sigma(t) = t - \sigma_0$ ,  $\tau_0, \sigma_0 \in \mathbb{R}_+$ , takes the form (see e.g., [14, 15])

$$\frac{d}{dt}[x(t) + cx(t - \tau_0)] + p(t)x(t - \sigma_0) = 0, \quad t \geq 0. \quad (E_3)$$

The search for the asymptotic behavior and, especially, for oscillation criteria and stability of neutral type (difference or differential) equations has received a great attention in the last few years. Hence, a large number of related papers have been published. See [4, 9, 10, 14–52] and the references cited therein. Most of these papers are concerning the special case of the delay difference equations (E<sub>1</sub>) and (E<sub>3</sub>) where the algebraic characteristic equation gives useful information about oscillation and stability.

The purpose of this paper is to investigate the convergence and divergence of the solutions of (E) in the case of a general delay argument  $\tau(n)$  and of a general deviated (retarded or advanced) argument  $\sigma(n)$ .

## 2. Some Preliminaries

Assume that  $(x(n))_{n \geq -k}$  is a nonoscillatory solution of (E). Then it is either eventually positive or eventually negative. As  $(-x(n))_{n \geq -k}$  is also a solution of (E), we may (and do) restrict ourselves only to the case where  $x(n) > 0$  for all large  $n$ . Let  $n_1 \geq -k$  be an integer such that  $x(n) > 0$  for all  $n \geq n_1$ . Then, there exists  $n_0 \geq n_1$  such that

$$x(\tau(n)), x(\sigma(n)) > 0 \quad \text{for every } n \geq n_0. \quad (2.1)$$

Set

$$z(n) = x(n) + cx(\tau(n)). \quad (2.2)$$

Then, in view of (E) and taking into account the fact that  $p(n) \geq p > 0$ , we have

$$\Delta z(n) + p(n)x(\sigma(n)) = 0 \quad (2.3)$$

or

$$\Delta z(n) = -p(n)x(\sigma(n)) \leq -px(\sigma(n)) < 0 \quad \forall n \geq n_0, \quad (2.4)$$

which means that the sequence  $(z(n))_{n \geq n_0}$  is strictly decreasing, regardless of the value of the real constant  $c$ .

Throughout this paper, we are going to use the following notation:

$$\tau \circ \tau = \tau^2, \quad \tau \circ \tau \circ \tau = \tau^3, \quad \text{and so on.} \quad (2.5)$$

Let the domain of  $\tau$  be the set  $D(\tau) = \mathbb{N}_{n_*} = \{n_*, n_* + 1, n_* + 2, \dots\}$ , where  $n_*$  is the smallest natural number that  $\tau$  is defined with. Then for every  $n > n_*$  it is clear that there exists a natural number  $m(n)$  such that

$$x(\tau^{m(n)}(n)) = x(\tau(n_*)), \quad \lim_{n \rightarrow \infty} m(n) = +\infty \quad (2.6)$$

since  $(m(n))$  is increasing and unbounded function of  $n$ .

The following lemma provides some tools which are useful for the main results.

**Lemma 2.1.** *Assume that  $(x(n))_{n \geq k}$  is a positive solution of (E). Then, one has the following.*

(i) *If  $c \neq 0$  and*

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = S_0 < +\infty, \quad (2.7)$$

*then*

$$\lim_{n \rightarrow \infty} z(n) = cL = c \lim_{n \rightarrow \infty} x(\tau(\sigma(n))). \quad (2.8)$$

(ii) *If*

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = +\infty, \quad (2.9)$$

*then*

$$z(n) < 0 \quad \text{eventually.} \quad (2.10)$$

(iii) *If  $c \geq -1$ , then*

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = S_0 < +\infty. \quad (2.11)$$

(iv) *If  $c < -1$ , then*

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = +\infty. \quad (2.12)$$

*Proof.* Summing up (2.3) from  $n_0$  to  $n$ , we obtain

$$z(n+1) - z(n_0) + \sum_{i=n_0}^n p(i)x(\sigma(i)) = 0 \quad (2.13)$$

or

$$z(n+1) = z(n_0) - \sum_{i=n_0}^n p(i)x(\sigma(i)). \quad (2.14)$$

For the above relation, there are only two possible cases:

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = S_0 < +\infty \quad (2.15a)$$

or

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = +\infty. \quad (2.15b)$$

Assume that (2.15a) holds. Since  $p(n) \geq p > 0$ , we have

$$+\infty > S_0 = \sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) \geq p \sum_{i=n_0}^{\infty} x(\sigma(i)). \quad (2.16)$$

The last inequality guarantees that

$$\sum_{i=n_0}^{\infty} x(\sigma(i)) < +\infty \quad (2.17)$$

and, consequently,

$$\lim_{n \rightarrow \infty} x(\sigma(n)) = 0. \quad (2.18)$$

Also, (2.15a) guarantees that  $\lim_{n \rightarrow \infty} z(n)$  exists as a real number. Now, assume that

$$\lim_{n \rightarrow \infty} z(n) = \ell \in \mathbb{R}, \quad c \neq 0. \quad (2.19)$$

Since  $(z(\sigma(n)))$  is a subsequence of  $(z(n))$ , it is obvious that

$$\lim_{n \rightarrow \infty} z(\sigma(n)) = \ell \quad (2.20)$$

or

$$\lim_{n \rightarrow \infty} [x(\sigma(n)) + cx(\tau(\sigma(n)))] = \ell, \quad (2.21)$$

and, in view of (2.18), we obtain

$$\lim_{n \rightarrow \infty} x(\tau(\sigma(n))) = \frac{\ell}{c} := L. \quad (2.22)$$

Hence,

$$\lim_{n \rightarrow \infty} z(n) = \ell = cL = c \lim_{n \rightarrow \infty} x(\tau(\sigma(n))). \quad (2.23)$$

The proof of part (i) of the lemma is complete.

Assume that (2.15b) holds. Then, by taking limits on both sides of (2.14) we obtain

$$\lim_{n \rightarrow \infty} z(n) = -\infty, \quad (2.24)$$

which, in view of the fact that the sequence  $(z(n))$  is strictly decreasing, means that

$$z(n) < 0 \quad \text{eventually.} \quad (2.25)$$

The proof of part (ii) of the lemma is complete.

Assume that  $-1 \leq c < 0$ , and suppose, for the sake of contradiction, that  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = +\infty$ . Then, in view of part (ii), we have  $z(n) < 0$  eventually. Thus,

$$x(n) < -cx(\tau(n)) \quad (2.26)$$

or

$$x(n) < -c[-cx(\tau^2(n))]. \quad (2.27)$$

Repeating the above procedure we obtain

$$x(n) < (-c)^{m(n)} x(\tau^{m(n)}(n)) = (-c)^{m(n)} x(\tau(n_*)). \quad (2.28)$$

If  $-1 < c < 0$ , clearly,  $(-c)^{m(n)} \rightarrow 0$  since  $m(n) \rightarrow \infty$  as  $n \rightarrow \infty$ . Since  $x(n) > 0$  for all large  $n$ , (2.28) guarantees that

$$\lim_{n \rightarrow \infty} x(n) = 0. \quad (2.29)$$

This implies that

$$\lim_{n \rightarrow \infty} z(n) = 0, \quad (2.30)$$

and consequently  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) < \infty$ , which contradicts our assumption.

If  $c = -1$ , by (2.28) we obtain

$$x(n) < x(\tau(n_*)), \quad (2.31)$$

which means that  $(x(n))$  is bounded and therefore  $(z(n))$  is bounded. Hence,  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) < \infty$ , which contradicts our assumption.

Assume that  $c \geq 0$  and that  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = \infty$ . In view of part (ii), (2.10) holds, that is,  $z(n) < 0$  eventually. This contradicts  $z(n) = x(n) + cx(\tau(n)) > 0$ . Therefore  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = S_0 < +\infty$ . The proof of part (iii) of the lemma is complete.

In the remainder of this proof, it will be assumed that  $c < -1$ .

If (2.15a) holds, that is,  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = S_0 < +\infty$ , then, in view of part (i), (2.8) is satisfied, that is,

$$\lim_{n \rightarrow \infty} z(n) = cL, \quad \text{where } L = \lim_{n \rightarrow \infty} x(\tau(\sigma(n))). \quad (2.32)$$

If  $L > 0$ , then

$$\lim_{n \rightarrow \infty} z(n) = \lim_{n \rightarrow \infty} (x(n) + cx(\tau(n))) = cL < 0. \quad (2.33)$$

Since  $(z(n))$  is strictly decreasing, we have

$$z(n) > cL, \quad (2.34)$$

$$x(n) > -cx(\tau(n)) + cL, \quad (2.35)$$

or

$$x(n) > -c \left[ -cx(\tau^2(n)) + cL \right] + cL. \quad (2.36)$$

Repeating this procedure  $m(n_\ell)$  times we have

$$\begin{aligned} x(n) &> -c \left[ -cx(\tau^2(n)) + cL \right] + cL \\ &= (-c)^2 x(\tau^2(n)) - c^2 L + cL \\ &> \dots > (-c)^{m(n_\ell)} x(\tau^{m(n_\ell)}(n)) + cL - c^2 L + \dots \pm c^{m(n_\ell)} L \\ &= (-c)^{m(n_\ell)} x(\tau(n_\lambda)) + cL \frac{(-c)^{m(n_\ell)} - 1}{-1 - c} \\ &= (-c)^{m(n_\ell)} x(\tau(n_\lambda)) - \frac{cL}{1 + c} \left[ (-c)^{m(n_\ell)} - 1 \right] \\ &= (-c)^{m(n_\ell)} \left[ x(\tau(n_\lambda)) - \frac{cL}{1 + c} \right] + \frac{cL}{1 + c}. \end{aligned} \quad (2.37)$$

Now, if for this index  $\lambda$  we have  $x(\tau(n_\lambda)) - (cL/(1+c)) > 0$ , then, as  $n \rightarrow \infty$ ,  $x(n) \rightarrow \infty$ , which contradicts (2.18). Therefore, there exists an index  $s$  such that

$$x(\tau(n_s)) = x\left(\tau^{m(n_\mu)}(n)\right) \quad (2.38)$$

with

$$x(\tau(n_s)) - \frac{cL}{1+c} < 0. \quad (2.39)$$

Then since  $\lim_{n \rightarrow \infty} z(n) = cL$ , for every  $\varepsilon > 0$  there exists  $n_2(\varepsilon)$  such that

$$z(n) < cL + \varepsilon \quad \text{for every } n \geq \max\{\tau(n_s), n_2\} = n_3, \quad (2.40)$$

where  $\tau(n_s)$  is satisfying the previous inequality.

Hence,

$$x(n) < -cx(\tau(n)) + cL + \varepsilon \quad \text{for every } n \geq n_3, \quad (2.41)$$

or

$$x(n) < -c\left[-cx(\tau^2(n)) + cL + \varepsilon\right] + cL + \varepsilon \quad \text{for every } n \geq n_3. \quad (2.42)$$

Repeating this procedure  $m(n_\mu)$  times we have

$$\begin{aligned} x(n) &< -c\left[-cx(\tau^2(n)) + cL + \varepsilon\right] + cL + \varepsilon \\ &= (-c)^2 x(\tau^2(n)) - c^2L - c\varepsilon + cL + \varepsilon \\ &< \dots < (-c)^{m(n_\mu)} x(\tau(n_s)) - \frac{cL}{1+c} \left[(-c)^{m(n_\mu)} - 1\right] + \frac{(-c)^{m(n_\mu)} - 1}{-1-c} \varepsilon \\ &= (-c)^{m(n_\mu)} \left[x(\tau(n_s)) - \frac{cL}{1+c} - \frac{\varepsilon}{1+c}\right] + \frac{cL}{1+c} + \frac{\varepsilon}{1+c} \end{aligned} \quad (2.43)$$

or

$$x(n) < (-c)^{m(n_\mu)} \left[x(\tau(n_s)) - \frac{cL}{1+c} - \frac{\varepsilon}{1+c}\right] + \frac{cL}{1+c} + \frac{\varepsilon}{1+c}. \quad (2.44)$$

Then, for sufficiently large  $n$  the above inequality gives

$$x(n) < 0, \quad (2.45)$$

which contradicts  $x(n) > 0$ .



If  $L = 0$ , then

$$\lim_{n \rightarrow \infty} z(n) = 0 \quad (2.46)$$

or

$$\lim_{n \rightarrow \infty} (x(n) + cx(\tau(n))) = 0, \quad (2.47)$$

and since  $(z(n))$  is strictly decreasing, it is obvious that  $z(n) > 0$  eventually or

$$x(n) > -cx(\tau(n)). \quad (2.48)$$

In view of (2.6), the last inequality becomes

$$x(n) > (-c)^{m(n)} x(\tau(n_*)), \quad (2.49)$$

and consequently

$$\lim_{n \rightarrow \infty} x(n) \geq \lim_{n \rightarrow \infty} [(-c)^{m(n)} x(\tau(n_*))] = +\infty, \quad (2.50)$$

which contradicts (2.18). Hence, it is clear that  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = +\infty$ . The proof of part (iv) of the lemma is complete.

The proof of the lemma is complete.  $\square$

### 3. Main Results

The asymptotic behavior of the solutions of the neutral type difference equation (E) is described by the following theorem.

**Theorem 3.1.** *For (E) one has the following.*

- (I) *Every nonoscillatory solution is unbounded if  $c < -1$ .*
- (II) *Every solution oscillates if  $c = -1$ .*
- (III) *Every nonoscillatory solution tends to zero if  $-1 < c < 1$ .*
- (IV) *Every nonoscillatory solution is bounded if  $c \geq 1$ .*

*Furthermore, if any solution of (E) is continuous with respect to  $c$ , one has the following.*

- (V) *Every solution is zero if  $c \leq -1$ .*
- (VI) *Every solution tends to zero if  $-1 < c \leq 1$ .*
- (VII) *If, additionally, any solution of (E) has continuous derivatives of any order and convergent Taylor series for every  $c \in \mathbb{R}$ , then the solution is zero.*

*Proof.* Assume that  $(x(n))_{n \geq -k}$  is a nonoscillatory solution of (E). Then it is either eventually positive or eventually negative. As  $(-x(n))_{n \geq -k}$  is also a solution of (E), we may (and do) restrict ourselves only to the case where  $x(n) > 0$  for all large  $n$ . Let  $n_1 \geq -k$  be an integer such that  $x(n) > 0$  for all  $n \geq n_1$ . Then, there exists  $n_0 \geq n_1$  such that

$$x(\tau(n)), x(\sigma(n)) > 0 \quad \forall n \geq n_0, \quad (3.1)$$

which, in view of the previous section, means that the sequence  $(z(n))_{n \geq n_0}$  is strictly decreasing, regardless of the value of the real constant  $c$ .

Assume that  $c < -1$ . Then, in view of part (iv) of Lemma 2.1, we have  $\sum_{i=n_0}^{\infty} p(i) x(\sigma(i)) = +\infty$ , and, consequently, in view of part (ii) of Lemma 2.1,  $z(n) < 0$  eventually. Therefore,

$$\lim_{n \rightarrow \infty} z(n) = -\infty \quad (3.2)$$

or

$$\lim_{n \rightarrow \infty} (x(n) + cx(\tau(n))) = -\infty. \quad (3.3)$$

Since  $c < -1$ , the last relation guarantees that

$$\lim_{n \rightarrow \infty} x(\tau(n)) = +\infty, \quad (3.4)$$

which means that  $(x(n))$  is unbounded. The proof of part (I) of the theorem is complete.

Assume that  $c = -1$ . Then, in view of part (iii) of Lemma 2.1, we have  $\sum_{i=n_0}^{\infty} p(i) x(\sigma(i)) < \infty$ , and, therefore, in view of part (i) of Lemma 2.1, we obtain

$$\lim_{n \rightarrow \infty} [x(n) - x(\tau(n))] = -L. \quad (3.5)$$

Assume that  $L > 0$ . Then there exists a natural number  $n_\lambda$  such that  $z(n) < 0$  for every  $n \geq n_\lambda$ , and therefore

$$x(n) < x(\tau(n)) < \dots < x(\tau^{m(n)}(n_\lambda)), \quad (3.6)$$

which means that  $(x(n))$  is bounded. Since  $(x(n))$  is bounded, let

$$M = \limsup x(n), \quad \text{where } M \geq L. \quad (3.7)$$

Then there exists a subsequence  $(x(\theta(n)))$  of  $(x(n))$  such that

$$\lim_{n \rightarrow \infty} x(\theta(n)) = M. \quad (3.8)$$

Thus,

$$\lim_{n \rightarrow \infty} z(\theta(n)) = -L, \quad (3.9)$$

$$\lim_{n \rightarrow \infty} [x(\theta(n)) - x(\tau(\theta(n)))] = -L, \quad (3.10)$$

or

$$\lim_{n \rightarrow \infty} x(\tau(\theta(n))) = M + L > M, \quad (3.11)$$

which contradicts (3.7). Therefore,  $L > 0$  is not valid. Hence,  $L = 0$  and consequently  $\lim_{n \rightarrow \infty} z(n) = 0$ . Furthermore, taking into account the fact that the sequence  $(z(n))$  is strictly decreasing we conclude that  $z(n) > 0$  or equivalently

$$x(n) > x(\tau(n)) > \dots > x(\tau(n_*)). \quad (3.12)$$

Since  $(x(n))$  has a lower bound greater than zero, it cannot have any subsequence that tends to zero. Thus,  $\lim_{n \rightarrow \infty} x(\sigma(n)) = 0$  is not valid, and therefore  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = \infty$  which contradicts our previous conclusions. Hence, if  $c = -1$ ,  $(x(n))$  oscillates. The proof of part (II) of the theorem is complete.

Assume that  $-1 < c < 0$ .

Assume that  $L > 0$ . Then there exists a natural number  $n_\lambda$  such that  $z(n) < 0$  for every  $n \geq n_\lambda$ , and therefore

$$x(n) < (-c)x(\tau(n)) < \dots < (-c)^{m(n_\epsilon)} x(\tau^{m(n_\epsilon)}(n_\lambda)), \quad (3.13)$$

which means that  $(x(n))$  tends to zero as  $n \rightarrow \infty$ . Therefore,  $(z(n))$  tends to zero as  $n \rightarrow \infty$ , that is,  $L = 0$ , which contradicts  $L > 0$ . Hence  $L = 0$ . Taking into account the fact that the sequence  $(z(n))$  is strictly decreasing, it is obvious that  $z(n) > 0$ . Hence, for every  $\epsilon > 0$  there exists a natural number  $n_4$  such that for every  $n \geq n_4$

$$x(n) + cx(\tau(n)) < \epsilon \quad (3.14)$$

or

$$x(n) < -cx(\tau(n)) + \epsilon < -c[-cx(\tau^2(n)) + \epsilon] + \epsilon = c^2x(\tau^2(n)) - c\epsilon + \epsilon. \quad (3.15)$$

For sufficiently large  $n$ , after  $m$ -steps we obtain

$$x(n) < c^{m+1}x(\tau^{m+1}(n)) + \epsilon - c\epsilon + \dots + (-c)^m\epsilon. \quad (3.16)$$

As  $n \rightarrow \infty$ , clearly  $m \rightarrow \infty$ , and therefore

$$\lim_{n \rightarrow \infty} x(n) \leq \lim_{m \rightarrow \infty} [\epsilon - c\epsilon + \dots + (-c)^m\epsilon] = \frac{\epsilon}{1+c}. \quad (3.17)$$

Since  $\epsilon$  is an arbitrary real positive number and, taking into account the fact that  $(x(n)) > 0$ , it is clear that

$$\lim_{n \rightarrow \infty} x(n) = 0. \quad (3.18)$$

Let  $c = 0$ . In view of part (iii) of Lemma 2.1, we have

$$\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) < +\infty. \quad (3.19)$$

This guarantees that  $(z(n))$  is bounded, and therefore, since  $z(n) = x(n)$ ,  $(x(n))$  is bounded. Also, since  $(z(n))$  is strictly decreasing,  $\lim_{n \rightarrow \infty} z(n)$  exists, that is,  $\lim_{n \rightarrow \infty} x(n)$  exists. In view of (2.18), we conclude that

$$\lim_{n \rightarrow \infty} x(n) = 0. \quad (3.20)$$

Assume that  $0 < c < 1$ , then

$$\lim_{n \rightarrow \infty} z(n) = cL, \quad \text{where} \quad \lim_{n \rightarrow \infty} x(\tau(\sigma(n))) = L \quad (3.21)$$

or

$$\lim_{n \rightarrow \infty} [x(n) + cx(\tau(n))] = cL. \quad (3.22)$$

Therefore,

$$\lim_{n \rightarrow \infty} z(\tau(\sigma(n))) = cL, \quad (3.23)$$

$$\lim_{n \rightarrow \infty} [x(\tau(\sigma(n))) + cx(\tau^2(\sigma(n)))] = cL, \quad (3.24)$$

or

$$\lim_{n \rightarrow \infty} x(\tau^2(\sigma(n))) = L - \frac{L}{c} \leq 0, \quad (3.25)$$

which means that  $L = 0$ . Hence,

$$\lim_{n \rightarrow \infty} z(n) = 0, \quad (3.26)$$

and since

$$z(n) \geq x(n) > 0, \quad (3.27)$$

we have

$$\lim_{n \rightarrow \infty} x(n) = 0. \quad (3.28)$$

The proof of part (III) of the theorem is complete.

Assume that  $c \geq 1$ . In view of Part (iii) of Lemma 2.1, it is clear that  $\sum_{i=n_0}^{\infty} p(i)x(\sigma(i)) = S_0 < +\infty$  which combined with (2.14) implies that  $(z(n))$  is bounded and, therefore  $(x(n))$  is bounded. The proof of part (IV) of the theorem is complete.

In the remainder of this proof, it will be assumed that  $x(n)$  is a continuous function with respect to  $c$ , and, therefore, instead of  $x(n)$  we will write  $x(c, n)$ .

Let  $c = -1$ . Then, in view of part (II),  $(x(-1, n))$  oscillates. On the other hand, since  $x(c, n)$  is continuous, we have

$$\lim_{c \rightarrow -1^-} x(c, n) = x(-1, n). \quad (3.29)$$

But  $x(c, n) > 0$  for all large  $n$ , and therefore its limit is always nonnegative. Thus,  $x(-1, n) > 0$  for all large  $n$  which contradicts that  $(x(-1, n))$  oscillates. Therefore,  $x(-1, n) = 0$  eventually.

Let  $c < -1$ . In view of part (I) we have  $(x(c, n))$  is unbounded, and therefore  $x(c, \tau(n)) \rightarrow \infty$ . Let  $M > 0$ , then there exists an index  $n_5$  such that, for every  $n \geq n_5$ ,  $x(c, \tau(n)) > M$ . Since the function  $x(c, n)$  is continuous,  $x(c, \tau(n))$  is continuous, and therefore

$$\lim_{c \rightarrow -1^-} x(c, \tau(n)) = x(-1, \tau(n)) = 0 \quad \text{for every } n \geq n_5. \quad (3.30)$$

Hence, there exists  $h > 0$  so that, if  $c > -1 - h$  then  $x(c, \tau(n)) < M$  for every  $n \geq n_5$ , which contradicts  $x(c, \tau(n)) > M$ . This implies that, there exists an interval  $(a, -1)$  such that

$$x(c, n) = 0 \quad \text{for every } c \in (a, -1) \text{ eventually.} \quad (3.31)$$

Let

$$A = \inf\{c \mid x(c, n) = 0\}. \quad (3.32)$$

Then, with the same argument as above, we conclude that

$$x(A, n) = 0 \text{ eventually.} \quad (3.33)$$

So, by a similar procedure we obtain an interval  $(B, A]$  such that

$$x(c, n) = 0 \quad \text{for every } c \in (B, A] \text{ eventually.} \quad (3.34)$$

This contradicts our assumption for  $A$ . Thus, if  $c < -1$ , we have  $x(c, n) = 0$  eventually. The proof of part (V) of the theorem is complete.

Assume that  $-1 < c \leq 1$ . Taking into account part (III), it is enough to show part (VI) only for  $c = 1$ . In view of parts (iii) and (i) of Lemma 2.1, we have

$$\lim_{n \rightarrow \infty} x(c, \tau(\sigma(n))) = L. \quad (3.35)$$

Assume, for the sake of contradiction, that  $L > 0$ . Taking into account the fact that  $\lim_{n \rightarrow \infty} x(c, n) = 0$  when  $0 < c < 1$ , there exists  $n_6 \in \mathbb{N}$  such that, for every  $n \geq n_6$ , we have

$$x(c, \tau(\sigma(n))) < \frac{L}{2}, \quad (3.36)$$

and, since  $x(c, \tau(\sigma(n)))$  is continuous,

$$\lim_{c \rightarrow 1^-} x(c, \tau(\sigma(n))) = x(1, \tau(\sigma(n))). \quad (3.37)$$

Hence,

$$x(1, \tau(\sigma(n))) \leq \frac{L}{2}, \quad \forall n \geq n_6 \quad (3.38)$$

which contradicts  $\lim_{n \rightarrow \infty} x(c, \tau(\sigma(n))) = L$ . Therefore,  $L > 0$  is impossible. Thus,  $L = 0$ , and, in view of (2.8), we conclude that

$$\lim_{n \rightarrow \infty} z(1, n) = 0, \quad (3.39)$$

which means that

$$\lim_{n \rightarrow \infty} x(1, n) = 0. \quad (3.40)$$

The proof of part (VI) of the theorem is complete.

Finally, since  $x(c, n)$  has convergent Taylor series, we have

$$x(c, n) = \sum_{m=0}^{\infty} \frac{x^{(m)}(a, n)}{m!} c^m \quad \text{for every } a \in \mathbb{R}. \quad (3.41)$$

Choose  $a < -1$ . Then  $x^{(m)}(a, n) = 0$  and, therefore,  $x(c, n) = 0$ . The proof of part (VII) of the theorem is complete.

The proof of the theorem is complete.  $\square$

By way of illustration and for purely pedagogical purposes, the asymptotic behavior of nonoscillatory solutions of (E) is presented in Figure 1.

|  |                           |   |     |  |
|--|---------------------------|---|-----|--|
| $c < -1$                                   | $-1$                      | $-1 < c < 1$                                | $1$ | $c \geq 1$                               |
| Every nonoscillatory solution is unbounded | Every solution oscillates | Every nonoscillatory solution tends to zero |     | Every nonoscillatory solution is bounded |

Figure 1

**Corollary 3.2.** For  $(E_1)$  one has the following.

- (i) Every nonoscillatory solution tends to  $\pm\infty$  if  $c < -1$ .
- (ii) Every solution oscillates if  $c = -1$ .
- (iii) Every nonoscillatory solution tends to zero if  $c > -1$ .

Furthermore, if any solution of  $(E_1)$  is continuous with respect to  $c$ , one has the following.

- (iv) Every solution is zero if  $c \leq -1$ .
- (v) If, additionally, any solution of  $(E_1)$  has continuous derivatives of any order and convergent Taylor series for every  $c \in \mathbb{R}$ , then the solution is zero.

*Proof.* In part (I) of Theorem 3.1 we have proved that

$$\lim_{n \rightarrow \infty} x(\tau(n)) = +\infty \tag{3.42}$$

and consequently

$$\lim_{n \rightarrow \infty} x(n - a) = +\infty, \tag{3.43}$$

which means that

$$\lim_{n \rightarrow \infty} x(n) = +\infty. \tag{3.44}$$

The proof of part (i) of the corollary is complete.

Part (ii) is direct from part (II) of Theorem 3.1.

As we have proved in parts (III) and (IV) of Theorem 3.1, if  $c > -1$ , then

$$\lim_{n \rightarrow \infty} x(\sigma(n)) = 0 \tag{3.45}$$

and consequently

$$\lim_{n \rightarrow \infty} x(n - b) = 0, \tag{3.46}$$

which means that

$$\lim_{n \rightarrow \infty} x(n) = 0. \quad (3.47)$$

The proof of part (iii) of the corollary is complete.

Parts (iv) and (v) are direct from parts (V) and (VII) of Theorem 3.1.

The proof of the corollary is complete.  $\square$

## References

- [1] R. D. Driver, "Existence and continuous dependence of solutions of a neutral functional-differential equation," *Archive for Rational Mechanics and Analysis*, vol. 19, pp. 149–166, 1965.
- [2] R. Bellman and K. L. Cooke, *Differential-Difference Equations*, Academic Press, New York, NY, USA, 1963.
- [3] J. Hale, *Theory of Functional Differential Equations*, Springer, New York, NY, USA, 2nd edition, 1977.
- [4] W. Snow, *Existence, Uniqueness, and Stability for Nonlinear Differential-Difference Equations in the Neutral Case*, New York: Courant Institute of Mathematical Sciences, New York University, New York, NY, USA, 1965, IMM-NYU 328.
- [5] R. P. Agarwal, M. Bohner, S. R. Grace, and D. O'D'Regan, *Discrete Oscillation Theory*, Hindawi Publishing Corporation, New York, NY, USA, 2005.
- [6] A. Bellen, N. Guglielmi, and A. E. Ruehli, "Methods for linear systems of circuit delay differential equations of neutral type," *IEEE Transactions on Circuits and Systems. I*, vol. 46, no. 1, pp. 212–216, 1999.
- [7] A. G. Balanov, N. B. Janson, P. V. E. McClintock, R. W. Tucker, and C. H. T. Wang, "Bifurcation analysis of a neutral delay differential equation modelling the torsional motion of a driven drill-string," *Chaos, Solitons and Fractals*, vol. 15, no. 2, pp. 381–394, 2003.
- [8] K. Gopalsamy, *Stability and Oscillations in Population Dynamics*, vol. 74, Kluwer Academic Publishers, Boston, Mass, USA, 1992.
- [9] X. H. Tang and S. S. Cheng, "Positive solutions of a neutral difference equation with positive and negative coefficients," *Georgian Mathematical Journal*, vol. 11, no. 1, pp. 177–185, 2004.
- [10] W. Xiong and J. Liang, "Novel stability criteria for neutral systems with multiple time delays," *Chaos, Solitons and Fractals*, vol. 32, no. 5, pp. 1735–1741, 2007.
- [11] J. Zhou, T. Chen, and L. Xiang, "Robust synchronization of delayed neural networks based on adaptive control and parameters identification," *Chaos, Solitons & Fractals*, vol. 27, pp. 905–913, 2006.
- [12] R. D. Driver, "A mixed neutral system," *Nonlinear Analysis*, vol. 8, no. 2, pp. 155–158, 1984.
- [13] R. K. Brayton and R. A. Willoughby, "On the numerical integration of a symmetric system of difference-differential equations of neutral type," *Journal of Mathematical Analysis and Applications*, vol. 18, pp. 182–189, 1967.
- [14] M. K. Grammatikopoulos, E. A. Grove, and G. Ladas, "Oscillations of first-order neutral delay differential equations," *Journal of Mathematical Analysis and Applications*, vol. 120, no. 2, pp. 510–520, 1986.
- [15] I. P. Stavroulakis, "Oscillations of mixed neutral equations," *Hiroshima Mathematical Journal*, vol. 19, no. 3, pp. 441–456, 1989.
- [16] J. Bařtinec, L. Berezansky, J. Diblík, and Z. Šmarda, "A final result on the oscillation of solutions of the linear discrete-delayed equation  $\Delta x(n) = -p(n)x(n - k)$  with a positive coefficient," *Abstract and Applied Analysis*, vol. 2011, Article ID 586328, 28 pages, 2011.
- [17] J. Bařtinec, J. Diblík, and Z. Šmarda, "Existence of positive solutions of discrete linear equations with a single delay," *Journal of Difference Equations and Applications*, vol. 16, no. 9, pp. 1047–1056, 2010.
- [18] J. Bařtinec, J. Diblík, and Z. Šmarda, "Oscillations of solutions of a linear second order discrete-delayed equation," *Advances in Difference Equations*, vol. 2010, Article ID 693867, 12 pages, 2010.
- [19] A. Bezeransky, J. Diblík, M. Ruzickova, and Z. Suta, "Asymptotic convergence of the solutions of a discrete equation with delays in the critical case," *Abstract and Applied Analysis*, vol. 2011, Article ID 709427, 15 pages, 2011.
- [20] G. E. Chatzarakis, G. L. Karakostas, and I. P. Stavroulakis, "Convergence of the positive solutions of a nonlinear neutral difference equation," *Nonlinear Oscillations*, vol. 14, no. 3, pp. 407–418, 2011.



- [21] M.-P. Chen, B. S. Lalli, and J. S. Yu, "Oscillation in neutral delay difference equations with variable coefficients," *Computers & Mathematics with Applications*, vol. 29, no. 3, pp. 5–11, 1995.
- [22] S. S. Cheng and Y.-Z. Lin, "Complete characterizations of an oscillatory neutral difference equation," *Journal of Mathematical Analysis and Applications*, vol. 221, no. 1, pp. 73–91, 1998.
- [23] J. Diblík and M. Ružičková, "Convergence of the solutions of the equation  $y'(t) = \beta(t)[y(t-\delta) - y(t-\tau)]$  in the critical case," *Journal of Mathematical Analysis and Applications*, vol. 331, pp. 1361–1370, 2007.
- [24] J. Diblík, M. Ružičková, and Z. Šmarda, "Ważewski's method for systems of dynamic equations on time scales," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 71, no. 12, pp. e1124–e1131, 2009.
- [25] J. Diblík, M. Ružičková, and Z. Šutá, "Asymptotical convergence of the solutions of a linear differential equation with delays," *Advances in Difference Equations*, vol. 2010, Article ID 749852, 12 pages, 2010.
- [26] J.-G. Dong, "Oscillation of solutions for first order neutral differential equations with distributed deviating arguments," *Computers & Mathematics with Applications*, vol. 58, no. 4, pp. 784–790, 2009.
- [27] Y. Domshlak, N. Partsvania, and I. P. Stavroulakis, "Oscillation properties of first order neutral differential equations near the critical states," *Nonlinear Functional Analysis and Applications*, vol. 9, no. 2, pp. 173–184, 2004.
- [28] Y. Gao and G. Zhang, "Oscillation of nonlinear first order neutral difference equations," *Applied Mathematics E-Notes*, vol. 1, pp. 5–10, 2001.
- [29] D. A. Georgiou, E. A. Grove, and G. Ladas, "Oscillations of neutral difference equations," *Applicable Analysis*, vol. 33, no. 3-4, pp. 243–253, 1989.
- [30] I. Györi and L. Horváth, "Asymptotic constancy in linear difference equations: limit formulae and sharp conditions," *Advances in Difference Equations*, vol. 2010, Article ID 789302, 20 pages, 2010.
- [31] X. Gong, X. Zhong, J. Jia, R. Quyang, and H. Han, "Oscillation of first-order neutral difference equation," *Modern Applied Science*, vol. 3, no. 8, pp. 90–94, 2009.
- [32] D. C. Huong, "Oscillation and Convergence for a neutral difference equation," *Journal of Science, Mathematics-Physics*, vol. 24, pp. 133–143, 2008.
- [33] J. Jia, X. Zhong, X. Gong, R. Quyang, and H. Han, "Nonoscillation of first-order neutral difference equation," *Modern Applied Science*, vol. 3, no. 11, pp. 30–33, 2009.
- [34] G. Ladas and Y. G. Sficas, "Oscillations of neutral delay differential equations," *Canadian Mathematical Bulletin*, vol. 29, no. 4, pp. 438–445, 1986.
- [35] B. S. Lalli, "Oscillation theorems for neutral difference equations," *Computers & Mathematics with Applications*, vol. 28, no. 1–3, pp. 191–202, 1994, Advances in difference equation.
- [36] B. S. Lalli and B. G. Zhang, "On existence of positive solutions and bounded oscillations for neutral difference equations," *Journal of Mathematical Analysis and Applications*, vol. 166, no. 1, pp. 272–287, 1992.
- [37] J. W. Li, Z. C. Wang, and H. Q. Zhang, "Oscillation of neutral delay difference equations," *Differential Equations and Dynamical Systems*, vol. 4, no. 1, pp. 113–121, 1996.
- [38] M. Migda and G. Zhang, "On unstable neutral difference equations with "maxima"," *Mathematica Slovaca*, vol. 56, no. 4, pp. 451–463, 2006.
- [39] S. L. Marian, M. P. Loganathan, and A. G. M. Selvan, "Oscillatory behavior of forced neutral difference equations with positive and negative coefficients," *International Journal of Computational and Applied Mathematics*, vol. 5, pp. 253–265, 2010.
- [40] O. Öcalan, "Oscillation criteria for forced neutral difference equation with positive and negative coefficients," *Dynamics of Continuous, Discrete & Impulsive Systems*, vol. 15, no. 2, pp. 151–161, 2008.
- [41] O. Öcalan and O. Duman, "Oscillation analysis of neutral difference equations with delays," *Chaos, Solitons and Fractals*, vol. 39, no. 1, pp. 261–270, 2009.
- [42] H. Péics, "Positive solutions of neutral delay difference equation," *Novi Sad Journal of Mathematics*, vol. 35, no. 2, pp. 111–122, 2005.
- [43] Y. G. Sficas and I. P. Stavroulakis, "Necessary and sufficient conditions for oscillations of neutral differential equations," *Journal of Mathematical Analysis and Applications*, vol. 123, no. 2, pp. 494–507, 1987.
- [44] W. Shan and G. Weigao, "Oscillation of neutral difference equations with positive and negative coefficients," *Computers & Mathematics with Applications*, vol. 47, no. 10-11, pp. 1647–1657, 2004.
- [45] X. H. Tang and X. Lin, "Necessary and sufficient conditions for oscillation of first-order nonlinear neutral difference equations," *Computers & Mathematics with Applications*, vol. 55, no. 6, pp. 1279–1292, 2008.
- [46] E. Thandapani, R. Arul, and P. S. Raja, "Oscillation of first order neutral delay difference equations," *Applied Mathematics E-Notes*, vol. 3, pp. 88–94, 2003.

- [47] E. Thandapani and P. M. Kumar, "Oscillation and nonoscillation of nonlinear neutral delay difference equations," *Tamkang Journal of Mathematics*, vol. 38, no. 4, pp. 323–333, 2007.
- [48] C.-J. Tian and S. S. Cheng, "Oscillation criteria for delay neutral difference equations with positive and negative coefficients," *Bulletin of Mathematical Society of Paraná*, vol. 21, no. 1-2, pp. 1–12, 2003.
- [49] J. S. Yu and Z. C. Wang, "Asymptotic behavior and oscillation in neutral delay difference equations," *Funkcialaj Ekvacioj*, vol. 37, no. 2, pp. 241–248, 1994.
- [50] G. Zhang, "Oscillation for nonlinear neutral difference equations," *Applied Mathematics E-Notes*, vol. 2, pp. 17–19, 2002.
- [51] G. Zhang and S. S. Cheng, "Oscillation criteria for a neutral difference equation with delay," *Applied Mathematics Letters*, vol. 8, no. 3, pp. 13–17, 1995.
- [52] Y. Zhou, "Oscillations of neutral difference equations," *Kyungpook Mathematical Journal*, vol. 39, pp. 283–291, 1999.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

