

Research Article

Hopf Bifurcation of a Differential-Algebraic Bioeconomic Model with Time Delay

Xiaojian Zhou,^{1,2} Xin Chen,¹ and Yongzhong Song¹

¹ Jiangsu Key Laboratory for NSLSCS, Institute of Mathematics, School of Mathematical Sciences, Nanjing Normal University, Nanjing 210023, China

² School of Science, Nantong University, Nantong 226008, China

Correspondence should be addressed to Yongzhong Song, yzsong@njnu.edu.cn

Received 7 September 2012; Accepted 26 September 2012

Academic Editor: Junjie Wei

Copyright © 2012 Xiaojian Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the dynamics of a differential-algebraic bioeconomic model with two time delays. Regarding time delay as a bifurcation parameter, we show that a sequence of Hopf bifurcations occur at the positive equilibrium as the delay increases. Using the theories of normal form and center manifold, we also give the explicit algorithm for determining the direction of the Hopf bifurcations and the stability of the bifurcating periodic solutions. Numerical tests are provided to verify our theoretical analysis.

1. Introduction

Predator-prey systems with delay play an important role in population dynamics since Vito Volterra and James Lotka proposed the seminal models of predator-prey in the mid of 1920s. In recent years, predator-prey systems with delay have been studied extensively due to their theoretical and practical significance.

In 1973, May [1] first proposed the delayed predator-prey system

$$\begin{aligned}\dot{x} &= x(r_1 - a_{11}x(t - \tau) - a_{12}y), \\ \dot{y} &= y(-r_2 + a_{21}x - a_{22}y),\end{aligned}\tag{1.1}$$

where x and y can be interpreted as the population densities of prey and predator at time t , respectively. $\tau > 0$ is the feedback time delay of prey species to the growth of species itself. $r_1 > 0$ denotes intrinsic growth rate of prey, and $r_2 > 0$ denotes the death rate of predator. The parameters a_{ij} ($i, j = 1, 2$) are all positive constants.

Recently, Song and Wei [2] further considered the existence of local Hopf bifurcations of system (1.1). They regarded the feedback time delay τ as a bifurcation parameter and investigated the stability and the direction of periodic solutions bifurcating from Hopf bifurcations, by applying the normal form theory and the center manifold reduction developed by Hassard et al. [3].

Considering the feedback time delay of predator species to the growth of species itself and also with the delay τ , Yan and Li [4] studied Hopf bifurcation and global periodic solutions of the following modified delayed predator-prey system:

$$\begin{aligned}\dot{x} &= x(r_1 - a_{11}x(t - \tau) - a_{12}y), \\ \dot{y} &= y(-r_2 + a_{21}x - a_{22}y(t - \tau)).\end{aligned}\tag{1.2}$$

They established the global existence results of periodic solutions bifurcating from Hopf bifurcations using a global Hopf bifurcation result due to Wu [5], which was different from that used in Song and Wei [2]. In [6], Yuan and Zhang also investigated the stability of the positive equilibrium and existence of Hopf bifurcation of the model (1.2).

Noting that, in real situations, the feedback time delay of the prey to the growth of the species itself and the feedback time delay of the predator to the growth of the species itself are different. If the time delay of the predator to the growth of the species itself is zero, (1.2) becomes (1.1). Meanwhile, if the time delay of the prey to the growth of the species itself is zero, (1.2) is simplified as

$$\begin{aligned}\dot{x} &= x(t)(r_1 - a_{11}x(t) - a_{12}y(t)), \\ \dot{y} &= y(t)(-r_2 + a_{21}x(t) - a_{22}y(t - \tau)).\end{aligned}\tag{1.3}$$

In 1954, Gordon [7] studied the effect of harvest effort on ecosystem from an economic perspective and proposed the following economic theory:

$$\text{Net Economic Revenue (NER)} = \text{Total Revenue (TR)} - \text{Total Cost (TC)}.\tag{1.4}$$

This provides theoretical fundament for the establishment of bioeconomic systems by differential-algebraic equations (DAEs).

Let E and y represent the harvest effort and the density of harvested population, respectively. Then $\text{TC} = cE$ and $\text{TR} = \alpha Ey$, where α is unit price of harvested population and c is the cost of harvest effort. Considering the economic interest m of the harvest effort on the predator, we can establish the following algebraic equation:

$$E(\alpha y - c) = m.\tag{1.5}$$

Recently, a class of differential-algebraic bioeconomic models were proposed and analyzed in [8–14]. For example, [8] discussed the problems of chaos and chaotic control for a differential-algebraic system. The existence and stability of equilibrium points, the sufficient conditions of existence for various bifurcations (transcritical bifurcation, singular induced bifurcation, and Hopf bifurcation) are investigated in [9–14]. However, the stability and the

direction of periodic solutions bifurcating from Hopf bifurcations have not been discussed there. Very recently, Zhang et al. [15] analyzed the stability and the Hopf bifurcation of a differential-algebraic bioeconomic system with a single time delay and Holling II type functional response and first investigated the direction of Hopf bifurcation and the stability of the bifurcating periodic solutions.

Based on the above economic theory and system (1.3), in this paper, we study the Hopf bifurcation of the following differential-algebraic bioeconomic system:

$$\begin{aligned}\dot{x} &= x(r_1 - a_{11}x(t) - a_{12}y), \\ \dot{y} &= y(-r_2 + a_{21}x - a_{22}y(t - \tau)) - Ey, \\ 0 &= E(\alpha y - c) - m.\end{aligned}\tag{1.6}$$

The rest of this paper is organized as follows. In Section 2, regarding time delay τ as bifurcation parameter, first, we study the stability of the equilibrium point of system based on the new normal form approach proposed by Chen et al. [16]. Then, following the normal form approach theory and the center manifold theory introduced by Hassard et al. [3], we compute the normal form for the Hopf bifurcation of system (1.6) and analyze the direction and stability of the bifurcating periodic orbits of the system. Numerical examples are given in Section 3 to illustrate our theoretical results.

2. Hopf Bifurcation

In this section, we analyze the stability and the Hopf bifurcation of the differential-algebraic bioeconomic system (1.6). Furthermore, we will derive the formula for determining the properties of the Hopf bifurcation.

2.1. Existence of Hopf Bifurcation

For $m > 0$, the system (1.6) has positive equilibrium point $P_* = (x_*, y_*, E_*)$, where $x_* = (r_1 - a_{12}y_*)/a_{11}$, $E_* = m/(\alpha y_* - c)$ and y_* is the positive root of the following equation:

$$\begin{aligned}\alpha(a_{11}a_{22} + a_{12}a_{21})y^2 - (ca_{11}a_{22} + ca_{12}a_{21} + \alpha r_1 a_{21} - \alpha a_{11}r_2)y \\ + (ma_{11} + cr_1 a_{21} - ca_{11}r_2) = 0.\end{aligned}\tag{2.1}$$

In order to guarantee the existence of the positive equilibrium point P_* , some restrictions must be satisfied also:

$$\frac{c}{\alpha} < y_* < \frac{r_1}{a_{12}}, \quad ca_{11}a_{22} + ca_{12}a_{21} + \alpha r_1 a_{21} > \alpha a_{11}r_2,\tag{2.2}$$

and $ma_{11} + cr_1 a_{21} - ca_{11}r_2 < 0$.

In order to analyze the local stability of the positive equilibrium point, we first use the linear transformation $(x, y, E)^T = Q \cdot (u, v, \bar{E})^T$, where

$$Q = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -\frac{\alpha E_*}{\alpha y_* - c} & 1 \end{bmatrix}. \quad (2.3)$$

Then the system (1.6) yields

$$\begin{aligned} \dot{u} &= u(r_1 - a_{11}u - a_{12}v), \\ \dot{v} &= v(-r_2 + a_{21}u - a_{22}v(t - \tau)) - \bar{E}v + \frac{\alpha E_* v^2}{\alpha y_* - c}, \\ 0 &= \left(\bar{E} - \frac{\alpha E_* v}{\alpha y_* - c} \right) (\alpha v - c) - m. \end{aligned} \quad (2.4)$$

According to the literature [16], considering the local parametrization of the third equation of the system (2.4), and we can obtain the following parametric system of it:

$$\begin{aligned} \dot{x}_1 &= (x_1 + u_*)(r_1 - a_{11}(x_1 + u_*) - a_{12}(x_2 + v_*)), \\ \dot{x}_2 &= (x_2 + v_*)(-r_2 + a_{21}(x_1 + u_*) - a_{22}(x_2(t - \tau) + v_*)) - (\bar{E}_* + h)(x_2 + v_*) + \frac{\alpha E_*(x_2 + v_*)^2}{\alpha y_* - c}, \end{aligned} \quad (2.5)$$

where

$$u = u_* + x_1, \quad v = v_* + x_2, \quad \bar{E} = \bar{E}_* + h(x_1, x_2), \quad (2.6)$$

$h : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a smooth mapping, and $(u_*, v_*, \bar{E}_*)^T$ satisfies

$$(x_*, y_*, E_*)^T = Q \cdot (u_*, v_*, \bar{E}_*)^T. \quad (2.7)$$

Thus the linearized system of parametric system (2.5) at $(0,0)$ is given as below:

$$\begin{aligned} \dot{x}_1 &= -a_{11}x_*x_1 - a_{12}x_*x_2, \\ \dot{x}_2 &= a_{21}y_*x_1 + \frac{\alpha E_* y_*}{\alpha y_* - c} x_2 - a_{22}y_*x_2(t - \tau). \end{aligned} \quad (2.8)$$

The characteristic equation of the linearized system of parametric system (2.8) at $(0,0)$ is given by

$$\lambda^2 + p\lambda + r + (s\lambda + q)e^{-\lambda\tau} = 0, \quad (2.9)$$

where $p = a_{11}x_* - \alpha y_* E_* / (\alpha y_* - c)$, $r = x_* y_* (a_{21}a_{12} - \alpha a_{11} E_* / (\alpha y_* - c))$, $s = a_{22}y_*$, and $q = a_{11}a_{22}x_* y_*$.

The second-degree transcendental polynomial equation (2.9) has been extensively studied and applied by many researchers [2, 17–20].

Let

$$(H1) \quad r_1 > \alpha y_* E_* / (\alpha y_* - c);$$

$$(H2) \quad a_{12} a_{21} + a_{11} a_{22} > \alpha a_{11} E_* / (\alpha y_* - c);$$

$$(H3) \quad \text{either } 2x_* y_* a_{21} a_{12} + a_{22}^2 y_*^2 < a_{11}^2 x_*^2 + (y_* E_* \alpha / (\alpha y_* - c))^2 \text{ and } (a_{21} a_{12} - a_{11} E_* \alpha / (\alpha y_* - c))^2 > (a_{11} a_{22})^2 \text{ or}$$

$$\left(2x_* y_* a_{21} a_{12} + a_{22}^2 y_*^2 - a_{11}^2 x_*^2 - \left(\frac{y_* E_* \alpha}{\alpha y_* - c} \right)^2 \right)^2 < 4x_*^2 y_*^2 \left(\left(a_{21} a_{12} - \frac{a_{11} E_* \alpha}{\alpha y_* - c} \right)^2 - a_{11}^2 a_{22}^2 \right); \quad (2.10)$$

$$(H4) \quad \text{either } (a_{21} a_{12} - a_{11} E_* \alpha / (\alpha y_* - c))^2 < (a_{11} a_{22})^2 \text{ or } 2x_* y_* a_{21} a_{12} + a_{22}^2 y_*^2 > a_{11}^2 x_*^2 + (y_* E_* \alpha / (\alpha y_* - c))^2, \text{ and}$$

$$\left(2x_* y_* a_{21} a_{12} + a_{22}^2 y_*^2 - a_{11}^2 x_*^2 - \left(\frac{y_* E_* \alpha}{\alpha y_* - c} \right)^2 \right)^2 = 4x_*^2 y_*^2 \left(\left(a_{21} a_{12} - \frac{a_{11} E_* \alpha}{\alpha y_* - c} \right)^2 - a_{11}^2 a_{22}^2 \right); \quad (2.11)$$

$$(H5) \quad (a_{21} a_{12} - a_{11} E_* \alpha / (\alpha y_* - c))^2 > (a_{11} a_{22})^2, 2x_* y_* a_{21} a_{12} + a_{22}^2 y_*^2 > a_{11}^2 x_*^2 + (y_* E_* \alpha / (\alpha y_* - c))^2, \text{ and}$$

$$\left(2x_* y_* a_{21} a_{12} + a_{22}^2 y_*^2 - a_{11}^2 x_*^2 - \left(\frac{y_* E_* \alpha}{\alpha y_* - c} \right)^2 \right)^2 > 4x_*^2 y_*^2 \left(\left(a_{21} a_{12} - \frac{a_{11} E_* \alpha}{\alpha y_* - c} \right)^2 - a_{11}^2 a_{22}^2 \right). \quad (2.12)$$

From Lemma 2.1 in [2], we easily obtain the following results about the stability of the positive equilibrium and the Hopf bifurcation of (1.6).

Theorem 2.1. *For system (1.6). One has the following.*

(i) *If (H1)–(H3) hold, then the equilibrium E_* of the system (1.6) is asymptotically stable for all $\tau \geq 0$.*

(ii) *If (H1), (H2), and (H4) hold, then the equilibrium E_* of the system (1.6) is asymptotically stable when $0 \leq \tau < \tau_0^+$ and unstable when $\tau > \tau_0^+$. System (1.6) undergoes a Hopf bifurcation at E_* when $\tau = \tau_j^+$.*

(iii) *If (H1), (H2), and (H5) hold, then there is a positive integer k , such that the equilibrium E_* switches k times from stability to instability to stability; that is, E_* is asymptotically stable when*

$$\tau \in [0, \tau_0^+) \cup (\tau_0^-, \tau_1^+) \cup \cdots \cup (\tau_{k-1}^-, \tau_k^+) \quad (2.13)$$

and unstable when

$$\tau \in (\tau_0^+, \tau_0^-) \cup (\tau_1^+, \tau_1^-) \cup \cdots \cup (\tau_{k-1}^+, \tau_{k-1}^-) \cup (\tau_{k-1}^+, +\infty). \quad (2.14)$$

Here,

$$\omega_{\pm} = \frac{\sqrt{2}}{2} \left[2x_*y_*a_{21}a_{12} + a_{22}^2y_*^2 - a_{11}^2x_*^2 - \left(\frac{y_*E_*\alpha}{\alpha y_* - c} \right)^2 \right. \\ \left. \pm \sqrt{\left(2x_*y_*a_{21}a_{12} + a_{22}^2y_*^2 - a_{11}^2x_*^2 - \left(\frac{y_*E_*\alpha}{\alpha y_* - c} \right)^2 \right)^2 + 4x_*^2y_*^2 \left(\left(a_{21}a_{12} - \frac{a_{11}E_*\alpha}{\alpha y_* - c} \right)^2 - (a_{11}a_{22})^2 \right)} \right]^{1/2}, \\ \tau_j^{\pm} = \frac{1}{\omega_{\pm}} \arccos \left(\frac{a_{22}(E_*\alpha / (\alpha y_* - c))\omega_{\pm}^2 - a_{11}a_{22}x_*^2(a_{21}a_{12} - a_{11}E_*\alpha / (\alpha y_* - c))}{a_{22}(\omega_{\pm}^2 + a_{11}^2x_*^2)} \right) + \frac{2j\pi}{\omega_{\pm}}, \\ j = 0, 1, 2, \dots \quad (2.15)$$

Denote $\lambda_j(\tau) = \alpha_j(\tau) + i\omega_j(\tau)$, $j = 0, 1, 2, \dots$; then the following transversality conditions hold:

$$\frac{d}{d\tau} \operatorname{Re} \lambda_j(\tau_j^+) > 0, \quad \frac{d}{d\tau} \operatorname{Re} \lambda_j(\tau_j^-) < 0. \quad (2.16)$$

2.2. Direction and Stability of the Hopf Bifurcation

In the previous section, we obtain the conditions for Hopf bifurcations at the positive equilibrium P_* when $\tau = \tau_n^{\pm}$, $n = 0, 1, 2, \dots$. In this section, we will derive the formulae determining the direction, stability, and period of these periodic solutions bifurcating from P_* at τ . The basic techniques used here are the normal form and the center manifold theory developed by Hassard et al. [3].

Assuming the system (1.6) undergoes Hopf bifurcations at the positive equilibrium P_* at $\tau = \tau_n$, we let $i\omega$ be the corresponding purely imaginary root of the characteristic equation at the positive equilibrium. In order to investigate the direction of Hopf bifurcation and the stability of the bifurcating periodic solutions of system (1.6), we consider the parametric system (2.5) of system (2.4).

Let $t = t\tau$, $\tau = \tau_n + \mu$; then the parametric system (2.5) is equivalent to the following functional differential equation system:

$$\dot{z}_1 = (\tau_n + \mu)(-a_{11}x_*z_1 - a_{12}x_*z_2), \\ \dot{z}_2 = (\tau_n + \mu) \left(a_{21}y_*z_1 + \frac{\alpha E_* y_*}{\alpha y_* - c} z_2 - a_{22}y_*z_2(t-1) \right), \quad (2.17)$$

where $z_1 = x_1(t\tau)$, $z_2 = x_2(t\tau)$.

For $\Phi \in C([-1, 0], \mathbb{R}^2)$, define a family of operators

$$L_\mu \Phi := (\tau_n + \mu) \begin{bmatrix} -a_{11}x_* & -a_{12}x_* \\ a_{21}y_* & \frac{\alpha E_* y_*}{\alpha y_* - c} \end{bmatrix} \Phi(0) + (\tau_n + \mu) \begin{bmatrix} 0 & 0 \\ 0 & -a_{22}y_* \end{bmatrix} \Phi(-1), \quad (2.18)$$

$$f(\mu, \Phi) := (\tau_n + \mu) \times \begin{bmatrix} -a_{11}\Phi_1^2(0) - a_{12}\Phi_1(0)\Phi_2(0) + \dots \\ a_{21}\Phi_1(0)\Phi_2(0) - a_{22}\Phi_2(0)\Phi_2(-1) - \frac{\alpha E_* c}{(\alpha y_* - c)^2} \Phi_2^2(0) + \dots \end{bmatrix},$$

where $\Phi = (\Phi_1, \Phi_2) \in C$. By the Riesz representation theorem, there exists a function of bounded variation for $\theta \in [-1, 0]$, such that

$$L_\mu \Phi = \int_{-1}^0 d\eta(\theta, \mu) \Psi(\theta), \quad \Psi \in C. \quad (2.19)$$

In fact, we can choose

$$\eta(\theta, \mu) = (\tau_n + \mu) \begin{bmatrix} -a_{11}x_* & -a_{12}x_* \\ a_{21}y_* & \frac{\alpha E_* y_*}{\alpha y_* - c} \end{bmatrix} \delta(\theta) + (\tau_n + \mu) \begin{bmatrix} 0 & 0 \\ 0 & a_{22}y_* \end{bmatrix} \delta(\theta + 1). \quad (2.20)$$

For $\Phi \in C$, define

$$A(\mu) \Phi := \begin{cases} \frac{d\Phi(\theta)}{d\theta}, & \theta \in [-1, 0), \\ \int_{-1}^0 d\eta(t, \mu) \Phi, & \theta = 0, \end{cases} \quad (2.21)$$

$$R(\mu) := \begin{cases} 0, & \theta \in [-1, 0), \\ f(t, \Phi), & \theta = 0. \end{cases}$$

Hence system (2.17) can be rewritten as

$$\dot{Z}_t = A(\mu) Z_t + R(\mu) Z_t. \quad (2.22)$$

For $\Psi \in C([-1, 0], (\mathbb{R}^2)^*)$, define

$$A^* \Psi(s) := \begin{cases} \frac{d\Psi(s)}{ds}, & s \in [-1, 0), \\ \int_{-1}^0 d\eta^T(s, 0) \Psi(-s), & s = 0, \end{cases} \quad (2.23)$$

and a bilinear inner product

$$\langle \Psi, \Phi \rangle = \bar{\Psi}^T(0)\Phi(0) - \int_{-1}^0 \int_0^\theta \bar{\Psi}^T(\xi - \theta) d\eta(\theta)\Phi(\xi) d\xi, \quad (2.24)$$

where $\eta(\theta) = \eta(\theta, 0)$. Then A^* and $A(0)$ are adjoint operators. By the discussion in Section 2.1, we know that $\pm i\omega$ are eigenvalues of $A(0)$. Thus they are also eigenvalues of A^* . Suppose that $q(\theta) = (1, q_1)^T e^{i\omega\tau\theta}$ and $\bar{q}^*(s) = D(q_1^*, 1) e^{i\omega\tau s}$ are eigenvectors of $A(0)$ and A^* corresponding to $i\omega\tau$ and $-i\omega\tau$, respectively. It is easy to obtain

$$\begin{aligned} q_1 &= -\frac{i\omega + a_{11}x_*}{a_{12}x_*}, \\ q_1^* &= \frac{i\omega - a_{22}y_* e^{i\omega\tau}}{a_{12}x_*}, \\ \bar{D} &= \left(q_1 + \bar{q}_1^* + \tau q_1 a_{22} y_* e^{-i\omega\tau} \right)^{-1}. \end{aligned} \quad (2.25)$$

Moreover, $\langle q^*(s), q(\theta) \rangle = 1$, $\langle q^*(s), \bar{q}(\theta) \rangle = 0$.

Now, we calculate the coordinates to describe the center manifold C_0 at $\mu = 0$. Using the same notations as in [3], we define

$$z = \langle q^*, Z_t \rangle, \quad W(t, \theta) = Z_t(\theta) - 2 \operatorname{Re}\{zq(\theta)\}. \quad (2.26)$$

On the center manifold C_0 , we have $W(t, \theta) = W(z, \bar{z}, \theta)$, where

$$W(z, \bar{z}, \theta) = W_{20}(\theta) \frac{z^2}{2} + W_{11}(\theta) z\bar{z} + W_{02}(\theta) \frac{\bar{z}^2}{2} + \dots, \quad (2.27)$$

and z and \bar{z} are local coordinates for C_0 in the direction of q^* and \bar{q}^* . Note that W is real if Z_t is real; we only consider real solutions. For $Z_t \in C_0$, we have

$$\dot{z} = i\omega\tau z + \bar{q}^*(0) f_0(z, \bar{z}) := i\omega\tau z + g(z, \bar{z}), \quad (2.28)$$

where

$$g(z, \bar{z}) = \bar{q}^*(0) f_0(z, \bar{z}) = \bar{q}^*(0) f(0, Z_t) = g_{20}(\theta) \frac{z^2}{2} + g_{11}(\theta) z\bar{z} + g_{02}(\theta) \frac{\bar{z}^2}{2} + \dots. \quad (2.29)$$

Moreover, we have

$$\begin{aligned}
g_{20} &= 2\bar{D}\tau \left(a_{21}q_1 - a_{11}\bar{q}_1^* - a_{12}\bar{q}_1^*q_1 - \frac{\alpha E_*c}{(\alpha y_* - c)^2}q_1^2 - a_{22}q_1^2 e^{-i\omega\tau\theta} \right), \\
g_{11} &= 2\bar{D}\tau \left(a_{21}\operatorname{Re}(q_1) - a_{11}\bar{q}_1^* - a_{12}\bar{q}_1^*\operatorname{Re}(q_1) - a_{22}\operatorname{Re}(q_1\bar{q}_1 e^{i\omega\tau\theta}) - \frac{\alpha E_*c}{(\alpha y_* - c)^2}q_1\bar{q}_1 \right), \\
g_{02} &= 2\bar{D}\tau \left(a_{21}\bar{q}_1 - a_{11}\bar{q}_1^* - a_{12}\bar{q}_1^*\bar{q}_1 - \frac{\alpha E_*c}{(\alpha y_* - c)^2}\bar{q}_1^2 - a_{22}\bar{q}_1^2 e^{i\omega\tau\theta} \right), \\
g_{21} &= 2\bar{D}\tau \left((a_{21}q_1 - 2a_{11}\bar{q}_1^* - a_{12}\bar{q}_1^*q_1)W_{11}^{(1)}(0) \right. \\
&\quad + \frac{1}{2}(a_{21}\bar{q}_1 - a_{12}\bar{q}_1^*\bar{q}_1 - 2a_{11}\bar{q}_1^*)W_{20}^{(1)}(0) \\
&\quad + \left(a_{21} - a_{22}q_1 e^{-i\omega\tau\theta} - a_{12}\bar{q}_1^* - 2\frac{\alpha E_*c}{(\alpha y_* - c)^2}q_1 \right)W_{11}^{(2)}(0) \\
&\quad + \frac{1}{2}\left(a_{21} - a_{22}\bar{q}_1 e^{i\omega\tau\theta} - a_{12}\bar{q}_1^* - 2\frac{\alpha E_*c}{(\alpha y_* - c)^2}\bar{q}_1 \right)W_{20}^{(2)}(0) \\
&\quad \left. - a_{22}q_1 W_{11}^{(2)}(-1) - \frac{1}{2}a_{22}\bar{q}_1 W_{20}^{(2)}(-1) \right).
\end{aligned} \tag{2.30}$$

Next, we will calculate $W_{11}(\theta)$ and $W_{20}(\theta)$. From (2.22) and (2.28), we have

$$\begin{aligned}
\dot{W} &= \dot{Z}_t - \dot{z}q - \dot{\bar{z}}\bar{q} = \begin{cases} AW - 2\operatorname{Re}\{\bar{q}^*(0)f(z, \bar{z})q(\theta)\}, & \theta \in [-1, 0), \\ AW - 2\operatorname{Re}\{\bar{q}^*(0)f(z, \bar{z})q(\theta)\} + f, & \theta = 0, \end{cases} \\
&:= AW + H(z, \bar{z}, \theta),
\end{aligned} \tag{2.31}$$

where $H(z, \bar{z}, \theta) = H_{20}(\theta)(z^2/2) + H_{11}(\theta)z\bar{z} + H_{02}(\theta)(\bar{z}^2/2) + \dots$.

For $\theta \in [-1, 0)$, we can get

$$(A - 2i\omega\tau)W_{20}(\theta) = -H_{20}(\theta), \quad AW_{11}(\theta) = -H_{11}(\theta). \tag{2.32}$$

On the other hand, $H(z, \bar{z}, \theta) = AW - 2\operatorname{Re}\{\bar{q}^*(0)f(z, \bar{z})q(\theta)\}$; hence we can obtain

$$H_{20}(\theta) = -g_{20}q(\theta) - \bar{g}_{02}\bar{q}(\theta), \quad H_{11}(\theta) = -g_{11}q(\theta) - \bar{g}_{11}\bar{q}(\theta). \tag{2.33}$$

It follows from (2.32) that

$$\begin{aligned} \dot{W}_{20}(\theta) &= 2i\omega W_{20}(\theta) + g_{20}q(\theta) + \bar{g}_{02}\bar{q}(\theta), \\ \dot{W}_{11}(\theta) &= g_{11}q(\theta) + \bar{g}_{11}\bar{q}(\theta). \end{aligned} \quad (2.34)$$

Solving it, we have

$$\begin{aligned} W_{20}(\theta) &= \frac{i g_{20}}{\omega \tau} q(0) e^{i\omega \tau \theta} + \frac{i \bar{g}_{02}}{3\omega \tau} \bar{q}(0) e^{-i\omega \tau \theta} + M e^{2i\omega \tau \theta}, \\ W_{11}(\theta) &= -\frac{i g_{11}}{\omega \tau} q(0) e^{i\omega \tau \theta} + \frac{i \bar{g}_{11}}{\omega \tau} \bar{q}(0) e^{-i\omega \tau \theta} + N. \end{aligned} \quad (2.35)$$

Now we will seek appropriate M and N . From (2.29) and (2.31), we have

$$\begin{aligned} H_{20}(\theta) &= -g_{20}q(0) - \bar{g}_{02}\bar{q}(0) + 2\tau H_1, \\ H_{11}(\theta) &= -g_{11}q(0) - \bar{g}_{11}\bar{q}(0) + 2\tau H_2, \end{aligned} \quad (2.36)$$

where

$$\begin{aligned} H_1 &= \begin{bmatrix} -a_{11} - a_{12}q_1 \\ a_{21}q_1 - \frac{\alpha E_* c}{(\alpha y_* - c)^2} q_1^2 - a_{22}q_1^2 e^{-i\omega \tau \theta} \end{bmatrix}, \\ H_2 &= \begin{bmatrix} -a_{11} - a_{12} \operatorname{Re}(q_1) \\ a_{21} \operatorname{Re}(q_1) - a_{22} \operatorname{Re}(q_1 \bar{q}_1 e^{i\omega \tau \theta}) - \frac{\alpha E_* c}{(\alpha y_* - c)^2} q_1 \bar{q}_1 \end{bmatrix}. \end{aligned} \quad (2.37)$$

Noting that

$$\begin{aligned} \left(i\omega \tau I - \int_{-1}^0 e^{i\omega \tau \theta} d\eta(\theta) \right) q(0) &= 0, \\ \left(-i\omega \tau I - \int_{-1}^0 e^{-i\omega \tau \theta} d\eta(\theta) \right) q(0) &= 0, \end{aligned} \quad (2.38)$$

we have the following linear equations:

$$\begin{aligned} \begin{bmatrix} 2i\omega + a_{11}x_* & a_{12}x_* \\ -a_{21}y_* & 2i\omega - a_{22}y_* - \frac{\alpha E_* y_*}{\alpha y_* - c} \end{bmatrix} M &= 2H_1, \\ \begin{bmatrix} a_{11}x_* & a_{12}x_* \\ -a_{21}y_* & -a_{22}y_* - \frac{\alpha E_* y_*}{\alpha y_* - c} \end{bmatrix} N &= 2H_2. \end{aligned} \quad (2.39)$$

It is easy to get M and N . Furthermore, we can get the following values:

$$\begin{aligned} C_1(0) &= \frac{i}{2\omega\tau} \left(g_{11}g_{20} - 2|g_{11}|^2 - \frac{|g_{02}|^2}{3} \right) + \frac{g_{21}}{2}, \\ \mu_2 &= -\frac{\operatorname{Re}(C_1(0))}{\operatorname{Re}(\lambda'(\tau))}, \\ \beta_2 &= 2 \operatorname{Re}(C_1(0)), \\ T_2 &= -\frac{\operatorname{Im}(C_1(0)) + \mu_2 \operatorname{Im}(\lambda'(\tau))}{\omega\tau}. \end{aligned} \quad (2.40)$$

It is well known that, at the critical value τ , the sign of μ_2 determines the direction of the Hopf bifurcation, β_2 determines the stability of bifurcated periodic solutions, while T_2 determines the period of bifurcated periodic solutions, respectively. In fact, we have the theorem as follows.

- Theorem 2.2.** (i) *The Hopf bifurcation is supercritical if $\mu_2 > 0$ and subcritical if $\mu_2 < 0$.*
(ii) *The bifurcated periodic solutions are unstable if $\beta_2 > 0$ and stable if $\beta_2 < 0$.*
(iii) *The period of bifurcated periodic solutions increases if $T_2 > 0$ and decreases if $T_2 < 0$.*

3. Numerical Examples

We now perform some simulations for better understanding of our analytical treatment.

Example 3.1. Consider the following differential-algebraic system:

$$\begin{aligned} \dot{x} &= x(6 - 2x(t) - 1.5y), \\ \dot{y} &= y(-0.5 + 2x - y(t - \tau)) - Ey, \\ 0 &= E(2y - 1) - 1.5. \end{aligned} \quad (3.1)$$

The only positive equilibrium point of the system (3.1) is $P^+ = (1.5, 2.0, 0.5)$. It is easy to get $p = 2.3333, q = 6, r = 7, s = 2$.




Figure 1: Dynamic behavior of the differential-algebraic system (3.1) with $\tau = 0.55 < \tau_0^+$, and the positive equilibrium point P^+ is stable.

When $\tau = 0$, from (2.9), the characteristic equation of the linearized system of parametric system of (3.1) at P^+ is given by

$$\lambda^2 + (p + s)\lambda + r + q = 0. \tag{3.2}$$

By simple calculating, $p + s = 4.3333 > 0, q + r = 13 > 0$. According to Routh-Hurwitz stability criterion, P^+ is stable.

When $\tau > 0$, we can further get $r^2 - q^2 = 13 > 0, s^2 - p^2 + 2r = 12.5556 > 0$, and $(s^2 - p^2 + 2r)^2 - 4(r^2 - q^2) = 105.6420 > 0$ which mean (H1), (H2), and (H5) hold. Furthermore, we have $\omega_- = 1.0671, \omega_+ = 3.3789$, and hence $\tau_0^+ = 0.5638 < \tau_1^+ = 2.4233 < \tau_0^- = 2.8880 < \tau_2^+ = 4.2829 < \tau_3^+ = 6.1424 < \tau_1^- = 8.7762 < \tau_2^- = 14.6645 < \tau_3^- = 20.5527 < \dots$. According to conclusion (iii) of Theorem 2.1, the equilibrium point P^+ is stable only for $\tau < \tau_0^+$, unstable for any $\tau > \tau_0^+$, and the Hopf bifurcation occurs at $\tau = \tau_0^+$.

By the aid of Mathematica, we can obtain the following values according to equalities in (2.40):

$$C_1(0) = 3.9725 - 4.7012i, \quad \lambda'(\tau_0^+) = 1.3427 - 2.9776i. \tag{3.3}$$

So we have $\mu_2 = -2.9586 < 0, \beta_2 = 7.9451 > 0$, and $T_2 = -2.1567 < 0$. Thus from Theorem 2.2, we can conclude that the Hopf bifurcation of system (3.1) is subcritical, the bifurcating periodic solution exists when τ crosses τ_0^+ to the left, and the bifurcating periodic solution is unstable and decreases.

Figure 1 shows the positive equilibrium point P^+ of system (3.1) is locally asymptotically stable when $\tau = 0.55 < \tau_0^+ = 0.5638$. The periodic solutions occur from P^+




Figure 2: Dynamic behavior of the differential-algebraic system (3.1) with $\tau = 0.563 < \tau_0^+$, periodic solutions bifurcating from the positive equilibrium point P^+ .




Figure 3: Dynamic behavior of the differential-algebraic system (3.1) with $\tau = 0.57 > \tau_0^+$, and the positive equilibrium point P^+ is unstable.

when $\tau = 0.563 < \tau_0^+ = 0.5638$ as is illustrated in Figure 2. When $\tau = 0.57 > \tau_0^+ = 0.5638$, the positive equilibrium point P^+ becomes unstable as shown in Figure 3.


Acknowledgments

The authors would like to thank the reviewers for their valuable comments which improved the paper. This work is supported by the Foundation for the Authors of the National Excellent Doctoral Thesis Award of China (200720), the "333 Project" Foundation of Jiangsu Province, and the Foundation for Innovative Program of Jiangsu Province(CXZZ11_0870, CXZZ12_0383).

References

- [1] R. M. May, "Time delay versus stability in population models with two and three trophic levels," *Ecology*, vol. 4, pp. 315–325, 1973.
- [2] Y. Song and J. Wei, "Local Hopf bifurcation and global periodic solutions in a delayed predator-prey system," *Journal of Mathematical Analysis and Applications*, vol. 301, no. 1, pp. 1–21, 2005.
- [3] B. D. Hassard, N. D. Kazarinoff, and Y. H. Wan, *Theory and Applications of Hopf Bifurcation*, vol. 41, Cambridge University Press, Cambridge, UK, 1981.
- [4] X.-P. Yan and W.-T. Li, "Hopf bifurcation and global periodic solutions in a delayed predator-prey system," *Applied Mathematics and Computation*, vol. 177, no. 1, pp. 427–445, 2006.
- [5] J. Wu, "Symmetric functional-differential equations and neural networks with memory," *Transactions of the American Mathematical Society*, vol. 350, no. 12, pp. 4799–4838, 1998.
- [6] S. Yuan and F. Zhang, "Stability and global Hopf bifurcation in a delayed predator-prey system," *Nonlinear Analysis*, vol. 11, no. 2, pp. 959–977, 2010.
- [7] H. S. Gordon, "The economic theory of a common property resource, the fishery," *Journal of Political Economy*, vol. 62, pp. 124–142, 1954.
- [8] Y. Zhang and Q. L. Zhang, "Chaotic control based on descriptor bioeconomic systems," *Control and Decision*, vol. 22, pp. 445–452, 2007.
- [9] Y. Zhang, Q. L. Zhang, and L. C. Zhao, "Bifurcations and control in singular biological economical model with stage structure," *Journal of Systems Engineering*, vol. 22, pp. 232–238, 2007.
- [10] X. Zhang, Q.-L. Zhang, C. Liu, and Z.-Y. Xiang, "Bifurcations of a singular prey-predator economic model with time delay and stage structure," *Chaos, Solitons and Fractals*, vol. 42, no. 3, pp. 1485–1494, 2009.
- [11] C. Liu, Q. Zhang, and X. Duan, "Dynamical behavior in a harvested differential-algebraic prey-predator model with discrete time delay and stage structure," *Journal of the Franklin Institute. Engineering and Applied Mathematics*, vol. 346, no. 10, pp. 1038–1059, 2009.
- [12] C. Liu, Q. Zhang, X. Zhang, and X. Duan, "Dynamical behavior in a stage-structured differential-algebraic prey-predator model with discrete time delay and harvesting," *Journal of Computational and Applied Mathematics*, vol. 231, no. 2, pp. 612–625, 2009.
- [13] Y. Feng, Q. Zhang, and C. Liu, "Dynamical behavior in a harvested differential-algebraic allelopathic phytoplankton model," *International Journal of Information & Systems Sciences*, vol. 5, no. 3-4, pp. 558–571, 2009.
- [14] Q. Zhang, C. Liu, and X. Zhang, *Complexity, Analysis and Control of Singular Bio-logical Systems*, Springer, 2012.
- [15] G. Zhang, L. Zhu, and B. Chen, "Hopf bifurcation in a delayed differential-algebraic biological economic system," *Nonlinear Analysis*, vol. 12, no. 3, pp. 1708–1719, 2011.
- [16] B. S. Chen, X. X. Liao, and Y. Q. Liu, "Normal forms and bifurcations for differential-algebraic systems," *Acta Mathematicae Applicatae Sinica*, vol. 23, no. 3, pp. 429–443, 2000 (Chinese).
- [17] K. L. Cooke and Z. Grossman, "Discrete delay, distributed delay and stability switches," *Journal of Mathematical Analysis and Applications*, vol. 86, no. 2, pp. 592–627, 1982.
- [18] Y. Kuang, *Delay Differential Equations with Applications in Population Dynamics*, vol. 191 of *Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1993.

- [19] S. Ruan, "Absolute stability, conditional stability and bifurcation in Kolmogorov-type predator-prey systems with discrete delays," *Quarterly of Applied Mathematics*, vol. 59, no. 1, pp. 159–173, 2001.
- [20] J. Wei and S. Ruan, "Stability and bifurcation in a neural network model with two delays," *Physica D*, vol. 130, no. 3-4, pp. 255–272, 1999.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

