

Research Article

Existence of Periodic Solutions for a Class of Asymptotically p -Linear Discrete Systems Involving p -Laplacian

Kai Chen¹ and Qiongfeng Zhang²

¹ Department of Information Engineering, Guilin University of Aerospace Technology, Guangxi, Guilin 541004, China

² College of Science, Guilin University of Technology, Guangxi, Guilin 541004, China

Correspondence should be addressed to Qiongfeng Zhang, qfzhangcsu@163.com

Received 20 July 2011; Accepted 16 January 2012

Academic Editor: K. S. Govinder

Copyright © 2012 K. Chen and Q. Zhang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By applying Mountain Pass Theorem in critical point theory, two existence results are obtained for the following asymptotically p -linear p -Laplacian discrete system $\Delta(|\Delta u(t-1)|^{p-2}\Delta u(t-1)) + \nabla[-K(t, u(t)) + W(t, u(t))] = 0$. The results obtained generalize some known works.

1. Introduction

Consider the periodic solutions of the following ordinary p -Laplacian discrete system

$$\Delta\left(|\Delta u(t-1)|^{p-2}\Delta u(t-1)\right) + \nabla F(t, u(t)) = 0, \quad (1.1)$$

where Δ is the forward difference operator defined by $\Delta u(t) = u(t+1) - u(t)$, $\Delta^2 u(t) = \Delta(\Delta u(t))$, $p \in (1, +\infty)$, $t \in \mathbb{Z}$, $u \in \mathbb{R}^N$, $F: \mathbb{Z} \times \mathbb{R}^N \rightarrow \mathbb{R}$, $F(t, x)$ is continuously differentiable in x for every $t \in \mathbb{Z}$ and T -periodic in t for all $x \in \mathbb{R}^N$, and F satisfies $(F) F(t, x) = -K(t, x) + W(t, x)$, and K and W are T -periodic in t for all $x \in \mathbb{R}^N$ with $T > 1$.

Difference equations provide a natural description of many discrete models in real world. Discrete models exist in various fields of science and technology such as statistics, computer science, electrical circuit analysis, biology, neural network, and optimal control; so it is very important to study the solutions of difference equations. For more details about difference equations, we refer the readers to the books [1–3].

In some recent papers [4–17], the authors investigated the existence of periodic solutions and subharmonic solutions of difference equations by applying critical point theory. These papers imply that the critical point theory is a useful method to the study of periodic solutions for difference equations. Motivated by the above papers and the paper [18], we will generalize the results of [18] to p -Laplacian systems (1.1). Here are our main results.

Theorem 1.1. *Assume that F satisfies (F) and K and W satisfy the following conditions:*

(A1) *there exist constants $a_1 > 0$ and $\gamma \in (p - 1, p]$ such that*

$$K(t, 0) = 0, \quad K(t, x) \geq a_1|x|^\gamma, \quad \forall (t, x) \in \mathbb{Z}[1, T] \times \mathbb{R}^N; \quad (1.2)$$

(A2) $(\nabla K(t, x), x) \leq pK(t, x)$ for all $(t, x) \in \mathbb{Z}[1, T] \times \mathbb{R}^N$, where $\mathbb{Z}[a, b] := \mathbb{Z} \cap [a, b]$ for every $a, b \in \mathbb{Z}$ with $a \leq b$;

(A3) $W(t, 0) = 0$, $\limsup_{|x| \rightarrow 0} (W(t, x)/|x|^p) < a_1$ uniformly for $t \in \mathbb{Z}[1, T]$;

(A4) *there exists a function $g \in l^1(\mathbb{Z}[1, T], \mathbb{R})$ such that*

$$(\nabla W(t, x), x) - pW(t, x) \geq g(t) \quad \text{for } (t, x) \in \mathbb{Z}[1, T] \times \mathbb{R}^N, \quad (1.3)$$

$$\lim_{|x| \rightarrow \infty} [(\nabla W(t, x), x) - pW(t, x)] = +\infty \quad \forall t \in \mathbb{Z}[1, T]; \quad (1.4)$$

(A5) *there exist constants $a_2 > 0$ and $d > 0$ such that*

$$W(t, x) \leq a_2|x|^p + d, \quad \forall (t, x) \in \mathbb{Z}[1, T] \times \mathbb{R}^N; \quad (1.5)$$

(A6) *there exists $x_0 \in \mathbb{R}^N$ such that*

$$\sum_{t=1}^T \left[K(t, x_0) - W(t, x_0) - \frac{g(t)}{p} \right] < 0. \quad (1.6)$$

Then problem (1.1) possesses one nontrivial periodic solution.

Corollary 1.2. *Assuming that F satisfies (F) and that K and W satisfy (A1)–(A4) and*

(A7) *there exists a function $V_\infty \in l^\infty(\mathbb{Z}[1, T], \mathbb{R})$ such that*

$$\lim_{|x| \rightarrow \infty} \frac{|\nabla W(t, x) - V_\infty(t)|x|^{p-2}x|}{|x|^{p-1}} = 0 \quad \text{uniformly for } t \in \mathbb{Z}[1, T], \quad (1.7)$$

$$\sum_{t=1}^T \left[\max_{|x|=1} K(t, x) - \frac{V_\infty(t)}{p} \right] < 0. \quad (1.8)$$

Then problem (1.1) possesses one nontrivial periodic solution.

Remark 1.3. As far as we know, similar results of discrete system (1.1) which satisfies (F) and is asymptotically p -linear at infinity cannot be found in the literature. From this point, our results are new.

2. Preliminaries

Let E_T be the Sobolev space defined by

$$E_T = \left\{ u : \mathbb{Z} \longrightarrow \mathbb{R}^N \mid u(t+T) = u(t), t \in \mathbb{Z} \right\}, \quad (2.1)$$

with the norm

$$\|u\| = \left(\sum_{t=1}^T (|\Delta u(t)|^p + |u(t)|^p) \right)^{1/p}, \quad u \in E_T, \quad (2.2)$$

where $|\cdot|$ denote the usual norm in \mathbb{R}^N . It is easy to see that $(E_T, \|\cdot\|)$ is a finite dimensional Banach space and linear homeomorphic to \mathbb{R}^{NT} . As usual, let

$$\|u\|_\infty = \sup\{|u(t)| : t \in \mathbb{Z}[1, T]\}, \quad \forall u \in l^\infty(\mathbb{Z}[1, T], \mathbb{R}^N). \quad (2.3)$$

Since E_T is finite dimensional Banach space, there exists a positive constant C_0 such that

$$\|u\|_\infty \leq C_0 \|u\|. \quad (2.4)$$

For any $u \in E_T$, let

$$\varphi(u) = \frac{1}{p} \sum_{t=1}^T |\Delta u(t)|^p + \sum_{t=1}^T [K(t, u(t)) - W(t, u(t))]. \quad (2.5)$$

We can compute the Fréchet derivative of (2.5) as

$$\frac{\partial \varphi(u)}{\partial u} = \Delta \left(|\Delta u(t-1)|^{p-2} \Delta u(t-1) \right) + \nabla [K(t, u(t)) - W(t, u(t))], \quad t \in \mathbb{Z}[1, T]. \quad (2.6)$$

Hence, u is a critical point of φ on E_T if and only if

$$\Delta \left(|\Delta u(t-1)|^{p-2} \Delta u(t-1) \right) + \nabla [K(t, u(t)) - W(t, u(t))] = 0, \quad t \in \mathbb{Z}[1, T], u \in \mathbb{R}^N. \quad (2.7)$$

So, the critical points of φ are classical solutions of (1.1). We will use the following lemma to prove our main results.

Lemma 2.1 (see [19]). Let E be a real Banach space and $\varphi \in C^1(E, \mathbb{R})$ satisfying the (PS) condition. Suppose $\varphi(0) = 0$ and

- (a) there exist constants $\rho, \alpha > 0$ such that $\varphi|_{\partial B_\rho(0)} \geq \alpha$;
- (b) there exists an $e \in E \setminus \overline{B_\rho(0)}$ such that $\varphi(e) \leq 0$.

Then φ possesses a critical value $c \geq \alpha$ which can be characterized as $c = \inf_{h \in \Gamma} \max_{s \in [0,1]} \varphi(h(s))$, where $\Gamma = \{h \in C([0,1], E) \mid h(0) = 0, h(1) = e\}$ and $B_\rho(0)$ is an open ball in E of radius ρ centered at 0.

It is well known that a deformation lemma can be proved with the weaker condition (C) replacing the usual (PS) condition. So Lemma 2.1 holds true under condition (C).

3. Proofs of Main Results

Proof of Theorem 1.1. The proof is divided into three steps.

Step 1. The functional φ satisfies condition (C). Let $\{u_n\} \subset E_T$ satisfying $(1 + \|u_n\|)\|\varphi'(u_n)\| \rightarrow 0$ as $n \rightarrow \infty$ and $\varphi(u_n)$ is bounded. Hence, there exists a positive constant C_1 such that

$$|\varphi(u_n)| \leq C_1, \quad (1 + \|u_n\|)\|\varphi'(u_n)\| \leq C_1. \quad (3.1)$$

We prove $\{u_n\}$ is bounded by contradiction. If $\{u_n\}$ is unbounded, without loss of generality, we can assume that $\|u_n\| \rightarrow \infty$ as $n \rightarrow \infty$. Let $z_n = u_n / \|u_n\|$, then we have $\|z_n\| = 1$. Going to a subsequence if necessary, we may assume that $z_n \rightharpoonup z$ weakly in E_T and so $z_n \rightarrow z$ strongly in $L^1(\mathbb{Z}[1, T], \mathbb{R})$. It follows from (3.1) and (A2) that

$$\begin{aligned} (p+1)C_1 &\geq p\varphi(u_n) - \langle \varphi'(u_n), u_n \rangle \\ &= \sum_{t=1}^T [(\nabla W(t, u_n), u_n) - pW(t, u_n)] + \sum_{t=1}^T [pK(t, u_n) - (\nabla K(t, u_n), u_n)] \\ &\geq \sum_{t=1}^T [(\nabla W(t, u_n), u_n) - pW(t, u_n)]. \end{aligned} \quad (3.2)$$

From (A1) and (A5), we obtain

$$\begin{aligned} \varphi(u_n) &= \frac{1}{p} \sum_{t=1}^T |\Delta u_n(t)|^p + \sum_{t=1}^T [K(t, u_n(t)) - W(t, u_n(t))] \\ &\geq \frac{1}{p} \sum_{t=1}^T |\Delta u_n(t)|^p - a_2 \sum_{t=1}^T |u_n(t)|^p - dT \\ &= \frac{1}{p} \|u_n\| - \left(\frac{1}{p} + a_2\right) \sum_{t=1}^T |u_n(t)|^p - dT. \end{aligned} \quad (3.3)$$

Hence, we have

$$\frac{\varphi(u_n)}{\|u_n\|^p} \geq \frac{1}{p} - \left(\frac{1}{p} + a_2\right) \sum_{t=1}^T |z_n(t)|^p - \frac{dT}{\|u_n\|^p}. \tag{3.4}$$

Passing to the limit in the above inequality, by using the fact that $\varphi(u_n)$ is bounded and $\{z_n(t)\}$ converges uniformly to $z(t)$ on $\mathbb{Z}[1, T]$, we obtain

$$\left(\frac{1}{p} + a_2\right) \sum_{t=1}^T |z(t)|^p \geq \frac{1}{p}, \tag{3.5}$$

which implies that $z \neq 0$. Let $\Omega \subset \mathbb{Z}[1, T]$ be the set on which $z \neq 0$, then the measure of Ω is positive. Moreover, $|u_n(t)| \rightarrow \infty$ as $n \rightarrow \infty$ for $t \in \Omega$. Thus, from (A4), we get

$$\begin{aligned} \sum_{t=1}^T [(\nabla W(t, u_n), u_n) - pW(t, u_n)] &= \sum_{\Omega} [(\nabla W(t, u_n), u_n) - pW(t, u_n)] \\ &\quad + \sum_{\mathbb{Z}[1, T] \setminus \Omega} [(\nabla W(t, u_n), u_n) - pW(t, u_n)] \\ &\geq \sum_{\Omega} [(\nabla W(t, u_n), u_n) - pW(t, u_n)] + \sum_{\mathbb{Z}[1, T] \setminus \Omega} g(t). \end{aligned} \tag{3.6}$$

It follows from Fatou’s lemma and (A4) that

$$\lim_{n \rightarrow \infty} \sum_{t=1}^T [(\nabla W(t, u_n), u_n) - pW(t, u_n)] = +\infty, \tag{3.7}$$

which contradicts with (3.2). Therefore, $\{u_n\}$ is bounded in E_T . Hence, there exists a subsequence, still denoted by $\{u_n\}$, such that

$$u_n \rightharpoonup u_0 \quad \text{weakly in } E_T. \tag{3.8}$$

Since E_T is finite dimensional space, we have $u_n \rightarrow u_0$ in E_T . Therefore, the functional φ satisfies condition (C).

Step 2. From (A3) and (A5), there exist constants $0 < \varepsilon < 1/p$, $q > p$ and $C_2 > 1/TC_0^p$ such that

$$W(t, u) \leq (a_1 - \varepsilon)|u|^p + C_2|u|^q \quad \text{for } u \in \mathbb{R}^N, t \in \mathbb{Z}[1, T]. \tag{3.9}$$

Let

$$\delta = \left(\frac{p\varepsilon}{qTC_2C_0^p}\right)^{1/(q-p)}. \tag{3.10}$$

Then $0 < \delta < 1$. If $u \in E_T$ and $\|u\| = \delta/C_0 := \rho$, then it follows from (2.4) that $|u(t)| \leq \delta$ for $t \in \mathbb{Z}[1, T]$. Set

$$\alpha = \frac{(q-p)\varepsilon}{q}\rho^p. \quad (3.11)$$

Then from (A1), (2.4), (3.9), (3.10), and (3.11), we have

$$\begin{aligned} \varphi(u) &= \frac{1}{p} \sum_{t=1}^T |\Delta u(t)|^p + \sum_{t=1}^T K(t, u(t)) - \sum_{t=1}^T W(t, u(t)) \\ &\geq \frac{1}{p} \sum_{t=1}^T |\Delta u(t)|^p + a_1 \sum_{t=1}^T |u(t)|^q - (a_1 - \varepsilon) \sum_{t=1}^T |u(t)|^p - C_2 \sum_{t=1}^T |u(t)|^q \\ &\geq \frac{1}{p} \sum_{t=1}^T |\Delta u(t)|^p + \varepsilon \sum_{t=1}^T |u(t)|^p - C_2 \sum_{t=1}^T |u(t)|^q \\ &\geq \varepsilon \left(\sum_{t=1}^T |\Delta u(t)|^p + \sum_{t=1}^T |u(t)|^p \right) - C_2 T \|u\|_\infty^q \geq \varepsilon \|u\|^p - C_2 T C_0^q \|u\|^q \\ &= \frac{(q-p)\varepsilon}{q} \rho^p = \alpha \quad \forall u \in E_T \text{ with } \|u\| = \rho. \end{aligned} \quad (3.12)$$

Step 3. Set $f(s) = s^{-p}W(t, sx_0)$ for $s > 0$. Then it follows from (A4) that

$$f'(s) = s^{-p-1}[-pW(t, sx_0) + (\nabla W(t, sx_0), sx_0)] \geq s^{-p-1}g(t) \quad \forall t \in \mathbb{Z}[1, T], s > 0. \quad (3.13)$$

Integrating the above inequality from 1 to $\xi > 1$, we have

$$W(t, \xi x_0) \geq \xi^p W(t, x_0) + \frac{g(t)}{p}(\xi^p - 1) \quad \forall t \in \mathbb{Z}[1, T], \xi > 1. \quad (3.14)$$

From (A2), it is easy to see that

$$K(t, \xi x_0) \leq \xi^p K(t, x_0) \quad \forall t \in \mathbb{Z}[1, T], \xi > 1. \quad (3.15)$$

From (3.14), (3.15), and (A6), we have

$$\begin{aligned} \varphi(\xi x_0) &= \sum_{t=1}^T [K(t, \xi x_0) - W(t, \xi x_0)] \leq \xi^p \sum_{t=1}^T \left[K(t, x_0) - W(t, x_0) - \frac{g(t)}{p} \right] + \frac{1}{p} \sum_{t=1}^T g(t) \\ &\leq 0 \quad \text{for large enough } \xi > 1. \end{aligned} \quad (3.16)$$

Choose $\xi_1 > 1$ such that $T^{1/p}|\xi_1 x_0| > \rho$ and $\varphi(\xi_1 x_0) \leq 0$. Let $e = \xi_1 x_0$, then $\|e\| = T^{1/p}|\xi_1 x_0| > \rho$ and $\varphi(e) \leq 0$. It is easy to see that $\varphi(0) = 0$. Hence, by Lemma 2.1, there exists $u \in E_T$ such that

$$\varphi(u) = c, \quad \varphi'(u) = 0. \quad (3.17)$$

Then the function u is a desired nontrivial T -periodic solution of (1.1). The proof is complete. \square

Proof of Corollary 1.2. Let $C_3 = -(1/3T) \sum_{t=1}^T [\max_{|x|=1} K(t, x) - (V_\infty(t)/p)]$. Then it follows from (A7) that $C_3 > 0$ and there exists a positive constant $C_4 > 0$ such that

$$\left| \nabla W(t, x) - V_\infty(t)|x|^{p-2}x \right| \leq C_3|x|^{p-1} \quad \text{for } t \in \mathbb{Z}[1, T], |x| \geq C_4. \quad (3.18)$$

For any $x \in \mathbb{R}^N \setminus \{0\}$, let $x_* = C_4 x/|x|$. Then it follows from (3.18) that for all $t \in \mathbb{Z}[1, T]$ and $x \in \mathbb{R}^N$ with $|x| > C_4$

$$\begin{aligned} W(t, x) - \frac{V_\infty(t)}{p}|x|^p &= W(t, x_*) - \frac{1}{p}V_\infty(t)|x_*|^p \\ &\quad + \int_0^1 (\nabla W(t, x_* + s(x - x_*)) - V_\infty(t)(x_* + s(x - x_*)), x - x_*) ds \\ &\leq W(t, x_*) - \frac{C_4^p}{p}V_\infty(t) + C_3 \int_0^1 |x_* + s(x - x_*)||x - x_*| ds \\ &\leq W(t, x_*) - \frac{C_4^p}{p}V_\infty(t) + 2C_3|x|^p, \end{aligned} \quad (3.19)$$

which implies that

$$W(t, x) \leq \left(\frac{V_\infty(t)}{p} + 2C_3 \right) |x|^p + W(t, x_*) - \frac{C_4^p}{p}V_\infty(t) \quad \text{for } t \in \mathbb{Z}[1, T], |x| > C_4, \quad (3.20)$$

which together with (A3) shows that (A5) holds. Similarly, we have

$$W(t, x) \geq \left(\frac{V_\infty(t)}{p} - 2C_3 \right) |x|^p + W(t, x_*) - \frac{C_4^p}{p}V_\infty(t) \quad \text{for } t \in \mathbb{Z}[1, T], |x| > C_4. \quad (3.21)$$

From (A2), it is easy to show that

$$K(t, x) \leq K\left(t, \frac{x}{|x|}\right)|x|^p \quad \text{for } t \in \mathbb{Z}[1, T], |x| > 1. \quad (3.22)$$

Choose $x_0 \in \mathbb{R}^N$ such that $|x_0| > C_4 + 1$ and

$$C_3 T |x_0|^p + \sum_{t=1}^T \left[\min_{|x|=C_4} W(t, x) + \frac{g(t)}{p} - \frac{C_4^p}{p} V_\infty(t) \right] > 0. \quad (3.23)$$

It follows from (3.21), (3.22), and (3.23) that

$$\begin{aligned} & \sum_{t=1}^T \left[K(t, x_0) - W(t, x_0) - \frac{g(t)}{p} \right] \\ & \leq |x_0|^p \sum_{t=1}^T \left[K\left(t, \frac{x_0}{|x_0|}\right) - \frac{V_\infty(t)}{p} + 2C_3 \right] - \sum_{t=1}^T \left[W(t, x_*) + \frac{g(t)}{p} - \frac{C_4^p}{p} V_\infty(t) \right] \\ & \leq |x_0|^p \sum_{t=1}^T \left[\max_{|x|=1} K(t, x) - \frac{V_\infty(t)}{p} + 2C_3 \right] - \sum_{t=1}^T \left[\min_{|x|=C_4} W(t, x) + \frac{g(t)}{p} - \frac{C_4^p}{p} V_\infty(t) \right] \\ & \leq -C_3 T |x_0|^p - \sum_{t=1}^T \left[\min_{|x|=C_4} W(t, x) + \frac{g(t)}{p} - \frac{C_4^p}{p} V_\infty(t) \right] \\ & < 0. \end{aligned} \quad (3.24)$$

This implies that (A6) holds. By Theorem 1.1, the conclusion of Corollary 1.2 holds true. The proof is complete. \square

4. An Example

In this section, we give an example to illustrate our results.

Example 4.1. In problem (1.1), let $p = 4/3$ and

$$W(t, x) = a(t)|x|^{4/3} \left(1 - \frac{1}{\ln(e + |x|)} \right), \quad K(t, x) = b|x|^\theta + c(t)|x|^\sigma, \quad (4.1)$$

where $b > 0$, $a, c \in l^1(\mathbb{Z}, [0, +\infty))$, $1 < \theta < \sigma \leq 4/3$, $a(t+T) = a(t)$, $c(t+T) = c(t)$. It is easy to check that (F), (A1)–(A3), and (A5) hold. On the one hand, we have

$$(\nabla W(t, x), x) - \frac{4}{3} W(t, x) = \frac{(4/3)a(t)|x|^{7/3}}{(e + |x|)(\ln(e + |x|))^2}. \quad (4.2)$$

Then, it is easy to check that condition (A4) holds. On the other hand, we have

$$\begin{aligned} \sum_{t=1}^T \left[K(t, x) - W(t, x) - \frac{g(t)}{p} \right] &= \sum_{t=1}^T \left[b|x|^\theta + c(t)|x|^\sigma - a(t)|x|^{4/3} \left(1 - \frac{1}{\ln(e + |x|)} \right) - \frac{g(t)}{p} \right] \\ &= bT|x|^\theta + |x|^\sigma \sum_{t=1}^T c(t) - \frac{\|g\|_{l^1}}{p} - |x|^{4/3} \left(1 - \frac{1}{\ln(e + |x|)} \right) \sum_{t=1}^T a(t), \end{aligned} \quad (4.3)$$

which implies that there exists $x_0 \in \mathbb{R}^N$ such that (A6) holds if

$$\sum_{t=1}^T a(t) > \sum_{t=1}^T c(t). \quad (4.4)$$

Hence, from Theorem 1.1, problem (1.1) with W and K as in (4.1) has one nontrivial T -periodic solution if (4.4) holds.

Acknowledgment

This work is partially supported by Scientific Research Foundation of Guilin University of Technology and Scientific Research Foundation of Guangxi Education Office of China (200911MS270).

References

- [1] R. P. Agarwal, *Difference Equations and Inequalities*, vol. 228, Marcel Dekker, New York, NY, USA, 2nd edition, 2000.
- [2] C. D. Ahlbrandt and A. C. Peterson, *Discrete Hamiltonian Systems*, vol. 16, Kluwer Academic Publishers Group, Dordrecht, The Netherlands, 1996.
- [3] S. N. Elaydi, *An Introduction to Difference Equations*, Springer, New York, NY, USA, 2nd edition, 1999.
- [4] R. P. Agarwal and J. Popenda, "Periodic solutions of first order linear difference equations," *Mathematical and Computer Modelling*, vol. 22, no. 1, pp. 11–19, 1995.
- [5] R. P. Agarwal, K. Perera, and D. O'Regan, "Multiple positive solutions of singular discrete p -Laplacian problems via variational methods," *Advances in Difference Equations*, no. 2, pp. 93–99, 2005.
- [6] Z. Guo and J. Yu, "Existence of periodic and subharmonic solutions for second-order superlinear difference equations," *Science in China. Series A*, vol. 46, no. 4, pp. 506–515, 2003.
- [7] Z. Guo and J. Yu, "Periodic and subharmonic solutions for superquadratic discrete Hamiltonian systems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 55, no. 7-8, pp. 969–983, 2003.
- [8] Z. Guo and J. Yu, "The existence of periodic and subharmonic solutions of subquadratic second order difference equations," *Journal of the London Mathematical Society Second Series*, vol. 68, no. 2, pp. 419–430, 2003.
- [9] H. Liang and P. Weng, "Existence and multiple solutions for a second-order difference boundary value problem via critical point theory," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 1, pp. 511–520, 2007.
- [10] Z. M. Luo and X. Y. Zhang, "Existence of nonconstant periodic solutions for a nonlinear discrete system involving the p -Laplacian," *Bulletin of the Malaysian Mathematical Science Society*. In press.
- [11] J. Rodriguez and D. L. Etheridge, "Periodic solutions of nonlinear second-order difference equations," *Advances in Difference Equations*, no. 2, pp. 173–192, 2005.
- [12] Y.-F. Xue and C.-L. Tang, "Existence of a periodic solution for subquadratic second-order discrete Hamiltonian system," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 67, no. 7, pp. 2072–2080, 2007.
- [13] J. Yu, Z. Guo, and X. Zou, "Periodic solutions of second order self-adjoint difference equations," *Journal of the London Mathematical Society Second Series*, vol. 71, no. 1, pp. 146–160, 2005.
- [14] J. Yu, Y. Long, and Z. Guo, "Subharmonic solutions with prescribed minimal period of a discrete forced pendulum equation," *Journal of Dynamics and Differential Equations*, vol. 16, no. 2, pp. 575–586, 2004.
- [15] J. Yu, X. Deng, and Z. Guo, "Periodic solutions of a discrete Hamiltonian system with a change of sign in the potential," *Journal of Mathematical Analysis and Applications*, vol. 324, no. 2, pp. 1140–1151, 2006.
- [16] Z. Zhou, J. Yu, and Z. Guo, "Periodic solutions of higher-dimensional discrete systems," *Proceedings of the Royal Society of Edinburgh Section A*, vol. 134, no. 5, pp. 1013–1022, 2004.

- [17] Q. Zhang, X. H. Tang, and Q.-M. Zhang, "Existence of periodic solutions for a class of discrete Hamiltonian systems," *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 463480, 14 pages, 2011.
- [18] X. H. Tang and J. Jiang, "Existence and multiplicity of periodic solutions for a class of second-order Hamiltonian systems," *Computers & Mathematics with Applications*, vol. 59, no. 12, pp. 3646–3655, 2010.
- [19] P. H. Rabinowitz, "Minimax methods in critical point theory with applications to differential equations," in *CBMS Regional Conference Series in Mathematics*, vol. 65, American Mathematical Society, Providence, RI, USA, 1986.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

