

Research Article

On Interpolation Functions of the Generalized Twisted (h, q) -Euler Polynomials

Kyoung Ho Park

Department of Mathematics, Sogang University, Seoul 121-742, South Korea

Correspondence should be addressed to Kyoung Ho Park, sagamath@yahoo.co.kr

Received 5 November 2008; Accepted 14 January 2009

Recommended by Vijay Gupta

The aim of this paper is to construct p -adic twisted two-variable Euler- (h, q) - L -functions, which interpolate generalized twisted (h, q) -Euler polynomials at negative integers. In this paper, we treat twisted (h, q) -Euler numbers and polynomials associated with p -adic invariant integral on \mathbb{Z}_p . We will construct two-variable twisted (h, q) -Euler-zeta function and two-variable (h, q) - L -function in Complex s -plane.

Copyright © 2009 Kyoung Ho Park. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Tsumura and Young treated the interpolation functions of the Bernoulli and Euler polynomials in [1, 2]. Kim and Simsek studied on p -adic interpolation functions of these numbers and polynomials [3–48]. In [49], Carlitz originally constructed q -Bernoulli numbers and polynomials. Many authors studied these numbers and polynomials [4, 28, 38, 41, 50]. After that, twisted (h, q) -Bernoulli and Euler numbers (polynomials) were studied by several authors [1–32, 32–65]. In [62], Whashington constructed one-variable p -adic- L -function which interpolates generalized classical Bernoulli numbers at negative integers. Fox introduced the two-variable p -adic L -functions [53]. Young defined p -adic integral representation for the two-variable p -adic L -functions [64]. Furthermore, Kim constructed the two-variable p -adic q - L -function, which is interpolation function of the generalized q -Bernoulli polynomials [8]. This function is the q -extension of the two-variable p -adic L -function. Kim constructed q -extension of the generalized formula for two-variable of Diamond and Ferrero and Greenberg formula for two-variable p -adic L -function in the terms of the p -adic gamma and log-gamma functions [8]. Kim and Rim introduced twisted q -Euler numbers and polynomials associated with basic twisted q - ℓ -functions [28]. Also, Jang et al. investigated the p -adic analogue twisted q - ℓ -function, which interpolates generalized twisted

q -Euler numbers $E_{n,q,\xi,\chi}$ attached to Dirichlet's character χ [55]. Kim et al. have studied two-variable p -adic L -functions, which interpolate the generalized Bernoulli polynomials at negative integers. In this paper, we will construct two-variable p -adic twisted Euler (h, q) - L -functions. These functions are interpolation functions of the generalized twisted (h, q) -Euler polynomials.

Let p be a fixed odd prime number. Throughout this paper \mathbb{Z} , \mathbb{Z}_p , \mathbb{Q}_p and \mathbb{C}_p will respectively denote the ring of rational integers, the ring of p -adic rational integers, the field of p -adic rational numbers and the completion of the algebraic closure of \mathbb{Q}_p . Let v_p be the normalized exponential valuation of \mathbb{C}_p such that $|p|_p = p^{-v_p(p)} = p^{-1}$. If $s \in \mathbb{C}$, then $|q| < 1$. If $q \in \mathbb{C}_p$, we normally assume $|1 - q|_p < p^{-(1/(p-1))}$, so that $q^x = \exp(\log q)$ for $|x|_p \leq 1$. Throughout this paper we use the following notations (cf. [1–32, 32–48, 50, 51, 54–65]):

$$[x]_q = [x : q] = \frac{1 - q^x}{1 - q}, \quad [x]_{-q} = \frac{1 - (-q)^x}{1 + q}. \quad (1.1)$$

Hence, $\lim_{q \rightarrow 1} [x]_q = x$, for any x with $|x|_p \leq 1$ in the present p -adic case.

For d a fixed positive integer with $(p, d) = 1$, set

$$\begin{aligned} X = X_d &= \lim_{\overline{N}} \frac{\mathbb{Z}}{dp^N \mathbb{Z}}, & X_1 &= \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0 < a < dp, \\ (a,p)=1}} (a + dp\mathbb{Z}_p), \\ a + dp^N \mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\}, \end{aligned} \quad (1.2)$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < dp^N$. The distribution is defined by

$$\mu_q(a + dp^N \mathbb{Z}_p) = \frac{q^a}{[dp^N]_q}. \quad (1.3)$$

We say that f is a uniformly differential function at a point $a \in \mathbb{Z}_p$, and we write $f \in UD(\mathbb{Z}_p)$, if the difference quotients, $F_f(x, y) = (f(x) - f(y))/(x - y)$ have a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$.

For $f \in UD(\mathbb{Z}_p)$, the p -adic invariant q -integral on \mathbb{Z}_p is defined as [4, 18]

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x. \quad (1.4)$$

The fermionic p -adic q -measures on \mathbb{Z}_p is defined as (cf. [14–16, 18, 22, 28])

$$\mu_{-q}(a + dp^N \mathbb{Z}_p) = \frac{(-q)^a}{[dp^N]_{-q}}, \quad (1.5)$$

for $f \in UD(\mathbb{Z}_p)$. For $f \in UD(\mathbb{Z}_p)$, the fermionic p -adic invariant q -integral on \mathbb{Z}_p is defined as

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x)(-q)^x, \tag{1.6}$$

which has a sense as we see readily that the limit is convergent. For $f \in UD(\mathbb{Z}_p, \mathbb{C}_p)$, we note that (cf. [14, 16, 18, 22, 28])

$$\int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \int_X f(x) d\mu_{-1}(x). \tag{1.7}$$

From the fermionic invariant integral on \mathbb{Z}_p , we derive the following integral equation (cf. [14, 35]):

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0), \tag{1.8}$$

where $f_1(x) = f(x + 1)$.

2. Twisted (h, q) -Euler Numbers and Polynomials

In this section, we will treat some properties of twisted (h, q) -Euler numbers and polynomials associated with p -adic invariant integral on \mathbb{Z}_p . From now on, we take $h \in \mathbb{Z}$ and $q \in \mathbb{C}_p$ with $|q - 1|_p < p^{-(1/(p-1))}$. Let C_{p^n} be the space of primitive p^n th root of unity,

$$C_{p^n} = \{w \in \mathbb{C}_{p^n} \mid w^{p^n} = 1\}. \tag{2.1}$$

Then, we denote

$$T_p = \lim_{n \rightarrow \infty} C_{p^n} = \bigcup_{n \geq 0} C_{p^n}. \tag{2.2}$$

Hence T_p is a p -adic locally constant space. For $\xi \in T_p$, we denote by $\phi_\xi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ defined by $\phi_\xi(x) = \xi^x$, the locally constant function. If we take $f(x) = \xi^x e^{xt}$, then we have (cf. [35])

$$E_{n,\xi} = \int_{\mathbb{Z}_p} x^n \xi^n d\mu_{-1}(x). \tag{2.3}$$

By induction in (1.8), Kim constructed the following useful identity (cf. [14, 28]):

$$I_{-1}(f_n) + (-1)^{n-1} I_{-1}(f) = 2 \sum_{\ell=0}^{n-1} (-1)^{n-1-\ell} f(\ell), \tag{2.4}$$

where $n \in \mathbb{N}$, $f_n = f(x+n)$. From (2.4), if n is odd, then we have

$$I_{-1}(f_n) + I_{-1}(f) = 2 \sum_{\ell=0}^{n-1} (-1)^\ell f(\ell). \quad (2.5)$$

If we replace n by d ($=$ odd) into (2.5), we obtain

$$I_{-1}(f_d) + I_{-1}(f) = 2 \sum_{\ell=0}^{d-1} (-1)^\ell f(\ell). \quad (2.6)$$

Let $\xi \in T_p$. Let χ be a Dirichlet's character of conductor d , which d is any multiple of p with $p \equiv 1 \pmod{2}$. By substituting $f(x) = \chi(x)\xi^x e^{xt}$ into (2.6), we have

$$I_{-1}(\chi(x)\xi^x e^{xt}) = \sum_{n=0}^{\infty} E_{n,\xi,\chi} \frac{t^n}{n!}. \quad (2.7)$$

Remark 2.1. In complex case, the generating function of the Euler numbers $E_{n,\xi,\chi}$ is given by (cf. [28])

$$\frac{2 \sum_{\ell=0}^{d-1} (-1)^\ell \chi(\ell) \xi^\ell e^{\ell t}}{\xi^d e^{dt} + 1} = \sum_{n=0}^{\infty} E_{n,\xi,\chi} \frac{t^n}{n!}, \quad |t| < \frac{\pi}{d}. \quad (2.8)$$

By using Taylor series of e^{xt} , then we can define the generalized twisted Euler numbers $E_{n,\xi,\chi}$ attached to χ as follows (cf. [55]):

$$E_{n,\xi,\chi} = \int_X \xi^n x^n \chi(x) d\mu_{-1}(x). \quad (2.9)$$

In [8], (h, q) -Euler numbers were defined by

$$E_{n,q}^{(h,1)}(x) = \int_{\mathbb{Z}_p} q^{(h-1)y} [x+y]_q^n d\mu_{-q}(y), \quad (2.10)$$

where $h \in \mathbb{Z}$ and $x \in \mathbb{Z}_p$. In particular, if we take $x = 0$, then $E_{n,q}^{(h,1)}(0) = E_{n,q}^{(h,1)}$. These numbers are called (h, q) -Euler numbers.

By using iterative method of p -adic invariant integral on \mathbb{Z}_p in the sense of fermionic, we define twisted (h, q) -Euler numbers as follows (cf. [55]):

$$E_{n,q,\xi}^{(h,1)}(x) = \int_{\mathbb{Z}_p} q^{(h-1)y} \phi_\xi(y) [x+y]_q^n d\mu_{-q}(y). \quad (2.11)$$

For $h \in \mathbb{Z}$ and $n \in \mathbb{N}$, we have that (cf. [55])

$$E_{n,q,\xi}^{(h,1)}(x) = \frac{1+q}{(1-q)^n} \sum_{i=0}^n \binom{n}{i} (-1)^i q^{xi} \frac{1}{1+\xi q^{h+i}}, \tag{2.12}$$

$$E_{n,q,\xi}^{(h,1)}(x) = \frac{1+q}{1+q^d} \sum_{a=0}^{d-1} (-1)^a q^{ha} \xi^a E_{n,\xi^d,q^d}^{(h,1)}\left(\frac{x+a}{d}\right) [d]_q^n, \tag{2.13}$$

where $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$.

Let $F_{q,\xi}^{(h,1)}(t, x)$ be the generating function of $E_{n,q,\xi}^{(h,1)}(x)$ in complex plane as follows (cf. [55]):

$$\begin{aligned} F_{q,\xi}^{(h,1)}(t, x) &= (1+q) \sum_{n=0}^{\infty} (-1)^n q^{hn} \xi^n e^{t[n+x]_q} \\ &= \sum_{n=0}^{\infty} E_{n,q,\xi}^{(h,1)}(x) \frac{t^n}{n!}. \end{aligned} \tag{2.14}$$

Let χ be the Dirichlet's character with conductor $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$. Then the generalized twisted (h, q) -Euler polynomials attached to χ is given by as follows:

For $n \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$,

$$E_{n,q,\xi,\chi}^{(h,1)}(x) = \int_X \chi(y) q^{(h-1)y} \xi^y [x+y]_q^n d\mu_{-q}(y), \tag{2.15}$$

where $h \in \mathbb{Z}$, d is any multiple of p with $p \equiv 1 \pmod{2}$ and $x \in \mathbb{C}_p$.

Then the distribution relation of the generalized twisted (h, q) -Euler polynomials is given by as follows (cf. [14]):

$$E_{n,q,\xi,\chi}^{(h,1)}(x) = \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a E_{n,q^d,\xi^d}^{(h,1)}\left(\frac{x+a}{d}\right) [d]_q^n. \tag{2.16}$$

3. Two-Variable Twisted (h, q) -Euler-Zeta Function and (h, q) -L-Function

In this section, we will construct two-variable twisted (h, q) -Euler-zeta function and two-variable (h, q) -L-function in Complex s -plane. We assume $q \in \mathbb{C}$ with $|q| < 1$.

Firstly, we consider twisted q -Euler numbers and polynomials in \mathbb{C} as follows (cf. [55]):

$$\begin{aligned} F_{q,\xi}^{(h,1)}(t, x) &= (1+q) \sum_{n=0}^{\infty} (-1)^n q^{hn} \xi^n e^{t[n+x]_q} \\ &= \sum_{n=0}^{\infty} E_{n,q,\xi}^{(h,1)}(x) \frac{t^n}{n!}, \end{aligned} \tag{3.1}$$

where $q, x \in \mathbb{C}$, $r \in \mathbb{Z}^+ = \mathbb{N} \cup \{0\}$ and ξ is r th root of unity. In particular, if we take $x = 0$, then we have $E_{n,q,\xi}^{(h,1)}(0) = E_{n,q,\xi}^{(h,1)}$. These numbers are called twisted Euler numbers. By using derivative operator, we have $(d^k/dt^k)F_{q,\xi}(t,x)|_{t=0} = E_{n,q,\xi}^{(h,1)}(x)$.

From (3.1), we can define Hurwitz-type twisted (h, q) -Euler-zeta function as follows (cf. [55]):

$$\zeta_{E,q,\xi}^{(h,1)}(s, x) = (1+q) \sum_{k=0}^{\infty} \frac{(-1)^k q^{hk} \xi^k}{[x+k]_q^s}, \quad (3.2)$$

where $q \in \mathbb{C}$, $|q| < 1$, $s \in \mathbb{C}$, $h \in \mathbb{Z}$ and $x \in \mathbb{R}$, $0 < x \leq 1$. Note that if $x = 1$ in (3.2), then we see that the twisted (h, q) -Euler-zeta function is defined by (cf. [28, 55])

$$\zeta_{E,q,\xi}^{(h,1)}(s) = (1+q) \sum_{k=1}^{\infty} \frac{(-1)^k q^{hk} \xi^k}{[k]_q^s}, \quad s \in \mathbb{C}, \operatorname{Re}(s) > 1. \quad (3.3)$$

For $n \in \mathbb{N}$, we know (cf. [28])

$$\zeta_{E,q,\xi}^{(h,1)}(-n, x) = E_{n,q,\xi}^{(h,1)}(x). \quad (3.4)$$

From now on, we will define the two-variable (h, q) - L -functions $L_{E,q,\xi}^{(h,1)}(s, x : \chi)$ which interpolates the generalized (h, q) -Euler polynomials.

Definition 3.1. Let χ be the Dirichlet's character with conductor d with $d \equiv 1 \pmod{2}$. For $s \in \mathbb{C}$, $h \in \mathbb{Z}$ and $x \in \mathbb{R}$, $0 < x \leq 1$, we define

$$L_{E,q,\xi}^{(h,1)}(s, x : \chi) = (1+q) \sum_{n=0}^{\infty} \frac{\chi(n) (-1)^n q^{hn} \xi^n}{[n+x]_q^s}. \quad (3.5)$$

By substituting $n = a + jd$, $d \equiv 1 \pmod{2}$, $1 \leq a \leq d$ and $n = 0, 1, 2, \dots$ into (3.5), then using (3.2), we have

$$\begin{aligned} L_{E,q,\xi}^{(h,1)}(s, x : \chi) &= (1+q) \sum_{a=1}^d \sum_{j=0}^{\infty} \frac{\chi(a+jd) (-1)^{a+jd} q^{h(a+jd)} \xi^{a+jd}}{[a+jd+x]_q^s} \\ &= (1+q) \sum_{a=1}^d \frac{\chi(a) (-1)^a q^{ha} \xi^a}{[d]_q^s} \sum_{j=0}^{\infty} \frac{(-1)^{jd} q^{hjd}}{[j + ((a+x)/d)]_{q^d}^s} \\ &= \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a \zeta_{E,q^d,\xi^d}^{(h,1)}\left(s, \frac{a+x}{d}\right) [d]_q^{-s}. \end{aligned} \quad (3.6)$$

Thus, we see the function $L_{E,q,\xi}^{(h,1)}(s, x : \chi)$ which interpolates the generalized (h, q) -Euler polynomials as follows.

Theorem 3.2. For $s \in \mathbb{C}$, $h \in \mathbb{Z}$, let χ be the Dirichlet's character with conductor d with $d \equiv 1 \pmod{2}$. Then one has

$$L_{E,q,\xi}^{(h,1)}(s, x : \chi) = \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a \zeta_{E,q^d,\xi^d}^{(h,1)} \left(s, \frac{a+x}{d} \right) [d]_q^{-s}. \tag{3.7}$$

By substituting $s = -n$ with $n > 0$, into (3.7), we obtain

$$\begin{aligned} L_{E,q,\xi}^{(h,1)}(-n, x : \chi) &= \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a \zeta_{E,q^d,\xi^d}^{(h,1)} \left(-n, \frac{a+x}{d} \right) [d]_q^n \\ &= \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a E_{n,q^d,\xi^d}^{(h,1)} \left(\frac{a+x}{d} \right) [d]_q^n \\ &= E_{n,q,\xi,\chi}^{(h,1)}(x), \end{aligned} \tag{3.8}$$

where $d \equiv 1 \pmod{2}$, $d \in \mathbb{N}$.

Thus, we have the following theorem.

Theorem 3.3. For $n \in \mathbb{N}$, let χ be the Dirichlet's character with conductor d with $d \equiv 1 \pmod{2}$. Then one has

$$L_{E,q,\xi}^{(h,1)}(-n, x : \chi) = E_{n,q,\xi,\chi}^{(h,1)}(x). \tag{3.9}$$

Remark 3.4. If we take $x = 1$ in (3.5), then we have (cf. [28, 55])

$$L_{E,q,\xi}^{(h,1)}(s, \chi) = (1+q) \sum_{n=1}^{\infty} \frac{\chi(n) (-1)^n q^{hn} \xi^n}{[n]_q^s}, \quad \text{for } s \in \mathbb{C}. \tag{3.10}$$

From (3.9) and (3.10), we have the following corollary.

Corollary 3.5. Let χ be the Dirichlet's character with conductor d with $d \equiv 1 \pmod{2}$. Then one has

$$E_{n,q,\xi,\chi}^{(h,1)}(x) = \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a E_{n,q^d,\xi^d}^{(h,1)} \left(\frac{a+x}{d} \right) [d]_q^n. \tag{3.11}$$

Secondly, we will define two-variable twisted Euler (h, q) -L-function as follows.

Definition 3.6. Let χ be the Dirichlet's character with conductor d with $d \equiv 1 \pmod{2}$, $d \in \mathbb{N}$. For $s \in \mathbb{C}$, $h \in \mathbb{Z}$, $x \in \mathbb{R}$, $0 < x \leq 1$ and $\xi^r = 1$ with $\xi \neq 1$, we define

$$L_{E,q,\xi}^{(h,1)}(s, x : \chi) = (1+q) \sum_{k=0}^{\infty} \frac{\chi(k) (-1)^k q^{hk} \xi^k}{[k+x]_q^s}. \tag{3.12}$$

We consider the well-known identity (cf. [44, 65])

$$\frac{1}{(1-x)^s} = \sum_{j=0}^{\infty} \binom{s+j-1}{j} x^j. \quad (3.13)$$

By using (3.12), we define two-variable twisted Euler (h, q) - L -function as follows:

$$L_{E,q,\xi}^{(h,1)}(s, x : \chi) = (1+q)(1-q)^s \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \binom{s+j-1}{j} \chi(k) (-1)^k \xi^k q^{hk+j(k+x)}. \quad (3.14)$$

We will investigate the relations between $L_{E,q,\xi}^{(h,1)}(s, x : \chi)$ and $L_{E,q,\xi}^{(h,1)}(s, \chi)$ as follows.

Substituting $k = a + jd$, $a = 1, 2, \dots, d$ with $d \equiv 1 \pmod{2}$, $j = 0, 1, 2, \dots$, into (3.12), we have

$$L_{E,q,\xi}^{(h,1)}(s, x : \chi) = (1+q) \sum_{a=1}^d \sum_{j=0}^{\infty} \frac{\chi(a+jd) (-1)^{a+jd} q^{h(a+jd)} \xi^{a+jd}}{[a+jd+x]_q^s}, \quad (3.15)$$

Thus we obtain the following theorem.

Theorem 3.7. For $s \in \mathbb{C}$ with $h \in \mathbb{Z}$, let χ be the Dirichlet character with conductor d with $d \equiv 1 \pmod{2}$ and $x \in \mathbb{R}$, $0 < x \leq 1$, $\xi^r = 1$ with $\xi \neq 1$. Then one has

$$L_{E,q,\xi}^{(h,1)}(s, x : \chi) = \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a \zeta_{E,q^d,\xi^d}^{(h,1)} \left(s, \frac{a+x}{d} \right) [d]_q^{-s}. \quad (3.16)$$

By substituting $s = -n$ with $n \in \mathbb{N}$ into (3.16) and using (3.4), we can obtain

$$\begin{aligned} L_{E,q,\xi}^{(h,1)}(-n, x : \chi) &= \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a \zeta_{E,q^d,\xi^d}^{(h,1)} \left(-n, \frac{a+x}{d} \right) [d]_q^n \\ &= \frac{1+q}{1+q^d} \sum_{a=1}^d \chi(a) (-1)^a q^{ha} \xi^a E_{n,q^d,\xi^d}^{(h,1)} \left(\frac{a+x}{d} \right) [d]_q^n \\ &= E_{n,q,\xi,\chi}^{(h,1)}(x). \end{aligned} \quad (3.17)$$

Thus, we see that the function $L_{E,q,\xi}^{(h,1)}(s, x : \chi)$ interpolates generalized (h, q) -Euler polynomials attached to χ at negative integer values of s as followings.

Theorem 3.8. For $n \in \mathbb{N}$, let χ be the Dirichlet's character with odd conductor d . Then one has

$$L_{E,q,\xi}^{(h,1)}(-n, x : \chi) = E_{n,q,\xi,\chi}^{(h,1)}(x). \quad (3.18)$$

Note that if we take $x = 1$, then Theorem 3.8 reduces to Theorem 3.3.

Let a and F be integers with $F \equiv 1 \pmod{2}$ and $0 < a < F$. For $s \in \mathbb{C}$, we define partial (h, q) -Hurwitz type zeta function $H_{E, q, \xi}^{(h, 1)}(s, a, x \mid F)$ as follows:

$$H_{E, q, \xi}^{(h, 1)}(s, a, x \mid F) = \sum_{\substack{m \equiv a \pmod{F}, \\ m > 0}} \frac{(-1)^m q^{hm} \xi^m}{[m + x]_q^s}. \tag{3.19}$$

By substituting $m = a + jF$, we have

$$\begin{aligned} H_{E, q, \xi}^{(h, 1)}(s, a, x \mid F) &= \sum_{j=0}^{\infty} \frac{(-1)^{a+jF} q^{h(a+jF)} \xi^{a+jF}}{[a + jF + x]_q^s} \\ &= (-1)^a q^{ha} \xi^a [F]_q^{-s} \sum_{j=0}^{\infty} \frac{(-1)^{jF} (q^F)^{hj} (\xi^F)^j}{[((a+x)/F) + j]_{q^F}^s} \\ &= [F]_q^{-s} (-1)^a (q)^{ha} \xi^a \frac{1}{1 + q^F} \sum_{j=0}^{\infty} \frac{(-1)^{jF} (q^F)^{hj} (\xi^F)^j}{[((a+x)/F) + j]_{q^F}^s} \\ &= [F]_q^{-s} \frac{(-1)^a (q)^{ha} \xi^a}{1 + q^F} \zeta_{E, q^F, \xi^F}^{(h, 1)}\left(s, \frac{a+x}{F}\right). \end{aligned} \tag{3.20}$$

By substituting (3.2), for $s = -n$, we get

$$H_{E, q, \xi}^{(h, 1)}(s, a, x \mid F) = [F]_q^n \frac{(-1)^a q^{ha} \xi^a}{1 + q^F} E_{n, q^F, \xi^F}^{(h, 1)}\left(\frac{a+x}{F}\right). \tag{3.21}$$

Equation (3.20) means that the function $H_{E, q, \xi}^{(h, 1)}(s, a, x \mid F)$ interpolates $E_{n, q, \xi}^{(h, 1)}(s, a, x \mid F)$ polynomials at negative integers.

From (3.16) and (3.20), we have the following theorem.

Theorem 3.9. For $s \in \mathbb{C}$, $\xi^r = 1$ with $\xi \neq 1$, let χ be the Dirichlet's character with conductor $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$ and $x \in \mathbb{R}$, $0 < x \leq 1$, F is any multiple of d . Then one has

$$L_{E, q, \xi}^{(h, 1)}(s, x : \chi) = (1 + q) \sum_{a=1}^F \chi(a) (-1)^a H_{E, q, \xi}^{(h, 1)}(s, a, x \mid F). \tag{3.22}$$

Remark 3.10. If we take $s = 0$ in (3.22), then we have

$$\begin{aligned} L_{E, q, \xi}^{(h, 1)}(0, x : \chi) &= (1 + q) \sum_{a=1}^F \chi(a) H_{E, q, \xi}^{(h, 1)}(0, a, x \mid F) \\ &= \frac{1 + q}{1 + q^F} \sum_{a=1}^F \chi(a) (-1)^a q^{ha} \xi^a E_{0, q^F, \xi^F}^{(h, 1)}\left(\frac{a+x}{F}\right). \end{aligned} \tag{3.23}$$

From (2.12), if we take $s = 0$, then we have the following corollary.

Corollary 3.11. For $s \in \mathbb{C}$, $\xi^r = 1$ with $\xi \neq 1$, let χ be the Dirichlet's character with conductor $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$ and $x \in \mathbb{R}$, $0 < x \leq 1$, F is any multiple of d . Then one has

$$L_{E,q,\xi}^{(h,1)}(0, x : \chi) = \frac{(1+q)^2}{(1+q^F)(1+\xi q^h)} \sum_{a=1}^F \chi(a)(-1)^a q^{ha} \xi^a. \quad (3.24)$$

4. p -Adic Twisted Two-Variable Euler (h, q) - L -Functions

In [62], Washington constructed one-variable p -adic- L -function which interpolates generalized classical Bernoulli numbers negative integers. Kim [22] investigated the p -adic analogues of two-variables Euler q - L -function. In this section, we will construct p -adic twisted two-variable Euler- (h, q) - L -functions, which interpolate generalized twisted (h, q) -Euler polynomials at negative integers. Our notations and methods are essentially due to Kim and Washington (cf. [22, 62]).

We assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-(1/(p-1))}$, so that $q^x = \exp(x \log q)$. Let p be an odd prime number. Let ω denote the Teichmüller character having conductor p . For an arbitrary character χ , we define $\chi_n = \chi \omega^{-n}$, where $n \in \mathbb{Z}$, in the sense of the product of characters. Let $\langle a \rangle = \langle a : q \rangle = \omega^{-1}(a)[a]_q = [a]_q / \omega(a)$. Then $\langle a \rangle \equiv 1 \pmod{p^{1+(1/(p-1))}}$. Hence we see that

$$\begin{aligned} \langle a + pt \rangle &= \omega^{-1}(a + pt)[a + pt]_q \\ &= \omega^{-1}(a)[a]_q + \omega^{-1}(a)q^a[pt]_q \\ &\equiv 1 \pmod{p^{1+(1/(p-1))}}, \end{aligned} \quad (4.1)$$

where $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, $(a, p) = 1$.

We denote the subset D of \mathbb{C}_p^* by (cf. [62])

$$D = \{s \in \mathbb{C}_p : |s|_p \leq p^{1-(1/(p-1))}\}. \quad (4.2)$$

Let

$$A_j(x) = \sum_{j=0}^{\infty} a_{n,j} x^n, \quad a_{n,j} \in \mathbb{C}_p, \quad j = 0, 1, 2, \dots, \quad (4.3)$$

be a sequence of power series, each of which converges in a fixed subset D such that

- (1) $a_{n,j} \rightarrow a_{n,0}$ as $j \rightarrow \infty$ for all n, j and
- (2) for each $s \in D$ and $\varepsilon > 0$, there exists $n_0 = n_0(s, \varepsilon)$ such that

$$\left| \sum_{n \geq n_0} a_{n,j} s^n \right|_p < \varepsilon, \quad \text{for } \forall j. \quad (4.4)$$

Then $\lim_{j \rightarrow \infty} A_j(s) = A_0(s)$ for all $s \in D$ (cf. [2, 22, 50, 51, 60, 62]).

Let χ be the Dirichlet's character with conductor d with $d \equiv 1 \pmod{2}$ and let F be a positive multiple of p and d .

Now we set

$$L_{E,p,q,\xi}^{(h,1)}(s, x : \chi) = \frac{1+q}{1+q^F} \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a)(-1)^a \xi^a \langle a+pt \rangle^{-s} \cdot \sum_{j=0}^{\infty} \binom{-s}{j} E_{j,q^F,\xi^F}^{(h,1)} q^{j(a+pt)} \left[\frac{F}{a+pt} \right]_{q^{a+pt}}^j. \tag{4.5}$$

Then $L_{E,p,q,\xi}^{(h,1)}(s, x : \chi)$ is analytic for $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, when $s \in D$. For $t \in \mathbb{C}_p$ with $|t|_p \leq 1$, we have

$$\sum_{j=0}^{\infty} \binom{-s}{j} E_{j,q^F,\xi^F}^{(h,1)} q^{j(a+pt)} \left[\frac{F}{a+pt} \right]_{q^{a+pt}}^j \tag{4.6}$$

is analytic for $s \in D$. It readily follows that

$$\langle a+pt \rangle^s = \omega^{-s}(a) [a+pt]_q^s = \langle a \rangle^s \sum_{m=0}^{\infty} \binom{s}{m} (q^a [a]_q^{-1} [pt]_q)^m \tag{4.7}$$

is analytic for $s \in \mathbb{C}_p$ with $|t|_p \leq 1$ when $s \in D$. Thus we see that

$$L_{E,p,q,\xi}^{(h,1)}(0, x : \chi) = \frac{1+q}{2} \sum_{a=1}^F (-1)^a \chi_n(a) \xi^a. \tag{4.8}$$

Let $n \in \mathbb{Z}_+$ and fixed $t \in \mathbb{C}_p$ with $|t|_p \leq 1$. Then we have that

$$E_{n,q,\xi,\chi_n}^{(h,1)}(pt) = [F]_q^n \frac{1+q}{1+q^F} \sum_{a=0}^F \chi_n(a) (-1)^a \xi^a E_{n,q^F,\xi^F}^{(h,1)} \left(\frac{a+pt}{F} \right). \tag{4.9}$$

If $\chi_n(p) \neq 0$, then $(p, d_{\chi_n}) = 1$, so F/p is a multiple of d_{χ_n} . Therefore, we have

$$\begin{aligned} & \chi_n(p) [p]_q^n E_{n,q^F,\xi^F,\chi_n}^{(h,1)}(t) \\ &= \chi_n(p) [p]_q^n \left\{ \left[\frac{F}{p} \right]_{q^p}^n \frac{1+q^p}{1+q^{pF/p}} \sum_{a=0}^{F/p-1} \chi_n(a) (-1)^a \xi^a E_{n,(q^p)^{F/p},(\xi^p)^{F/p}}^{(h,1)} \left(\frac{a+t}{F/p} \right) \right\} \\ &= [F]_q^n \frac{1+q^p}{1+q^F} \sum_{\substack{a=0 \\ p \nmid a}}^F \chi_n(a) (-1)^a \xi^a E_{n,q^F,\xi^F}^{(h,1)} \left(\frac{a+pt}{F} \right). \end{aligned} \tag{4.10}$$

Then we note that

$$\frac{1+q}{1+q^p} \chi_n(p) [p]_q^n E_{n,q^F, \xi^F, \chi_n}^{(h,1)}(t) = \frac{1+q}{1+q^F} [F]_q^n \sum_{\substack{a=0 \\ p \nmid a}}^F \chi_n(a) (-1)^a \xi^a E_{n,q^F, \xi^F}^{(h,1)}\left(\frac{a+pt}{F}\right). \quad (4.11)$$

The difference of these equations yields

$$E_{n,q, \xi, \chi_n}^{(h,1)}(pt) - \frac{1+q}{1+q^p} \chi_n(p) [p]_q^n E_{n,q^F, \xi^F, \chi_n}^{(h,1)}(t) = \frac{1+q}{1+q^F} [F]_q^n \sum_{\substack{a=0 \\ p \nmid a}}^F \chi_n(a) (-1)^a \xi^a E_{n,q^F, \xi^F}^{(h,1)}\left(\frac{a+pt}{F}\right). \quad (4.12)$$

Using distribution for (h, q) -Euler polynomials, we easily see that

$$E_{n,q^F, \xi^F}^{(h,1)}\left(\frac{a+pt}{F}\right) = [F]_q^{-n} [a+pt]_q^n \sum_{k=0}^n \binom{n}{k} q^{(a+pt)k} \xi^a \left[\frac{F}{a+pt}\right]_{q^{a+pt}}^k E_{k,q^F, \xi^F}^{(h,1)}. \quad (4.13)$$

Since $\chi_n(a) = \chi(a)\omega^{-n}(a)$, for $(a, p) = 1$, and $t \in \mathbb{C}_p$, with $|t|_p \leq 1$, we have

$$\begin{aligned} & E_{n,q, \xi, \chi_n}^{(h,1)}(pt) - \frac{1+q}{1+q^p} \chi_n(p) [p]_q^n E_{n,q^F, \xi^F, \chi_n}^{(h,1)}(t) \\ &= \frac{1+q}{1+q^F} \sum_{a=0}^{F-1} \chi_n(a) (-1)^a \xi^a E_{n,q^F, \xi^F}^{(h,1)}\left(\frac{a+pt}{F}\right) \\ &= \frac{1+q}{1+q^p} \sum_{\substack{a=0, \\ p \nmid a}}^{F-1} \chi_n(a) (-1)^a \xi^a (a+pt)^n \sum_{k=0}^n \binom{n}{k} q^{(a+pt)k} \left[\frac{F}{a+pt}\right]_{q^{a+pt}}^k E_{k,q^F, \xi^F}^{(h,1)}. \end{aligned} \quad (4.14)$$

From (4.5)–(4.14), we can derive that

$$E_{n,q, \xi, \chi_n}^{(h,1)}(pt) - \frac{1+q}{1+q^p} \chi_n(p) [p]_q^n E_{n,q^F, \xi^F, \chi_n}^{(h,1)}(t) = L_{E,p,q, \xi}^{(h,1)}(-n, t : \chi). \quad (4.15)$$

Therefore we obtain the following theorem.

Theorem 4.1. Let F be a positive integral multiple of p and $d (= d_\chi)$ with $F \equiv 1 \pmod{2}$, and let

$$L_{E,p,q, \xi}^{(h,1)}(s, t : \chi) = \frac{1+q}{1+q^d} \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a) (-1)^a \xi^a (a+pt)^{-s} \sum_{m=0}^{\infty} \binom{-s}{m} q^{(a+pt)m} \left[\frac{F}{a+pt}\right]_{q^{a+pt}}^m E_{m,q^F, \xi^F}^{(h,1)}. \quad (4.16)$$

Then $L_{E,p,q,\xi}^{(h,1)}(s, t : \chi)$ is analytic for $t \in \mathbb{C}_p, |t|_p \leq 1$, provides $s \in D$ when $\chi = 1$. Furthermore, for each $n \in \mathbb{Z}_+$, we have

$$L_{E,p,q,\xi}^{(h,1)}(-n, t : \chi) = E_{n,q,\xi,\chi^n}^{(h,1)}(pt) - \frac{1+q}{1+q^p} \chi^n(p) [p]_q^n E_{n,q^p,\xi^p,\chi^n}^{(h,1)}(t). \tag{4.17}$$

Thus we note that $L_{E,p,q,\xi}^{(h,1)}(s, 0 : \chi) = L_{E,p,q,\xi}^{(h,1)}(s, \chi)$ for all $s \in D$, where $L_{E,p,q,\xi}^{(h,1)}(s, \chi)$ is twisted p -adic Euler (h, q) - L -function, (cf. [15, 22]).

We now generalized to two-variable p -adic Euler (h, q) - L -function, $L_{E,p,q,\xi}^{(h,1)}(s, t : \chi)$ which is first defined by the interpolation function

$$H_{E,p,q,\xi}^{(h,1)}(s, a, x | F) = \frac{(-1)^a}{1+q^F} q^{ha} \xi^a \langle a+pt \rangle^{-s} \cdot \sum_{j=0}^{\infty} \binom{-s}{j} q^{j(a+pt)} \left(\frac{[F]_q}{[a+pt]_q} \right)^j E_{j,q^F,\xi^F}^{(h,1)} \tag{4.18}$$

for $s \in \mathbb{Z}_p$.

From (4.18), we have that

$$\begin{aligned} H_{E,p,q,\xi}^{(h,1)}(-n, a, x | F) &= \frac{(-1)^a}{1+q^F} \xi^a q^{ha} \langle a+pt \rangle^n \sum_{j=0}^a \binom{n}{j} q^{(a+pt)j} \left(\frac{[F]_q}{[a]_q} \right)^j E_{j,q^F,\xi^F}^{(h,1)} \\ &= \frac{(-1)^a}{1+q^F} q^{ha} \xi^a \omega^{-n}(a) [F]_q^n E_{n,q^F,\xi^F} \left(\frac{a}{F} \right) \\ &= \omega^{-n}(a) H_{E,q,\xi}^{(h,1)}(-n, a, x | F). \end{aligned} \tag{4.19}$$

By using the definition of $H_{E,p,q,\xi}^{(h,1)}(s, a, x | F)$, we can express $L_{E,p,q,\xi}^{(h,1)}(s, t : \chi)$ for all $a \in \mathbb{Z}, (a, p) = 1$ and $t \in \mathbb{C}_p$ with $|t| \leq 1$ as follows:

$$L_{E,p,q,\xi}^{(h,1)}(s, t : \chi) = \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a) H_{E,p,q,\xi}^{(h,1)}(s, a+pt | F). \tag{4.20}$$

We know that $H_{E,p,q,\xi}^{(h,1)}(s, a+pt | F)$ is analytic for $t \in \mathbb{C}_p, |t| \leq 1$, when $s \in D$. The value of $(\partial/\partial s)L_{E,p,q,\xi}^{(h,1)}(s, t : \chi)$ is the coefficients of s in the expansion of $L_{E,p,q,\xi}^{(h,1)}(s, t : \chi)$ at $s = 0$. Using the Taylor expansion at $s = 0$, we see that

$$\langle a+pt \rangle^{-s} = 1 - s \log \langle a+pt \rangle + \dots, \quad \binom{-s}{m} = \frac{(-1)^m}{m} s + \dots. \tag{4.21}$$

The p -adic logarithmic function, \log_p , is the unique function $\mathbb{C}_p^* \rightarrow \mathbb{C}_p$ that satisfies

$$\begin{aligned}\log_p(1+x) &= \sum_{n=1}^{\infty} \frac{(-1)^n}{n} x^n, \quad |x|_p < 1, \\ \log_p(xy) &= \log_p(x) + \log_p(y), \quad \forall x, y \in \mathbb{C}_p^*, \\ \log_p(p) &= 0.\end{aligned}\tag{4.22}$$

By employing these expansion and some algebraic manipulations, we evaluate the derivative $(\partial/\partial s)L_{E,p,q,\xi}^{(h,1)}(0, t : \chi)$. It follows from the definition of $L_{E,p,q,\xi}(s, t : \chi)$ that

$$\begin{aligned}L_{E,p,q,\xi}^{(h,1)}(s, t : \chi) &= \frac{1+q}{1+q^F} \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a) (-1)^a \xi^a \langle a+pt \rangle^{-s} \\ &\quad \cdot \sum_{m=0}^{\infty} \binom{-s}{m} q^{(a+pt)m} \left[\frac{F}{a+pt} \right]_{q^{a+pt}}^m E_{m,q^F,\xi^F}^{(h,1)}.\end{aligned}\tag{4.23}$$

Thus, we have

$$\begin{aligned}\frac{\partial}{\partial s} L_{E,p,q,\xi}^{(h,1)}(s, t : \chi)|_{s=0} &= \frac{1+q}{1+q^F} \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a) (-1)^a \xi^a \\ &\quad \cdot \left(-\log(a+pt) E_{0,q^F,\xi^F}^{(h,1)} + \sum_{m=1}^{\infty} \frac{(-1)^m}{m} q^{(a+pt)m} \left[\frac{F}{a+pt} \right]_{q^{a+pt}}^m E_{m,q^F,\xi^F}^{(h,1)} \right).\end{aligned}\tag{4.24}$$

Since $\omega(a)$ is a root of unity for $(a, p) = 1$, we have

$$\log_p \langle a+pt \rangle = \log_p(a+pt) + \log_p \omega^{-1}(a) = \log_p(a+pt).\tag{4.25}$$

Thus we have the following theorem.

Theorem 4.2. *Let χ be a primitive Dirichlet's character with odd conductor d , $d \in \mathbb{N}$ and let F be a odd positive integral multiple of p and d . Then for any $t \in \mathbb{C}_p$ with $|t| \leq 1$, one has*

$$\begin{aligned}\frac{\partial}{\partial s} L_{E,p,q,\xi}^{(h,1)}(s, t : \chi) &= \frac{1+q}{1+q^F} \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a) (-1)^a \xi^a \sum_{m=1}^{\infty} \frac{(-1)^m}{m} q^{(a+pt)m} \left(\frac{[F]_q}{[a+pt]_q} \right)^m E_{m,q^F,\xi^F}^{(h,1)} \\ &\quad - \frac{1+q}{2} \sum_{\substack{a=1, \\ p \nmid a}}^F \chi(a) (-1)^a \xi^a \log(a+pt).\end{aligned}\tag{4.26}$$

References

- [1] H. Tsumura, "On a p -adic interpolation of the generalized Euler numbers and its applications," *Tokyo Journal of Mathematics*, vol. 10, no. 2, pp. 281–293, 1987.
- [2] P. T. Young, "Congruences for Bernoulli, Euler, and Stirling numbers," *Journal of Number Theory*, vol. 78, no. 2, pp. 204–227, 1999.
- [3] T. Kim, "On a q -analogue of the p -adic log gamma functions and related integrals," *Journal of Number Theory*, vol. 76, no. 2, pp. 320–329, 1999.
- [4] T. Kim, " q -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [5] T. Kim, "On Euler-Barnes multiple zeta functions," *Russian Journal of Mathematical Physics*, vol. 10, no. 3, pp. 261–267, 2003.
- [6] T. Kim, " q -Riemann zeta function," *International Journal of Mathematics and Mathematical Sciences*, vol. 2004, no. 12, pp. 599–605, 2004.
- [7] T. Kim, "Analytic continuation of multiple q -zeta functions and their values at negative integers," *Russian Journal of Mathematical Physics*, vol. 11, no. 1, pp. 71–76, 2004.
- [8] T. Kim, "Power series and asymptotic series associated with the q -analog of the two-variable p -adic L -function," *Russian Journal of Mathematical Physics*, vol. 12, no. 2, pp. 186–196, 2005.
- [9] T. Kim, " q -generalized Euler numbers and polynomials," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 293–298, 2006.
- [10] T. Kim, "A new approach to p -adic q - L -functions," *Advanced Studies in Contemporary Mathematics*, vol. 12, no. 1, pp. 61–72, 2006.
- [11] T. Kim, "A note on p -adic invariant integral in the rings of p -adic integers," *Advanced Studies in Contemporary Mathematics*, vol. 13, no. 1, pp. 95–99, 2006.
- [12] T. Kim, "Multiple p -adic L -function," *Russian Journal of Mathematical Physics*, vol. 13, no. 2, pp. 151–157, 2006.
- [13] T. Kim, " q -extension of the Euler formula and trigonometric functions," *Russian Journal of Mathematical Physics*, vol. 14, no. 3, pp. 275–278, 2007.
- [14] T. Kim, "On the analogs of Euler numbers and polynomials associated with p -adic q -integral on \mathbb{Z}_p at $q = -1$," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 2, pp. 779–792, 2007.
- [15] T. Kim, "On p -adic q - l -functions and sums of powers," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 2, pp. 1472–1481, 2007.
- [16] T. Kim, "On the q -extension of Euler and Genocchi numbers," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 1458–1465, 2007.
- [17] T. Kim, "A note on p -adic q -integral on \mathbb{Z}_p associated with q -Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 15, no. 2, pp. 133–137, 2007.
- [18] T. Kim, " q -Euler numbers and polynomials associated with p -adic q -integrals," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 15–27, 2007.
- [19] T. Kim, "Euler numbers and polynomials associated with zeta functions," *Abstract and Applied Analysis*, vol. 2008, Article ID 581582, 11 pages, 2008.
- [20] T. Kim, " q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients," *Russian Journal of Mathematical Physics*, vol. 15, no. 1, pp. 51–57, 2008.
- [21] T. Kim, "Note on the Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 17, no. 2, pp. 109–115, 2008.
- [22] T. Kim, "On p -adic interpolating function for q -Euler numbers and its derivatives," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 1, pp. 598–608, 2008.
- [23] T. Kim, "A note on q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 161–170, 2008.
- [24] T. Kim, "The modified q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 161–170, 2008.
- [25] T. Kim, "On a p -adic interpolation function for the q -extension of the generalized Bernoulli polynomials and its derivative," *Discrete Mathematics*, vol. 309, no. 6, pp. 1593–1602, 2009.
- [26] T. Kim, "Note on the Euler q -zeta functions," *Journal of Number Theory*. In press.
- [27] T. Kim, J. Y. Choi, and J. Y. Sug, "Extended q -Euler numbers and polynomials associated with fermionic p -adic q -integral on \mathbb{Z}_p ," *Russian Journal of Mathematical Physics*, vol. 14, no. 2, pp. 160–163, 2007.
- [28] T. Kim and S.-H. Rim, "On the twisted q -Euler numbers and polynomials associated with basic q - l -functions," *Journal of Mathematical Analysis and Applications*, vol. 336, no. 1, pp. 738–744, 2007.

- [29] H. Ozden, I. N. Cangul, and Y. Simsek, "Remarks on sum of products of (h, q) -twisted Euler polynomials and numbers," *Journal of Inequalities and Applications*, vol. 2008, Article ID 816129, 8 pages, 2008.
- [30] H. Ozden, Y. Simsek, and I. N. Cangul, "Euler polynomials associated with p -adic q -Euler measure," *General Mathematics*, vol. 15, no. 2, pp. 24–37, 2007.
- [31] H. Ozden and Y. Simsek, "A new extension of q -Euler numbers and polynomials related to their interpolation functions," *Applied Mathematics Letters*, vol. 21, no. 9, pp. 934–939, 2008.
- [32] I. N. Cangul, H. Ozden, and Y. Simsek, "Generating functions of the (h, q) extension of twisted Euler polynomials and numbers," *Acta Mathematica Hungarica*, vol. 120, no. 3, pp. 281–299, 2008.
- [33] H. Ozden, I. N. Cangul, and Y. Simsek, "Multivariate interpolation functions of higher-order q -Euler numbers and their applications," *Abstract and Applied Analysis*, vol. 2008, Article ID 390857, 16 pages, 2008.
- [34] Y. Simsek, O. Yurekli, and V. Kurt, "On interpolation functions of the twisted generalized Frobenius-Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 15, no. 2, pp. 187–194, 2007.
- [35] S.-H. Rim and T. Kim, "A note on q -Euler numbers associated with the basic q -zeta function," *Applied Mathematics Letters*, vol. 20, no. 4, pp. 366–369, 2007.
- [36] Y. Simsek, "Theorems on twisted L -function and twisted Bernoulli numbers," *Advanced Studies in Contemporary Mathematics*, vol. 11, no. 2, pp. 205–218, 2005.
- [37] Y. Simsek, " q -analogue of twisted l -series and q -twisted Euler numbers," *Journal of Number Theory*, vol. 110, no. 2, pp. 267–278, 2005.
- [38] Y. Simsek, "On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 340–348, 2006.
- [39] Y. Simsek, "Hardy character sums related to Eisenstein series and theta functions," *Advanced Studies in Contemporary Mathematics*, vol. 12, no. 1, pp. 39–53, 2006.
- [40] Y. Simsek, "Remarks on reciprocity laws of the Dedekind and Hardy sums," *Advanced Studies in Contemporary Mathematics*, vol. 12, no. 2, pp. 237–246, 2006.
- [41] Y. Simsek, "Twisted (h, q) -Bernoulli numbers and polynomials related to twisted (h, q) -zeta function and L -function," *Journal of Mathematical Analysis and Applications*, vol. 324, no. 2, pp. 790–804, 2006.
- [42] Y. Simsek, "On twisted q -Hurwitz zeta function and q -two-variable L -function," *Applied Mathematics and Computation*, vol. 187, no. 1, pp. 466–473, 2007.
- [43] Y. Simsek, "The behavior of the twisted p -adic (h, q) - L -functions at $s = 0$," *Journal of the Korean Mathematical Society*, vol. 44, no. 4, pp. 915–929, 2007.
- [44] Y. Simsek, "Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 251–278, 2008.
- [45] Y. Simsek, D. Kim, and S.-H. Rim, "On the two-variable Dirichlet q - L -series," *Advanced Studies in Contemporary Mathematics*, vol. 10, no. 2, pp. 131–142, 2005.
- [46] Y. Simsek and A. Mehmet, "Remarks on Dedekind eta function, theta functions and Eisenstein series under the Hecke operators," *Advanced Studies in Contemporary Mathematics*, vol. 10, no. 1, pp. 15–24, 2005.
- [47] H. Ozden, I. N. Cangul, and Y. Simsek, "On the behavior of two variable twisted p -adic Euler q - l -functions," *Nonlinear Analysis*. In press.
- [48] Y. Simsek and S. Yang, "Transformation of four Titchmarsh-type infinite integrals and generalized Dedekind sums associated with Lambert series," *Advanced Studies in Contemporary Mathematics*, vol. 9, no. 2, pp. 195–202, 2004.
- [49] L. Carlitz, " q -Bernoulli and Eulerian numbers," *Transactions of the American Mathematical Society*, vol. 76, pp. 332–350, 1954.
- [50] M. Cenkci and M. Can, "Some results on q -analogue of the Lerch zeta function," *Advanced Studies in Contemporary Mathematics*, vol. 12, no. 2, pp. 213–223, 2006.
- [51] M. Cenkci, Y. Simsek, and V. Kurt, "Further remarks on multiple p -adic q - l -function of two variables," *Advanced Studies in Contemporary Mathematics*, vol. 14, no. 1, pp. 49–68, 2007.
- [52] A. Dąbrowski, "A note on p -adic q - ζ -functions," *Journal of Number Theory*, vol. 64, no. 1, pp. 100–103, 1997.
- [53] G. J. Fox, "A p -adic L -function of two variables," *L'Enseignement Mathématique, IIe Série*, vol. 46, no. 3–4, pp. 225–278, 2000.
- [54] L.-C. Jang, S.-D. Kim, D.-W. Park, and Y.-S. Ro, "A note on Euler number and polynomials," *Journal of Inequalities and Applications*, vol. 2006, Article ID 34602, 5 pages, 2006.

- [55] L.-C. Jang, V. Kurt, Y. Simsek, and S. H. Rim, " q -analogue of the p -adic twisted l -function," *Journal of Concrete and Applicable Mathematics*, vol. 6, no. 2, pp. 169–176, 2008.
- [56] N. Koblitz, "On Carlitz's q -Bernoulli numbers," *Journal of Number Theory*, vol. 14, no. 3, pp. 332–339, 1982.
- [57] K. H. Park and Y.-H. Kim, "On some arithmetical properties of the Genocchi numbers and polynomials," *Advances in Difference Equations*, 14 pages, 2008.
- [58] S.-H. Rim, K. H. Park, and E. J. Moon, "On Genocchi numbers and polynomials," *Abstract and Applied Analysis*, vol. 2008, Article ID 898471, 7 pages, 2008.
- [59] W. H. Schikhof, *Ultrametric Calculus: An Introduction to p -Adic Analysis*, vol. 4 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, UK, 1984.
- [60] K. Shiratani and S. Yamamoto, "On a p -adic interpolation function for the Euler numbers and its derivatives," *Memoirs of the Faculty of Science, Kyushu University. Series A*, vol. 39, no. 1, pp. 113–125, 1985.
- [61] H. M. Srivastava, T. Kim, and Y. Simsek, " q -Bernoulli numbers and polynomials associated with multiple q -zeta functions and basic L -series," *Russian Journal of Mathematical Physics*, vol. 12, no. 2, pp. 241–268, 2005.
- [62] L. C. Washington, *Introduction to Cyclotomic Fields*, vol. 83 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 2nd edition, 1997.
- [63] C. F. Woodcock, "Special p -adic analytic functions and Fourier transforms," *Journal of Number Theory*, vol. 60, no. 2, pp. 393–408, 1996.
- [64] P. T. Young, "On the behavior of some two-variable p -adic L -functions," *Journal of Number Theory*, vol. 98, no. 1, pp. 67–88, 2003.
- [65] J. Zhao, "Multiple q -zeta functions and multiple q -polylogarithms," *Ramanujan Journal*, vol. 14, no. 2, pp. 189–221, 2007.