

Research Article

Weak and Strong Convergence Theorems for Equilibrium Problems and Countable Strict Pseudocontractions Mappings in Hilbert Space

Rudong Chen,¹ Xilin Shen,² and Shujun Cui¹

¹ Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, China

² Department of Mathematics, Xinxiang College, Xinxiang, Henan 453002, China

Correspondence should be addressed to Rudong Chen, tjcrd@yahoo.com.cn

Received 27 August 2009; Accepted 10 January 2010

Academic Editor: Jong Kim

Copyright © 2010 Rudong Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce two iterative sequence for finding a common element of the set of solutions of an equilibrium problem and the set of fixed points of a countable family of strict pseudocontractions in Hilbert Space. Then we study the weak and strong convergence of the sequences.

1. Introduction

Let C be a nonempty closed convex subset of a Hilbert space H and let T be a self-mapping of C . Then T is said to be a strict pseudocontraction mappings if for all $x, y \in C$, there exists a constant $0 \leq \kappa < 1$ such that

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + \kappa\|(I - T)x - (I - T)y\|^2 \quad (1.1)$$

(if (1.1) holds, we also say that T is a κ -strict pseudocontraction). We use $F(T)$ to denote the set of fixed points of T , \rightharpoonup (\rightarrow) to denote weak(strong) convergence, and $W_w(x_n) = \{x : \exists x_{n_k} \rightharpoonup x\}$ to denote the W -limit set of $\{x_n\}$.

Let $f : C \times C \rightarrow R$ be a bifunction where R is the set of real numbers. Then, we consider the following equilibrium problem:

$$\text{Find } z \in C \text{ such that } f(z, y) \geq 0, \quad \forall y \in C. \quad (1.2)$$

The set of such $z \in C$ is denoted by $EP(f)$. Numerous problems in physics, optimization, and economics can be reduced to find a solution of (1.2). Some methods have been proposed to solve the equilibrium problem (see [1–3]). Recently, S. Takahashi and W. Takahashi [4] introduced an iterative scheme by the viscosity approximation method for finding a common element of the set of solutions of the equilibrium problem and the set of fixed points of a nonexpansive mapping in Hilbert spaces. They also studied the strong convergence of the sequences generated by their algorithm for a solution of the EP which is also a fixed point of a nonexpansive mapping defined on a closed convex subset of a Hilbert space.

In this paper, thanks to the condition introduced by Aoyama et al. [5], We introduce two iterative sequence for finding a common element of the set of solutions of an equilibrium problems and the set of fixed points of a countable family of strict pseudocontractions mappings in Hilbert Space. Then we study the weak and strong convergence of the sequences. The additional condition is inspired by Marino and Xu [6] and Kim and Xu [7].

2. Preliminaries

For solving the equilibrium problem, let us assume that the bifunction f satisfies the following conditions (see [3]):

- (A1) $f(x, x) = 0$ for all $x \in C$;
- (A2) f is monotone, that is, $f(x, y) + f(y, x) \leq 0$ for any $x, y \in C$;
- (A3) f is upper-hemicontinuous, that is, for each $x, y, z \in C$, $\limsup_{t \rightarrow 0^+} f(tz + (1 - t)x, y) \leq f(x, y)$;
- (A4) $f(x, \cdot)$ is convex and lower semicontinuous for each $x \in C$.

Let H be a real Hilbert space. Then there hold the following well-known results:

$$\begin{aligned} \|tx + (1 - t)y\|^2 &= t\|x\|^2 + (1 - t)\|y\|^2 - t(1 - t)\|x - y\|^2, \quad \forall x, y \in H, \forall t \in [0, 1]; \\ \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2\langle x, y \rangle, \quad \forall x, y \in H. \end{aligned} \quad (2.1)$$

If $\{x_n\}$ is a sequence in H weakly convergent to z , then

$$\limsup_{n \rightarrow \infty} \|x_n - y\|^2 = \limsup_{n \rightarrow \infty} \|x_n - z\|^2 + \|z - y\|^2 \quad \forall y \in H. \quad (2.2)$$

Recall that the nearest point projection P_C from H onto C assigns to each $x \in H$ its nearest point denoted by $P_C x$ in C ; that is, $P_C x$ is the unique point in C with the property

$$\|x - P_C x\| \leq \|x - y\|, \quad \forall y \in C. \quad (2.3)$$

Given $x \in H$ and $z \in C$, then $z = P_C x$ if and only if there holds the following relation:

$$\langle x - z, y - z \rangle \leq 0, \quad \forall y \in C. \quad (2.4)$$

Lemma 2.1 (see [6]). Let C be a nonempty closed convex subset of a real Hilbert space H . Let $T: C \rightarrow C$ be a κ -strict pseudocontraction such that $F(T) \neq \emptyset$.

- (1) (Demi-closed principle) T is demi-closed on C , that is, if $x_n \rightarrow x \in C$ and $x_n - Tx_n \rightarrow 0$, then $x = Tx$.
- (2) T satisfies the Lipschitz condition

$$\|Tx - Ty\| \leq L\|x - y\| = \frac{1 + \kappa}{1 - \kappa}\|x - y\| \quad \forall x, y \in C. \quad (2.5)$$

- (3) The fixed point set $F(T)$ of T is closed and convex so that the projection $P_{F(T)}$ is well defined.

Lemma 2.2 (see [5]). Let C be a nonempty closed convex subset of a Banach space and let $\{T_n\}$ be a sequence of mapping of C into itself. Suppose $\sum_{n=1}^{\infty} \sup_{x \in C} \|T_{n+1}x - T_nx\| < \infty$. Then, for each $y \in C$, $\{T_n y\}$ converges strongly to some point of C . Moreover, let T be a mapping of C into itself defined by

$$Ty = \lim_{n \rightarrow \infty} T_n y \quad \forall y \in C. \quad (2.6)$$

Then $\lim_{n \rightarrow \infty} \sup_{x \in C} \|T_n x - Tx\| = 0$.

Lemma 2.3 (see [8]). Let C be a closed convex subset of H . Let $\{x_n\}$ be a sequence in H and $u \in H$. Let $q = P_C u$. If $\{x_n\}$ is such that $W_w(x_n) \subset C$ and satisfies the condition

$$\|x_n - u\| \leq \|u - q\| \quad \forall n, \quad (2.7)$$

then $x_n \rightarrow q$.

Lemma 2.4 (see [9]). Let C be a nonempty closed convex subset of H . Let f be a bifunction from $C \times C$ into \mathbb{R} satisfying (A1), (A2), (A3), and (A4). Then, for any $\lambda > 0$ and $x \in H$, there exists $z \in C$ such that

$$f\langle z, y \rangle + \frac{1}{\lambda} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C. \quad (2.8)$$

Further, if $T_\lambda x = \{z \in C : f\langle z, y \rangle + (1/\lambda) \langle y - z, z - x \rangle \geq 0, \forall y \in C\}$, then the following holds:

- (1) $T_\lambda x$ is single-valued;
- (2) $T_\lambda x$ is firmly nonexpansive, that is,

$$\|T_\lambda x - T_\lambda y\|^2 \leq \langle T_\lambda x - T_\lambda y, x - y \rangle, \quad \forall x, y \in H; \quad (2.9)$$

- (3) $F(T_\lambda) = EP(f)$;
- (4) $EP(f)$ is closed and convex.

3. Weak Convergence Theorems

Theorem 3.1. *Let C be a nonempty closed convex subset of a Hilbert space H and let $\{T_n\}$ be a sequence of κ_n -strict pseudocontractions mappings on C into itself with $0 \leq \kappa_n < 1$. Assume that $\kappa = \max\{\kappa_n : n \geq 1\}$. Let $f : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A1), (A2), (A3), (A4), and $\text{EP}(f) \cap \bigcap_{n=1}^{\infty} F(T_n) \neq \emptyset$. Let $\{x_n\}$ and $\{z_n\}$ be sequence generated by $x_1 \in C$ and*

$$\begin{aligned} f(z_n, y) + \frac{1}{r_n} \langle y - z_n, z_n - x_n \rangle &\geq 0, \quad \forall y \in C, \\ x_{n+1} &= \alpha_n z_n + (1 - \alpha_n) T_n z_n, \quad \forall n \geq 1. \end{aligned} \quad (3.1)$$

Assume that $\{\alpha_n\} \subset [0, 1]$ with $\kappa + \delta < \alpha_n < 1 - \delta$ for all n , where $\delta \in (0, 1)$ is a small enough constant, and $\{r_n\}$ is a sequence in $(0, \infty)$ with $\liminf_{n \rightarrow \infty} r_n > 0$ and $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$. Let $\sum_{n=1}^{\infty} \sup_{x \in B} \|T_{n+1}x - T_nx\| < \infty$ for any bounded subset B of C and let T be a mapping of C into itself defined by $Tx = \lim_{n \rightarrow \infty} T_nx$ for all $x \in C$ and suppose that $F(T) = \bigcap_{n=1}^{\infty} F(T_n)$. Then the sequences $\{x_n\}$ and $\{z_n\}$ converge weakly to an element of $F(T) \cap \text{EP}(f)$.

Proof. Pick $p \in F(T) \cap \text{EP}(f)$. Then from the definition of T_r in Lemma 2.4, we have $z_n = T_{r_n}x_n$, and therefore $\|z_n - p\| = \|T_{r_n}x_n - T_{r_n}p\| \leq \|x_n - p\|$. It follows from (3.1) that

$$\begin{aligned} \|x_{n+1} - p\|^2 &= \|(1 - \alpha_n)(T_n z_n - p) + \alpha_n(z_n - p)\|^2 \\ &= \alpha_n \|z_n - p\|^2 + (1 - \alpha_n) \|T_n z_n - p\|^2 - \alpha_n(1 - \alpha_n) \|z_n - T_n z_n\|^2 \\ &\leq \alpha_n \|z_n - p\|^2 + (1 - \alpha_n) \left(\|z_n - p\|^2 + \kappa \|z_n - T_n z_n\|^2 \right) \\ &\quad - \alpha_n(1 - \alpha_n) \|z_n - T_n z_n\|^2 \\ &= \|z_n - p\|^2 - (\alpha_n - \kappa)(1 - \alpha_n) \|z_n - T_n z_n\|^2 \\ &\leq \|x_n - p\|^2 - (\alpha_n - \kappa)(1 - \alpha_n) \|z_n - T_n z_n\|^2. \end{aligned} \quad (3.2)$$

Since $\kappa + \delta < \alpha_n < 1 - \delta$ for all n , we get $\|x_{n+1} - p\| \leq \|x_n - p\|$; that is, the sequence $\{\|x_n - p\|\}$ is decreasing. Hence $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. In particular, $\{x_n\}$ is bounded. Since T_r is firmly nonexpensive, $\{z_n\}$ is also bounded. Also (3.2) implies that

$$\|z_n - T_n z_n\|^2 \leq \frac{1}{\delta^2} \left(\|x_n - p\|^2 - \|x_{n+1} - p\|^2 \right). \quad (3.3)$$

Taking the limit as $n \rightarrow \infty$ yields that

$$\lim_{n \rightarrow \infty} \|z_n - T_n z_n\| = 0. \quad (3.4)$$

Since $\{z_n\}$ is bounded, it follows that

$$\sum_{n=1}^{\infty} \sup_{x \in \{z_n\}} \|T_{n+1}x - T_nx\| < \infty. \quad (3.5)$$

We apply Lemma 2.2 to get

$$\begin{aligned} \|z_n - Tz_n\| &\leq \|z_n - T_n z_n\| + \|T_n z_n - Tz_n\| \\ &\leq \|z_n - T_n z_n\| + \sup\{\|T_n z - Tz\| : z \in \{z_n\}\} \longrightarrow 0. \end{aligned} \quad (3.6)$$

Next, we claim that $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$. Indeed, let p be an arbitrary element of $F(T) \cap EP(f)$. Then as above

$$\begin{aligned} \|z_n - p\|^2 &= \|T_r x_n - T_r p\|^2 \\ &\leq \langle T_r x_n - T_r p, x_n - p \rangle \\ &= \langle z_n - T_r p, x_n - p \rangle \\ &= \frac{1}{2} \left(\|z_n - p\|^2 + \|x_n - p\|^2 - \|x_n - z_n\|^2 \right), \end{aligned} \quad (3.7)$$

and hence

$$\|z_n - p\|^2 \leq \|x_n - p\|^2 - \|x_n - z_n\|^2. \quad (3.8)$$

Therefore, from (3.2), we have

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq \|z_n - p\|^2 - (\alpha_n - \kappa)(1 - \alpha_n) \|z_n - T_n z_n\|^2 \\ &\leq \|z_n - p\|^2 \\ &\leq \|x_n - p\|^2 - \|x_n - z_n\|^2, \end{aligned} \quad (3.9)$$

and hence

$$\|x_n - z_n\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2. \quad (3.10)$$

So, from the existence of $\lim_{n \rightarrow \infty} \|x_n - p\|$, we have

$$\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0. \quad (3.11)$$

Next, we claim that $W_w(x_n) \subset F(T) \cap EP(f)$. since $\{x_n\}$ is bounded and H is reflexive, $W_w(x_n)$ is nonempty. Let $w \in W_w(x_n)$ be an arbitrary element. Then a subsequence x_{n_i} of $\{x_n\}$ converges weakly to w . Hence, from (3.11) we know that $z_{n_i} \rightharpoonup w$. As $\|z_n - Tz_n\| \rightarrow 0$, we obtain that $Tz_{n_i} \rightharpoonup w$. Let us show $W_w(x_n) \subset EP(f)$. Since $z_n = T_{r_n} x_n$, we have

$$f(z_n, y) + \frac{1}{r_n} \langle y - z_n, z_n - x_n \rangle \geq 0, \quad \forall y \in C. \quad (3.12)$$

By (A2), we have

$$\frac{1}{r_n} \langle y - z_n, z_n - x_n \rangle \geq f(y, z_n), \quad (3.13)$$

and hence

$$\left\langle y - z_{n_i}, \frac{z_{n_i} - x_{n_i}}{r_{n_i}} \right\rangle \geq f(y, z_{n_i}). \quad (3.14)$$

From (A4), we have

$$0 \geq f(y, w) \quad \forall y \in C. \quad (3.15)$$

Then, for $t \in (0, 1]$ and $y \in C$, from (A1), and (A4), we also have

$$\begin{aligned} 0 &= f(ty + (1-t)w, ty + (1-t)w) \\ &\leq tf(ty + (1-t)w, y) + (1-t)f(ty + (1-t)w, w) \\ &\leq tf(ty + (1-t)w, y), \end{aligned} \quad (3.16)$$

Taking $t \rightarrow 0^+$ and using (A3), we get

$$f(w, y) \geq 0 \quad \forall y \in C, \quad (3.17)$$

and hence $w \in \text{EP}(f)$. Since T is a strict pseudocontraction mapping, by Lemma 2.1(1) we know that the mapping T is demiclosed at zero. Note that $\|z_n - Tz_n\| \rightarrow 0$ and $z_{n_i} \rightharpoonup w$. Thus, $w \in F(T)$. Consequently, we deduce that $w \in F(T) \cap \text{EP}(f)$. Since w is an arbitrary element, we conclude that $W_w(x_n) \subset F(T) \cap \text{EP}(f)$.

To see that $\{x_n\}$ and $\{z_n\}$ are actually weakly convergent, we take $\bar{x}, \tilde{x} \in W_w(x_n)$ ($x_{n_i} \rightharpoonup \bar{x}, x_{m_j} \rightharpoonup \tilde{x}$). Since $\lim_{n \rightarrow \infty} \|x_n - p\|$ exist for every $p \in F(T)$, by (2.2), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - \tilde{x}\|^2 &= \lim_{i \rightarrow \infty} \|(x_{n_i} - \tilde{x})\|^2 \\ &= \lim_{i \rightarrow \infty} \|x_{n_i} - \bar{x}\|^2 + \|\bar{x} - \tilde{x}\|^2 \\ &= \lim_{j \rightarrow \infty} \|x_{m_j} - \bar{x}\|^2 + \|\bar{x} - \tilde{x}\|^2 \\ &= \lim_{j \rightarrow \infty} \|x_{m_j} - \tilde{x}\|^2 + 2\|\bar{x} - \tilde{x}\|^2 \\ &= \lim_{n \rightarrow \infty} \|x_n - \tilde{x}\|^2 + 2\|\bar{x} - \tilde{x}\|^2. \end{aligned} \quad (3.18)$$

Hence $\tilde{x} = \bar{x}$ and proof is completed. \square

4. Strong Convergence Theorems

Theorem 4.1. Let C be a closed convex subset of a real Hilbert space H . Let $\{T_n\}$ be a sequence of κ_n -strict pseudocontractions mappings on C into itself with $0 \leq \kappa_n < 1$. Assume that $\kappa = \max\{\kappa_n : n \geq 1\}$. Let $f : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A1), (A2), (A3), (A4) and $\text{EP}(f) \cap \bigcap_{n=1}^{\infty} F(T_n) \neq \emptyset$. For $C_1 = C$ and $x_1 = P_{C_1}x_0$, let $\{x_n\}$ and $\{z_n\}$ be sequence generated by $x_0 \in C$ and

$$\begin{aligned} y_n &= \alpha_n x_n + (1 - \alpha_n) T_n x_n, \\ z_n &\in C \text{ such that } f(z_n, y) + \frac{1}{r_n} \langle y - z_n, z_n - y_n \rangle \geq 0, \quad \forall y \in C, \\ C_{n+1} &= \{v \in C : \|z_n - v\| \leq \|x_n - v\|\}, \\ x_{n+1} &= P_{C_{n+1}} x_0, \quad n \geq 1. \end{aligned} \tag{4.1}$$

Assume that $\{\alpha_n\} \subset [0, 1]$ with $\kappa + \delta < \alpha_n < 1 - \delta$ for all n , where $\delta \in (0, 1)$ is a small enough constant, and $\{r_n\}$ is a sequence in $(0, \infty)$ with $\liminf_{n \rightarrow \infty} r_n > 0$ and $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$. Let $\sum_{n=1}^{\infty} \sup_{x \in B} \|T_{n+1}x - T_nx\| < \infty$ for any bounded subset B of C and let T be a mapping of C into itself defined by $Tx = \lim_{n \rightarrow \infty} T_nx$ for all $x \in C$. Suppose that $F(T) = \bigcap_{n=1}^{\infty} F(T_n)$. Then, $\{x_n\}$ converges strongly to $P_{F(T) \cap \text{EP}(f)} x_0$.

Proof. First, we show that C_n is closed and convex. It is obvious that $C_1 = C$ is closed and convex. Suppose that C_k is closed and convex for some $k \geq 1$. For $z \in C_k$, we know that $\|z_k - z\| \leq \|x_k - z\|$ is equivalent to

$$\|z_k - x_k\|^2 + 2\langle z_k - x_k, x_k - z \rangle \leq 0. \tag{4.2}$$

So C_{k+1} is closed and convex. Then, C_n is closed and convex.

Next, we show by induction that $F(T) \cap \text{EP}(f) \subset C_n$ for all $n \geq 1$. $F(T) \cap \text{EP}(f) \subset C_1$ is obvious. Suppose that $F(T) \cap \text{EP}(f) \subset C_k$ for some $k \geq 1$. Let $p \in F(T) \cap \text{EP}(f) \subset C_k$. Putting $z_n = T_{r_n} y_n$ for all n , we know from (4.1) that

$$\begin{aligned} \|z_n - p\|^2 &= \|T_{r_n} y_n - p\|^2 \\ &\leq \|y_n - p\|^2 \\ &= \|(1 - \alpha_n)(T_n x_n - p) + \alpha_n(x_n - p)\|^2 \\ &= \alpha_n \|x_n - p\|^2 + (1 - \alpha_n) \|T_n x_n - p\|^2 - \alpha_n(1 - \alpha_n) \|x_n - T_n x_n\|^2 \\ &\leq \alpha_n \|x_n - p\|^2 + (1 - \alpha_n) (\|x_n - p\|^2 + \kappa \|x_n - T_n x_n\|^2) \\ &\quad - \alpha_n(1 - \alpha_n) \|x_n - T_n x_n\|^2 \end{aligned}$$

$$\begin{aligned}
&= \|x_n - p\|^2 - (\alpha_n - \kappa)(1 - \alpha_n)\|x_n - T_n x_n\|^2 \\
&\leq \|x_n - p\|^2 - \delta^2 \|x_n - T_n x_n\|^2 \\
&\leq \|x_n - p\|^2,
\end{aligned} \tag{4.3}$$

and hence $p \in C_{k+1}$. This implies that $F(T) \cap \text{EP}(f) \subset C_n$ for all $n \geq 1$.

This implied that $\{x_n\}$ is well defined.

From $x_n = P_{C_n} x_0$, we have

$$\|x_0 - x_n\| \leq \|x_0 - y\| \quad \forall y \in C_n. \tag{4.4}$$

Using $F(T) \cap \text{EP}(f) \subset C_n$, we have

$$\|x_0 - x_n\| \leq \|x_0 - u\| \quad \forall u \in F(T) \cap \text{EP}(f), \quad n \geq 1. \tag{4.5}$$

Then, $\{x_n\}$ is bounded. So are $\{y_n\}$ and $\{z_n\}$. In particular,

$$\|x_0 - x_n\| \leq \|x_0 - p\| \quad \text{where } p = P_{F(T) \cap \text{EP}(f)} x_0. \tag{4.6}$$

From $x_n = P_{C_n} x_0$ and $x_{n+1} = P_{C_{n+1}} x_0 \in C_{n+1} \subset C_n$, we have

$$\|x_0 - x_n\| \leq \|x_0 - x_{n+1}\|. \tag{4.7}$$

Since $\{\|x_n - x_0\|\}$ is bounded, $\lim_{n \rightarrow \infty} \|x_n - x_0\|$ exists. From $x_n = P_{C_n} x_0$ and $x_{n+1} = P_{C_{n+1}} x_0 \in C_{n+1} \subset C_n$, we also have

$$\langle x_0 - x_n, x_n - x_{n+1} \rangle \geq 0. \tag{4.8}$$

In fact, from (4.8), we have

$$\begin{aligned}
\|x_n - x_{n+1}\| &= \|x_n - x_0 + x_0 - x_{n+1}\|^2 \\
&= \|x_0 - x_{n+1}\|^2 - \|x_0 - x_n\|^2 - 2\langle x_0 - x_n, x_n - x_{n+1} \rangle \\
&\leq \|x_0 - x_{n+1}\|^2 - \|x_0 - x_n\|^2.
\end{aligned} \tag{4.9}$$

Since $\lim_{n \rightarrow \infty} \|x_n - x_0\|$ exists, we have that $\|x_n - x_{n+1}\| \rightarrow 0$. On the other hand $x_{n+1} \in C_{n+1} \subset C_n$ implies that

$$\|z_n - x_{n+1}\| \leq \|x_n - x_{n+1}\| \rightarrow 0. \tag{4.10}$$

Further, we have

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \quad (4.11)$$

From (4.3), we have

$$\|x_n - T_n x_n\|^2 \leq \frac{1}{\delta^2} \left(\|x_n - p\|^2 - \|z_n - p\|^2 \right). \quad (4.12)$$

On the other hand, we have

$$\begin{aligned} \|x_n - p\|^2 - \|z_n - p\|^2 &= \|x_n\|^2 - \|z_n\|^2 + 2\langle z_n - x_n, p \rangle \\ &\leq \|x_n - z_n\|(\|x_n\| + \|z_n\|) + 2\|p\|\|x_n - z_n\|. \end{aligned} \quad (4.13)$$

Then, we have

$$\lim_{n \rightarrow \infty} \|x_n - p\|^2 - \|z_n - p\|^2 = 0. \quad (4.14)$$

Therefore, we have

$$\lim_{n \rightarrow \infty} \|x_n - T_n x_n\| = 0. \quad (4.15)$$

We apply Lemma 2.2 to get

$$\begin{aligned} \|x_n - T x_n\| &\leq \|x_n - T_n x_n\| + \|T_n x_n - T x_n\| \\ &\leq \|x_n - T_n x_n\| + \sup\{\|T_n x - T x\| : x \in \{x_n\}\} \longrightarrow 0. \end{aligned} \quad (4.16)$$

Lastly, we show that the sequence $\{x_n\}$ converges to $P_{F(T) \cap EP(f)} x_0$. Since $\{x_n\}$ is bounded and H is reflexive, $W_w(x_n)$ is nonempty. Let $w \in W_w(x_n)$ be an arbitrary element. Then a subsequence x_{n_i} of $\{x_n\}$ converges weakly to w . From Lemma 2.1 and (4.16), we obtain that $\omega_w(x_n) \subset F(T)$. Next, we show $W_w(x_n) \subset EP(f)$. Let p be an arbitrary element of $F(T) \cap EP(f)$. From $z_n = T_{r_n} y_n$ and $\|y_n - p\| \leq \|x_n - p\|$, we have

$$\begin{aligned} \|z_n - p\|^2 &\leq \|T_{r_n} y_n - T_{r_n} p\|^2 \\ &\leq \langle T_{r_n} y_n - T_{r_n} p, y_n - p \rangle \\ &= \langle z_n - T_{r_n} p, y_n - p \rangle \\ &= \frac{1}{2} \left(\|z_n - p\|^2 + \|y_n - p\|^2 - \|y_n - z_n\|^2 \right) \\ &= \frac{1}{2} \left(\|z_n - p\|^2 + \|x_n - p\|^2 - \|y_n - z_n\|^2 \right), \end{aligned} \quad (4.17)$$

and hence

$$\|y_n - z_n\|^2 \leq \|x_n - p\|^2 - \|z_n - p\|^2. \quad (4.18)$$

Therefore, we have

$$\lim_{n \rightarrow \infty} \|y_n - z_n\| = 0. \quad (4.19)$$

As in the proof of Theorem 3.1, we have

$$f(z_n, y) + \frac{1}{r_n} \langle y - z_n, z_n - y_n \rangle \geq 0, \quad \forall y \in C. \quad (4.20)$$

By (A2), we have

$$\frac{1}{r_n} \langle y - z_n, z_n - y_n \rangle \geq f(y, z_n), \quad (4.21)$$

and hence

$$\langle y - z_{n_i}, \frac{z_{n_i} - y_{n_i}}{r_{n_i}} \rangle \geq f(y, z_{n_i}). \quad (4.22)$$

From (A4), we have

$$0 \geq f(y, w) \quad \forall y \in C. \quad (4.23)$$

Then, for $t \in (0, 1]$ and $y \in C$, from (A1) and (A4), we also have

$$\begin{aligned} 0 &= f(ty + (1-t)w, ty + (1-t)w) \\ &\leq tf(ty + (1-t)w, y) + (1-t)f(ty + (1-t)w, w) \\ &\leq tf(ty + (1-t)w, y). \end{aligned} \quad (4.24)$$

Taking $t \rightarrow 0^+$ and using (A3), we get

$$f(w, y) \geq 0 \quad \forall y \in C, \quad (4.25)$$

and hence $w \in EP(f)$. Lemma 2.3 and (4.6) ensure the strong convergence of $\{x_n\}$ to $P_{F(T) \cap EP(f)}x_0$. This completes the proof. \square

Acknowledgment

This work is supported by the National Science Foundation of China, Grant 10771050.

References

- [1] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [2] S. D. Flam and A. S. Antipin, "Equilibrium programming using proximal-like algorithms," *Mathematical Programming*, vol. 78, no. 1, pp. 29–41, 1997.
- [3] A. Moudafi and M. Thera, "Proximal and dynamical approaches to equilibrium problems," in *Ill-Posed Variational Problems and Regularization Techniques (Trier, 1998)*, vol. 477 of *Lecture Notes in Economics and Mathematical Systems*, pp. 187–201, Springer, New York, NY, USA, 1999.
- [4] S. Takahashi and W. Takahashi, "Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 1, pp. 506–515, 2007.
- [5] K. Aoyama, Y. Kimura, W. Takahashi, and M. Toyoda, "Approximation of common fixed points of a countable family of nonexpansive mappings in a Banach space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 8, pp. 2350–2360, 2007.
- [6] G. Marino and H.-K. Xu, "Weak and strong convergence theorems for strict pseudo-contractions in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 336–346, 2007.
- [7] T.-H. Kim and H.-K. Xu, "Strong convergence of modified Mann iterations for asymptotically nonexpansive mappings and semigroups," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 5, pp. 1140–1152, 2006.
- [8] C. Martinez-Yanes and H.-K. Xu, "Strong convergence of the CQ method for fixed point iteration processes," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 11, pp. 2400–2411, 2006.
- [9] A. Tada and W. Takahashi, "Weak and strong convergence theorems for a nonexpansive mapping and an equilibrium problem," *Journal of Optimization Theory and Applications*, vol. 133, no. 3, pp. 359–370, 2007.