

Research Article

New Iterative Schemes for Asymptotically Quasi-Nonexpansive Mappings

Yonghong Yao¹ and Yeong-Cheng Liou²

¹ Department of Mathematics, Tianjin Polytechnic University, Tianjin 300160, China

² Department of Information Management, Cheng Shiu University, Kaohsiung 833, Taiwan

Correspondence should be addressed to Yonghong Yao, yaoyonghong@yahoo.cn

Received 20 December 2009; Accepted 15 January 2010

Academic Editor: Yeol Je Cho

Copyright © 2010 Y. Yao and Y.-C. Liou. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider an iterative scheme for approximating the common fixed points of two asymptotically quasi-nonexpansive mappings in the intermediate sense in Banach spaces. The present results improve and extend some recent corresponding results of Lan (2006) and many others.

1. Introduction

Let C be a nonempty subset of a real Banach space E . Let $T : C \rightarrow C$ be a mapping. We use $F(T)$ to denote the set of fixed points of T . Recall that a mapping $T : C \rightarrow C$ is said to be generalized asymptotically quasi-nonexpansive with respect to $\{\sigma_n\}$ and $\{\delta_n\}$ if there exists the sequences $\{\sigma_n\}$ and $\{\delta_n\} \subset [0, 1)$ with $\sigma_n \rightarrow 0$ and $\delta_n \rightarrow 0$ as $n \rightarrow \infty$ such that

$$\|T^n(x) - p\| \leq (1 + \sigma_n)\|x - p\| + \delta_n\|x - T^n(x)\| \quad (1.1)$$

for all $x \in C$, $p \in F(T)$ and $n \geq 1$. It is clear that if $F(T)$ is nonempty, then the asymptotically nonexpansive mapping, the asymptotically quasi-nonexpansive mapping, and the generalized quasi-nonexpansive mapping are all the generalized asymptotically quasi-nonexpansive mapping.

Recall also that a mapping $T : C \rightarrow C$ is said to be asymptotically quasi-nonexpansive in the intermediate sense provided that T is uniformly continuous and

$$\limsup_{n \rightarrow \infty} \sup_{x \in C, p \in F(T)} (\|T^n(x) - p\| - \|x - p\|) \leq 0. \quad (1.2)$$

Remark 1.1. From the above definition, if $F(T)$ is nonempty, it is easy to see that the generalized asymptotically quasi-nonexpansive mapping must be the asymptotically quasi-nonexpansive mapping in the intermediate sense.

It is well known that the concept of asymptotically nonexpansive mapping, which is closely related to the theory of fixed points in Banach spaces, is introduced by Goebel and Kirk [1]. An early fundamental result due to Goebel and Kirk [1] proved that every asymptotically nonexpansive mapping of a nonempty closed bounded and convex subset of a uniformly convex Banach space has a fixed point. Since 1972, the weak and strong convergence problems of iterative sequences (with errors) for nonexpansive mappings, asymptotically nonexpansive mappings in the setting of Hilbert space or Banach space, have been studied by many authors; please see, for example, [1–29] and the references therein. Recently, Zhou et al. [30] introduced a class of new generalized asymptotically nonexpansive mappings and gave a sufficient and necessary condition for the modified Ishikawa and Mann iterative sequences to converge to fixed points for the class of mappings. Define the Ishikawa iterative process involving the generalized asymptotically nonexpansive mappings in a Banach space E as follows.

$$\begin{aligned}x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T(y_n), \\y_n &= (1 - \beta_n)x_n + \beta_n T(x_n), \quad n = 1, 2, \dots\end{aligned}\tag{1.3}$$

where $\{\alpha_n\}_{n=0}^{\infty}$ and $\{\beta_n\}_{n=0}^{\infty}$ are two real sequences in $[0, 1]$ satisfying some conditions. For details, we can refer to [31–33]. Very recently, Lan [3] introduced a new class of iterative procedures as follows:

Let $T_i : C \rightarrow C$ ($i = 1, 2$) be given mappings. Then, for arbitrary $\omega \in C$ and $x_1 \in C$, the sequence $\{x_n\}$ in C defined by

$$\begin{aligned}y_n &= (1 - \beta_n)\omega + \beta_n T_2^n(x_n), \\x_{n+1} &= (1 - \alpha_n)\omega + \alpha_n T_1^n(y_n), \quad n = 1, 2, \dots,\end{aligned}\tag{1.4}$$

is called the generalized modified Ishikawa iterative sequence.

Further, Lan [3] remarked that the above iterative processes include many iterative processes as special cases and he gave a sufficient and necessary condition for the iterative sequence to converge to the common fixed points for two generalized asymptotically quasi-nonexpansive mappings.

It is our purpose in this paper that we will extend the above iterative processes to the more general iterative processes and give a sufficient and necessary condition for two asymptotically quasi-nonexpansive mapping in the intermediate sense. Our result extends the corresponding results of Lan [3], Zhou et al. [30], and many others.

2. Preliminaries

Let C be a nonempty closed convex subset of a real Banach space E . Let $T_i : C \rightarrow C$ ($i = 1, 2$) be given mappings. For given $x_1 \in C$, the sequence $\{x_n\}$ in C defined iteratively by

$$\begin{aligned} y_n &= (1 - \beta_n)\omega_n + \beta_n T_2^n(x_n), \\ x_{n+1} &= (1 - \alpha_n)v_n + \alpha_n T_1^n(y_n), \quad n = 1, 2, \dots \end{aligned} \quad (2.1)$$

is called the more general modified Ishikawa iterative sequence, where $\{\omega_n\}$ and $\{v_n\}$ are sequences in C , and $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$ satisfying some conditions.

If we replace ω_n and v_n in all the iteration steps by ω , then the sequence $\{x_n\}$ defined by (2.1) becomes to (1.4) which is studied by Lan [3].

If we replace ω_n and v_n in all the iteration steps by x_n and ω , respectively, then the sequence $\{x_n\}$ defined by (2.1) becomes to

$$\begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_n T_2^n(x_n), \\ x_{n+1} &= (1 - \alpha_n)\omega + \alpha_n T_1^n(y_n), \quad n = 1, 2, \dots \end{aligned} \quad (2.2)$$

If we replace ω_n and v_n in all the iteration steps by x_n , then the sequence $\{x_n\}$ defined by (2.1) becomes to

$$\begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_n T_2^n(x_n), \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T_1^n(y_n), \quad n = 1, 2, \dots \end{aligned} \quad (2.3)$$

If $\beta_n = v_n = 0$ in (2.1), then the sequence $\{x_n\}$ defined by

$$x_{n+1} = (1 - \alpha_n)v_n + \alpha_n T_1^n(x_n), \quad n = 1, 2 \quad (2.4)$$

is called the more general modified Mann iterative sequence.

It is clear that the iterative processes (2.1) include many iterative processes as special cases.

In the sequel, we need the following lemmas for the main results in this paper.

Lemma 2.1 (see [32]). *Let $\{a_n\}$, $\{b_n\}$, and $\{\delta_n\}$ be sequences of nonnegative real numbers satisfying the inequality*

$$a_{n+1} \leq (1 + \delta_n)a_n + b_n. \quad (2.5)$$

If $\sum_{n=1}^{\infty} \delta_n < \infty$ and $\sum_{n=1}^{\infty} b_n < \infty$, then $\lim_{n \rightarrow \infty} a_n$ exists. In particular, if $\{a_n\}$ has a subsequence converging to zero, then $\lim_{n \rightarrow \infty} a_n = 0$.

3. Main Results

Theorem 3.1. Let C be a nonempty closed convex subset of a real Banach space E . Let $T_i : C \rightarrow C$ ($i = 1, 2$) be asymptotically quasi-nonexpansive mappings in the intermediate sense such that $F(T_1) \cap F(T_2) \neq \emptyset$. Let $\omega_n \in C, \nu_n \in C$ be two bounded sequences. For any given $x_1 \in C$, let the sequences $\{x_n\}$ and $\{y_n\}$ be defined by (2.1). Put

$$G_n = \max \left\{ \sup_{p \in F(T_1) \cap F(T_2), n \geq 1} (\|T_2^n(x_n) - p\| - \|x_n - p\|) \vee 0, \right. \\ \left. \sup_{p \in F(T_1) \cap F(T_2), n \geq 1} (\|T_1^n(y_n) - p\| - \|y_n - p\|) \vee 0 \right\}. \quad (3.1)$$

Assume that $\sum_{n=1}^{\infty} G_n < \infty$, $\sum_{n=1}^{\infty} (1 - \alpha_n) < \infty$ and $\sum_{n=1}^{\infty} (1 - \beta_n) < \infty$.

Then the sequence $\{x_n\}$ converges strongly to a common fixed point p^* of T_1 and T_2 if and only if

$$\liminf_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2)) = 0, \quad (3.2)$$

where $d(x, F(T_1) \cap F(T_2))$ denotes the distance between x and the set $F(T_1) \cap F(T_2)$.

Proof. The necessity is obvious and so it is omitted.

Now, we prove the sufficiency. For any $p \in F(T_1) \cap F(T_2)$, from (2.1), we have

$$\begin{aligned} \|y_n - p\| &= \|(1 - \beta_n)\omega_n + \beta_n T_2^n(x_n) - p\| \\ &\leq (1 - \beta_n)\|\omega_n - p\| + \beta_n \|T_2^n(x_n) - p\| \\ &= (1 - \beta_n)\|\omega_n - p\| + \beta_n (\|T_2^n(x_n) - p\| - \|x_n - p\|) + \beta_n \|x_n - p\| \\ &\leq \beta_n \|x_n - p\| + \beta_n G_n + (1 - \beta_n)\|\omega_n - p\|, \end{aligned} \quad (3.3)$$

$$\begin{aligned} \|x_{n+1} - p\| &= \|(1 - \alpha_n)\nu_n + \alpha_n T_1^n(y_n) - p\| \\ &\leq (1 - \alpha_n)\|\nu_n - p\| + \alpha_n \|T_1^n(y_n) - p\| \\ &= (1 - \alpha_n)\|\nu_n - p\| + \alpha_n (\|T_1^n(y_n) - p\| - \|y_n - p\|) + \alpha_n \|y_n - p\| \\ &\leq \alpha_n \|y_n - p\| + \alpha_n G_n + (1 - \alpha_n)\|\nu_n - p\|. \end{aligned} \quad (3.4)$$

Substituting (3.3) into (3.4) and simplifying, we have

$$\begin{aligned} \|x_{n+1} - p\| &\leq \alpha_n \beta_n \|x_n - p\| + \alpha_n (1 + \beta_n) G_n \\ &\quad + \alpha_n (1 - \beta_n)\|\omega_n - p\| + (1 - \alpha_n)\|\nu_n - p\| \\ &\leq \|x_n - p\| + b_n, \end{aligned} \quad (3.5)$$

where

$$b_n = \alpha_n(1 + \beta_n)G_n + \alpha_n(1 - \beta_n)\|\omega_n - p\| + (1 - \alpha_n)\|v_n - p\|. \quad (3.6)$$

We note that $\sum_{n=1}^{\infty} G_n < \infty$, $\sum_{n=1}^{\infty} (1 - \beta_n) < \infty$, $\sum_{n=1}^{\infty} (1 - \alpha_n) < \infty$ and $\|\omega_n - p\|, \|v_n - p\|$ are bounded; therefore, we have $\sum_{n=1}^{\infty} b_n < \infty$. Then, from (3.5), we have

$$d(x_{n+1}, F(T_1) \cap F(T_2)) \leq d(x_n, F(T_1) \cap F(T_2)) + b_n. \quad (3.7)$$

By Lemma 2.1, we know that $\lim_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2))$ exists. Because $\liminf_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2)) = 0$, then

$$\lim_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2)) = 0. \quad (3.8)$$

Next we prove that $\{x_n\}$ is a Cauchy sequence in C .

It follows from (3.5) that for any $m \geq 1$, for all $n \geq n_0$, for all $p \in F(T_1) \cap F(T_2)$,

$$\begin{aligned} \|x_{n+m} - p\| &\leq \|x_{n+m-1} - p\| + b_{n+m-1} \leq \|x_{n+m-2} - p\| + (b_{n+m-1} + b_{n+m-2}) \\ &\leq \dots \leq \|x_n - p\| + \sum_{k=n}^{n+m-1} b_k. \end{aligned} \quad (3.9)$$

So we have

$$\|x_{n+m} - x_n\| \leq \|x_{n+m} - p\| + \|x_n - p\| \leq 2\|x_n - p\| + \sum_{k=n}^{\infty} b_k. \quad (3.10)$$

Then, we have

$$\|x_{n+m} - x_n\| \leq 2d(x_n, F(T_1) \cap F(T_2)) + \sum_{k=n}^{\infty} b_k, \quad \forall n \geq n_0. \quad (3.11)$$

For any given $\epsilon > 0$, there exists a positive integer $n_1 \geq n_0$ such that for any $n \geq n_1$, $d(x_n, F(T_1) \cap F(T_2)) < \epsilon/4$ and $\sum_{k=n}^{\infty} b_k < \epsilon/2$. Thus when $n \geq n_1$, $\|x_{n+m} - x_n\| < \epsilon$. So we have that

$$\lim_{n \rightarrow \infty} \|x_{n+m} - x_n\| = 0. \quad (3.12)$$

This implies that $\{x_n\}$ is a Cauchy sequence in E . Thus, the completeness of E implies that $\{x_n\}$ must be convergent. Assume that $x_n \rightarrow p^*$ as $n \rightarrow \infty$.

Now we have to prove that p^* is a common fixed point of T_1 and T_2 . Indeed, we know that the set $F(T_1) \cap F(T_2)$ is closed. From the continuity of $d(x, F(T_1) \cap F(T_2)) = 0$

with $\lim_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2)) = 0$ and $\lim_{n \rightarrow \infty} x_n = p^*$, we get

$$d(p^*, F(T_1) \cap F(T_2)) = 0, \quad (3.13)$$

and so $p^* \in F(T_1) \cap F(T_2)$. This completes the proof. \square

We can conclude immediately Theorem 3.1 in [3], which can be reviewed as a corollary of Theorem 3.1.

Corollary 3.2. *Let C be a nonempty closed convex subset of a real Banach space E and for $i = 1, 2$, let $T_i : C \rightarrow C$ be a generalized asymptotically quasi-nonexpansive mapping with respect to $\{\sigma_{in}\}$ and $\{\delta_{in}\}$ such that $F(T_1) \cap F(T_2) \neq \emptyset$ in C , and $\sum_{n=1}^{\infty} (\sigma_n + 2\delta_n) / (1 - \delta_n) < \infty$, where $\sigma_n = \max\{\sigma_{1n}, \sigma_{2n}\}$ and $\delta_n = \max\{\delta_{1n}, \delta_{2n}\}$. Assume that $\sum_{n=1}^{\infty} (1 - \alpha_n) < \infty$ and $\sum_{n=1}^{\infty} (1 - \beta_n) < \infty$.*

Then the iterative sequence $\{x_n\}$ defined by (1.4) converges strongly to a common fixed point p^ of T_1 and T_2 if and only if*

$$\liminf_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2)) = 0, \quad (3.14)$$

where $d(x, F(T_1) \cap F(T_2))$ denotes the distance between x and the set $F(T_1) \cap F(T_2)$.

Proof. We note that condition $\sum_{n=1}^{\infty} (\sigma_n + 2\delta_n) / (1 - \delta_n) < \infty$ implies $\sum_{n=1}^{\infty} \sigma_n < \infty$ and $\sum_{n=1}^{\infty} \delta_n < \infty$. From the boundedness of $\{x_n\}$, $\{y_n\}$, and (1.1), we can obtain $\sum_{n=1}^{\infty} G_n < \infty$. It is easy to see that all conditions of Theorem 3.1 are satisfied; it follows from Theorem 3.1; we can conclude our desired result. This completes the proof. \square

Theorem 3.3. *Let C be a nonempty closed convex subset of a real Banach space E . Let $T_i : C \rightarrow C$ ($i = 1, 2$) be asymptotically quasi-nonexpansive mappings in the intermediate sense such that $F(T_1) \cap F(T_2) \neq \emptyset$. For any given $x_1 \in C$, let the sequences $\{x_n\}$ and $\{y_n\}$ be defined by (2.3). Assume that $\sum_{n=1}^{\infty} \alpha_n < \infty$.*

Then the sequence $\{x_n\}$ converges strongly to a common fixed point p^ of T_1 and T_2 if and only if*

$$\liminf_{n \rightarrow \infty} d(x_n, F(T_1) \cap F(T_2)) = 0. \quad (3.15)$$

Proof. The necessity is obvious and so it is omitted.

Now, we prove the sufficiency. For any $p \in F(T_1) \cap F(T_2)$, it follows from (1.2) that for any given $\epsilon > 0$, there exists a positive integer n_1 such that for $n \geq n_1$, we have

$$\max \left\{ \begin{array}{l} \sup_{p \in F(T_1) \cap F(T_2), n \geq n_1} (\|T_2^n(x_n) - p\| - \|x_n - p\|), \\ \sup_{p \in F(T_1) \cap F(T_2), n \geq n_1} (\|T_1^n(y_n) - p\| - \|y_n - p\|) < \epsilon \end{array} \right\}. \quad (3.16)$$

From (2.3), we have

$$\begin{aligned} \|y_n - p\| &\leq (1 - \beta_n)\|x_n - p\| + \beta_n(\|T_2^n(x_n) - p\| - \|x_n - p\|) + \beta_n\|x_n - p\| \\ &\leq \|x_n - p\| + \beta_n\epsilon, \end{aligned} \quad (3.17)$$

$$\begin{aligned} \|x_{n+1} - p\| &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n(\|T_1^n(y_n) - p\| - \|y_n - p\|) + \alpha_n\|y_n - p\| \\ &\leq (1 - \alpha_n)\|x_n - p\| + \alpha_n\epsilon + \alpha_n\|y_n - p\|. \end{aligned} \quad (3.18)$$

Substituting (3.17) into (3.18), we have

$$\|x_{n+1} - p\| \leq \|x_n - p\| + \alpha_n(1 + \beta_n)\epsilon = \|x_n - p\| + b_n, \quad (3.19)$$

where $\sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} \alpha_n(1 + \beta_n)\epsilon < \infty$. The rest proof follows as those of Theorem 3.1 and therefore is omitted. This completes the proof. \square

From Theorem 3.1, we can obtain the following results.

Theorem 3.4. *Let C be a nonempty closed convex subset of a real Banach space E . Let $T : C \rightarrow C$ be asymptotically quasi-nonexpansive mappings in the intermediate sense such that $F(T) \neq \emptyset$. Let $\{v_n\} \subset C$ be bounded sequence. For any given $x_1 \in C$, let the sequence $\{x_n\}$ be defined by (2.4). Put*

$$G_n = \sup_{p \in F(T), n \geq 1} (\|T^n(x_n) - p\| - \|x_n - p\|) \vee 0. \quad (3.20)$$

Assume that $\sum_{n=1}^{\infty} G_n < \infty$ and $\sum_{n=1}^{\infty} (1 - \alpha_n) < \infty$.

Then the sequence $\{x_n\}$ converges strongly to a fixed point p^* of T if and only if

$$\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0. \quad (3.21)$$

Remark 3.5. Constructing iterative algorithms for approximating (common) fixed points of some nonlinear operators has been studied extensively. It is worth mentioning that our iterative scheme (2.1) appears to be a new one, which includes many iterative schemes as special cases. Our results improve and extend the corresponding results of Lan [3], Chang et al. [4], Xu and Noor [7], Zhou et al. [30], and many others.

Acknowledgments

The authors are extremely grateful to three anonymous referees for their useful comments and suggestions which improved the manuscript. The second author was supported in part by NSC 98-2622-E-230-006-CC3 and NSC 98-2923-E-110-003-MY3.

References

- [1] K. Goebel and W. A. Kirk, "A fixed point theorem for asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 35, pp. 171–174, 1972.
- [2] Q. Liu, "Iterative sequences for asymptotically quasi-nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 259, no. 1, pp. 1–7, 2001.
- [3] H.-Y. Lan, "Common fixed-point iterative processes with errors for generalized asymptotically quasi-nonexpansive mappings," *Computers & Mathematics with Applications*, vol. 52, no. 10-11, pp. 1403–1412, 2006.
- [4] S. S. Chang, J. K. Kim, and S. M. Kang, "Approximating fixed points of asymptotically quasi-nonexpansive type mappings by the Ishikawa iterative sequences with mixed errors," *Dynamic Systems and Applications*, vol. 13, no. 2, pp. 179–186, 2004.
- [5] Y. J. Cho, H. Zhou, and G. Guo, "Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mappings," *Computers & Mathematics with Applications*, vol. 47, no. 4-5, pp. 707–717, 2004.
- [6] K.-K. Tan and H. K. Xu, "Fixed point iteration processes for asymptotically nonexpansive mappings," *Proceedings of the American Mathematical Society*, vol. 122, no. 3, pp. 733–739, 1994.
- [7] B. Xu and M. A. Noor, "Fixed-point iterations for asymptotically nonexpansive mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 267, no. 2, pp. 444–453, 2002.
- [8] Z. Huang, "Mann and Ishikawa iterations with errors for asymptotically nonexpansive mappings," *Computers & Mathematics with Applications*, vol. 37, no. 3, pp. 1–7, 1999.
- [9] L.-C. Ceng, H.-K. Xu, and J.-C. Yao, "The viscosity approximation method for asymptotically nonexpansive mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 4, pp. 1402–1412, 2008.
- [10] L.-C. Ceng, A. Petruşel, and J.-C. Yao, "Strong convergence of modified implicit iterative algorithms with perturbed mappings for continuous pseudocontractive mappings," *Applied Mathematics and Computation*, vol. 209, no. 2, pp. 162–176, 2009.
- [11] Y. Yao, Y.-C. Liou, and J.-C. Yao, "Convergence theorem for equilibrium problems and fixed point problems of infinite family of nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2007, Article ID 64363, 12 pages, 2007.
- [12] T. Honda, W. Takahashi, and J. C. Yao, "Nonexpansive retractions onto closed convex cones in Banach spaces," to appear in *Taiwanese Journal of Mathematics*.
- [13] Y.-C. Liou, Y. Yao, and R. Chen, "Iteration scheme with perturbed mapping for common fixed points of a finite family of nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2007, Article ID 29091, 10 pages, 2007.
- [14] L. C. Ceng, A. Petrusel, and J. C. Yao, "Iterative approximation of fixed points for asymptotically strict pseudocontractive type mappings in the intermediate sense," to appear in *Taiwanese Journal of Mathematics*.
- [15] X. Wu, J.-C. Yao, and L.-C. Zeng, "Uniform normal structure and strong convergence theorems for asymptotically pseudocontractive mappings," *Journal of Nonlinear and Convex Analysis*, vol. 6, no. 3, pp. 453–463, 2005.
- [16] L. C. Ceng, N. C. Wong, and J. C. Yao, "Implicit predictor-corrector iteration process for finitely many asymptotically (quasi-)nonexpansive mappings," *Journal of Inequalities and Applications*, vol. 2006, Article ID 65983, 11 pages, 2006.
- [17] J.-C. Yao and L.-C. Zeng, "Strong convergence of averaged approximants for asymptotically pseudocontractive mappings in Banach spaces," *Journal of Nonlinear and Convex Analysis*, vol. 8, no. 3, pp. 451–462, 2007.
- [18] Y. Yao, Y. C. Liou, and G. Marino, "Strong convergence of two iterative algorithms for nonexpansive mappings in Hilbert spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 279058, 7 pages, 2009.
- [19] L.-C. Ceng, S. Al-Homidan, Q. H. Ansari, and J.-C. Yao, "An iterative scheme for equilibrium problems and fixed point problems of strict pseudo-contraction mappings," *Journal of Computational and Applied Mathematics*, vol. 223, no. 2, pp. 967–974, 2009.
- [20] D. R. Sahu, H.-K. Xu, and J.-C. Yao, "Asymptotically strict pseudocontractive mappings in the intermediate sense," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 10, pp. 3502–3511, 2009.

- [21] J.-W. Peng and J.-C. Yao, "Some new iterative algorithms for generalized mixed equilibrium problems with strict pseudo-contractions and monotone mappings," *Taiwanese Journal of Mathematics*, vol. 13, no. 5, pp. 1537–1582, 2009.
- [22] Y.-C. Liou and Y. Yao, "Iterative algorithms for nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2008, Article ID 384629, 10 pages, 2008.
- [23] L. C. Ceng, D. S. Shyu, and J. C. Yao, "Relaxed composite implicit iteration process for common fixed points of a finite family of strictly pseudocontractive mappings," *Fixed Point Theory and Applications*, vol. 2009, Article ID 402602, 16 pages, 2009.
- [24] L. C. Ceng, A. Petrusel, and J. C. Yao, "A hybrid method for Lipschitz continuous monotone mappings and asymptotically strict pseudocontractive mappings in the intermediate sense," *Journal of Nonlinear and Convex Analysis*, vol. 11, no. 1, 2010.
- [25] Y. Yao, R. Chen, and H. Zhou, "Strong convergence to common fixed points of nonexpansive mappings without commutativity assumption," *Fixed Point Theory and Applications*, vol. 2006, Article ID 89470, 8 pages, 2006.
- [26] L. C. Ceng, D. R. Sahu, and J. C. Yao, "Implicit iterative algorithms for asymptotically nonexpansive mappings in the intermediate sense and Lipschitz continuous monotone mappings," *Journal of Computational and Applied Mathematics*, vol. 233, no. 11, pp. 2902–2915, 2010.
- [27] Y.-C. Liou, Y. Yao, and K. Kimura, "Strong convergence to common fixed points of a finite family of nonexpansive mappings," *Journal of Inequalities and Applications*, vol. 2007, Article ID 37513, 10 pages, 2007.
- [28] L. C. Ceng, Q. H. Ansari, and J. C. Yao, "Strong and weak convergence theorems for asymptotically strict pseudocontractive mappings in intermediate sense," to appear in *Journal of Nonlinear and Convex Analysis*.
- [29] W. Takahashi and J. C. Yao, "Weak and strong convergence theorems for positively homogenous nonexpansive mappings in Banach spaces," to appear in *Taiwanese Journal of Mathematics*.
- [30] H. Zhou, Y. J. Cho, and M. Grabiec, "Iterative processes for generalized asymptotically nonexpansive mappings in Banach spaces," *PanAmerican Mathematical Journal*, vol. 13, no. 4, pp. 99–107, 2003.
- [31] Y.-C. Lin, N.-C. Wong, and J.-C. Yao, "Strong convergence theorems of Ishikawa iteration process with errors for fixed points of Lipschitz continuous mappings in Banach spaces," *Taiwanese Journal of Mathematics*, vol. 10, no. 2, pp. 543–552, 2006.
- [32] L.-C. Zeng, N.-C. Wong, and J.-C. Yao, "Strong convergence theorems for strictly pseudocontractive mappings of Browder-Petryshyn type," *Taiwanese Journal of Mathematics*, vol. 10, no. 4, pp. 837–849, 2006.
- [33] Y.-C. Lin, "Three-step iterative convergence theorems with errors in Banach spaces," *Taiwanese Journal of Mathematics*, vol. 10, no. 1, pp. 75–86, 2006.