


KWOK KEI CHONG

Department of Applied Mathematics,  
The Hong Kong Polytechnic University,  
Hung Hom, Kowloon, Hong Kong  
The People's Republic of China.

*E*Mail: [makkchon@inet.polyu.edu.hk](mailto:makkchon@inet.polyu.edu.hk)



---

volume 2, issue 2, article 19,  
2001.

*Received 03 October, 2000;*  
*accepted 02 February, 2001.*

*Communicated by: F. Qi*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



## Abstract

It is well-known that inequalities between means play a very important role in many branches of mathematics. Please refer to [1, 3, 7], etc. The main aims of the present article are:

- (i) to show that there are monotonic and continuous functions  $H(t)$ ,  $K(t)$ ,  $P(t)$  and  $Q(t)$  on  $[0, 1]$  such that for all  $t \in [0, 1]$ ,

$$H_n \leq H(t) \leq G_n \leq K(t) \leq A_n \quad \text{and}$$

$$H_n/(1 - H_n) \leq P(t) \leq G_n/G'_n \leq Q(t) \leq A_n/A'_n,$$


where  $A_n, G_n$  and  $H_n$  are respectively the weighted arithmetic, geometric and harmonic means of the positive numbers  $x_1, x_2, \dots, x_n$  in  $(0, 1/2]$ , with positive weights  $\alpha_1, \alpha_2, \dots, \alpha_n$ ; while  $A'_n$  and  $G'_n$  are respectively the weighted arithmetic and geometric means of the numbers  $1 - x_1, 1 - x_2, \dots, 1 - x_n$  with the same positive weights  $\alpha_1, \alpha_2, \dots, \alpha_n$ ;

- (ii) to present more general monotonic refinements for the Ky Fan inequality as well as some inequalities involving means; and  
(iii) to present some generalized and new inequalities in this connection.

**2000 Mathematics Subject Classification:** 26D15, 26A48.

**Key words:** Ky Fan inequality, monotonic refinements of inequalities, arithmetic, geometric and harmonic means.

The author would like to thank the referees for their invaluable comments and suggestions.



---

### Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

---

Title Page

Contents



Go Back


Close

Quit

Page 2 of 24

# Contents

1	Introduction .....	4
2	Some Generalizations .....	8
3	Monotonic Refinements of the Ky Fan Inequality .....	15
4	Second Proof of Theorem 2.1 .....	21
References		



---

## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 24

# 1. Introduction

Let  $n$  be a positive integer. To two given sequences of positive numbers  $x_1, x_2, \dots, x_n$  and  $\alpha_1, \alpha_2, \dots, \alpha_n$  such that  $\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$ , we denote by  $A_n, G_n$  and  $H_n$  respectively the weighted arithmetic, geometric and harmonic means, that is,

$$A_n = \sum_{i=1}^n \alpha_i x_i,$$

$$G_n = \prod_{i=1}^n x_i^{\alpha_i},$$


$$H_n = \left( \sum_{i=1}^n \alpha_i / x_i \right)^{-1}.$$

We use the symbols  $\mathbf{a}_n, \mathbf{g}_n$  and  $\mathbf{h}_n$  to denote the corresponding unweighted arithmetic, geometric and harmonic means of the  $n$  positive numbers  $x_1, x_2, \dots, x_n$ . The following well-known inequality has been proved, using many different methods: (Please refer to [3].)

$$(1.1) \quad H_n \leq G_n \leq A_n$$

Let the real numbers  $x_i$  be such that  $0 < x_i \leq 1/2$ , for all  $i = 1, 2, \dots, n$ . We denote by  $A'_n, G'_n$  and  $H'_n$  the weighted arithmetic, geometric and harmonic means of the numbers  $1 - x_1, 1 - x_2, \dots, 1 - x_n$ , namely,

$$A'_n = \sum_{i=1}^n \alpha_i (1 - x_i),$$



## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents




Go Back

Close

Quit

Page 4 of 24



**Monotonic Refinements of a Ky Fan Inequality**

K. K. Chong

Title Page

Contents



Go Back

Close

Quit

Page 5 of 24

$$G'_n = \prod_{i=1}^n (1 - x_i)^{\alpha_i},$$

$$H'_n = \left( \sum_{i=1}^n \alpha_i / (1 - x_i) \right)^{-1}.$$

Also, let  $\mathbf{a}'_n$ ,  $\mathbf{g}'_n$  and  $\mathbf{h}'_n$  denote the corresponding unweighted arithmetic, geometric and harmonic means of the numbers  $1 - x_1, 1 - x_2, \dots, 1 - x_n$  respectively. In recent years many interesting inequalities involving these mean values have been published, in particular, the following well-known Ky Fan and Wang-Wang inequalities :

$$(1.2) \quad \frac{H_n}{H'_n} \leq \frac{G_n}{G'_n} \leq \frac{A_n}{A'_n}$$

with equality holding if and only if  $x_1 = \dots = x_n$ . Please refer to the following papers by H. Alzer [1] – [2] and Wang-Wang [10] or [7], etc. The right-hand inequality of (1.2) is the famous Ky Fan inequality; the left-hand inequality for the unweighted case was first discovered by Wang-Wang in 1984 [10]. The main purpose of this paper is to present some new monotonic continuous functions  $H(\lambda)$ ,  $K(\lambda)$ ,  $P(\lambda)$  and  $Q(\lambda)$  on  $[0, 1]$  such that

$$H_n \leq H(\lambda) \leq G_n \leq K(\lambda) \leq A_n$$

and

$$H_n / (1 - H_n) \leq P(\lambda) \leq G_n / G'_n \leq Q(\lambda) \leq A_n / A'_n.$$

In fact, our theorems here generalize results of Wang and Yang in [9] and a theorem of K.M. Chong [6]. In Section 2, we shall generalize refinements

of inequalities between means. In Section 3, we shall present generalizations to refinements of the Ky Fan inequality, with some new inequalities deduced. Finally, in Section 4, we shall show that Theorem 3.1 can be used to deduce many other established refinements of the Ky Fan inequality.

In a recent paper of Wang and Yang [9], the following two interesting theorems were put forward. In fact, they are refinements of inequalities (1.1) and (1.2) in Section 1, for the discrete unweighted case. They are restated here without proof. For the details of the proof, please refer to [9].

**Theorem 1.1.** *Given a sequence  $\{x_1, x_2, \dots, x_n\}$  of positive numbers, which are not all equal:*

(a) *For any  $t$  in  $[0, 1/n]$ , let*


$$(1.3) \quad h(t) = \prod_{i=1}^n \left[ \frac{1}{x_i} + t \sum_{j=1}^n \left( \frac{1}{x_j} - \frac{1}{x_i} \right) \right]^{-1/n}$$

*Then,  $h(t)$  is continuous, strictly decreasing and  $\mathbf{h}_n = h(1/n) \leq h(t) \leq h(0) = \mathbf{g}_n$  on  $[0, 1/n]$ .*

(b) *For any  $t$  in  $[0, 1/n]$ , let*

$$(1.4) \quad k(t) = \prod_{i=1}^n \left[ x_i + t \sum_{j=1}^n (x_j - x_i) \right]^{1/n}$$

*Then,  $k(t)$  is continuous, strictly increasing and  $\mathbf{g}_n = k(0) \leq k(t) \leq k(1/n) = \mathbf{a}_n$  on  $[0, 1/n]$ .*



## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents



Go Back

Close

Quit

Page 6 of 24

**Theorem 1.2.** Given a sequence  $\{x_1, x_2, \dots, x_n\}$  with  $x_i$  in  $(0, 1/2]$ ,  $i = 1, 2, \dots, n$ , which are not all equal:

(a) For any  $t$  in  $[0, 1/n]$ , let


$$(1.5) \quad p(t) = \prod_{i=1}^n \left[ \frac{1}{x_i} + t \sum_{j=1}^n \left( \frac{1}{x_j} - \frac{1}{x_i} \right) - 1 \right]^{-1/n}$$

Then,  $p(t)$  is continuous, strictly decreasing, and  $\mathbf{h}_n/(1 - \mathbf{h}_n) = p(1/n) \leq p(t) \leq p(0) = \mathbf{g}_n/\mathbf{g}'_n$  on  $[0, 1/n]$ .

(b) For any  $t$  in  $[0, 1/n]$ , let

$$(1.6) \quad q(t) = \frac{\prod_{i=1}^n \left[ x_i + t \sum_{j=1}^n (x_j - x_i) \right]^{1/n}}{\prod_{i=1}^n \left[ 1 - x_i - t \sum_{j=1}^n (x_j - x_i) \right]^{1/n}}$$

Then,  $q(t)$  is continuous, strictly increasing and  $\mathbf{g}_n/\mathbf{g}'_n = q(0) \leq q(t) \leq q(1/n) = \mathbf{a}_n/\mathbf{a}'_n$  on  $[0, 1/n]$ .



### Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents



Go Back

Close

Quit

Page 7 of 24

## 2. Some Generalizations

In this section, we are going to present and prove a generalization of Theorem 1.1 and Theorem 1.2(a), in particular, to the case for weighted means. Its statement runs as follows:

**Theorem 2.1.** *Let  $a_1, a_2, \dots, a_n$ , and  $\alpha_1, \alpha_2, \dots, \alpha_n$  be two sequences of positive numbers, with  $a_i$  not all equal and  $\sum_{i=1}^n \alpha_i = 1$ . Let  $a$  be any positive number such that  $A_n \leq a$ , where  $A_n = \sum_{i=1}^n \alpha_i a_i$ , and  $k$  is a constant such that  $k < a_i$ , for all  $i = 1, 2, \dots, n$ . Let*

$$(2.1) \quad F(\lambda) = \prod_{i=1}^n [\lambda a + (1 - \lambda) a_i - k]^{\alpha_i}$$


$$(2.2) \quad G(\lambda) = \prod_{i=1}^n [\lambda a + (1 - \lambda) a_i - k]^{-\alpha_i}$$

Then,

- (i)  $F(\lambda)$  is continuous and strictly increasing on  $[0, 1]$ ;
- (ii)  $G(\lambda)$  is continuous and strictly decreasing on  $[0, 1]$ .

*Proof.* (i) Taking the logarithm of  $F(\lambda)$ , we have,

$$\ln F(\lambda) = \sum_{i=1}^n \alpha_i \ln [\lambda a + (1 - \lambda) a_i - k]$$



---

### Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

---

Title Page

Contents




Go Back

Close

Quit

Page 8 of 24





Title Page

Contents



Go Back

Close

Quit

Page 9 of 24

Differentiating the last expression with respect to  $\lambda$ , we have:

$$\frac{F'(\lambda)}{F(\lambda)} = \sum_{i=1}^n \frac{\alpha_i (a - a_i)}{[\lambda a + (1 - \lambda) a_i - k]}$$

Differentiating again, we obtain:

$$(2.3) \quad \left[ \frac{F'(\lambda)}{F(\lambda)} \right]' = - \sum_{i=1}^n \frac{\alpha_i (a - a_i)^2}{[\lambda a + (1 - \lambda) a_i - k]^2} < 0$$

for all  $\lambda$  in  $[0, 1]$ , as the  $a_i$  are not all equal. Hence,  $F'(\lambda)/F(\lambda)$  is strictly decreasing on  $[0, 1]$ . Also, as  $A_n \leq a$  and  $k < a$ , we have :

$$(2.4) \quad \frac{F'(1)}{F(1)} = \sum_{i=1}^n \frac{\alpha_i (a - a_i)}{a - k} = \frac{a - A_n}{a - k} \geq 0$$

Therefore,  $F'(\lambda)/F(\lambda) > 0$ , for all  $\lambda$  in  $[0, 1)$ . As  $F(\lambda)$  is positive for all  $\lambda$  in  $[0, 1]$ ,  $F'(\lambda) > 0$  for  $\lambda$  in  $[0, 1)$ . Hence,  $F(\lambda)$  is strictly increasing on  $[0, 1]$ . The continuity of  $F(\lambda)$  on  $[0, 1]$  is obvious.

- (ii) As  $F(\lambda)$  is positive for all  $\lambda$  in  $[0, 1]$  and  $G(\lambda) = 1/F(\lambda)$ ,  $G(\lambda)$  is continuous and strictly decreasing on  $[0, 1]$ . Hence, the proof of Theorem 2.1 is complete.

□

Now, we use Theorem 2.1 to deduce some established theorems.

**Remark 2.1.** (i) From Theorem 2.1, we have, for all  $\lambda \in (0, 1)$ ,

$$F(0) < F(\lambda) < F(1),$$

which yields for not all equal  $a_i$ ,

$$(2.5) \quad \prod_{i=1}^n (a_i - k)^{\alpha_i} < a - k.$$

In particular, if  $a = A_n$ , for not all equal  $a_i$  we have,

$$(2.6) \quad \prod_{i=1}^n (a_i - k)^{\alpha_i} < A_n - k,$$

which is a generalization of the weighted arithmetic-geometric means inequality.


(ii) Again, in Theorem 2.1, we let  $k = 0, a = \sum_{i=1}^n \alpha_i a_i = A_n$ . Then,  $F(\lambda)$  will reduce to, say

$$(2.7) \quad K(\lambda) = \prod_{i=1}^n [\lambda A_n + (1 - \lambda) a_i]^{\alpha_i}$$

It is clear that  $K(\lambda)$  is continuous and strictly increasing on  $[0, 1]$ , and for all  $\lambda \in (0, 1)$ ,

$$(2.8) \quad K(0) = G_n < K(\lambda) < K(1) = A_n.$$

This is a refinement of the weighted arithmetic-geometric means inequality.



## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents



Go Back

Close

Quit

Page 10 of 24

(iii) Furthermore, if we put  $\lambda = nt, \alpha_i = \frac{1}{n}, i = 1, 2, \dots, n$ , into  $K(\lambda)$ , we obtain for all  $t \in [0, 1/n]$ ,

$$K(nt) = \prod_{i=1}^n [ntA_n + (1 - nt)a_i]^{1/n} = \prod_{i=1}^n \left[ a_i + t \sum_{j=1}^n (a_j - a_i) \right]^{1/n}.$$

The last expression is in fact the function  $k(t)$  of Theorem 1.1(b). Hence, we have shown that Theorem 1.1(b) is a particular case of Theorem 2.1.

**Remark 2.2.** If, in Theorem 2.1, we let  $k = 0, a_i = \frac{1}{x_i}, i = 1, 2, \dots, n, a = \frac{1}{H_n} = \frac{\alpha_1}{x_1} + \dots + \frac{\alpha_n}{x_n}$ , then  $G(\lambda)$  will reduce to,

$$(2.9) \quad H(\lambda) = \prod_{i=1}^n \left[ \frac{\lambda}{H_n} + (1 - \lambda) \frac{1}{x_i} \right]^{-\alpha_i}.$$


Then,  $H(\lambda)$  is continuous and strictly decreasing on  $[0, 1]$ , and for all  $\lambda \in (0, 1)$ ,

$$(2.10) \quad H(1) = H_n < H(\lambda) < H(0) = G_n.$$

(2.10) is a refinement of the weighted means inequality. Furthermore, if we put  $\lambda = nt, \alpha_i = \frac{1}{n}, a_i = 1/x_i, i = 1, 2, \dots, n$ , into  $H(\lambda)$ , we obtain for all  $t$  in  $[0, 1/n]$ ,

$$H(nt) = \prod_{i=1}^n \left[ nta + (1 - nt) \frac{1}{x_i} \right]^{-1/n} = \prod_{i=1}^n \left[ \frac{1}{x_i} + t \sum_{j=1}^n \left( \frac{1}{x_j} - \frac{1}{x_i} \right) \right]^{-1/n}$$

This is the function  $h(t)$  in Theorem 1.1(a). Hence, we have deduced Theorem 1.1(a) as a particular case of Theorem 2.1.



## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents




Go Back

Close

Quit

Page 11 of 24



**Theorem 2.2.** Let  $x_1, x_2, \dots, x_n$  be  $n$  positive numbers, not all equal, with  $x_i \in (0, 1/2]$  for all  $i = 1, 2, \dots, n$ . Let  $\alpha_1, \alpha_2, \dots, \alpha_n$  be the corresponding weights, i.e.,  $\alpha_i > 0$ ,  $i = 1, 2, \dots, n$  and  $\alpha_1 + \dots + \alpha_n = 1$ . Let  $\gamma$  be a constant such that  $\gamma < \frac{1}{x_i}$ , for all  $i = 1, 2, \dots, n$ . We define  $P(\lambda)$  as :

$$(2.11) \quad P(\lambda) = \prod_{i=1}^n \left[ \frac{\lambda}{H_n} + (1-\lambda) \frac{1}{x_i} - \gamma \right]^{-\alpha_i}$$

Then,

- (i)  $P(\lambda)$  is continuous and strictly decreasing on  $[0, 1]$ ;
- (ii) for all  $\lambda \in (0, 1)$ , we have,

$$(2.12) \quad P(1) = \frac{H_n}{1-\gamma H_n} < P(\lambda) < P(0) = \prod_{i=1}^n \left( \frac{x_i}{1-\gamma x_i} \right)^{\alpha_i}.$$

*Proof.* (i)  $P(\lambda)$  is continuous and strictly decreasing on  $[0, 1]$ , as we get  $P(\lambda)$  from the continuous and strictly decreasing function  $G(\lambda)$ , by putting  $k = \gamma$ ,  $a_i = 1/x_i$  with  $x_i \in (0, 1/2]$ ,  $i = 1, 2, \dots, n$ ,  $a = 1/H_n = \alpha_1/x_1 + \dots + \alpha_n/x_n$  into  $G(\lambda)$  of Theorem 2.1.

(ii) We have :  $P(0) = G(0) = \prod_{i=1}^n \left( \frac{x_i}{1-\gamma x_i} \right)^{\alpha_i}$ ,

$$P(1) = G(1) = H_n/(1-\gamma H_n).$$

Hence, for all  $\lambda \in (0, 1)$ ,

$$H_n/(1-\gamma H_n) < P(\lambda) < \prod_{i=1}^n \left( \frac{x_i}{1-\gamma x_i} \right)^{\alpha_i}$$

This completes the proof of Theorem 2.2. □

**Remark 2.3.** If we put  $\lambda = nt$ ,  $\alpha_i = 1/n$ ,  $i = 1, 2, \dots, n$ ,  $\gamma = 1$  into  $P(\lambda)$  of Theorem 2.2, we obtain for any  $t$  in  $[0, 1/n]$ ,

$$\begin{aligned} P(nt) &= \prod_{i=1}^n \left[ \frac{nt}{h_n} + (1 - nt) a_i - 1 \right]^{-1/n} \\ &= \prod_{i=1}^n \left[ \frac{1}{x_i} + t \sum_{j=1}^n \left( \frac{1}{x_j} - \frac{1}{x_i} \right) - 1 \right]^{-1/n} \end{aligned}$$


This is the function  $p(t)$  of Theorem 1.2(a), and we have deduced Theorem 1.2(a) as a particular case of Theorem 2.1.

The only part in Section 1, which is not yet dealt with, is Theorem 1.2(b). Its proof is postponed to the next section, with some additional theorems. We end this section by considering another similar theorem. In [6], K.M. Chong presented the following theorem:

**Theorem 2.3.** Let  $a_1, a_2, \dots, a_n$  be positive numbers and  $\alpha_1, \alpha_2, \dots, \alpha_n$  be their corresponding weights, i.e.  $\alpha_i > 0$ ,  $i = 1, 2, \dots, n$  and  $\sum_{i=1}^n \alpha_i = 1$ . Let  $f(\lambda)$  be defined as:

$$(2.13) \quad f(\lambda) = \prod_{i=1}^n \left[ \lambda \sum_{j=1}^n \alpha_j a_j + (1 - \lambda) a_i \right]^{\alpha_i}$$

Then  $f(\lambda)$  is a strictly increasing function of  $\lambda$  for  $\lambda \in [0, 1]$ , unless  $a_1 = a_2 = \dots = a_n$ ; in which case  $f(0) = G_n = A_n = f(1)$ .




**Monotonic Refinements of a Ky Fan Inequality**

K. K. Chong

Title Page	
Contents	
◀◀	▶▶
◀	▶
Go Back	
Close	
Quit	
Page 13 of 24	

*Proof.* It is obvious that when  $k = 0$  and  $a = A_n$  in Theorem 2.1 we obtain K.M. Chong's theorem at once.  $\square$



---

**Monotonic Refinements of a Ky Fan Inequality**

K. K. Chong

---

Title Page

Contents



Go Back

Close

Quit

Page 14 of 24

### 3. Monotonic Refinements of the Ky Fan Inequality

In the previous section, we have seen that Theorem 2.1 is a generalization of various theorems. In this section, we shall present a refinement of the well-known Ky Fan inequality, which is a generalization of Theorem 1.2(b).


**Theorem 3.1.** *Let  $x_1, x_2, \dots, x_n$  be  $n$  positive numbers, not all equal, such that  $x_i \in (0, 1/2]$  for all  $i = 1, 2, \dots, n$  and let  $\alpha_1, \alpha_2, \dots, \alpha_n$  be their corresponding weights i.e.  $\alpha_i > 0, i = 1, 2, \dots, n$  and  $\sum_{i=1}^n \alpha_i = 1$ . Let  $\beta$  and  $\delta$  be two constants such that  $\delta \leq \beta$  and  $\beta < x_i$ , for all  $i = 1, 2, \dots, n$ . Let  $r(\lambda)$  be defined as :*

$$(3.1) \quad r(\lambda) = \frac{\prod_{i=1}^n [\lambda z + (1 - \lambda)x_i - \beta]^{\alpha_i}}{\prod_{i=1}^n [\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta]^{\alpha_i}}$$

for any  $\lambda \in [0, 1]$ , and any real number  $z$  such that  $\sum_{i=1}^n \alpha_i x_i \leq z \leq 1/2$ .  
Then,

- (i)  $r(\lambda)$  is continuous and strictly increasing on  $[0, 1]$ ; and
- (ii) when  $\beta = \delta = 0$ , we have, for all  $\lambda \in (0, 1)$ ,

$$(3.2) \quad G_n/G'_n = r(0) < r(\lambda) < r(1) = \frac{z}{1 - z}.$$



#### Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents



Go Back

Close

Quit

Page 15 of 24

*Proof.* (i) Taking the logarithm, we have,


$$\ln\{r(\lambda)\} = \sum_{i=1}^n \alpha_i \ln[\lambda z + (1 - \lambda)x_i - \beta] - \sum_{i=1}^n \alpha_i \ln[\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta]$$

Differentiating with respect to  $\lambda$ , we have :

$$\begin{aligned} \frac{r'(\lambda)}{r(\lambda)} &= \sum_{i=1}^n \alpha_i \frac{z - x_i}{\lambda z + (1 - \lambda)x_i - \beta} - \sum_{i=1}^n \alpha_i \frac{(1 - z) - (1 - x_i)}{\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta} \\ &= \sum_{i=1}^n \alpha_i \frac{z - x_i}{\lambda z + (1 - \lambda)x_i - \beta} + \sum_{i=1}^n \alpha_i \frac{z - x_i}{\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta}. \end{aligned}$$

Let  $u(\lambda) = \frac{r'(\lambda)}{r(\lambda)}$ .

We are going to show that  $\ln\{r(\lambda)\}$  and hence  $r(\lambda)$  are both strictly increasing by showing that  $u(\lambda) > 0$  for all  $\lambda \in [0, 1)$ .



## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents




Go Back

Close

Quit

Page 16 of 24





Title Page

Contents



Go Back

Close

Quit

Page 17 of 24

Differentiating  $u(\lambda)$  with respect to  $\lambda$ , we have :

$$u'(\lambda) = - \sum_{i=1}^n \frac{\alpha_i(z - x_i)^2}{[\lambda z + (1 - \lambda)x_i - \beta]^2} + \sum_{i=1}^n \frac{\alpha_i(z - x_i)^2}{[\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta]^2} < 0$$

as

$$\frac{1}{[\lambda z + (1 - \lambda)x_i - \beta]^2} > \frac{1}{[\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta]^2}$$

$i = 1, 2, \dots, n$ , unless  $z = x_1 = x_2 = \dots = x_n = 1/2$ , and  $\beta = \delta$ .

Hence  $u(\lambda)$  is strictly decreasing on  $[0, 1]$ .

$$\begin{aligned} u(1) &= \sum_{i=1}^n \alpha_i \frac{z - x_i}{z - \beta} + \sum_{i=1}^n \alpha_i \frac{z - x_i}{1 - z - \delta} \\ &= \frac{1}{z - \beta} \sum_{i=1}^n \alpha_i(z - x_i) + \frac{1}{1 - z - \delta} \sum_{i=1}^n \alpha_i(z - x_i) \\ &= \left( z - \sum_{i=1}^n \alpha_i x_i \right) \frac{1 - \beta - \delta}{(z - \beta)(1 - z - \delta)} \geq 0, \text{ for } \sum_{i=1}^n \alpha_i x_i \leq z \leq 1/2. \end{aligned}$$

Hence,  $u(\lambda) = \frac{r'(\lambda)}{r(\lambda)} > 0$ , for all  $\lambda \in [0, 1)$ .

As  $r(\lambda)$  is always positive, we have  $r'(\lambda) > 0$  for all  $\lambda \in [0, 1)$  and  $r(\lambda)$  is strictly increasing on  $[0, 1]$ .

(ii) It is easy to see that when  $\beta = \delta = 0$ ,  $r(0) = G_n/G'_n$ ,  $r(1) = \frac{z}{1-z}$  and  $r(0) < r(\lambda) < r(1)$  on  $(0, 1)$ . □

It is remarked, that if  $\beta = \delta = 0$ ,  $z = \sum_{i=1}^n \alpha_i x_i$  in Theorem 3.1, then the chain of inequalities in (3.2), with  $r(\lambda)$  replaced by  $Q(\lambda)$ , will become : for any  $\lambda \in (0, 1)$ ,

$$(3.3) \quad \frac{G_n}{G'_n} = Q(0) < Q(\lambda) < Q(1) = \frac{z}{1-z} = \frac{A_n}{A'_n}$$

This is a refinement of the Ky Fan inequality.


**Remark 3.1.** (3.3) is a refinement of the weighted Ky Fan inequality and we have  $r(0) < r(\lambda) < r(1)$ , unless  $x_1 = x_2 = \dots = x_n$ . In general, (3.2) yields a generalization of the Ky Fan inequality as follows :

For  $A_n \leq z \leq 1/2$  and  $\delta \leq \beta < x_i \in (0, 1/2]$ , for all  $i = 1, 2, \dots, n$ , we have,

$$(3.4) \quad r(0) = \frac{\prod_{i=1}^n [x_i - \beta]^{\alpha_i}}{\prod_{i=1}^n [1 - x_i - \delta]^{\alpha_i}} \leq \frac{z - \beta}{1 - z - \delta} = r(1),$$

with equality if and only if  $x_1 = x_2 = \dots = x_n$ .

If in (3.4) we let  $z = A_n$  the weighted arithmetic mean of  $x_1, x_2, \dots, x_n$ , we




---

### Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 24

obtain a generalization of the weighted Ky Fan inequality :

$$(3.5) \quad \frac{\prod_{i=1}^n [x_i - \beta]^{\alpha_i}}{\prod_{i=1}^n [1 - x_i - \delta]^{\alpha_i}} \leq \frac{A_n - \beta}{A'_n - \delta},$$


with equality if and only if  $x_1 = x_2 = \dots = x_n$ .

**Remark 3.2.** If we put  $\alpha_i = 1/n, i = 1, 2, \dots, n, z = \sum_{i=1}^n \alpha_i x_i, \beta = \delta = 0$  and  $\lambda = nt$  into (3.1), we then obtain after simplification,

$$\begin{aligned} r(nt) &= \frac{\prod_{i=1}^n \left[ nt \sum_{j=1}^n \frac{1}{n} x_j + (1 - nt)x_i \right]^{1/n}}{\prod_{i=1}^n \left[ nt \left( 1 - \sum_{j=1}^n \frac{1}{n} x_j \right) + (1 - nt)(1 - x_i) \right]^{1/n}} \\ &= \frac{\prod_{i=1}^n \left[ x_i + t \sum_{j=1}^n (x_j - x_i) \right]^{1/n}}{\prod_{i=1}^n \left[ 1 - x_i - t \sum_{j=1}^n (x_j - x_i) \right]^{1/n}} = q(t), \quad \text{for } t \in [0, 1/n]. \end{aligned}$$

This is the function  $q(t)$  in Theorem 1.2(b), showing that Theorem 3.1 is a generalization of Theorem 1.2(b).

**Remark 3.3.** In [5], we have the following theorem, which can be easily seen to follow as a particular case of Theorem 3.1, when  $\beta = 0$  and  $\delta = 0$  :



## Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents




Go Back

Close

Quit

Page 19 of 24




---

**Monotonic Refinements of a Ky Fan Inequality**

K. K. Chong

---

Title Page

Contents



Go Back

Close

Quit

Page 20 of 24

---

**Theorem 3.2.** Let  $x_1, x_2, \dots, x_n$  be  $n$  positive numbers, such that  $x_i \in (0, 1/2]$ , for all  $i = 1, 2, \dots, n$ , and let  $\alpha_1, \alpha_2, \dots, \alpha_n$  be their corresponding positive weights, with  $\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$ . Let  $z$  be a constant such that  $A_n \leq z \leq 1/2$ , where  $A_n = \sum_{i=1}^n \alpha_i x_i$ . We define  $w(\lambda)$ , for any  $\lambda \in [0, 1]$ , to be the function:

$$(3.6) \quad w(\lambda) = \frac{\prod_{i=1}^n [\lambda z + (1 - \lambda)x_i]^{\alpha_i}}{\prod_{i=1}^n [\lambda(1 - z) + (1 - \lambda)(1 - x_i)]^{\alpha_i}}.$$

Then,

- (i)  $w(\lambda)$  is continuous and strictly increasing on  $[0, 1]$ , unless  $x_1 = x_2 = \dots = x_n$ ;
- (ii)  $G_n/G_n' = w(0) \leq w(\lambda) \leq w(1) = \frac{z}{1-z}$ , for  $\lambda \in [0, 1]$ .

## 4. Second Proof of Theorem 2.1

In this section, we shall show that Theorem 3.1 is not only a generalization of Theorem 1.2(b), but also it can be used to deduce some elementary theorems.

*Second proof of Theorem 2.1.* Suppose the  $n$  numbers  $x_1, x_2, \dots, x_n$  are not all equal.

For  $x_i \in (0, 1/2]$ ,  $i = 1, 2, \dots, n$ ,  $z$  lying between  $\sum_{i=1}^n \alpha_i x_i$  and  $1/2$ , the function  $r(\lambda)$  of Theorem 3.1 is strictly increasing on  $[0, 1]$ , where for  $\lambda \in [0, 1]$ ,  $r(\lambda)$  is defined as :


$$(4.1) \quad r(\lambda) = \frac{\prod_{i=1}^n [\lambda z + (1 - \lambda)x_i - \beta]^{\alpha_i}}{\prod_{i=1}^n [\lambda(1 - z) + (1 - \lambda)(1 - x_i) - \delta]^{\alpha_i}}.$$

Now, we put  $x_i = \frac{a_i}{l}$ ,  $i = 1, 2, \dots, n$ , where  $l$  is a large positive number, and let  $z = \frac{a}{l}$ ,  $\beta = \frac{k}{l}$  with  $\delta = \beta$ . Then, the function  $r(\lambda)$  becomes :

$$(4.2) \quad r(\lambda) = \frac{\frac{1}{l} \prod_{i=1}^n [\lambda a + (1 - \lambda)a_i - k]^{\alpha_i}}{\prod_{i=1}^n \left[ \lambda \left(1 - \frac{a}{l}\right) + (1 - \lambda) \left(1 - \frac{a_i}{l}\right) - \beta \right]^{\alpha_i}}.$$

By Theorem 3.1,

$$(4.3) \quad v(\lambda) = \frac{\prod_{i=1}^n [\lambda a + (1 - \lambda)a_i - k]^{\alpha_i}}{\prod_{i=1}^n \left[ \lambda \left(1 - \frac{a}{l}\right) + (1 - \lambda) \left(1 - \frac{a_i}{l}\right) - \frac{k}{l} \right]^{\alpha_i}}$$



Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

Title Page

Contents



Go Back

Close

Quit

Page 21 of 24

is strictly increasing as  $\lambda$  increases from 0 to 1.


We let  $l$  tend to  $+\infty$ , the denominator tends to 1 and we have shown that the function in Theorem 2.1

$$(4.4) \quad F(\lambda) = \prod_{i=1}^n [\lambda a + (1 - \lambda)a_i - k]^{\alpha_i}$$

is an increasing function on  $[0, 1]$ .

Differentiation calculations as in Theorem 2.1 easily reveal that in fact  $F(\lambda)$  is strictly increasing on  $[0, 1]$ . This completes the proof of Theorem 2.1.  $\square$

**Remark 4.1.** *From the discussions in Section 2, it can be seen that Theorem 2.1 generalizes Theorem 1.1, Theorem 1.2(a), Theorem 2.2 and Theorem 2.3. From the discussions of the last two sections, it can be seen that Theorem 3.1 generalizes Theorem 1.2(b), Theorem 3.2 and Theorem 2.1. As a whole, we have shown that Theorem 3.1 is a generalization of all other refinements of inequalities (1.1) and (1.2), appearing in this paper.*




---

**Monotonic Refinements of a Ky Fan Inequality**

K. K. Chong

---

Title Page

Contents



Go Back


Close

Quit

Page 22 of 24

## References

- [1] H. ALZER, Inequalities for arithmetic, geometric and harmonic means, *Bull. London Math. Soc.*, **22** (1990), 362–366.
- [2] H. ALZER, The inequality of Ky Fan and related results, *Acta Appl. Math.*, **38** (1995), 305–354.
- [3] P.S. BULLEN, D.S. MITRINOVIĆ AND J.E. PEČARIĆ, *Means and Their Inequalities*, Reiddel Dordrecht, 1988.
- [4] K.K. CHONG, On a Ky Fan's inequality and some related inequalities between means, *Southeast Asian Bull. Math.*, **22** (1998), 363–372.
- [5] K.K. CHONG, On some generalizations and refinements of a Ky Fan inequality, *Southeast Asian Bull. Math.*, **24** (2000), 355–364.
- [6] K.M. CHONG, On the arithmetic-mean-geometric-mean inequality, *Nanta Mathematica*, **10**(1) (1977), 26–27.
- [7] A.M. FINK, J.E. PEČARIĆ AND D.S. MITRINOVIĆ, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993.
- [8] N. LEVINSON, Generalization of an inequality of Ky Fan, *J. Math. Anal. Appl.*, **3** (1964), 133–134.
- [9] C.S. WANG AND G.S. YANG, Refinements on an inequality of Ky Fan, *J. Math. Anal. Appl.*, **201** (1996), 955–965.
- [10] W.L. WANG AND P.F. WANG, A class of inequalities for the symmetric functions, *Acta Math. Sinica*, **27** (1984), 485–497 [In Chinese].



---

### Monotonic Refinements of a Ky Fan Inequality

K. K. Chong

---

Title Page

Contents




Go Back

Close

Quit

Page 23 of 24

- [11] W.L. WANG, Some inequalities involving means and their converses, *J. Math. Anal. Appl.*, **238** (1999), 567–579.



---

**Monotonic Refinements of a Ky Fan Inequality**

K. K. Chong

---

Title Page

Contents



Go Back

Close

Quit

Page 24 of 24