

NOTES ON CERTAIN SUBCLASS OF p -VALENTLY BAZILEVIČ FUNCTIONS



p-Valently Bazilevič Functions
Zhi-Gang Wang and Yue-Ping Jiang
vol. 9, iss. 3, art. 70, 2008

ZHI-GANG WANG

School of Mathematics and Computing Science
Changsha University of Science and Technology
Changsha 410076, Hunan,
People's Republic of China
EMail: zhigangwang@foxmail.com

YUE-PING JIANG

School of Mathematics and Econometrics
Hunan University
Changsha 410082, Hunan,
People's Republic of China
EMail: ypjiang731@163.com

Received: 04 February, 2007

Accepted: 07 July, 2008

Communicated by: A. Sofo

2000 AMS Sub. Class.: Primary 30C45.



Key words: Analytic functions, Multivalent functions, Bazilevič functions, subordination between analytic functions, Briot-Bouquet differential subordination.

Abstract: In the present paper, we discuss a subclass $\mathcal{M}_p(\lambda, \mu, A, B)$ of p -valently Bazilevič functions, which was introduced and investigated recently by Patel [5]. Such results as inclusion relationship, coefficient inequality and radius of convexity for this class are proved. The results presented here generalize and improve some earlier results. Several other new results are also obtained.

Acknowledgements: The present investigation was supported by the *National Natural Science Foundation* under Grant 10671059 of People's Republic of China. The first-named author would like to thank Professors Chun-Yi Gao and Ming-Sheng Liu for their continuous support and encouragement. The authors would also like to thank the referee for his careful reading and making some valuable comments which have essentially improved the presentation of this paper.

Title Page

Contents



Page 1 of 15

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Preliminary Results	6
3	Main Results	8



p-Valently Bazilevič Functions

Zhi-Gang Wang and Yue-Ping Jiang

vol. 9, iss. 3, art. 70, 2008

Title Page

Contents



Page 2 of 15

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

1. Introduction

Let \mathcal{A}_p denote the class of functions of the form:

$$f(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n \quad (p \in \mathbb{N} := \{1, 2, 3, \dots\}),$$

which are *analytic* in the *open* unit disk

$$\mathbb{U} := \{z : z \in \mathbb{C} \text{ and } |z| < 1\}.$$

For simplicity, we write

$$\mathcal{A}_1 =: \mathcal{A}.$$

For two functions f and g , analytic in \mathbb{U} , we say that the function f is subordinate to g in \mathbb{U} , and write

$$f(z) \prec g(z) \quad (z \in \mathbb{U}),$$

if there exists a Schwarz function ω , which is analytic in \mathbb{U} with

$$\omega(0) = 0 \quad \text{and} \quad |\omega(z)| < 1 \quad (z \in \mathbb{U})$$

such that

$$f(z) = g(\omega(z)) \quad (z \in \mathbb{U}).$$

Indeed it is known that

$$f(z) \prec g(z) \quad (z \in \mathbb{U}) \implies f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subset g(\mathbb{U}).$$

Furthermore, if the function g is univalent in \mathbb{U} , then we have the following equivalence:

$$f(z) \prec g(z) \quad (z \in \mathbb{U}) \iff f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subset g(\mathbb{U}).$$



Title Page

Contents



Page 3 of 15

Go Back

Full Screen

Close



[Title Page](#)

[Contents](#)

◀▶

◀ ▶

Page 4 of 15

[Go Back](#)

[Full Screen](#)

[Close](#)

Let $\mathcal{M}_p(\lambda, \mu, A, B)$ denote the class of functions in \mathcal{A}_p satisfying the following subordination condition:

$$(1.1) \quad \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \lambda \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right] \prec p \frac{1+Az}{1+Bz}$$

$$(-1 \leq B < A \leq 1; z \in \mathbb{U})$$

for some real μ ($\mu \geq 0$), λ ($\lambda \geq 0$) and $g \in \mathcal{S}_p^*$, where \mathcal{S}_p^* denotes the usual class of *p*-valently starlike functions in \mathbb{U} .

For simplicity, we write

$$\mathcal{M}_p \left(\lambda, \mu, 1 - \frac{2\alpha}{p}, -1 \right) = \mathcal{M}_p(\lambda, \mu, \alpha)$$

$$:= \left\{ f(z) \in \mathcal{A}_p : \Re \left(\frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \lambda \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right] \right) > \alpha \right\},$$

for some α ($0 \leq \alpha < p$) and $z \in \mathbb{U}$.

The class $\mathcal{M}_p(\lambda, \mu, A, B)$ was introduced and investigated recently by Patel [5]. The author obtained some interesting properties for this class in the case $\lambda > 0$, he also proved the following result:

Theorem 1.1. *Let*

$$\mu \geq 0, \lambda > 0 \quad \text{and} \quad -1 \leq B < A \leq 1.$$

If $f \in \mathcal{M}_p(\lambda, \mu, A, B)$, then

$$(1.2) \quad \frac{zf'(z)}{pf^{1-\mu}(z)g^\mu(z)} \prec \frac{\lambda}{pQ(z)} = q(z) \prec \frac{1+Az}{1+Bz} \quad (z \in \mathbb{U}),$$

where

$$Q(z) = \begin{cases} \int_0^1 s^{\frac{p}{\lambda}-1} \left(\frac{1+Bsz}{1+Bz} \right)^{\frac{p(A-B)}{\lambda B}} ds & (B \neq 0), \\ \int_0^1 s^{\frac{p}{\lambda}-1} \exp\left(\frac{p}{\lambda}(s-1)Az\right) ds & (B = 0), \end{cases}$$

and $q(z)$ is the best dominant of (1.2).

In the present paper, we shall derive such results as inclusion relationship, coefficient inequality and radius of convexity for the class $\mathcal{M}_p(\lambda, \mu, A, B)$ by making use of the techniques of Briot-Bouquet differential subordination. The results presented here generalize and improve some known results. Several other new results are also obtained.



p-Valently Bazilevič Functions

Zhi-Gang Wang and Yue-Ping Jiang

vol. 9, iss. 3, art. 70, 2008

Title Page

Contents



Page 5 of 15

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 6 of 15

[Go Back](#)

[Full Screen](#)

[Close](#)

2. Preliminary Results

In order to prove our main results, we shall require the following lemmas.

Lemma 2.1. *Let*

$$\mu \geq 0, \lambda \geq 0 \quad \text{and} \quad -1 \leq B < A \leq 1.$$

Then

$$\mathcal{M}_p(\lambda, \mu, A, B) \subset \mathcal{M}_p(0, \mu, A, B).$$

Proof. Suppose that $f \in \mathcal{M}_p(\lambda, \mu, A, B)$. By virtue of (1.2), we know that

$$\frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} \prec_p \frac{1+Az}{1+Bz} \quad (z \in \mathbb{U}),$$

which implies that $f \in \mathcal{M}_p(0, \mu, A, B)$. Therefore, the assertion of Lemma 2.1 holds true. \square

Lemma 2.2 (see [3]). *Let*

$$-1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1.$$

Then

$$\frac{1+A_2z}{1+B_2z} \prec \frac{1+A_1z}{1+B_1z}.$$

Lemma 2.3 (see [4]). *Let F be analytic and convex in \mathbb{U} . If*

$$f, g \in \mathcal{A} \quad \text{and} \quad f, g \prec F,$$

then

$$\lambda f + (1-\lambda)g \prec F \quad (0 \leq \lambda \leq 1).$$

Lemma 2.4 (see [6]). *Let*

$$f(z) = \sum_{k=0}^{\infty} a_k z^k$$

be analytic in \mathbb{U} and

$$g(z) = \sum_{k=0}^{\infty} b_k z^k$$

be analytic and convex in \mathbb{U} . If $f \prec g$, then

$$|a_k| \leq |b_1| \quad (k \in \mathbb{N}).$$



***p*-Valently Bazilevič Functions**

Zhi-Gang Wang and Yue-Ping Jiang

vol. 9, iss. 3, art. 70, 2008

Title Page

Contents

◀▶

◀▶

Page 7 of 15

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



3. Main Results

We begin by stating our first inclusion relationship given by Theorem 3.1 below.

Theorem 3.1. *Let*

$$\mu \geq 0, \lambda_2 \geq \lambda_1 \geq 0 \quad \text{and} \quad -1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1.$$

Then

$$\mathcal{M}_p(\lambda_2, \mu, A_2, B_2) \subset \mathcal{M}_p(\lambda_1, \mu, A_1, B_1).$$

Proof. Suppose that $f \in \mathcal{M}_p(\lambda_2, \mu, A_2, B_2)$. We know that

$$\frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \lambda_2 \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right] \prec p \frac{1+A_2z}{1+B_2z}.$$

Since

$$-1 \leq B_1 \leq B_2 < A_2 \leq A_1 \leq 1,$$

it follows from Lemma 2.2 that

$$(3.1) \quad \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \lambda_2 \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right] \prec p \frac{1+A_1z}{1+B_1z}.$$

that is, that $f \in \mathcal{M}_p(\lambda_2, \mu, A_1, B_1)$. Thus, the assertion of Theorem 3.1 holds true for

$$\lambda_2 = \lambda_1 \geq 0.$$

If

$$\lambda_2 > \lambda_1 \geq 0,$$

by virtue of Lemma 2.1 and (3.1), we know that $f \in \mathcal{M}_p(0, \mu, A_1, B_1)$, that is

$$(3.2) \quad \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} \prec p \frac{1+A_1z}{1+B_1z}.$$

[Title Page](#)

[Contents](#)



Page 8 of 15

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)



Page 9 of 15

[Go Back](#)

[Full Screen](#)

[Close](#)

At the same time, we have

$$(3.3) \quad \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \lambda_1 \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right]$$

$$= \left(1 - \frac{\lambda_1}{\lambda_2} \right) \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \frac{\lambda_1}{\lambda_2} \left\{ \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} \right.$$

$$\left. + \lambda_2 \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right] \right\}.$$

It is obvious that

$$h_1(z) = \frac{1 + A_1z}{1 + B_1z}$$

is analytic and convex in \mathbb{U} . Thus, we find from Lemma 2.3, (3.1), (3.2) and (3.3) that

$$\frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} + \lambda_1 \left[1 + \frac{zf''(z)}{f'(z)} - (1-\mu)\frac{zf'(z)}{f(z)} - \mu\frac{zg'(z)}{g(z)} \right] \prec p \frac{1 + A_1z}{1 + B_1z},$$

that is, that $f \in \mathcal{M}_p(\lambda_1, \mu, A_1, B_1)$. This implies that

$$\mathcal{M}_p(\lambda_2, \mu, A_2, B_2) \subset \mathcal{M}_p(\lambda_1, \mu, A_1, B_1).$$

□

Remark 1. Setting $A_1 = A_2 = A$ and $B_1 = B_2 = B$ in Theorem 3.1, we get the corresponding result obtained by Guo and Liu [2].

Corollary 3.2. *Let*

$$\mu \geq 0, \lambda_2 \geq \lambda_1 \geq 0 \quad \text{and} \quad p > \alpha_2 \geq \alpha_1 \geq 0.$$

Then

$$\mathcal{M}_p(\lambda_2, \mu, \alpha_2) \subset \mathcal{M}_p(\lambda_1, \mu, \alpha_1).$$



Title Page

Contents



Page 10 of 15

Go Back

Full Screen

Close

Theorem 3.3. If $f \in \mathcal{A}_p$ satisfies the following conditions:

$$\Re \left(\frac{f(z)}{z^p} \right) > 0 \quad \text{and} \quad \left| \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} - p \right| < \nu p \quad \left(0 \leq \mu < \frac{1}{2}; 0 < \nu \leq 1; z \in \mathbb{U} \right)$$

for $g \in \mathcal{S}_p^*$, then f is p -valently convex (univalent) in $|z| < R(p, \mu, \nu)$, where

$$R(p, \mu, \nu) = \frac{2p\mu + 2\mu - \nu - 2 + \sqrt{(2p\mu + 2\mu - \nu - 2)^2 + 4(\nu + p)(p - 2p\mu)}}{2(\nu + p)}.$$

Proof. Suppose that

$$h(z) := \frac{zf'(z)}{pf^{1-\mu}(z)g^\mu(z)} - 1 \quad (z \in \mathbb{U}).$$

Then h is analytic in \mathbb{U} with

$$h(0) = 0 \quad \text{and} \quad |h(z)| < 1 \quad (z \in \mathbb{U}).$$

Thus, by applying Schwarz's Lemma, we get

$$h(z) = \nu z\psi(z) \quad (0 < \nu \leq 1),$$

where ψ is analytic in \mathbb{U} with

$$|\psi(z)| \leq 1 \quad (z \in \mathbb{U}).$$

Therefore, we have

$$(3.4) \quad zf'(z) = pf^{1-\mu}(z)g^\mu(z)(1 + \nu z\psi(z)).$$

Differentiating both sides of (3.4) with respect to z logarithmically, we obtain

$$(3.5) \quad 1 + \frac{zf''(z)}{f'(z)} = (1 - \mu) \frac{zf'(z)}{f(z)} + \mu \frac{zg'(z)}{g(z)} + \frac{\nu z(\psi(z) + z\psi'(z))}{1 + \nu z\psi(z)}.$$



Title Page

Contents



Page 11 of 15

Go Back

Full Screen

Close

We now suppose that

$$(3.6) \quad \phi(z) := \frac{f(z)}{z^p} = 1 + c_1z + c_2z^2 + \dots,$$

by hypothesis, we know that

$$(3.7) \quad \Re(\phi(z)) > 0 \quad (z \in \mathbb{U}).$$

It follows from (3.6) that

$$(3.8) \quad \frac{zf'(z)}{f(z)} = p + \frac{z\phi'(z)}{\phi(z)}.$$

Upon substituting (3.8) into (3.5), we get

$$(3.9) \quad 1 + \frac{zf''(z)}{f'(z)} = (1 - \mu)p + (1 - \mu)\frac{z\phi'(z)}{\phi(z)} + \mu\frac{zg'(z)}{g(z)} + \frac{\nu z(\psi(z) + z\psi'(z))}{1 + \nu z\psi(z)}.$$

Now, by using the following well known estimates (see [1]):

$$\Re\left(\frac{z\phi'(z)}{\phi(z)}\right) \geq -\frac{2r}{1-r^2}, \quad \Re\left(\frac{zg'(z)}{g(z)}\right) \geq -p\frac{1-r}{1+r},$$

and

$$\Re\left(\frac{z(\psi(z) + z\psi'(z))}{1 + z\psi(z)}\right) \geq -\frac{r}{1-r}$$

for $|z| = r < 1$ in (3.9), we obtain

$$\begin{aligned} \Re\left(1 + \frac{zf''(z)}{f'(z)}\right) &\geq (1 - \mu)p - \frac{2(1 - \mu)r}{1 - r^2} - \frac{p\mu(1 - r)}{1 + r} - \frac{\nu r}{1 - \nu r} \\ &\geq (1 - \mu)p - \frac{2(1 - \mu)r}{1 - r^2} - \frac{p\mu(1 - r)}{1 + r} - \frac{\nu r}{1 - r} \end{aligned}$$

$$\geq -\frac{(\nu + p)r^2 - (2p\mu + 2\mu - \nu - 2)r - (p - 2p\mu)}{1 - r^2},$$

which is certainly positive if $r < R(p, \mu, \nu)$. □

Putting $\nu = 1$ in Theorem 3.3, we get the following result.

Corollary 3.4. *If $f \in \mathcal{A}_p$ satisfies the following conditions:*

$$\Re \left(\frac{f(z)}{z^p} \right) > 0 \quad \text{and} \quad \left| \frac{zf'(z)}{f^{1-\mu}(z)g^\mu(z)} - p \right| < p \quad \left(0 \leq \mu < \frac{1}{2}; z \in \mathbb{U} \right)$$

for $g \in \mathcal{S}_p^*$, then f is p -valently convex (univalent) in $|z| < R(p, \mu)$, where

$$R(p, \mu) = \frac{2p\mu + 2\mu - 3 + \sqrt{(2p\mu + 2\mu - 3)^2 + 4(1+p)(p - 2p\mu)}}{2(1+p)}.$$

Remark 2. Corollary 3.4 corrects the mistakes of Theorem 3.8 which was obtained by Patel [5].

Theorem 3.5. *Let*


$$\mu \geq 0, \lambda \geq 0 \quad \text{and} \quad -1 \leq B < A \leq 1.$$

If

$$f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k \quad (z \in \mathbb{U})$$

satisfies the following subordination condition:

$$(3.10) \quad f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \lambda \left[\frac{zf''(z)}{f'(z)} + (1-\mu) \left(1 - \frac{zf'(z)}{f(z)} \right) \right] \prec \frac{1 + Az}{1 + Bz},$$



[Title Page](#)

[Contents](#)



Page 12 of 15

[Go Back](#)

[Full Screen](#)

[Close](#)



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 13 of 15

[Go Back](#)

[Full Screen](#)

[Close](#)

then

$$(3.11) \quad |a_{n+1}| \leq \frac{A - B}{(1 + n\lambda)(n + \mu)} \quad (n \in \mathbb{N}).$$

Proof. Suppose that

$$f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k \quad (z \in \mathbb{U})$$

satisfies (3.10). It follows that

$$(3.12) \quad f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \lambda \left[\frac{z f''(z)}{f'(z)} + (1 - \mu) \left(1 - \frac{z f'(z)}{f(z)} \right) \right] \\ = 1 + (1 + n\lambda)(n + \mu)a_{n+1}z^n + \cdots < \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}).$$

Therefore, we find from Lemma 2.4, (3.12) and $-1 \leq B < A \leq 1$ that

$$(3.13) \quad |(1 + n\lambda)(n + \mu)a_{n+1}| \leq A - B.$$

The assertion (3.11) of Theorem 3.5 can now easily be derived from (3.13). \square

Taking $A = 1 - 2\alpha$ ($0 \leq \alpha < 1$) and $B = -1$ in Theorem 3.5, we get the following result.

Corollary 3.6. *Let*

$$\mu \geq 0, \lambda \geq 0 \quad \text{and} \quad 0 \leq \alpha < 1.$$

If

$$f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k$$



satisfies the following inequality:

$$\Re \left(f'(z) \left(\frac{f(z)}{z} \right)^{\mu-1} + \lambda \left[\frac{zf''(z)}{f'(z)} + (1-\mu) \left(1 - \frac{zf'(z)}{f(z)} \right) \right] \right) > \alpha,$$

then

$$|a_{n+1}| \leq \frac{2(1-\alpha)}{(1+n\lambda)(n+\mu)} \quad (n \in \mathbb{N}).$$

Remark 3. Corollary 3.6 provides an extension of the corresponding result obtained by Guo and Liu [2].

p-Valently Bazilevič Functions

Zhi-Gang Wang and Yue-Ping Jiang

vol. 9, iss. 3, art. 70, 2008

Title Page

Contents



Page 14 of 15

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] W.M. CAUSEY AND E.P. MERKES, Radii of starlikeness for certain classes of analytic functions, *J. Math. Anal. Appl.*, **31** (1970), 579–586.
- [2] D. GUO AND M.-S. LIU, On certain subclass of Bazilevič functions, *J. Inequal. Pure Appl. Math.*, **8**(1) (2007), Art. 12. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=825>].
- [3] M.-S. LIU, On a subclass of p -valent close-to-convex functions of order β and type α , *J. Math. Study*, **30** (1997), 102–104 (in Chinese).
- [4] M.-S. LIU, On certain subclass of analytic functions, *J. South China Normal Univ.*, **4** (2002), 15–20 (in Chinese).
- [5] J. PATEL, On certain subclass of p -valently Bazilevič functions, *J. Inequal. Pure Appl. Math.*, **6**(1) (2005), Art. 16. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=485>].
- [6] W. ROGOSINSKI, On the coefficients of subordinate functions, *Proc. London Math. Soc. (Ser. 2)*, **48** (1943), 48–82.



p -Valently Bazilevič Functions

Zhi-Gang Wang and Yue-Ping Jiang

vol. 9, iss. 3, art. 70, 2008

Title Page

Contents



Page 15 of 15

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756