

HYERS-ULAM STABILITY OF THE GENERALIZED QUADRATIC FUNCTIONAL EQUATION IN AMENABLE SEMIGROUP

BOUIKHALENE BELAID

Department of Mathematics, Faculty of Sciences,
University of Ibn Tofail, Kenitra, Morocco.
EMail: bbouikhalene@yahoo.fr

ELQORACHI ELHOUCIEN AND REDOUANI AHMED

Laboratory LAMA, Harmonic Analysis and Functional Equations Team
Department of Mathematics
Faculty of Sciences, University Ibn Zohr
Agadir, Morocco
EMail: elqorachi@hotmail.com and redouani_ahmed@yahoo.fr

Received: 6 March, 2007

Accepted: 26 April, 2007

Communicated by: [Th.M. Rassias](#)

2000 AMS Sub. Class.: 39B82, 39B52.



Key words: Hyers-Ulam stability, Quadratic functional equation,
Amenable semigroup, Morphism of semigroup.



Hyers-Ulam Stability
Bouikhalene Belaid, Elqorachi Elhoucien
and Redouani Ahmed
vol. 8, iss. 2, art. 47, 2007

[Title Page](#)

[Contents](#)



Page 1 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Abstract:

In this paper we derive the Hyers-Ulam stability of the quadratic functional equation

$$f(xy) + f(x\sigma(y)) = 2f(x) + 2f(y), \quad x, y \in G,$$

respectively the functional equation

$$f(xy) + g(x\sigma(y)) = f(x) + g(y), \quad x, y \in G,$$

where G is an amenable semigroup, σ is a morphism of G such that $\sigma \circ \sigma = I$, respectively where G is an amenable semigroup and σ is an homomorphism of G such that $\sigma \circ \sigma = I$.

Hyers-Ulam Stability

Bouikhalene Belaid, Elqorachi Elhoucien
and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

Title Page

Contents



Page 2 of 36

Go Back

Full Screen


Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	4
2	Stability of Equation (1.4) in Amenable Semigroups	6
3	Stability of Equation (1.5) in Amenable Semigroups	22



Hyers-Ulam Stability
Bouikhalene Belaid, Elqorachi Elhoucien
and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

Title Page

Contents



Page 3 of 36


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 4 of 36

Go Back

Full Screen

Close

1. Introduction

In 1940, Ulam [21] raised a question concerning the stability problem of group homomorphisms:

Given a group G_1 , a metric group (G_2, d) , a number $\varepsilon > 0$ and a mapping $f : G_1 \rightarrow G_2$ which satisfies the inequality $d(f(xy), f(x)f(y)) < \varepsilon$ for all $x, y \in G_1$, does there exist a homomorphism $h : G_1 \rightarrow G_2$ and a constant $k > 0$, depending only on G_1 and G_2 such that $d(f(x), h(x)) \leq k\varepsilon$ for all x in G_1 ?

The case of approximately additive mappings was solved by D. H. Hyers [8] under the assumption that G_1 and G_2 are Banach spaces.

In 1978, Th. M. Rassias [16] gave a remarkable generalization of the Hyers's result which allows the Cauchy difference to be unbounded. Since then, several mathematicians have been attracted to the results of Hyers and Rassias and investigated a number of stability problems of different functional equations. See for example the monographs of the following references [5, 6, 9, 10, 11, 16].

The quadratic functional equation

$$(1.1) \quad f(x + y) + f(x - y) = 2f(x) + 2f(y), \quad x, y \in G$$

has been much studied. It was generalized by Stetkær [19] to the more general equation

$$(1.2) \quad f(x + y) + f(x + \sigma(y)) = 2f(x) + 2f(y), \quad x, y \in G,$$

where σ is an automorphism of the abelian group G such that $\sigma^2 = I$, (I denotes the identity).

A stability result for the quadratic functional equation (1.1) was derived by Skof [18], Cholewa [3] and by Czerwik [4].



Recently, Bouikhalene, Elqorachi and Rassias stated the stability theorem of equation (1.2), see [1] and [2].

Székelyhidi [20] extended the Hyers's result to amenable semigroups. He replaced the original proof given by Hyers by a new one based on the use of invariant means.

In [22] Yang obtained the stability of the quadratic functional equation

$$(1.3) \quad f(xy) + f(xy^{-1}) = 2f(x) + 2f(y), \quad x, y \in G,$$

in amenable groups.

The purpose of the present paper is a joint treatment of the functional equations (1.1) and (1.3) and their generalization, where the unifying object is a morphism like the one introduced in (1.2).

New features of the paper:

1. In comparison with [20] and [22], we work with a general morphism σ .
2. In contrast to [1] and [2] we here allow the (semi)group G to be non-abelian.

In Section 2, we obtain the stability of the quadratic functional equation

$$(1.4) \quad f(xy) + f(x\sigma(y)) = 2f(x) + 2f(y), \quad x, y \in G,$$

where σ is a morphism of G such that $\sigma \circ \sigma = I$. The result of this section can be compared with the ones of Yang [22] because we formulate them in the same way by using some ideas from [22].

In Section 3, we obtain the stability of the generalized quadratic functional equation

$$(1.5) \quad f(xy) + g(x\sigma(y)) = f(x) + g(y), \quad x, y \in G,$$

where σ is an automorphism of G such that $\sigma \circ \sigma = I$.

Title Page

Contents



Page 5 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



2. Stability of Equation (1.4) in Amenable Semigroups

In this section we investigate the Hyers-Ulam stability of the quadratic functional equation

$$(2.1) \quad f(xy) + f(x\sigma(y)) = 2f(x) + 2f(y), \quad x, y \in G,$$

where G is an amenable semigroup with unit element e and $\sigma : G \rightarrow G$ is a morphism of G , i.e. σ is an antiautomorphism: $\sigma(xy) = \sigma(y)\sigma(x)$ for all $x, y \in G$ or σ is an automorphism: $\sigma(xy) = \sigma(x)\sigma(y)$ for all $x, y \in G$. Furthermore, we assume that σ satisfies $(\sigma \circ \sigma)(x) = x$, for all $x \in G$.

We recall that a semigroup G is said to be amenable if there exists an invariant mean on the space of the bounded complex functions defined on G . We refer to [7] for the definition and properties of invariant means.

Throughout this paper, as in [5], we use the following definition.

Definition 2.1. *Let G be a semigroup and B a Banach space. We say that the equation*

$$(2.2) \quad f(xy) + f(x\sigma(y)) = 2f(x) + 2f(y), \quad x, y \in G$$

is stable for the pair (G, B) if for every function $f : G \rightarrow B$ such that

$$(2.3) \quad \left\| \frac{1}{2}[f(xy) + f(x\sigma(y))] - f(x) - f(y) \right\| \leq \delta, \quad x, y \in G \text{ for some } \delta \geq 0,$$

there exists a solution q of equation (2.2) and a constant $\gamma \geq 0$ dependent only on δ such that

$$(2.4) \quad \|f(x) - q(x)\| \leq \gamma \text{ for all } x \in G.$$

Hyers-Ulam Stability

Bouikhalene Belaid, Elqorachi Elhoucien

and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

[Title Page](#)

[Contents](#)



Page 6 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 7 of 36

Go Back

Full Screen

Close

Proposition 2.1. Let σ be an antiautomorphism of the semigroup G such that $\sigma \circ \sigma = I$. Let B a Banach space. Suppose that $f : G \rightarrow B$ satisfies the inequality (2.3). Then for every $x \in G$, the limit

$$(2.5) \quad g(x) = \lim_{n \rightarrow +\infty} 2^{-2n} \left[f(x^{2^n}) + \sum_{k=1}^n 2^{k-1} f((x^{2^{n-k}} \sigma(x)^{2^{n-k}})^{2^{k-1}}) \right]$$

exists. Moreover, $g : G \rightarrow \mathbb{C}$ is a unique function satisfying

$$(2.6) \quad \|f(x) - g(x)\| \leq \delta, \text{ and } g(x^2) + g(x\sigma(x)) = 4g(x) \text{ for all } x \in G.$$

Proof. Assume that $f : G \rightarrow \mathbb{C}$ satisfies the inequality (2.3) and define by induction the sequence function $f_0(x) = f(x)$ and $f_n(x) = \frac{1}{2}[f_{n-1}(x^2) + f_{n-1}(x\sigma(x))]$ for $n \geq 1$. By direct computation, we obtain

$$f_n(x) = 2^{-n} \left[f(x^{2^n}) + \sum_{k=1}^n 2^{k-1} f((x^{2^{n-k}} \sigma(x)^{2^{n-k}})^{2^{k-1}}) \right]$$

for all $n \geq 1$.

By letting $x = y$, in (2.3) we get


$$(2.7) \quad \left\| \frac{1}{2}[f(x^2) + f(x\sigma(x))] - 2f(x) \right\| \leq \delta,$$

so

$$(2.8) \quad \|f_1(x) - 2f_0(x)\| \leq \delta \text{ for all } x \in G.$$

In the following, we prove by induction the inequalities

$$(2.9) \quad \|f_n(x) - 2f_{n-1}(x)\| \leq \delta$$



$$(2.10) \quad \|f_n(x) - 2^n f(x)\| \leq (2^n - 1)\delta$$

for all $n \in \mathbb{N}$ and $x \in G$. It is clear that (2.8) is (2.9) for $n = 1$. The inductive step must now be demonstrated to hold true for the integer $n + 1$, that is

$$(2.11) \quad \begin{aligned} & \|f_{n+1}(x) - 2f_n(x)\| \\ &= \left\| \frac{1}{2}[f_n(x^2) + f_n(x\sigma(x))] - 2\frac{1}{2}[f_{n-1}(x^2) + f_{n-1}(x\sigma(x))] \right\| \\ &\leq \frac{1}{2}[\|f_n(x^2) - 2f_{n-1}(x^2)\|] + \left\| \frac{1}{2}[f_n(x\sigma(x)) - 2f_{n-1}(x\sigma(x))] \right\| \\ &\leq \frac{1}{2}[\delta + \delta] = \delta. \end{aligned}$$

This proves that (2.9) is true for any natural number n .

Now, by using the inequality

$$(2.12) \quad \|f_n(x) - 2^n f(x)\| \leq \|f_n(x) - 2f_{n-1}(x)\| + 2\|f_{n-1}(x) - 2^{n-1}f(x)\|$$

we check that (2.10) holds true for any $n \in \mathbb{N}$.

Let us define

$$(2.13) \quad g_n(x) = \frac{f_n(x)}{2^n} = 2^{-2n} \left[f(x^{2^n}) + \sum_{k=1}^n 2^{k-1} f((x^{2^{n-k}} \sigma(x)^{2^{n-k}})^{2^{k-1}}) \right]$$

for any positive integer n and $x \in G$.

Now, by using (2.11) and (2.13), we get

$$(2.14) \quad \|g_{n+1}(x) - g_n(x)\| \leq \frac{\delta}{2^{n+1}}.$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 8 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



It easily follows that $\{g_n(x)\}$ is a Cauchy sequence for all $x \in G$. Since B is complete, we can define $g(x) = \lim_{n \rightarrow +\infty} g_n(x)$ for any $x \in G$. From (2.10), one can verify that g satisfies the first assertion of (2.6). Now, we will show that g satisfies the second assertion of (2.6). By induction one proves that the sequence $f_n(x)$ satisfies

$$(2.15) \quad \left\| \frac{1}{2}[f_n(x^2) + f_n(x\sigma(x))] - f_n(x) - f_n(x) \right\| \leq \delta$$

for all $n \in \mathbb{N}$.

For $n = 1$, we have

$$\begin{aligned}
 (2.16) \quad & \left\| \frac{1}{2}[f_1(x^2) + f_1(x\sigma(x))] - f_1(x) - f_1(x) \right\| \\
 &= \left\| \frac{1}{2} \left[\frac{1}{2} \left[f(x^2) + f(x^2\sigma(x)^2) + f((x\sigma(x))^2) + f((x\sigma(x))^2) \right] \right. \right. \\
 &\quad \left. \left. - \frac{1}{2}[f(x^2) + f(x\sigma(x))] - \frac{1}{2}[f(x^2) + f(x\sigma(x))] \right] \right\| \\
 &\leq \left\| \frac{1}{2} \left[\frac{1}{2} \left[f(x^2) + f(x^2\sigma(x)^2) - 2f(x^2) \right] \right. \right. \\
 &\quad \left. \left. + \frac{1}{2} \left[\frac{1}{2} \left[f((x\sigma(x))^2) + f((x\sigma(x))^2) - 2f(x\sigma(x))] \right] \right] \right\| \\
 &\leq \frac{1}{2}[\delta + \delta] = \delta.
 \end{aligned}$$

So (2.15) is true for $n = 1$. We assume then that (2.15) holds for n and we prove

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 9 of 36

Go Back

Full Screen

Close

journal of **inequalities**
 in pure and applied
 mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 10 of 36

Go Back

Full Screen

Close

that (2.15) is true for $n + 1$.

$$\begin{aligned}
 (2.17) \quad & \left\| \frac{1}{2} [f_{n+1}(x^2) + f_{n+1}(x\sigma(x))] - f_{n+1}(x) - f_{n+1}(x) \right\| \\
 &= \left\| \frac{1}{2} \left[\frac{1}{2} [f_n(x^2) + f_n(x^2\sigma(x)^2) + f_n((x\sigma(x))^2) + f_n((x\sigma(x))^2)] \right. \right. \\
 &\quad \left. \left. - \frac{1}{2} [f_n(x^2) + f_n(x\sigma(x))] - \frac{1}{2} [f_n(x^2) + f_n(x\sigma(x))] \right] \right\| \\
 &\leq \left\| \frac{1}{2} \left[\frac{1}{2} [f_n(x^2) + f_n(x^2\sigma(x)^2) - 2f_n(x^2)] \right] \right\| \\
 &\quad + \left\| \frac{1}{2} \left[\frac{1}{2} [f_n((x\sigma(x))^2) + f_n((x\sigma(x))^2) - 2f_n(x\sigma(x))] \right] \right\| \\
 &\leq \frac{1}{2} [\delta + \delta] = \delta.
 \end{aligned}$$

Consequently, the sequence $g_n(x)$ satisfies

$$(2.18) \quad \left\| \frac{1}{2} [g_n(x^2) + g_n(x\sigma(x))] - g_n(x) - g_n(x) \right\| \leq \frac{\delta}{2^n}$$

for all $n \in \mathbb{N}$ and then by letting $n \rightarrow +\infty$ we get the desired result.

Assume now that there exists another mapping $h : G \rightarrow B$ which satisfies

$$\|f(x) - h(x)\| \leq \delta \text{ and } h(x^2) + h(x\sigma(x)) = 4h(x) \text{ for all } x \in G.$$

First, we will prove by mathematical induction that

$$(2.19) \quad \|f(x) - 2^n h(x)\| \leq \delta \text{ for all } x \in G.$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 11 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)

For $n = 1$, we have

$$\begin{aligned}
 (2.20) \quad \|f_1(x) - 2h(x)\| &= \left\| \frac{1}{2}[f(x^2) + f(x\sigma(x))] - \frac{1}{2}[h(x^2) + h(x\sigma(x))] \right\| \\
 &\leq \frac{1}{2}[\|f(x^2) - h(x^2)\| + \|f(x\sigma(x)) - h(x\sigma(x))\|] \\
 &\leq \frac{1}{2}[\delta + \delta] = \delta.
 \end{aligned}$$

Suppose (2.19) is true for n and we will prove it for $n + 1$. Hence, we have

$$\begin{aligned}
 (2.21) \quad \|f_{n+1}(x) - 2^{n+1}h(x)\| &= \left\| \frac{1}{2}[f_n(x^2) + f_n(x\sigma(x))] - 2^n \frac{1}{2}[h(x^2) + h(x\sigma(x))] \right\| \\
 &\leq \frac{1}{2}[\|f_n(x^2) - 2^n h(x^2)\| + \|f_n(x\sigma(x)) - 2^n h(x\sigma(x))\|] \\
 &\leq \frac{1}{2}[\delta + \delta] = \delta.
 \end{aligned}$$

This proves that (2.19) is true for all $n \in \mathbb{N}$. From (2.19), we obtain $\left\| \frac{f_n(x)}{2^n} - h(x) \right\| \leq \frac{\delta}{2^n}$, so by letting $n \rightarrow +\infty$ and by using the definition of g we get $g = h$. This completes the proof of the theorem. \square

By using the precedent proof we easily obtain the following result.

Proposition 2.2. *Let σ be an automorphism of the semigroup G such that $\sigma \circ \sigma = I$. Let B a Banach space. Suppose that $f : G \rightarrow B$ satisfies the inequality (2.3). Then for every $x \in G$, the limit*

$$g(x) = \lim_{n \rightarrow +\infty} 2^{-n} f_n(x)$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 12 of 36

Go Back

Full Screen

Close

exists, and g is a unique function satisfying

$$\|f(x) - g(x)\| \leq \delta, \text{ and } g(x^2) + g(x\sigma(x)) = 4g(x) \text{ for all } x \in G,$$

where the sequence of functions f_n is defined on G by the formulas $f_0(x) = f(x)$ and $f_n(x) = \frac{1}{2}[f_{n-1}(x^2) + f_{n-1}(x\sigma(x))]$ for $n \geq 1$.

The following result shows that in this context the only property of B (Definition 2.1) involved is the completeness. For the proof, we refer to the one used by Yang in [22].

Theorem 2.3. *Let σ be a morphism of the semigroup G such that $\sigma \circ \sigma = I$. Suppose that equation (2.2) is stable for the pair (G, \mathbb{C}) (resp. (G, \mathbb{R})) Then for every complex (resp. real) Banach space B , (2.2) is stable for the pair (G, B) .*

The main result of the present section is the following

Theorem 2.4. *Let σ be an antiautomorphism of the amenable semigroup G such that $\sigma \circ \sigma = I$. Then equation (2.2) is stable for the pair (G, \mathbb{C}) .*

First, we prove the following useful lemma

Lemma 2.5. *Let σ be an antiautomorphism of the semigroup G such that $\sigma \circ \sigma = I$. Let B be a Banach space. Suppose that $f : G \rightarrow B$ satisfies the inequality*


$$(2.22) \quad \left\| \frac{1}{2}(f(xy) + f(x\sigma(y))) - f(x) - f(y) \right\| \leq \delta, \text{ for some } \delta \geq 0.$$

Then for every $x \in G$, the limit

$$(2.23) \quad q(x) = \lim_{n \rightarrow +\infty} 2^{-2n} \left\{ f(x^{2^n}) + (2^n - 1)f(x^{2^{n-1}}\sigma(x)^{2^{n-1}}) \right\}$$

exists. Moreover, the mapping q satisfies the inequality

$$(2.24) \quad \|f(x) - q(x)\| \leq 7\delta \text{ for all } x \in G.$$



Title Page

Contents



Page 13 of 36

Go Back

Full Screen

Close

Proof. Assume that $f : G \rightarrow B$ satisfies the inequality (2.22). We will prove by induction that

$$(2.25) \quad \left\| f(x) - \frac{1}{2^{2n}} \{f(2^n x) + (2^n - 1)f(2^{n-1}x + 2^{n-1}\sigma(x))\} \right\| \leq 2 \left(\frac{7}{2} + \frac{3}{2^{2n-1}} - \frac{19}{2^{n+1}} \right) \delta$$

for some positive integer n . By letting $y = x$, in (2.22) we get

$$(2.26) \quad \|f(x^2) + (2 - 1)f(x\sigma(x)) - 2^2 f(x)\| \leq 2\delta,$$

so

$$(2.27) \quad \left\| f(x) - \frac{1}{2^2} \{f(x^2) + (2 - 1)f(x\sigma(x))\} \right\| \leq \delta \left(1 - \frac{1}{2} \right) \quad \text{for all } x \in G.$$

This proves (2.25) for $n = 1$. The inductive step must now be demonstrated to hold true for the integer $n + 1$, that is

$$\begin{aligned} & \left\| f(x) - \frac{1}{2^{2(n+1)}} \left\{ f(x^{2^{n+1}}) + (2^{n+1} - 1)f(x^{2^n} \sigma(x)^{2^n}) \right\} \right\| \\ & \leq \frac{1}{2^{2(n+1)}} \left\| f(x^{2^{n+1}}) + f(x^{2^n} \sigma(x)^{2^n}) - 4f(x^{2^n}) \right\| \\ & \quad + \frac{1}{2^{2(n+1)}} \left\| 2(2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}} x^{2^{n-1}} \sigma(x)^{2^{n-1}}) - 4(2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right\| \\ & \quad + \frac{1}{2^{2(n+1)}} \left\| 4f(x^{2^n}) + 4(2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}}) - 2^{2(n+1)} f(x) \right\| \\ & \quad + \frac{2(2^n - 1)}{2^{2(n+1)}} \left\| f(x^{2^n} \sigma(x)^{2^n}) - f(x^{2^{n-1}} \sigma(x)^{2^{n-1}} x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right\| \end{aligned}$$



$$\leq \frac{2\delta}{2^{2(n+1)}} + \frac{(2^n - 1)2\delta}{2^{2(n+1)}} + \left(\frac{7}{2} + \frac{3}{2^{2n-1}} - \frac{19}{2^{n+1}} \right) 2\delta \\ + \frac{2(2^n - 1)}{2^{2(n+1)}} \left\| f(x^{2^n} \sigma(x)^{2^n}) - f\left(x^{2^{n-1}} \sigma(x)^{2^{n-1}} x^{2^{n-1}} \sigma(x)^{2^{n-1}}\right) \right\|.$$

To complete the proof of the induction assumption (2.25), we need the following inequalities. Let $x = y = e$ in (2.22) to get $\|2f(e)\| \leq 2\delta$. Putting $x = e$ in (2.22), gives

$$\|f(y) + f(\sigma(y)) - 2f(e) - 2f(y)\| \leq 2\delta.$$

Consequently,

$$(2.28) \quad \|f(y) - f(\sigma(y))\| \leq 4\delta.$$

By interchanging x by y in (2.22), we obtain

$$(2.29) \quad \|f(yx) + f(y\sigma(x)) - 2f(x) - 2f(y)\| \leq 2\delta.$$

By using (2.22), (2.28), (2.29) and the triangle inequality, we deduce that

$$(2.30) \quad \|f(xy) - f(yx)\| \leq 8\delta.$$

Now, from (2.22), we obtain

$$(2.31) \quad \left\| 2f\left(x^{2^{n-1}} \sigma(x)^{2^n} x^{2^{n-1}}\right) - 2f\left(x^{2^{n-1}} \sigma(x)^{2^{n-1}}\right) - 2f\left(\sigma(x)^{2^{n-1}} x^{2^{n-1}}\right) \right\| \\ \leq 2\delta.$$

Since

$$(2.32) \quad \|f(x^{2^n} \sigma(x)^{2^n}) - f(\sigma(x)^{2^n} x^{2^n})\| \leq 8\delta$$

Title Page

Contents



Page 14 of 36


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



and

$$(2.33) \quad \left\| f\left(x^{2^{n-1}}\sigma(x)^{2^n}x^{2^{n-1}}\right) - f\left(x^{2^n}\sigma(x)^{2^n}\right) \right\| \leq 8\delta$$

then

$$(2.34) \quad \left\| 2f\left(x^{2^{n-1}}\sigma(x)^{2^n}x^{2^{n-1}}\right) - 4f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) \right\| \leq 18\delta$$

and



$$(2.35) \quad \begin{aligned} & \left\| f\left(x^{2^n}\sigma(x)^{2^n}\right) - 2f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) \right\| \\ & \leq \left\| f\left(x^{2^{n-1}}\sigma(x)^{2^n}x^{2^{n-1}}\right) - f\left(x^{2^n}\sigma(x)^{2^n}\right) \right\| \\ & \quad + \left\| f\left(x^{2^{n-1}}\sigma(x)^{2^n}x^{2^{n-1}}\right) - f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) \right\| \\ & \leq 8\delta + \frac{18\delta}{2} = 17\delta. \end{aligned}$$

Finally, we deduce that

$$(2.36) \quad \begin{aligned} & \left\| f\left(x^{2^n}\sigma(x)^{2^n}\right) - f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) \right\| \\ & \leq \left\| f\left(x^{2^n}\sigma(x)^{2^n}\right) - 2f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) \right\| \\ & \quad + \left\| 2f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) - f\left(x^{2^{n-1}}\sigma(x)^{2^{n-1}}x^{2^{n-1}}\sigma(x)^{2^{n-1}}\right) \right\| \\ & \leq 17\delta + \delta = 18\delta \end{aligned}$$

Title Page

Contents



Page 15 of 36


Go Back

Full Screen

Close



journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 16 of 36

Go Back

Full Screen

Close

journal of **inequalities**
 in pure and applied
 mathematics

issn: 1443-5756

and consequently,

$$\begin{aligned} & \left\| f(x) - \frac{1}{2^{2(n+1)}} \left\{ f(x^{2^{n+1}}) + (2^{n+1} - 1)f(x^{2^n} \sigma(x)^{2^n}) \right\} \right\| \\ & \leq \frac{2\delta}{2^{2(n+1)}} + \frac{2(2^n - 1)\delta}{2^{2(n+1)}} + \left(\frac{7}{2} + \frac{3}{2^{2n-1}} - \frac{19}{2^{n+1}} \right) 2\delta + \frac{2(2^n - 1)}{2^{2(n+1)}} 18\delta \\ & = 2 \left(\frac{7}{2} + \frac{3}{2^{2n+1}} - \frac{19}{2^{n+2}} \right) \delta. \end{aligned}$$

This proves the validity of the inequality (2.25).


Let us define

$$(2.37) \quad q_n(x) = \frac{1}{2^{2n}} \left\{ f(x^{2^n}) + (2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right\}$$

for any positive integer n and $x \in G$.

Then $\{q_n(x)\}$ is a Cauchy sequence for every $x \in G$. In fact by using (2.22), (2.36) and (2.37), we get

$$\begin{aligned} & \|q_{n+1}(x) - q_n(x)\| \\ & \leq \frac{1}{2^{2(n+1)}} \left\| f(x^{2^{n+1}}) + f(x^{2^n} \sigma(x)^{2^n}) - 4f(x^{2^n}) \right\| \\ & \quad + \frac{1}{2^{2(n+1)}} \left\| 2(2^n - 1)f(x^{2^n} \sigma(x)^{2^n}) - 4(2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right\| \\ & \leq \frac{\delta}{2^{2(n+1)}} + \frac{1}{2^{2(n+1)}} \left\| 2(2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}} x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right. \\ & \quad \left. - 4(2^n - 1)f(x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right\| \\ & \quad + \frac{2(2^n - 1)}{2^{2(n+1)}} \left\| f(x^{2^n} \sigma(x)^{2^n}) - f(x^{2^{n-1}} \sigma(x)^{2^{n-1}} x^{2^{n-1}} \sigma(x)^{2^{n-1}}) \right\| \end{aligned}$$



$$\leq \frac{2\delta}{2^{2(n+1)}} + \frac{2(2^n - 1)\delta}{2^{2(n+1)}} + \frac{36\delta(2^n - 1)}{2^{2(n+1)}} \leq \frac{40\delta}{2^n}.$$

It easily follows that $\{q_n(x)\}$ is a Cauchy sequence for all $x \in G$. Since B is complete, we can define $q(x) = \lim_{n \rightarrow +\infty} q_n(x)$ for any $x \in G$ and, in view of (2.25) one can verify that q satisfies the inequality (2.24). This completes the proof of Lemma 2.5. \square

Proof of Theorem 2.4. We follow the ideas and the computations used in [22]. By using (2.30) one derives the functional inequality

$$(2.38) \quad |f((xy)^n) - f((yx)^n)| \leq 8\delta.$$

In addition, from (2.3), (2.38) and the triangle inequality we deduce that

$$(2.39) \quad |f((xy)^{2^n}(\sigma(xy))^{2^n}) - f((yx)^{2^n}(\sigma(yx))^{2^n})| \\ \leq |f((xy)^{2^n}(\sigma(xy))^{2^n}) + f((xy)^{2^n}(xy)^{2^n}) - 4f((xy)^{2^n})| \\ + | - f((yx)^{2^n}(yx)^{2^n}) - f((yx)^{2^n}(\sigma(yx))^{2^n}) + 4f((yx)^{2^n})| \\ + |f((xy)^{2^n}(xy)^{2^n}) - f((yx)^{2^n}(yx)^{2^n})| + 4|f((xy)^{2^n}) - f((yx)^{2^n})| \\ \leq 2\delta + 2\delta + 8\delta + 32\delta = 44\delta.$$

From Lemma 2.5, for every $x \in G$, the limit

$$(2.40) \quad q(x) = \lim_{n \rightarrow +\infty} 2^{-2n} \left\{ f(x^{2^n}) + (2^n - 1)f\left(x^{2^n-1}\sigma(x)^{2^n-1}\right) \right\}$$

exists and

$$(2.41) \quad |f(x) - q(x)| \leq 7\delta.$$

Furthermore, in view of (2.38) – (2.39) q satisfies the relation

$$(2.42) \quad q(xy) = q(yx), \quad x, y \in G$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 17 of 36

Go Back

Full Screen

Close



and by (2.41) – (2.22) q satisfies the inequality equation

$$(2.43) \quad |q(xy) + q(x\sigma(y)) - 2q(x) - 2q(y)| \leq 44\delta$$

for all $x, y \in G$.

Consequently, for any fixed $y \in G$ the function

$$x \longmapsto q(xy) + q(x\sigma(y)) - 2q(x)$$

is bounded. Since G is amenable, there exists an invariant mean m_x on the space of bounded, complex-functions on G . With the help of m_x we define the following function on G

$$(2.44) \quad \psi(y) = m_x\{q_y + q_{\sigma(y)} - 2q\}$$

for all $y \in G$, where $q_y(z) = q(z)y$, $z \in G$.

Furthermore, by using (2.29) and (2.44), we get

$$(2.45) \quad \begin{aligned} \psi(z)y + \psi(\sigma(z)y) &= m_x\{q_{zy} + q_{\sigma(y)\sigma(z)} - 2q\} + m_x\{q_{\sigma(z)y} + q_{\sigma(y)z} - 2q\} \\ &= m_x\{zyq + q_{\sigma(y)\sigma(z)} - 2q\} + m_x\{\sigma(z)yq + q_{\sigma(y)z} - 2q\} \\ &= m_x\{zyq + \sigma(z)yq - 2(yq)\} + m_x\{q_{\sigma(y)\sigma(z)} + q_{\sigma(y)z} - 2q_{\sigma(y)}\} \\ &\quad + m_x\{2q_y + 2q_{\sigma(y)} - 4q\} \\ &= 2\psi(z) + 2\psi(y). \end{aligned}$$

So, $Q(y) = \frac{\psi(y)}{2}$ satisfies equation (2.2) and the following inequality

$$(2.46) \quad \begin{aligned} |Q(y) - q(y)| &= \frac{1}{2} |m_x\{q_y + q_{\sigma(y)} - 2q - 2q(y)\}| \\ &\leq \sup_{x \in G} \frac{1}{2} | \{q(xy) + q(x\sigma(y)) - 2q(x) - 2q(y)\} | \end{aligned}$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 18 of 36

Go Back

Full Screen

Close

$$\leq \frac{44}{2}\delta = 22\delta.$$

Consequently, there exists a mapping Q which satisfies the functional equation (2.2) and the inequality $|f(y) - Q(y)| \leq 29\delta$. This completes the proof of the theorem. \square

Theorem 2.6. *Let σ be an automorphism of the amenable semigroup G such that $\sigma \circ \sigma = I$. Then equation (2.2) is stable for the pair (G, \mathbb{C}) .*

Proof. From inequality (2.3), we deduce that for any fixed $y \in G$ the function $x \mapsto f(xy) + f(x\sigma(y)) - 2f(x)$ is bounded. Since G is amenable, then we can define

$$(2.47) \quad \phi(y) = m_x \{f_y + f_{\sigma(y)} - 2f\}$$

for all $y \in G$.


We have

$$(2.48) \quad \begin{aligned} \phi(yz) + \phi(y\sigma(z)) &= m_x \{f_{yz} + f_{\sigma(y)\sigma(z)} - 2f\} + m_x \{f_{y\sigma(z)} + f_{\sigma(y)z} - 2f\} \\ &= m_x \{f_{yz} + f_{y\sigma(z)} - 2f_y\} + m_x \{f_{\sigma(y)\sigma(z)} + f_{\sigma(y)z} - 2f_{\sigma(y)}\} \\ &\quad + 2m_x \{f_y + f_{\sigma(y)} - 2f\} \\ &= \phi(z) + \phi(\sigma(z)) + 2\phi(y) = 2\phi(z) + 2\phi(y), \end{aligned}$$

so ϕ is a solution of equation (2.2). Moreover, we have



$$(2.49) \quad \begin{aligned} \left| f(y) - \frac{\phi(y)}{2} \right| &= \frac{1}{2} |m_x \{f_y + f_{\sigma(y)} - 2f - 2f(y)\}| \\ &\leq \frac{1}{2} \sup_{x \in G} |f(xy) + f(x\sigma(y)) - 2f(x) - 2f(y)| \leq \delta. \end{aligned}$$

This completes the proof of theorem. \square



Title Page

Contents



Page 19 of 36


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



By using Theorem 2.4 and the proof of Proposition 2.1 we get the following corollaries.

In the first corollary, the estimate improves the ones obtained in the proof of Theorem 2.4.

Corollary 2.7. *Let σ be an antiautomorphism of the amenable semigroup G such that $\sigma \circ \sigma = I$. Let B a Banach space. Suppose that $f : G \rightarrow B$ satisfies the inequality (2.3). Then for every $x \in G$, the limit*

$$(2.50) \quad Q(x) = \lim_{n \rightarrow +\infty} 2^{-2n} \left[f(x^{2^n}) + \sum_{k=1}^n 2^{k-1} f \left((x^{2^{n-k}} \sigma(x)^{2^{n-k}})^{2^{k-1}} \right) \right]$$

exists. Moreover, Q is the unique solution of equation (1.4) satisfying

$$(2.51) \quad \|f(x) - Q(x)\| \leq \delta \text{ for all } x \in G.$$

Corollary 2.8. *Let σ be a morphism of the amenable semigroup G such that $\sigma \circ \sigma = I$. Then for every Banach space B , equation (2.2) is stable for the pair (G, B) .*

Corollary 2.9 ([20]). *Let $\sigma = I$. Let G be an amenable semigroup. Then for every Banach space B , equation*

$$(2.52) \quad f(xy) = f(x) + f(y), \quad x, y \in G$$

is stable for the pair (G, B) .

Corollary 2.10 ([22]). *Let $\sigma(x) = x^{-1}$. Let G be an amenable group. Then for every Banach space B , equation*

$$(2.53) \quad f(xy) + f(xy^{-1}) = 2f(x) + 2f(y), \quad x, y \in G$$

is stable for the pair (G, B) .

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 20 of 36


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Corollary 2.11 ([2]). *Let σ be an automorphism of the vector space G such that $\sigma \circ \sigma = I$. Then for every Banach space B , the equation*

$$(2.54) \quad f(x + y) + f(x + \sigma(y)) = 2f(x) + 2f(y), \quad x, y \in G$$

is stable for the pair (G, B) .

Hyers-Ulam Stability

Bouikhalene Belaid, Elqorachi Elhoucien

and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

Title Page

Contents



Page 21 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 22 of 36

Go Back

Full Screen

Close

3. Stability of Equation (1.5) in Amenable Semigroups

In this section we investigate the Hyers-Ulam stability of the functional equation

$$(3.1) \quad f(xy) + g(x\sigma(y)) = f(x) + g(y), \quad x, y \in G,$$

where G is an amenable semigroup with element unity e and $\sigma : G \rightarrow G$ is an automorphism of G such that $\sigma \circ \sigma = I$.

The stability of equation (3.1) was studied by several authors in the case where G is an abelian group and $\sigma = -I$. For more information, see for example [12].

First we establish some results which will be instrumental in proving our main results.

In the following lemma, we will present a Hyers-Ulam stability result for Jensen's functional equation:

$$(3.2) \quad f(xy) + f(x\sigma(y)) = 2f(x), \quad x, y \in G.$$

Lemma 3.1. *Let G be an amenable semigroup. Let σ be an homomorphism of G such that $\sigma \circ \sigma = I$ and let $f : G \rightarrow \mathbb{C}$ be a function. Assume that there exists $\delta \geq 0$ such that*

$$(3.3) \quad |f(xy) + f(x\sigma(y)) - 2f(x)| \leq \delta$$

for all $x, y \in G$. Then, there exists a solution $J : G \rightarrow \mathbb{C}$ of Jensen's functional equation (3.2) such that

$$(3.4) \quad |f(x) - J(x) - f(e)| \leq \delta$$

for all $x \in G$.

Proof. Let us denote by $f^e(x) = \frac{f(x)+f(\sigma(x))}{2}$ the even part of f and by $f^o(x) = \frac{f(x)-f(\sigma(x))}{2}$ the odd part of f .



Title Page

Contents



Page 23 of 36

Go Back

Full Screen

Close

By replacing x by $\sigma(x)$ and y by $\sigma(y)$ in (3.3), we get

$$(3.5) \quad |f(\sigma(x)\sigma(y)) + f(\sigma(x)y) - 2f(\sigma(x))| \leq \delta.$$

Now, if we add (subtract) the argument of the inequality (3.3) to (from) inequality (3.5), we deduce that the functions f^e and f^o satisfy the following inequalities

$$(3.6) \quad |f^e(xy) + f^e(x\sigma(y)) - 2f^e(x)| \leq \delta$$

$$(3.7) \quad |f^o(xy) + f^o(x\sigma(y)) - 2f^o(x)| \leq \delta$$

for all $x, y \in G$.

By putting $x = e$ in (3.6), we obtain

$$(3.8) \quad |f^e(y) - f(e)| \leq \frac{\delta}{2}.$$

The inequality (3.7) can be written as follows

$$(3.9) \quad |f^o(yx) - f^o(\sigma(y)x) - 2f^o(y)| \leq \delta.$$

This implies that for fixed $y \in G$, the function $x \mapsto f^o(yx) - f^o(\sigma(y)x)$ is bounded. Since G is amenable, let m_x be an invariant mean on the space of complex bounded functions on G and define the mapping:

$$(3.10) \quad \psi(y) = m_x \{y f^o -_{\sigma(y)} f^o\} \quad \text{for all } y \in G.$$



Consequently from (3.10), we obtain that ψ satisfies the Jensen's functional equation

$$\begin{aligned}(3.11) \quad & \psi(yz) + \psi(y\sigma(z)) \\ &= m_x \{ yz f^o - \sigma(y)\sigma(z) f^o \} + m_x \{ y\sigma(z) f^o - \sigma(y)z f^o \} \\ &= m_x \{ yz f^o - \sigma(y)z f^o \} + m_x \{ y\sigma(z) f^o - \sigma(y)\sigma(z) f^o \} \\ &= m_x \{ z [y f^o - \sigma(y) f^o] \} + m_x \{ \sigma(z) [y f^o - \sigma(y) f^o] \} \\ &= 2\psi(y).\end{aligned}$$

The function $J(y) = \frac{\psi(y)}{2}$ satisfies the Jensen's functional equation (3.2) and the following inequality

$$(3.12) \quad |J(y) - f^o(y)| \leq \frac{1}{2} \sup_{x \in G} |f^o(yx) - f^o(\sigma(y)x) - 2f^o(y)| \leq \frac{\delta}{2}.$$

Finally, we obtain

$$\begin{aligned}(3.13) \quad & |f(y) - J(y) - f(e)| = |f^e(y) + f^o(y) - J(y) - f(e)| \\ & \leq |f^e(y) - f(e)| + |f^o(y) - J(y)| \leq \delta.\end{aligned}$$

This completes the proof of Lemma 3.1. □

By using the proof of the preceding lemma, we get the stability of the Jensen function equation

$$(3.14) \quad f(yx) + f(\sigma(y)x) = 2f(x), \quad x, y \in G.$$

Lemma 3.2. *Let G be an amenable semigroup. Let σ be a homomorphism of G such that $\sigma \circ \sigma = I$ and let $f : G \rightarrow \mathbb{C}$ be a function. Assume that there exists $\delta \geq 0$ such that*

$$(3.15) \quad |f(yx) + f(\sigma(y)x) - 2f(x)| \leq \delta$$

[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 24 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



for all $x, y \in G$. Then, there exists a solution $J : G \rightarrow \mathbb{C}$ of Jensen's functional equation (3.14) such that

$$(3.16) \quad |f(x) - J(x) - f(e)| \leq \delta$$

for all $x \in G$. More precisely, J is given by the formula

$$(3.17) \quad J(y) = m_x \{f_y^o - f_{\sigma(y)}^o\} \text{ for all } y \in G.$$

In the following lemma, we obtain a partial stability theorem for the Pexider's functional equation

$$(3.18) \quad f_1(xy) + f_2(x\sigma(y)) = f_3(x) + f_4(y), \quad x, y \in G$$

that includes the functional equation (1.5) and the Drygas's functional equation:

$$(3.19) \quad f(xy) + f(x\sigma(y)) = 2f(x) + f(y) + f(\sigma(y)), \quad x, y \in G$$

as special cases.

Lemma 3.3. *Let G be an amenable semigroup. Let σ be an automorphism of G such that $\sigma \circ \sigma = I$. If the functions $f_1, f_2, f_3, f_4 : G \rightarrow \mathbb{C}$ satisfy the inequality*

$$(3.20) \quad |f_1(xy) + f_2(x\sigma(y)) - f_3(x) - f_4(y)| \leq \delta$$

for all $x, y \in G$, then there exists a unique function $q : G \rightarrow \mathbb{C}$, a solution of equation (1.4). Also, there exists a solution J_1 , (resp. J_2): $G \rightarrow \mathbb{C}$ of Jensen's functional equation (3.14), (resp. (3.2)) such that,

$$(3.21) \quad |f_3(x) - J_2(x) - q(x) - f_3(e)| \leq 16\delta,$$

$$(3.22) \quad |f_4(x) - J_1(x) - q(x) - f_4(e)| \leq 16\delta,$$

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 25 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)



$$(3.23) \quad \left| f_1^e(x) + f_2^e(x) - q(x) - \frac{1}{2}f_1(e) - \frac{1}{2}f_2(e) \right| \leq 6\delta,$$

$$(3.24) \quad |(f_1^e - f_2^e)(xy) - (f_1^e - f_2^e)(x\sigma(y))| \leq 12\delta,$$

$$\left| f_1^o(x) - \frac{1}{2}J_1(x) - \frac{1}{2}J_2(x) \right| \leq 10\delta$$

and

$$\left| f_2^o(x) - \frac{1}{2}J_2(x) + \frac{1}{2}J_1(x) \right| \leq 10\delta$$

for all $x, y \in G$.

Proof. In the present proof, we follow the computations used in the papers [1], [12], and [23].

For any function $f : G \rightarrow \mathbb{C}$, we define $F(x) = f(x) - f(e)$.

By putting $x = y = e$ in (3.20), we get

$$(3.25) \quad |f_1(e) + f_2(e) - f_3(e) - f_4(e)| \leq \delta.$$

Consequently, if we subtract the inequality (3.20) from the new inequality (3.25), we obtain



$$(3.26) \quad |F_1(xy) + F_2(x\sigma(y)) - F_3(x) - F_4(y)| \leq 2\delta.$$

Now, by replacing x by $\sigma(x)$ and y by $\sigma(y)$ in (3.26) and if we add (subtract) the inequality obtained to (3.26), we deduce that

$$(3.27) \quad |F_1^e(xy) + F_2^e(x\sigma(y)) - F_3^e(x) - F_4^e(y)| \leq 2\delta,$$

Title Page

Contents



Page 26 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



and

$$(3.28) \quad |F_1^o(xy) + F_2^o(x\sigma(y)) - F_3^o(x) - F_4^o(y)| \leq 2\delta$$

for all $x, y \in G$. Hence, if we replace y by e , and x by e respectively in (3.27), we get

$$(3.29) \quad |F_1^e(x) + F_2^e(x) - F_3^e(x)| \leq 2\delta$$

and

$$(3.30) \quad |F_1^e(y) + F_2^e(y) - F_4^e(y)| \leq 2\delta.$$

So, in view of (3.27), (3.29) and (3.30), we obtain

$$(3.31) \quad \begin{aligned} &|F_1^e(xy) + F_2^e(x\sigma(y)) - (F_1^e + F_2^e)(x) - (F_1^e + F_2^e)(y)| \\ &\leq |F_1^e(xy) + F_2^e(x\sigma(y)) - F_3^e(x) - F_4^e(y)| \\ &\quad + |F_1^e(x) + F_2^e(x) - F_3^e(x)| + |F_1^e(y) + F_2^e(y) - F_4^e(y)| \\ &\leq 6\delta. \end{aligned}$$

By replacing y by $\sigma(y)$ in (3.31), we get the following

$$(3.32) \quad |F_1^e(x\sigma(y)) + F_2^e(xy) - (F_1^e + F_2^e)(x) - (F_1^e + F_2^e)(y)| \leq 6\delta.$$

If we add (subtract) the inequality (3.31) to (3.32), we get

$$(3.33) \quad |(F_1^e + F_2^e)(xy) + (F_1^e + F_2^e)(x\sigma(y)) - 2(F_1^e + F_2^e)(x) - 2(F_1^e + F_2^e)(y)| \leq 12\delta,$$

$$(3.34) \quad |(F_1^e - F_2^e)(xy) - (F_1^e - F_2^e)(x\sigma(y))| \leq 12\delta$$

Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 27 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



for all $x, y \in E_1$. Hence, in view of Theorem 2.6, there exists a unique function q , a solution of equation (1.4) such that

$$(3.35) \quad |(F_1^e + F_2^e)(x) - q(x)| \leq 6\delta \text{ for all } x \in G.$$

Consequently, from (3.29), (3.30) and (3.35), we deduce that

$$(3.36) \quad |F_3^e(x) - q(x)| \leq 8\delta$$

and

$$(3.37) \quad |F_4^e(x) - q(x)| \leq 8\delta$$

for all $x \in G$.

On the other hand, from (3.28) we get

$$(3.38) \quad |F_3^o(x) - F_1^o(x) - F_2^o(x)| \leq 2\delta$$

and

$$(3.39) \quad |F_4^o(x) - F_1^o(x) + F_2^o(x)| \leq 2\delta,$$

for all $x \in G$. Hence, we obtain

$$(3.40) \quad |2F_1^o(x) - F_3^o(x) - F_4^o(x)| \leq 4\delta$$

and

$$(3.41) \quad |2F_2^o(x) - F_3^o(x) + F_4^o(x)| \leq 4\delta$$

Title Page

Contents



Page 28 of 36


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

Page 29 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)

for all $x \in G$ and consequently, we have

$$\begin{aligned} (3.42) \quad & |F_3^o(xy) + F_3^o(x\sigma(y)) - 2F_3^o(x)| \\ & \leq |F_3^o(xy) - F_1^o(xy) - F_2^o(xy)| \\ & \quad + |F_3^o(x\sigma(y)) - F_1^o(x\sigma(y)) - F_2^o(x\sigma(y))| \\ & \quad + |F_1^o(xy) + F_2^o(x\sigma(y)) - F_3^o(x) - F_4^o(y)| \\ & \quad + |F_1^o(x\sigma(y)) + F_2^o(xy) - F_3^o(x) - F_4^o(\sigma(y))| \\ & \leq 8\delta \end{aligned}$$

and

$$\begin{aligned} (3.43) \quad & |F_4^o(yx) + F_4^o(\sigma(y)x) - 2F_4^o(x)| \\ & \leq |F_4^o(yx) - F_1^o(yx) + F_2^o(yx)| \\ & \quad + |F_4^o(\sigma(y)x) - F_1^o(\sigma(y)x) + F_2^o(\sigma(y)x)| \\ & \quad + |F_1^o(yx) + F_2^o(y\sigma(x)) - F_3^o(y) - F_4^o(x)| \\ & \quad + |F_1^o(\sigma(y)x) + F_2^o(\sigma(y)\sigma(x)) - F_3^o(\sigma(y)) - F_4^o(x)| \\ & \leq 8\delta \end{aligned}$$

for all $x, y \in G$.

Now, from Lemma 3.1 and Lemma 3.2 there exist two solutions of Jensen's functional equation (3.14) and (3.2), $J_1, J_2 : G \rightarrow \mathbb{C}$ such that

$$(3.44) \quad |F_4^o(x) - J_1(x)| \leq 8\delta.$$

and

$$(3.45) \quad |F_3^o(x) - J_2(x)| \leq 8\delta$$

for all $x \in G$. Now, by small computations, we obtain the rest of the proof. \square



[Title Page](#)

[Contents](#)

[⏪](#) [⏩](#)

[◀](#) [▶](#)

Page 30 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)

By using the previous lemmas, we may deduce our main result.

Theorem 3.4. *Let G be an amenable semigroup. Let σ be an automorphism of G such that $\sigma \circ \sigma = I$. If the functions $f, g : G \rightarrow \mathbb{C}$ satisfy the inequality*

$$(3.46) \quad |f(xy) + g(x\sigma(y)) - f(x) - g(y)| \leq \delta,$$

for all $x, y \in G$, then there exists a unique function $q : G \rightarrow \mathbb{C}$, a solution of equation (1.4). Also, there exist two solutions J_1 , (resp. J_2): $G \rightarrow \mathbb{C}$ of Jensen's functional equation (3.14), (resp. (3.2)) such that

$$(3.47) \quad |f(x) - J_2(x) - q(x) - f(e)| \leq 16\delta$$

and

$$(3.48) \quad |g(x) - J_1(x) - q(x) - g(e)| \leq 16\delta,$$

for all $x \in G$.

The stability of the Drygas's functional equation (3.19) is a consequence of the preceding theorem.

Theorem 3.5. *Let G be an amenable semigroup. Let σ be an automorphism of G such that $\sigma \circ \sigma = I$. Let the function $f : G \rightarrow \mathbb{C}$ satisfy the inequality*

$$(3.49) \quad |f(xy) + f(x\sigma(y)) - 2f(x) - f(y) - f(\sigma(y))| \leq \delta,$$

for all $x, y \in G$. Then there exists a unique function $q : G \rightarrow \mathbb{C}$, a solution of equation (1.4), and a solution $J : G \rightarrow \mathbb{C}$ of Jensen's functional equation (3.2) such that

$$(3.50) \quad |f(x) - J(x) - q(x) - f(e)| \leq 16\delta$$

for all $x \in G$.



Corollary 3.6. *Let G be an amenable semigroup with a unity element. Let σ be an automorphism of G such that $\sigma \circ \sigma = I$. If the functions $f_1, f_2, f_3, f_4 : G \rightarrow \mathbb{C}$ satisfy the functional equation*

$$(3.51) \quad f_1(xy) + f_2(x\sigma(y)) = f_3(x) + f_4(y)$$

for all $x, y \in G$, then there exists a quadratic function $q : G \rightarrow \mathbb{C}$. There also exists a function $\nu : G \rightarrow \mathbb{C}$, a solution of

$$(3.52) \quad \nu(xy) = \nu(x\sigma(y)), \quad x, y \in G.$$

In addition, there exist $\alpha, \beta, \gamma, \delta \in \mathbb{C}$ and two solutions J_1 , (resp. J_2) of Jensen's equation (3.14) (resp. (3.2)) such that

$$(3.53) \quad f_1(x) = \frac{1}{2}J_1(x) + \frac{1}{2}J_2(x) + \frac{1}{2}\nu(x) + \frac{1}{2}q(x) + \alpha,$$

$$(3.54) \quad f_2(x) = -\frac{1}{2}J_1(x) + \frac{1}{2}J_2(x) - \frac{1}{2}\nu(x) + \frac{1}{2}q(x) + \beta,$$

$$(3.55) \quad f_3(x) = J_2(x) + q(x) + \gamma$$

and

$$(3.56) \quad f_4(x) = J_1(x) + q(x) + \delta$$

for all $x \in G$.

From Lemma 3.3, we can deduce the results obtained in [12].

Title Page

Contents



Page 31 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Corollary 3.7. Let G be a vector space. If the functions $f_1, f_2, f_3, f_4 : G \rightarrow \mathbb{C}$ satisfy the inequality

$$(3.57) \quad |f_1(x+y) + f_2(x-y) - f_3(x) - f_4(y)| \leq \delta$$

for all $x, y \in G$, then there exists a unique function $q : G \rightarrow \mathbb{C}$, a solution of equation (1.2). Also, $\alpha \in \mathbb{C}$ exists, and there are exactly two additive functions $a_1, a_2 : G \rightarrow \mathbb{C}$ such that

$$(3.58) \quad \left| f_1(x) - \frac{1}{2}a_1(x) - \frac{1}{2}a_2(x) - \frac{1}{2}q(x) - f_1(0) - \alpha \right| \leq 19\delta,$$

$$(3.59) \quad \left| f_2(x) + \frac{1}{2}a_1(x) - \frac{1}{2}a_2(x) - \frac{1}{2}q(x) - f_2(0) + \alpha \right| \leq 19\delta,$$

$$(3.60) \quad |f_3(x) - a_2(x) - q(x) - f_3(0)| \leq 16\delta$$

and

$$(3.61) \quad |f_4(x) - a_1(x) - q(x) - f_4(0)| \leq 16\delta$$

for all $x \in G$.

The following corollary follows from Lemma 3.3. This result is well known in the commutative case, see for example [15].

Corollary 3.8. Let G be an amenable semigroup. If the functions $f_1, f_2, f_3 : G \rightarrow \mathbb{C}$ satisfy the inequality

$$(3.62) \quad |f_1(xy) - f_2(x) - f_3(y)| \leq \delta$$

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)


[◀](#) [▶](#)

Page 32 of 36

[Go Back](#)

[Full Screen](#)

[Close](#)



for all $x, y \in G$, then there exists a unique additive function $a : G \rightarrow \mathbb{C}$ such that

$$(3.63) \quad |f_1(x) - a(x) - f_1(e)| \leq 38\delta,$$

$$(3.64) \quad |f_2(x) - a(x) - f_2(e)| \leq 16\delta$$

and

$$(3.65) \quad |f_3(x) - a(x) - f_3(e)| \leq 16\delta$$

for all $x \in G$.

Hyers-Ulam Stability

Bouikhalene Belaid, Elqorachi Elhoucien

and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

Title Page

Contents



Page 33 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



References

- [1] B. BOUIKHALENE, E. ELQORACHI AND Th. M. RASSIAS, On the Hyers-Ulam stability of approximately Pexider mappings, *Math. Inequalities. and Appl.*, (accepted for publication).
- [2] B. BOUIKHALENE, E. ELQORACHI AND Th. M. RASSIAS, On the generalized Hyers-Ulam stability of the quadratic functional equation with a general involution, *Nonlinear Funct. Anal. Appl.*, (accepted for publication).
- [3] P.W. CHOLEWA, Remarks on the stability of functional equations, *Aequationes Math.*, **27** (1984), 76–86.
- [4] S. CZERWIK, On the stability of the quadratic mapping in normed spaces, *Abh. Math. Sem. Univ. Hamburg*, **62** (1992), 59–64.
- [5] G.L. FORTI, Hyers-Ulam stability of functional equations in several variables, *Aequationes Math.*, **50** (1995), 143–190.
- [6] Z. GAJDA, On stability of additive mappings, *Internat. J. Math. Sci.*, **14** (1991), 431–434.
- [7] F.P. GREENLEAF, *Invariant Means on Topological Groups*, [Van Nostrand Mathematical Studies V. 16], Van Nostrand, New York-Toronto-London-Melbourne, 1969.
- [8] D.H. HYERS, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci. U.S.A.* **27** (1941), 222–224.
- [9] D.H. HYERS AND Th.M. RASSIAS, Approximate homomorphisms, *Aequationes Math.*, **44** (1992), 125–153.



Title Page

Contents



Page 34 of 36

Go Back

Full Screen

Close



- [10] D.H. HYERS, G. I. ISAC AND Th.M. RASSIAS, *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [11] S.-M. JUNG, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press Inc., Palm Harbor, Florida, 2003.
- [12] S.-M. JUNG, Hyers-Ulam-Rassias stability of Jensen's equation and its application, *Proc. Amer. Math. Soc.* **126** (1998), 3137–3143.
- [13] S.-M. JUNG AND P.K. SAHOO, Hyers-Ulam stability of the quadratic equation of Pexider type, *J. Korean Math. Soc.*, **38**(3) (2001), 645–656.
- [14] S.-M. JUNG AND P.K. SAHOO, Stability of a functional equation of Drygas, *Aequationes Math.*, **64**(3) (2002), 263–273.
- [15] Y.-H. LEE AND K.-W. JUN, A note on the Hyers-Ulam-Rassias stability of Pexider equation, *J. Korean Math. Soc.*, **37**(1) (2000), 111–124.
- [16] Th.M. RASSIAS, On the stability of linear mapping in Banach spaces, *Proc. Amer. Math. Soc.*, **72** (1978), 297–300.
- [17] Th.M. RASSIAS AND J. TABOR, *Stability of Mappings of Hyers-Ulam Type*, Hadronic Press Inc., Palm Harbor, Florida, 1994.
- [18] F. SKOF, Local properties and approximations of operators, *Rend. Sem. Math. Fis. Milano*, **53** (1983), 113–129.
- [19] H. STETKÆR, Functional equations on abelian groups with involution, *Aequationes Math.*, **54** (1997), 144–172.
- [20] L. SZÉKELYHIDI, Note on a stability theorem, *Canad. Math. Bull.*, **25** (1982), 500–501.

Hyers-Ulam Stability

Bouikhalene Belaid, Elqorachi Elhoucien
and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

Title Page

Contents



Page 35 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [21] S.M. ULAM, *A Collection of Mathematical Problems*, Interscience Publ. New York, 1961. *Problems in Modern Mathematics*, Wiley, New York 1964.
- [22] *Contributions to the Theory of Functional Equations*, PhD Thesis, University of Waterloo, Waterloo, Ontario, Canada, 2006.
- [23] D. YANG, D. YANG, Remarks on the stabilities of Drygas' equation and Pexider-quadratic equation, *Aequationes Math.*, **68** (2004), 108–116.

Hyers-Ulam Stability

Bouikhalene Belaid, Elqorachi Elhoucien
and Redouani Ahmed

vol. 8, iss. 2, art. 47, 2007

Title Page

Contents



Page 36 of 36

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756