

# Journal of Inequalities in Pure and Applied Mathematics


## SOME INEQUALITIES EXHIBITING CERTAIN PROPERTIES OF SOME SUBCLASSES OF MULTIVALENTLY ANALYTIC FUNCTIONS

HÜSEYİN IRMAK AND R.K. RAINA

Department Of Mathematics  
Faculty Of Education  
Baskent University  
Tr-06530, Baglica Campus  
Baglica, Etimesgut- Ankara, Turkey.  
EMail: [hisimya@baskent.edu.tr](mailto:hisimya@baskent.edu.tr)

Department Of Mathematics  
College Of Technology and Engineering  
M.P. University Of Agriculture And Technology  
Udaipur- 31300, India  
EMail: [rainark\\_7@hotmail.com](mailto:rainark_7@hotmail.com)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
088-04



---

volume 6, issue 1, article 9,  
2005.

*Received 01 May, 2004;  
accepted 27 October, 2004.*

*Communicated by: N.K. Govil*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)



## Abstract

This paper introduces a new subclass and investigates the sufficiency conditions for a function to belong to this subclass. Certain types of inequalities are also studied exhibiting the well-known geometric properties of multivalently analytic functions in the unit disk. Several interesting consequences of the main results are also mentioned.

*2000 Mathematics Subject Classification:* 30C45, 30C50, 30A10.


*Key words:* Open unit disk, analytic, multivalently analytic functions, multivalently starlike, multivalently convex, rational and complex inequalities, rational functions with complex variable and Jack's Lemma.

This paper is funded, in part, by TÜBİTAK (The Scientific and Technical Research Council of Turkey) and Başkent University (Ankara, Turkey). The first author would like to acknowledge and express his thanks to Professor (Dr.) Mehmet Haberal, Rector of Başkent University, who generously supports scientific researches in all aspects.

The second author's work was supported by All India Council of Technical Education (Govt. of India), New Delhi.

## Contents

1	Introduction and Definition .....	3
2	Main Results .....	5
3	Some Consequences of Main Results .....	11
	References	



### Some Inequalities Exhibiting Certain Properties Of Some Subclasses Of Multivalently Analytic Functions

Hüseyin İrmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

# 1. Introduction and Definition

Let  $\mathcal{T}(p)$  denote the class of functions  $f(z)$  of the form:

$$(1.1) \quad f(z) = z^p + \sum_{k=p+1}^{\infty} a_k z^k \quad (p \in \mathcal{N} = \{1, 2, 3, \dots\}),$$

which are *analytic* and *multivalent* in the open disk  $\mathcal{U} = \{z : z \in \mathcal{C} \text{ and } |z| < 1\}$ . A function  $f(z)$  belonging to  $\mathcal{T}(p)$  is said to be *multivalently starlike order*  $\alpha$  in  $\mathcal{U}$  if it satisfies the inequality:


$$(1.2) \quad \Re \left\{ \frac{z f'(z)}{f(z)} \right\} > \alpha \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$$

and, a function  $f(z) \in \mathcal{T}(p)$  is said to be *multivalently convex of order*  $\alpha$  in  $\mathcal{U}$  if it satisfies the inequality:

$$(1.3) \quad \Re \left\{ 1 + \frac{z f''(z)}{f'(z)} \right\} > \alpha \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}).$$

For the aforementioned definitions, one may refer to [1] (see also [11]). Further, a function  $f(z) \in \mathcal{T}(p)$  is said to be in the subclass  $\mathcal{TSK}_{\lambda}^{\delta}(p; \alpha)$  if it satisfies the inequality:

$$(1.4) \quad \Re \left\{ \left( \frac{z f'(z) + \lambda z^2 f''(z)}{(1 - \lambda) f(z) + \lambda z f'(z)} \right)^{\delta} \right\} > \alpha, \\ (z \in \mathcal{U}; \delta \neq 0; 0 \leq \lambda \leq 1; 0 \leq \alpha < p; p \in \mathcal{N}).$$



Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions

Hüseyin Irmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

Here, and throughout this paper, the value of expressions like

$$\left( \frac{zf'(z) + \lambda z^2 f''(z)}{(1 - \lambda)f(z) + \lambda z f'(z)} \right)^\delta,$$

is considered to be its principal value. We mention below some of the subclasses of the functions  $\mathcal{T}(p)$  from the families of functions  $\mathcal{TSK}_\lambda^\delta(p; \alpha)$  (defined above). Indeed, we have

$$(1.5) \quad \mathcal{TS}^\delta(p; \alpha) \equiv \mathcal{TSK}_0^\delta(p; \alpha) \quad (\delta \neq 0, 0 \leq \alpha < p, p \in \mathcal{N}),$$

$$(1.6) \quad \mathcal{TK}^\delta(p; \alpha) \equiv \mathcal{TSK}_1^\delta(p; \alpha) \quad (\delta \neq 0, 0 \leq \alpha < p, p \in \mathcal{N}),$$

$$(1.7) \quad \mathcal{T}_\lambda(p; \alpha) \equiv \mathcal{TSK}_\lambda^1(p; \alpha) \quad (0 \leq \lambda \leq 1, 0 \leq \alpha < p, p \in \mathcal{N}) \text{ (see [5]).}$$

The important subclasses in Geometric Function Theory such as multivalently starlike functions  $\mathcal{S}_p(\alpha)$  of order  $\alpha$  ( $0 \leq \alpha < p; p \in \mathcal{N}$ ) in  $\mathcal{U}$ , multivalently convex functions  $\mathcal{K}_p(\alpha)$  of order  $\alpha$  ( $0 \leq \alpha < p; p \in \mathcal{N}$ ) in  $\mathcal{U}$ , multivalently starlike functions  $\mathcal{S}_p$  in  $\mathcal{U}$ , multivalently convex functions  $\mathcal{K}_p$  in  $\mathcal{U}$ , starlike functions  $\mathcal{S}(\alpha)$  of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathcal{U}$ , convex functions  $\mathcal{K}(\alpha)$  of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathcal{U}$ , starlike functions  $\mathcal{S}$  in  $\mathcal{U}$  and convex functions  $\mathcal{K}$  in  $\mathcal{U}$ , are seen to be easily identifiable with the aforementioned classes ([1], [5] and [11]).

By introducing a subclass  $\mathcal{TSK}_\lambda^\delta(p; \alpha)$  of functions  $f(z) \in \mathcal{T}(p)$  satisfying the inequality (1.4), our motive in this paper is to obtain sufficient conditions for a function to belong to the above subclass. The other results investigated include certain inequalities for multivalent functions depicting the properties of starlikeness, close-to-convexity and convexity in the open unit disk. Several corollaries are deduced as worthwhile consequences of our main results.



**Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalent  
Analytic Functions**

Hüseyin İrmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

## 2. Main Results

Before stating and proving our main results, we require the following assertion (popularly known as Jack's Lemma).

**Lemma 2.1 ([7]).** *Let the function  $w(z)$  be non-constant and regular in the unit disc  $\mathcal{U}$  such that  $w(0) = 0$ . If  $|w(z)|$  attains its maximum value on the circle  $|z| = r < 1$  at the point  $z_0$ , then*

$$(2.1) \quad z_0 w'(z_0) = c w(z_0) \quad (c \geq 1).$$

We begin now to prove the following:

**Theorem 2.2.** *Let  $\delta \in \mathbb{R} \setminus \{0\}$ ,  $0 \leq \alpha < p$ ,  $p \in \mathcal{N}$  and  $f(z) \in \mathcal{T}(p)$ . If a function  $F(z)$  defined by*


$$(2.2) \quad F(z) = (1 - \lambda)f(z) + \lambda z f'(z) \quad (0 \leq \lambda \leq 1),$$

*satisfies the inequality:*

$$(2.3) \quad \Re \left\{ \frac{1 + z \left( \frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right)}{1 - p^\delta \left( \frac{z F'(z)}{F(z)} \right)^{-\delta}} \right\} \left\{ \begin{array}{l} < \frac{1}{\delta} \text{ when } \delta > 0 \\ > \frac{1}{\delta} \text{ when } \delta < 0 \end{array} \right\} \quad (z \in \mathcal{U}),$$

*then  $f(z) \in \mathcal{TSK}_\lambda^\delta(p; \beta)$ , where  $\beta = p^\delta - (p - \alpha)^\delta$ .*

*Proof.* Let  $f(z) \in \mathcal{T}(p)$  and  $F(z)$  be defined by (2.2). From (1.1) and (2.2), we



Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions

Hüseyin Irmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 5 of 16

have

$$(2.4) \quad \frac{zF'(z)}{F(z)} = \frac{zf'(z) + \lambda z^2 f''(z)}{(1-\lambda)f(z) + \lambda zf'(z)}$$

$$= \frac{p + \sum_{k=p+1}^{\infty} \frac{k[1+\lambda(k-1)]}{1+\lambda(p-1)} a_k z^{k-p}}{1 + \sum_{k=p+1}^{\infty} \frac{1+\lambda(k-1)}{1+\lambda(p-1)} a_k z^{k-p}}.$$

$(z \in \mathcal{U}; 0 \leq \lambda \leq 1; p \in \mathcal{N})$

Now, define a function  $w(z)$  by

$$(2.5) \quad \left( \frac{zF'(z)}{F(z)} \right)^{\delta} - p^{\delta} = (p-\alpha)^{\delta} w(z), \quad (z \in \mathcal{U}; \delta \neq 0; 0 \leq \alpha < p; p \in \mathcal{N}),$$

then the function  $w(z)$  is analytic in  $\mathcal{U}$  and  $w(0) = 0$ . Differentiation of (2.5) gives


$$(2.6) \quad 1 + \frac{zF''(z)}{F'(z)} - \frac{zF'(z)}{F(z)} = \left( \frac{(p-\alpha)^{\delta}}{p^{\delta} + (p-\alpha)^{\delta} w(z)} \right) \frac{zw'(z)}{\delta}.$$

Hence, (2.5) and (2.6) yields

$$(2.7) \quad \frac{1 + z \left( \frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right)}{1 - p^{\delta} \left( \frac{zF'(z)}{F(z)} \right)^{-\delta}} = \frac{zw'(z)}{\delta w(z)}.$$

We claim that  $|w(z)| < 1$  in  $\mathcal{U}$ . For otherwise (by Jack's Lemma), there exists a point  $z_0 \in \mathcal{U}$  such that

$$z_0 w'(z_0) = c w(z_0), \quad \text{where } |w(z_0)| = 1 \quad (c \geq 1).$$



**Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions**

Hüseyin İrmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

Therefore, (2.7) yields

$$(2.8) \quad \Re \left\{ \frac{1 + z \left( \frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right)}{1 - p^\delta \left( \frac{zF'(z)}{F(z)} \right)^{-\delta}} \right\} \Bigg|_{z=z_0} = \frac{1}{\delta} \Re \left\{ \frac{z_0 w'(z_0)}{w(z_0)} \right\}$$

$$= \frac{c}{\delta} \begin{cases} \geq \frac{1}{\delta} & \text{when } \delta > 0 \\ \leq \frac{1}{\delta} & \text{when } \delta < 0, \end{cases}$$

which contradicts our assumption (2.3). Therefore,  $|w(z)| < 1$  holds true for all  $z \in \mathcal{U}$ , and we conclude from (2.5) that

$$(2.9) \quad \left| \left( \frac{zF'(z)}{F(z)} \right)^\delta - p^\delta \right| = (p - \alpha)^\delta |w(z)| < (p - \alpha)^\delta,$$


which evidently implies that

$$(2.10) \quad \Re \left\{ \left( \frac{zF'(z)}{F(z)} \right)^\delta \right\} > p^\delta - (p - \alpha)^\delta,$$

and hence  $f(z) \in \mathcal{TSK}_\lambda^\delta(p; \alpha)$ . □

**Theorem 2.3.** Let  $\delta \in \mathbb{R} \setminus \{0\}$ ;  $0 \leq \alpha < p$ ;  $n, m, p \in \mathbb{N}$ ;  $q = n - m$ ;  $f(z) \in \mathcal{T}(n)$  and  $g(z) \in \mathcal{T}(m)$ . If  $f(z)$  satisfies the inequality:

$$(2.11) \quad \Re \left( \frac{zf'(z)}{f(z)} \right) \begin{cases} < q + \alpha + \frac{1}{2\delta} & \text{when } \delta > 0 \text{ and } g(z) \in \mathcal{S}_m(\alpha) \\ > q + \alpha + \frac{1}{2\delta} & \text{when } \delta < 0 \text{ and } g(z) \notin \mathcal{S}_m(\alpha), \end{cases}$$



Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions

Hüseyin İrmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

then

$$(2.12) \quad \Re \left\{ \left( z^{-q} \frac{f(z)}{g(z)} \right)^\delta \right\} > 0,$$

where the value of  $\left( z^{-q} \frac{f(z)}{g(z)} \right)^\delta$  is taken to be its principle value.

*Proof.* Let  $f(z) \in \mathcal{T}(n)$  and  $g(z) \in \mathcal{T}(m)$  with  $n - m \in \mathcal{N}$ . Since

$$\frac{f(z)}{g(z)} = z^q + c_1 z^{q+1} + c_2 z^{q+2} + \dots \in \mathcal{T}(q) \quad (q = n - m \in \mathcal{N}),$$

we define  $w(z)$  by

$$(2.13) \quad \left( z^{-q} \frac{f(z)}{g(z)} \right)^\delta = 1 + w(z) \quad (z \in \mathcal{U}; \delta \neq 0).$$


It is clear that the function  $w(z)$  is an analytic function in  $\mathcal{U}$  and  $w(0) = 0$ .

Differentiating (2.13), we have

$$(2.14) \quad \frac{z f'(z)}{f(z)} = q + \frac{z w'(z)}{\delta (1 + w(z))} + \frac{z g'(z)}{g(z)}.$$

If we suppose that there exists a point  $z_0 \in \mathcal{U}$  such that  $z_0 w'(z_0) = c w(z_0)$  where  $|w(z_0)| = 1$  ( $c \geq 1$ ), i.e.  $w(z_0) = e^{i\theta}$  ( $\theta \in [0, 2\pi) - \{\pi\}$ ), then

$$(2.15) \quad \begin{aligned} \Re \left\{ \frac{z_0 f'(z_0)}{f(z_0)} \right\} &= q + \frac{1}{\delta} \Re \left\{ \frac{z_0 w'(z_0)}{1 + w(z_0)} + \frac{\delta z_0 g'(z_0)}{g(z_0)} \right\} \\ &= q + \frac{1}{\delta} \Re \left\{ \frac{c e^{i\theta}}{1 + e^{i\theta}} \right\} + \Re \left\{ \frac{z_0 g'(z_0)}{g(z_0)} \right\}. \end{aligned}$$



**Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions**

Hüseyin İrmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16



From (2.15) it follows that

$$(2.16) \quad \Re \left\{ \frac{z_0 f'(z_0)}{f(z_0)} \right\} \geq q + \alpha + \frac{1}{2\delta} \quad (\delta > 0),$$

provided that

$$\Re \left\{ \frac{z_0 g'(z_0)}{g(z_0)} \right\} > \alpha,$$

and

$$(2.17) \quad \Re \left\{ \frac{z_0 f'(z_0)}{f(z_0)} \right\} \leq q + \alpha + \frac{1}{2\delta} \quad (\delta < 0),$$

provided that

$$\Re \left\{ \frac{z_0 g'(z_0)}{g(z_0)} \right\} \leq \alpha.$$

But the inequalities in (2.16) and (2.17) contradict the inequalities in (2.11).


Hence  $|w(z)| < 1$ , for all  $z \in \mathcal{U}$ , and therefore (2.13) yields

$$(2.18) \quad \left| \left( z^{-q} \frac{f(z)}{g(z)} \right)^\delta - 1 \right| = |w(z)| < 1,$$

which evidently implies (2.12), and this completes the proof of Theorem 2.3.  $\square$

**Theorem 2.4.** Let  $\delta \in \mathbb{R} \setminus \{0\}$ ;  $0 \leq \alpha < p$ ;  $n, m, p \in \mathcal{N}$ ;  $q = n - m$ ;  $f(z) \in \mathcal{T}(n)$ , and  $g(z) \in \mathcal{T}(m)$ . If  $f(z)$  satisfies the inequality:

$$(2.19) \quad \Re \left( 1 + \frac{z f''(z)}{f'(z)} \right) \begin{cases} < q + \alpha + \frac{1}{2\delta} \text{ when } \delta > 0 \text{ and } g(z) \in \mathcal{K}_m(\alpha) \\ > q + \alpha + \frac{1}{2\delta} \text{ when } \delta < 0 \text{ and } g(z) \notin \mathcal{K}_m(\alpha), \end{cases}$$



Some Inequalities Exhibiting Certain Properties Of Some Subclasses Of Multivalently Analytic Functions

Hüseyin İrmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 9 of 16

then

$$(2.20) \quad \Re \left\{ \left( z^{-q} \frac{m f'(z)}{n g'(z)} \right)^\delta \right\} > 0,$$

where the value of  $\left( z^{-q} \frac{m f'(z)}{n g'(z)} \right)^\delta$  is taken its principle value.


*Proof.* Let  $f(z) \in \mathcal{T}(n)$  and  $g(z) \in \mathcal{T}(m)$  with  $n - m \in \mathcal{N}$ . Since

$$\frac{m f'(z)}{n g'(z)} = z^q + k_1 z^{q+1} + k_2 z^{q+2} + \dots \in \mathcal{T}(q) \quad (q = n - m \in \mathcal{N}),$$

and if we define  $w(z)$  by

$$(2.21) \quad \left( z^{-q} \frac{m f'(z)}{n g'(z)} \right)^\delta = 1 + w(z) \quad (z \in \mathcal{U}),$$

then by appealing to the same technique as in the proof of Theorem 2.3, we arrive at the assertion (2.20) of Theorem 2.4 under the conditions stated with (2.19).  $\square$




**Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalent  
Analytic Functions**

Hüseyin Irmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

### 3. Some Consequences of Main Results

Among the various interesting and important consequences of Theorems 2.2 – 2.4, we mention now some of the corollaries relating to the classes  $\mathcal{T}_\lambda(p; \alpha)$ ,  $\mathcal{T}_\lambda(\alpha)$ ,  $\mathcal{S}_p(\alpha)$ ,  $\mathcal{K}_p(\alpha)$ ,  $\mathcal{S}_p$ ,  $\mathcal{K}_p$ ,  $\mathcal{S}(\alpha)$ ,  $\mathcal{K}(\alpha)$ , which are easily deducible from the main results. Inequalities concerning analytic and multivalent functions were also studied in [2] – [6], and in [8] – [10].

Firstly, if we take  $\delta = 1$ , then Theorem 2.2 by virtue of (1.7) gives the following:

**Corollary 3.1.** *Let a function  $F(z)$  defined by (2.2) satisfy the condition:*

$$(3.1) \quad \Re \left\{ \frac{1 + z \left( \frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right)}{1 - p \left( \frac{F(z)}{zF'(z)} \right)} \right\} < 1,$$

$$(z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}; f(z) \in \mathcal{T}(p))$$

then  $f(z) \in \mathcal{T}_\lambda(p; \alpha)$ .

Next, if we take  $\delta - 1 = \lambda = 0$  in Theorem 2.2, so that  $F(z) = f(z)$ , then we get

**Corollary 3.2.** *If  $F(z) = f(z)$  satisfies the condition in (3.1), then  $f(z) \in \mathcal{S}_p(\alpha)$ , i.e.  $f(z)$  is  $p$ -valent starlike of order  $\alpha$  ( $0 \leq \alpha < p; p \in \mathcal{N}$ ) in  $\mathcal{U}$ .*

If we take  $\delta = \lambda = 1$  in Theorem 2.2, so that  $F(z) = z f'(z)$ , then we obtain the following:



Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions

Hüseyin İrmak and R.K. Raina

Title Page

Contents




Go Back

Close

Quit

Page 11 of 16



Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

**Corollary 3.3.** If  $f(z)$  satisfies the condition

$$(3.2) \quad \Re \left\{ \frac{1 + z \left( \frac{(zf'(z))''}{(zf'(z))'} - \frac{(zf'(z))'}{zf'(z)} \right)}{1 - p \left( \frac{zf'(z)}{(zf'(z))'} \right)} \right\} < 1 \quad (z \in \mathcal{U}; 0 \leq \alpha < p; p \in \mathcal{N}),$$

then  $f(z) \in \mathcal{K}_p(\alpha)$ , that is  $f(z)$  is  $p$ -valent convex of the order  $\alpha$  ( $0 \leq \alpha < p; p \in \mathcal{N}$ ) in  $\mathcal{U}$ .

For  $p = 1$  in Corollaries 3.1 – 3.3 give the following:

**Corollary 3.4.** Let a function  $F(z)$  defined by (2.2) satisfy the condition

$$(3.3) \quad \Re \left\{ \frac{1 + z \left( \frac{F''(z)}{F'(z)} - \frac{F'(z)}{F(z)} \right)}{1 - \frac{F(z)}{zF'(z)}} \right\} < 1,$$

$(z \in \mathcal{U}; 0 \leq \alpha < 1; f(z) \in \mathcal{T})$

then  $f(z) \in \mathcal{T}_\lambda(\alpha)$ .

**Corollary 3.5.** If  $F(z) = f(z)$  satisfies the condition (3.3), then  $f(z) \in \mathcal{S}(\alpha)$ , i.e.  $f(z)$  is starlike of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathcal{U}$ .

**Corollary 3.6.** If  $f(z)$  satisfies the condition

$$(3.4) \quad \Re \left\{ \frac{1 + z \left( \frac{(zf'(z))''}{(zf'(z))'} - \frac{(zf'(z))'}{zf'(z)} \right)}{1 - \frac{zf'(z)}{(zf'(z))'}} \right\} < 1 \quad (z \in \mathcal{U}; 0 \leq \alpha < 1),$$

then  $f(z) \in \mathcal{K}(\alpha)$ , i.e.,  $f(z)$  is convex of order  $\alpha$  ( $0 \leq \alpha < 1$ ) in  $\mathcal{U}$ .

Let us take  $\delta = 1$  in Theorems 2.3 and 2.4, then we get the following:

**Corollary 3.7.** *Let  $z \in \mathcal{U}$ ;  $0 \leq \alpha < p$ ;  $n, m, p \in \mathcal{N}$ ;  $f(z) \in \mathcal{T}(n)$  and a function  $g(z) \in \mathcal{T}(m)$  belong to the class  $\mathcal{S}_m(\alpha)$  with  $q = n - m \in \mathcal{N}$ . If  $f(z)$  satisfies the inequality:*

$$(3.5) \quad \Re \left\{ \frac{z f'(z)}{f(z)} \right\} < q + \alpha + \frac{1}{2},$$

then

$$(3.6) \quad \Re \left\{ z^{-q} \frac{f(z)}{g(z)} \right\} > 0.$$

**Corollary 3.8.** *Let  $z \in \mathcal{U}$ ;  $0 \leq \alpha < p$ ;  $n, m, p \in \mathcal{N}$ ;  $f(z) \in \mathcal{T}(n)$  and a function  $g(z)$  in  $\mathcal{T}(m)$  belong to the class  $\mathcal{K}_m(\alpha)$  with  $q = n - m \in \mathcal{N}$ . If  $f(z)$  satisfies the inequality:*

$$(3.7) \quad \Re \left\{ 1 + \frac{z f''(z)}{f'(z)} \right\} < q + \alpha + \frac{1}{2},$$

then

$$(3.8) \quad \Re \left\{ z^{-q} \frac{m f'(z)}{n g'(z)} \right\} > 0.$$

Lastly, setting  $\delta = -1$  in Theorems 2.3 and 2.4, we obtain the following:



**Some Inequalities Exhibiting Certain Properties Of Some Subclasses Of Multivalently Analytic Functions**

Hüseyin Irmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 13 of 16

**Corollary 3.9.** Let  $z \in \mathcal{U}$ ;  $0 \leq \alpha < p$ ;  $n, m, p \in \mathcal{N}$ ;  $f(z) \in \mathcal{T}(n)$  and suppose a function  $g(z) \in \mathcal{T}(m)$  does not belong to the class  $\mathcal{S}_m(\alpha)$  with  $q = n - m \in \mathcal{N}$ . If  $f(z)$  satisfies the inequality:

$$(3.9) \quad \Re \left\{ \frac{z f'(z)}{f(z)} \right\} > q + \alpha - \frac{1}{2},$$

then

$$(3.10) \quad \Re \left\{ z^q \frac{g(z)}{f(z)} \right\} > 0.$$

**Corollary 3.10.** Let  $z \in \mathcal{U}$ ;  $0 \leq \alpha < p$ ;  $n, m, p \in \mathcal{N}$ ;  $f(z) \in \mathcal{T}(n)$  and suppose a function  $g(z) \in \mathcal{T}(m)$  does not belong to the class  $\mathcal{K}_m(\alpha)$  with  $q = n - m \in \mathcal{N}$ . If  $f(z)$  satisfies the inequality:

$$(3.11) \quad \Re \left\{ 1 + \frac{z f''(z)}{f'(z)} \right\} > q + \alpha - \frac{1}{2},$$

then

$$(3.12) \quad \Re \left\{ z^q \frac{n g'(z)}{m f'(z)} \right\} > 0.$$



Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions

Hüseyin Irmak and R.K. Raina

Title Page

Contents



Go Back

Close

Quit

Page 14 of 16

## References

- [1] P.L. DUREN, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften 259, Springer-Verlag, New York Berlin, Heidelberg, and Tokyo, 1983.
- [2] H. IRMAK AND S. OWA, Certain inequalities involving analytic and univalent functions, *Far East J. Math. Sci.*, **10** (2003), 353–358.
- [3] H. IRMAK AND S. OWA, Certain inequalities for multivalent and meromorphically multivalent starlike functions, *Bull. Inst. Math. Acad. Sinica*, **31** (2003), 11–21.
- [4] H. IRMAK, R.K. RAINA AND S. OWA, A certain for multivalent starlike and meromorphically multivalent starlike functions, *Inter. J. Appl. Math.*, **12** (2003), 93–98.
- [5] H. IRMAK AND R.K. RAINA, The starlikeness and convexity of multivalent function involving certain inequalities, *Rev. Mat. Complut.*, **16** (2003), 391–398.
- [6] H. IRMAK, R.K. RAINA AND S. OWA, Certain results involving inequalities on analytic and univalent functions, *Far East J. Math. Sci.*, **10** (2003), 359–366.
- [7] I.S. JACK, Functions starlike and convex of the order  $\alpha$ , *J. London Math. Soc.*, **3** (1971), 469–474.
- [8] S. OWA, M. NUNOKAWA AND H. SAITOH, Some inequalities involving multivalent functions, *Ann. Polon. Math.*, **60** (1994), 159–162.



---

Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions

Hüseyin Irmak and R.K. Raina

---

Title Page

Contents




Go Back

Close

Quit

Page 15 of 16

- [9] S. OWA, M. NUNOKAWA AND S. FUKUI, A criterion for  $p$ -valent starlike functions, *Intern. J. Math. & Math. Sci.*, **17** (1994), 205–207.
- [10] S. OWA, H.M. SRIVASTAVA, F.-Y. REN AND W.-Q. YANG, The starlikeness of a certain class of integral operators, *Complex Variables*, **27** (1995), 185–191.
- [11] H.M. SRIVASTAVA AND S. OWA, (EDITORS), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London, and Hong Kong, 1992.



---

**Some Inequalities Exhibiting  
Certain Properties Of Some  
Subclasses Of Multivalently  
Analytic Functions**

Hüseyin Irmak and R.K. Raina

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16