

PIROSKA LAKATOS AND LÁSZLÓ LOSONCZI

Institute of Mathematics,  
Debrecen University,  
4010 Debrecen, pf.12, Hungary.

*EMail:* [lapi@math.klte.hu](mailto:lapi@math.klte.hu)

*EMail:* [losi@math.klte.hu](mailto:losi@math.klte.hu)



---

volume 4, issue 3, article 60,  
2003.

*Received 12 December, 2002;  
accepted 30 July, 2003.*

*Communicated by: A. Sofo*

---

[Abstract](#)

[Contents](#)




[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)




---

**On Zeros of Reciprocal  
Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 18

### Abstract

The first author [1] proved that all zeros of the reciprocal polynomial

$$P_m(z) = \sum_{k=0}^m A_k z^k \quad (z \in \mathbb{C}),$$

of degree  $m \geq 2$  with real coefficients  $A_k \in \mathbb{R}$  (i.e.  $A_m \neq 0$  and  $A_k = A_{m-k}$  for all  $k = 0, \dots, [\frac{m}{2}]$ ) are on the unit circle, provided that

$$|A_m| \geq \sum_{k=0}^m |A_k - A_m| = \sum_{k=1}^{m-1} |A_k - A_m|.$$

Moreover, the zeros of  $P_m$  are near to the  $m+1$ st roots of unity (except the root 1). A. Schinzel [3] generalized the first part of Lakatos' result for self-inversive polynomials i.e. polynomials

$$P_m(z) = \sum_{k=0}^m A_k z^k$$

for which  $A_k \in \mathbb{C}$ ,  $A_m \neq 0$  and  $\epsilon \bar{A}_k = A_{m-k}$  for all  $k = 0, \dots, m$  with a fixed  $\epsilon \in \mathbb{C}$ ,  $|\epsilon| = 1$ . He proved that all zeros of  $P_m$  are on the unit circle, provided that

$$|A_m| \geq \inf_{c,d \in \mathbb{C}, |d|=1} \sum_{k=0}^m |cA_k - d^{m-k} A_m|.$$

If the inequality is strict the zeros are single. The aim of this paper is to show that for real reciprocal polynomials of odd degree Lakatos' result remains valid even if

$$|A_m| \geq \cos^2 \frac{\pi}{2(m+1)} \sum_{k=1}^{m-1} |A_k - A_m|.$$

We conjecture that Schinzel's result can also be extended similarly: all zeros of  $P_m$  are on the unit circle if  $P_m$  is self-inversive and


$$|A_m| \geq \cos \frac{\pi}{2(m+1)} \inf_{c,d \in \mathbb{C}, |d|=1} \sum_{k=0}^m |cA_k - d^{m-k} A_m|.$$

*2000 Mathematics Subject Classification:* Primary 30C15, Secondary 12D10, 42C05  
*Key words:* Reciprocal, Semi-reciprocal polynomials, Chebyshev transform, Zeros on the unit circle.

The authors had their research supported by Hungarian NFSR grant No.s T 043034 and T 043080 respectively.

## Contents

1	Introduction .....	4
2	The Main Result .....	7
3	Remarks on Schinzel's Theorem .....	13
References		



---

**On Zeros of Reciprocal  
Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczi

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 18

# 1. Introduction

Studying the spectral properties of the Coxeter transformation Lakatos [1] found that all zeros of the reciprocal polynomial

$$P_m(z) = \sum_{k=0}^m A_k z^k \quad (z \in \mathbb{C})$$

of degree  $m \geq 2$  with real coefficients, i.e.

$$(1.1) \quad A_m \neq 0, \quad A_k \in \mathbb{R}, \quad \text{and} \quad A_k = A_{m-k} \quad (k = 0, \dots, \lfloor \frac{m}{2} \rfloor)$$

are on the unit circle, provided that

$$(1.2) \quad |A_m| \geq \sum_{k=1}^{m-1} |A_k - A_m|.$$


She used Chebyshev transformation to prove this result.

The manuscript on this was sent to A. Schinzel for his comments. He generalized the above theorem [3] for self-inversive polynomials by proving that all zeros of the polynomial  $P_m(z) = \sum_{k=0}^m A_k z^k$  where

$$(1.3) \quad A_m \neq 0, \quad A_k \in \mathbb{C}, \quad \text{and} \quad \epsilon \bar{A}_k = A_{m-k} \quad (k = 0, \dots, m) \quad \text{with} \quad \epsilon \in \mathbb{C}, \quad |\epsilon| = 1$$

are on the unit circle, provided that

$$(1.4) \quad |A_m| \geq \inf_{c, d \in \mathbb{C}, |d|=1} \sum_{k=0}^m |c A_k - d^{m-k} A_m|$$



**On Zeros of Reciprocal Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 4 of 18

holds.

Schinzel's proof was based on a theorem of Cohn [4] and on the estimate

$$(1.5) \quad \min_{z \in \mathbb{C}, |z|=1} \left| \sum_{k=1}^m k z^{m-k} \right| \geq \frac{m}{2}.$$

Learning of Schinzel's generalization, the first author made an attempt to improve her result. Although the method of Chebyshev transformation does not seem to work for self-inversive polynomials, it can be used to obtain information about the location of the zeros on the unit circle. She could prove [1] that condition (1.2) ensures that the distribution of the zeros  $e^{iu_j}$  ( $j = 1, \dots, m$ ) of  $P_m(z) = \sum_{k=0}^m A_k z^k$  satisfying (1.1), (1.2) is quite regular. They can be arranged such that

$$(1.6) \quad |\epsilon_j - e^{iu_j}| < \frac{\pi}{m+1} \quad (j = 1, \dots, m)$$

holds, where

$$\epsilon_j = e^{i \frac{j}{m+1} 2\pi} \quad (j = 1, 2, \dots, m)$$


are the  $m+1$ st roots of unity except 1.

If  $m = 2n+1$  is odd then  $-1 = e^{iu_{n+1}}$  is always a zero and all zeros of  $P_{2n+1}$  are single.

If  $m = 2n$  is even, (1.2) holds with equality and

$$\operatorname{sgn} A_{2n} = \operatorname{sgn}(-1)^{k+1}(A_k - A_{2n})$$

for all  $k = 1, 2, \dots, n$  with  $A_k - A_{2n} \neq 0$ , then  $u_n = u_{n+1} = \pi$ , the number  $-1 = e^{iu_n} = e^{iu_{n+1}}$  is a double zero of  $P_{2n}$ . Otherwise all zeros of  $P_{2n}$  are single.



**On Zeros of Reciprocal Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents




Go Back

Close

Quit

Page 5 of 18

The aim of this paper is to show that for polynomials of odd degree both results can be improved. For even degree polynomials this is not possible. Since, for first degree reciprocal or self-inversive polynomials the only zero of the polynomial has modulus one we may assume that the degree  $m \geq 2$ .



---

**On Zeros of Reciprocal  
Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 18

## 2. The Main Result

The theorem of Lakatos can be improved as follows.

**Theorem 2.1.** *All zeros of the reciprocal polynomial*

$$(2.1) \quad P_{2n+1}(z) = \sum_{k=0}^{2n+1} A_k z^k \quad (z \in \mathbb{C})$$

*of odd degree  $2n + 1 \geq 3$  with real coefficients i.e.*

$$(2.2) \quad A_{2n+1} \neq 0, \quad A_k \in \mathbb{R}, \quad \text{and} \quad A_k = A_{2n+1-k} \quad (k = 0, \dots, n)$$

*are on the unit circle, provided that*

$$(2.3) \quad |A_{2n+1}| \geq \cos^2 \frac{\pi}{2(2n+2)} \sum_{k=1}^{2n} |A_k - A_{2n+1}|.$$


*Moreover, if (2.2), (2.3) hold then all zeros  $e^{iu_j}$  ( $j = 1, 2, \dots, 2n+1$ ) of  $P_{2n+1}$  are single,  $-1 = e^{iu_{n+1}}$  is always a zero and the zeros can be arranged such that*

$$(2.4) \quad |\epsilon_j - e^{iu_j}| < \frac{\pi}{2n+2} \quad (j = 1, \dots, 2n+1),$$

*where*

$$\epsilon_j = e^{i \frac{j}{2n+2} 2\pi} \quad (j = 1, 2, \dots, 2n+1)$$

*are the  $2n + 2$ nd roots of unity except 1.*



**On Zeros of Reciprocal  
Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 7 of 18

*Proof.* With the notation

$$l := A_{2n+1} = A_0, a_1 := A_{2n} - l = A_1 - l, \dots, a_n := A_{n+1} - l = A_n - l$$

we have

$$\begin{aligned} h_{2n+1}(z) &:= l(z^{2n+1} + z^{2n} + \dots + z + 1) + \sum_{k=1}^n a_k (z^{2n+1-k} + z^k) \\ &= P_{2n+1}(z) = \sum_{k=0}^{2n+1} A_k z^k \quad (z \in \mathbb{C}). \end{aligned}$$

(2.2) goes over into

$$l \neq 0, l, a_k \in R \quad (k = 1, \dots, n)$$

while (2.3) goes over into


$$(2.5) \quad |l| \geq 2 \cos^2 \frac{\pi}{2(2n+2)} \sum_{k=1}^n |a_k|.$$

We have  $h_{2n+1}(z) = (z+1)\bar{h}_{2n}(z)$  with

$$\bar{h}_{2n}(z) = l\bar{v}_{2n}(z) + \sum_{k=1}^n a_k e_k(z) \bar{w}_{2n-2k}(z)$$

where

$$\begin{aligned} \bar{v}_{2n}(z) &= z^{2n} + z^{2n-2} + \dots + z^2 + 1 \\ \bar{w}_{2n-2k}(z) &= \frac{z^{2n+1-2k} + 1}{z + 1}, \quad e_k(z) = z^k. \end{aligned}$$



**On Zeros of Reciprocal Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 8 of 18



The Chebyshev transform  $\mathcal{T}\bar{h}_{2n}$  of  $\bar{h}_{2n}$  was calculated in [1]:

$$\begin{aligned}\mathcal{T}\bar{h}_{2n}(x) &= l\mathcal{T}\bar{v}_{2n}(x) + \sum_{k=1}^n a_k \mathcal{T}(e_k \cdot \bar{w}_{2n-2k})(x) \\ &= l U_n\left(\frac{x}{2}\right) + \sum_{k=1}^n a_k \left[ U_{n-k}\left(\frac{x}{2}\right) - U_{n-k-1}\left(\frac{x}{2}\right) \right],\end{aligned}$$

where  $U_n$  is the Chebyshev polynomial of degree  $n$  of the second kind defined by  $U_n(\cos x) := \frac{\sin(n+1)x}{\sin x}$  ( $n = 1, 2, \dots$ ) and  $U_{-1}(x) := 0$ . Evaluating of  $\mathcal{T}\bar{h}_{2n}$  at the points


$$x_j = 2 \cos y_j \quad \text{with} \quad y_j = \frac{j + \frac{1}{2}}{2n + 2} 2\pi \quad (j = 0, \dots, n)$$

of the open interval  $] - 2, 2[$  gives that (see [1])

$$\begin{aligned}\mathcal{T}\bar{h}_{2n}(x_j) &= \frac{l(-1)^j}{\sin y_j} + \sum_{k=1}^n a_k \frac{\cos \frac{2n-2k+1}{2} y_j}{\cos \frac{y_j}{2}} \\ &= \frac{l(-1)^j + 2 \sum_{k=1}^n a_k \sin \frac{y_j}{2} \cos \frac{2n-2k+1}{2} y_j}{\sin y_j}.\end{aligned}$$

We have for  $j = 0, \dots, n - 1$

$$(2.6) \quad (0 <) \sin \frac{y_j}{2} < \sin \frac{y_n}{2} = \sin \left( \frac{\pi}{2} - \frac{\pi}{2(2n+2)} \right) = \cos \frac{\pi}{2(2n+2)},$$



**On Zeros of Reciprocal  
Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 9 of 18

while for  $j = n$  there is equality here. The absolute value of the factor

$$\begin{aligned} & \cos \frac{2n - 2k + 1}{2} y_j \\ &= \cos \frac{2(n - k)j + j + (n - k) + \frac{1}{2}}{2n + 2} \pi \quad (k = 1, \dots, n; j = 0, \dots, n) \end{aligned}$$

takes its maximum if the fraction

$$\frac{2(n - k)j + j + (n - k) + \frac{1}{2}}{2n + 2}$$

is nearest to an integer. Clearly the nearest possible value of this fraction to an integer is  $\frac{1}{2(2n+2)}$  (this value is attained at  $j = 0, k = n$ ). Thus we have shown that


$$(2.7) \quad \left| \cos \frac{2n - 2k + 1}{2} y_j \right| \leq \cos \frac{\pi}{2(2n + 2)} \quad (k = 1, \dots, n; j = 0, \dots, n).$$

Let for  $j = 0, \dots, n$

$$S_j := 2 \left| \sum_{k=1}^n a_k \sin \frac{y_j}{2} \cos \frac{2n - 2k + 1}{2} y_j \right|.$$

Then, for  $j = 0, \dots, n - 1$  by (2.6), (2.7) we have

$$S_j \leq 2 \sin \frac{y_j}{2} \sum_{k=1}^n |a_k| \cos \frac{\pi}{2(2n + 2)} < 2 \cos^2 \frac{\pi}{2(2n + 2)} \sum_{k=1}^n |a_k|$$



On Zeros of Reciprocal  
Polynomials of Odd Degree

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 10 of 18

unless  $\sum_{k=1}^n |a_k| = 0$ .

Thus, by (2.3) or (2.5) we have

$$S_j < 2 \cos^2 \frac{\pi}{2(2n+2)} \sum_{k=1}^n |a_k| \leq |l|$$

and the resulting inequality

$$S_j < |l|$$


remains valid even if  $\sum_{k=1}^n |a_k| = 0$ .

For  $j = n$  we have

$$\begin{aligned} S_n &\leq 2 \sin \frac{y_n}{2} \sum_{k=1}^n |a_k| \left| \cos \frac{2(n-k)n + n + (n-k) + \frac{1}{2}}{2n+2} \pi \right| \\ &= 2 \cos \frac{\pi}{2(2n+2)} \sum_{k=1}^n |a_k| \left| \cos \left( n-k + \frac{k + \frac{1}{2}}{2n+2} \right) \pi \right| \\ &= 2 \cos \frac{\pi}{2(2n+2)} \sum_{k=1}^n |a_k| \cos \frac{k + \frac{1}{2}}{2n+2} \pi \\ &< 2 \cos^2 \frac{\pi}{2(2n+2)} \sum_{k=1}^n |a_k| \end{aligned}$$

unless  $\sum_{k=1}^n |a_k| = 0$ . Thus, by (2.3) or (2.5) we have

$$S_n < 2 \cos^2 \frac{\pi}{2(2n+2)} \sum_{k=1}^n |a_k| \leq |l|.$$



**On Zeros of Reciprocal Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 11 of 18

and the inequality

$$S_n < |l|$$

remains valid even if  $\sum_{k=1}^n |a_k| = 0$ .


Looking again at the Chebyshev transform we can see that by the inequalities  $S_j < |l|$  ( $j = 0, \dots, n$ ) we have

$$\operatorname{sgn}(\mathcal{T}\bar{h}_{2n}(x_j)) = \operatorname{sgn} l \operatorname{sgn}(-1)^j \quad (j = 0, 1, \dots, n)$$

therefore  $\mathcal{T}\bar{h}_{2n}$  has  $n$  different zeros in  $] -2, 2[$ . Writing these zeros in the form  $2 \cos v_j$  with  $0 \leq v_1 \leq v_2 \leq \dots \leq v_n \leq \pi$  and applying Lemma 1 of [1] we conclude that all zeros of  $\bar{h}_{2n}$  are single, and of the form  $e^{\pm i v_j}$ , where

$$(2.8) \quad y_{j-1} < v_j < y_j \quad (j = 1, \dots, n).$$

Let  $u_j := v_j$  for  $j = 1, \dots, n$ ,  $u_{n+1} := \pi$  and  $u_{n+1+j} := 2\pi - u_{n+1-j}$  ( $j = 1, \dots, n$ ), then we obtain that all zeros of  $P_{2n+1} = h_{2n+1}$  are  $e^{i u_j}$  ( $j = 1, \dots, 2n+1$ ) and by (2.8) the condition (2.4) holds.  $\square$



**On Zeros of Reciprocal  
Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 12 of 18

### 3. Remarks on Schinzel's Theorem

Schinzel's result can be generalized as follows.

**Theorem 3.1.** *Let  $P_m(z) = \sum_{k=0}^m A_k z^k$  be a self-inversive polynomial of degree  $m$ , i.e. let  $A_k \in \mathbb{C}$ ,  $A_m \neq 0$  and  $\epsilon \bar{A}_k = A_{m-k}$  for all  $k = 0, \dots, m$  with a fixed  $\epsilon \in \mathbb{C}$ ,  $|\epsilon| = 1$ . If*

$$(3.1) \quad |A_m| \geq \frac{m}{2\mu_m} \inf_{c,d \in \mathbb{C}, |d|=1} \sum_{k=0}^m |cA_k - d^{m-k}A_m|,$$

where

$$(3.2) \quad \mu_m := \min_{|z| \leq 1} \left| \sum_{k=1}^m k z^{m-k} \right|$$


then all zeros of  $P_m$  are on the unit circle. If the inequality is strict the zeros are single.

Apart from minor changes Schinzel's proof [3] is valid for this more general result hence we omit the proof.

By Cohn's theorem [4] any self-inversive polynomial  $P_m$  and the polynomial

$$z^{m-1} P'_m(z^{-1}) = \sum_{k=1}^m k A_k z^{m-k} \quad (z \in \mathbb{C})$$

have the same number of zeros inside the unit circle. Applying this for the polynomial  $\sum_{k=0}^m z^k$  we obtain that  $\sum_{k=1}^m k z^{m-k}$  has no zeros inside the unit



On Zeros of Reciprocal  
Polynomials of Odd Degree

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 13 of 18

circle, thus the modulus of the latter takes its positive minimum in the unit disk on the unit circle:

$$\mu_m = \min_{|z|=1} \left| \sum_{k=1}^m kz^{m-k} \right| > 0.$$

By some known identities (see [2, Part 6, Problems 16, 18]) for trigonometric sums we easily get that

$$\begin{aligned} D_m(t) &:= \left| \sum_{k=1}^m kz^{m-k} \right|_{z=e^{it}} \\ &= \sqrt{\left[ \sum_{k=1}^m k \cos(m-k)t \right]^2 + \left[ \sum_{k=1}^m k \sin(m-k)t \right]^2} \\ &= \sqrt{\left[ \frac{m}{2} + \frac{1}{2} \left( \frac{\sin \frac{mt}{2}}{\sin \frac{t}{2}} \right)^2 \right]^2 + \left[ \frac{m \sin t - \sin mt}{4 \sin^2 \frac{t}{2}} \right]^2}. \end{aligned}$$

From this it follows that  $\mu_m = \min_{t \in [0, 2\pi]} D_m(t) \geq \frac{m}{2}$  and for even  $m = 2n$  we have equality here, since for  $t = \pi$

$$D_m(\pi) = \frac{m}{2}.$$

This means that for even  $m$  Theorem 3.1 coincides with Schinzel's result. For odd  $m$  however

$$\mu_m > \frac{m}{2}.$$



On Zeros of Reciprocal Polynomials of Odd Degree

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 14 of 18

Let for  $t \in [0, 2\pi]$

$$x_m(t) := \frac{m}{2} + \frac{1}{2} \left( \frac{\sin \frac{mt}{2}}{\sin \frac{t}{2}} \right)^2,$$

$$y_m(t) := \frac{m \sin t - \sin mt}{4 \sin^2 \frac{t}{2}},$$

$$z_m(t) := x_m(t) + iy_m(t),$$

then  $D_m^2(t) = |z_m(t)|^2 = x_m(t)^2 + y_m(t)^2$ . As  $D_m(\pi + t) = D_m(\pi - t)$  it is enough to consider  $D_m$  on the interval  $[0, \pi]$ .

The next figure shows the graph of  $z_9$  in the complex plane and an enlargement of the part which is nearest to the origin. In the latter the distance of the origin from the graph of  $z_9$  is also shown. The point of the graph with  $t = t_m = \frac{m-1}{m}\pi$  is distinguished by a small circle.


Our numerical experiments give base to the following conjecture.

**Conjecture 1.** *For odd  $m$  we have*

$$\mu_m = \min_{t \in [0, 2\pi]} D_m(t) \geq \frac{m}{2} \sec \frac{\pi}{2m+2}.$$

A simple calculation shows that for odd  $m = 2n + 1$

$$D_m(t_m) = \frac{m}{2} \sec \frac{\pi}{2m}.$$



**On Zeros of Reciprocal Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 15 of 18




Figure 1: Graph of  $z_0$ .


It is clear that  $\mu_m$  is the *distance of the graph of  $z_m$  from the origin*. The minimum of  $D_m$  is attained near to  $t_m$ . It is relatively easy to show that the minimum is attained in the interval  $[t_m, \pi]$ . Numerical calculations seem to justify that the minimum point is in the smaller interval

$$\left[ t_m, t_m + \frac{2\pi}{m^2} \right].$$

Theorem 3.1 and Conjecture 1 give

**Conjecture 2.** *All zeros of the self-inversive polynomial*

$$P_{2n+1}(z) = \sum_{k=0}^{2n+1} A_k z^k \quad (z \in \mathbb{C})$$



On Zeros of Reciprocal Polynomials of Odd Degree

Piroska Lakatos and  
László Losonczy

Title Page

Contents



Go Back

Close

Quit

Page 16 of 18




of odd degree  $2n + 1$ , i.e.

$A_{2n+1} \neq 0$ ,  $A_k \in \mathbb{C}$ , and  $\epsilon \bar{A}_k = A_{2n+1-k}$  ( $k = 0, \dots, 2n+1$ ) with  $\epsilon \in \mathbb{C}$ ,  $|\epsilon| = 1$

are on the unit circle, provided that

$$|A_{2n+1}| \geq \cos \frac{\pi}{2(2n+2)} \inf_{c,d \in \mathbb{C}, |d|=1} \sum_{k=0}^{2n+1} |cA_k - d^{2n+1-k} A_{2n+1}|.$$

holds. If the inequality is strict here then the zeros are single.




---

**On Zeros of Reciprocal Polynomials of Odd Degree**

Piroska Lakatos and  
László Losonczi

---

Title Page

Contents



Go Back


Close

Quit

Page 17 of 18

## References

- [1] P. LAKATOS, On zeros of reciprocal polynomials, *Publ. Math.* (Debrecen) **61** (2002), 645–661.
- [2] G. PÓLYA AND G. SZEGŐ, *Problems and Theorems in Analysis*, Vol. II, Springer 1976.
- [3] A. SCHINZEL, Self-inversive polynomials with all zeros on the unit circle, *Ramanujan Journal*, to appear.
- [4] A. COHN, Über die Anzahl der Wurzeln einer algebraischen Gleichung in einem Kreise, *Math. Zeit.*, **14** (1922), 110–148.



---

On Zeros of Reciprocal  
Polynomials of Odd Degree

Piroska Lakatos and  
László Losonczy

---

Title Page

Contents



Go Back

Close

Quit

Page 18 of 18