

Journal of Inequalities in Pure and Applied Mathematics

REVERSES OF SCHWARZ, TRIANGLE AND BESSEL INEQUALITIES IN INNER PRODUCT SPACES


S.S. DRAGOMIR

School of Computer Science and Mathematics
Victoria University of Technology
PO Box 14428, MCMC 8001
Victoria, Australia.

EMail: sever.dragomir@vu.edu.au

URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>

©2000 Victoria University
ISSN (electronic): 1443-5756
171-03



volume 5, issue 3, article 76,
2004.

*Received 12 December, 2003;
accepted 11 July, 2004.*

Communicated by: B. Mond

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


Reverses of the Schwarz, triangle and Bessel inequalities in inner product spaces that improve some earlier results are pointed out. They are applied to obtain new Grüss type inequalities. Some natural integral inequalities are also stated.

2000 Mathematics Subject Classification: Primary 26D15, 46C05.

Key words: Schwarz's inequality, Triangle inequality, Bessel's inequality, Grüss type inequalities, Integral inequalities.

Contents

1	Introduction	3
2	Some Reverses of Schwarz's Inequality	9
3	Reverses of the Triangle Inequality	16
4	Some Grüss Type Inequalities	19
5	Reverses of Bessel's Inequality	23
6	Some Grüss Type Inequalities for Orthonormal Families	29
7	Integral Inequalities	33
	References	



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 2 of 40

1. Introduction

Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} . The following inequality is known in the literature as *Schwarz's inequality*:

$$(1.1) \quad |\langle x, y \rangle|^2 \leq \|x\|^2 \|y\|^2, \quad x, y \in H;$$

where $\|z\|^2 = \langle z, z \rangle$, $z \in H$. The equality occurs in (1.1) if and only if x and y are linearly dependent.

In [7], the following *reverse* of Schwarz's inequality has been obtained:

$$(1.2) \quad 0 \leq \|x\|^2 \|y\|^2 - |\langle x, y \rangle|^2 \leq \frac{1}{4} |A - a|^2 \|y\|^4,$$

provided $x, y \in H$ and $a, A \in \mathbb{K}$ are so that either


$$(1.3) \quad \operatorname{Re} \langle Ay - x, x - ay \rangle \geq 0,$$

or, equivalently,

$$(1.4) \quad \left\| x - \frac{a + A}{2} \cdot y \right\| \leq \frac{1}{2} |A - a| \|y\|,$$

holds. The constant $\frac{1}{4}$ is best possible in (1.2) in the sense that it cannot be replaced by a smaller quantity.

If x, y, A, a satisfy either (1.3) or (1.4), then the following reverse of Schwarz's



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 3 of 40

inequality also holds [8]

$$(1.5) \quad \|x\| \|y\| \leq \frac{1}{2} \cdot \frac{\operatorname{Re} [A\overline{\langle x, y \rangle} + \bar{a}\langle x, y \rangle]}{[\operatorname{Re}(\bar{a}A)]^{\frac{1}{2}}} \\ \leq \frac{1}{2} \cdot \frac{|A| + |a|}{[\operatorname{Re}(\bar{a}A)]^{\frac{1}{2}}} |\langle x, y \rangle|,$$

provided that, the complex numbers a and A satisfy the condition $\operatorname{Re}(\bar{a}A) > 0$. In both inequalities in (1.5), the constant $\frac{1}{2}$ is best possible.

An additive version of (1.5) may be stated as well (see also [9])

$$(1.6) \quad 0 \leq \|x\|^2 \|y\|^2 - |\langle x, y \rangle|^2 \\ \leq \frac{1}{4} \cdot \frac{(|A| - |a|)^2 + 4[|Aa| - \operatorname{Re}(\bar{a}A)]}{\operatorname{Re}(\bar{a}A)} |\langle x, y \rangle|^2.$$

In this inequality, $\frac{1}{4}$ is the best possible constant.

It has been proven in [10], that


$$(1.7) \quad 0 \leq \|x\|^2 - |\langle x, y \rangle|^2 \leq \frac{1}{4} |\phi - \varphi|^2 - \left| \frac{\phi + \varphi}{2} - \langle x, e \rangle \right|^2;$$

provided, either

$$(1.8) \quad \operatorname{Re} \langle \phi e - x, x - \varphi e \rangle \geq 0,$$

or, equivalently,

$$(1.9) \quad \left\| x - \frac{\phi + \varphi}{2} e \right\| \leq \frac{1}{2} |\phi - \varphi|,$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 4 of 40

where $e \in H$, $\|e\| = 1$. The constant $\frac{1}{4}$ in (1.7) is also best possible.

If we choose $e = \frac{y}{\|y\|}$, $\phi = \Gamma \|y\|$, $\varphi = \gamma \|y\|$ ($y \neq 0$), $\Gamma, \gamma \in \mathbb{K}$, then by (1.8) and (1.9) we have,

$$(1.10) \quad \operatorname{Re} \langle \Gamma y - x, x - \gamma y \rangle \geq 0,$$

or, equivalently,

$$(1.11) \quad \left\| x - \frac{\Gamma + \gamma}{2} y \right\| \leq \frac{1}{2} |\Gamma - \gamma| \|y\|,$$

imply the following reverse of Schwarz's inequality:


$$(1.12) \quad 0 \leq \|x\|^2 \|y\|^2 - |\langle x, y \rangle|^2 \leq \frac{1}{4} |\Gamma - \gamma|^2 \|y\|^4 - \left| \frac{\Gamma + \gamma}{2} \|y\|^2 - \langle x, y \rangle \right|^2.$$

The constant $\frac{1}{4}$ in (1.12) is sharp.

Note that this inequality is an improvement of (1.2), but it might not be very convenient for applications.

Now, let $\{e_i\}_{i \in I}$ be a finite or infinite family of orthonormal vectors in the inner product space $(H; \langle \cdot, \cdot \rangle)$, i.e., we recall that

$$\langle e_i, e_j \rangle = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}, \quad i, j \in I.$$



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 5 of 40

In [11], we proved that, if $\{e_i\}_{i \in I}$ is as above, $F \subset I$ is a finite part of I such that either

$$(1.13) \quad \operatorname{Re} \left\langle \sum_{i \in F} \phi_i e_i - x, x - \sum_{i \in F} \varphi_i e_i \right\rangle \geq 0,$$

or, equivalently,


$$(1.14) \quad \left\| x - \sum_{i \in F} \frac{\phi_i + \varphi_i}{2} e_i \right\| \leq \frac{1}{2} \left(\sum_{i \in F} |\phi_i - \varphi_i|^2 \right)^{\frac{1}{2}},$$

holds, where $(\phi_i)_{i \in I}$, $(\varphi_i)_{i \in I}$ are real or complex numbers, then we have the following reverse of *Bessel's inequality*:

$$(1.15) \quad \begin{aligned} 0 &\leq \|x\|^2 - \sum_{i \in F} |\langle x, e_i \rangle|^2 \\ &\leq \frac{1}{4} \cdot \sum_{i \in F} |\phi_i - \varphi_i|^2 - \operatorname{Re} \left\langle \sum_{i \in F} \phi_i e_i - x, x - \sum_{i \in F} \varphi_i e_i \right\rangle \\ &\leq \frac{1}{4} \cdot \sum_{i \in F} |\phi_i - \varphi_i|^2. \end{aligned}$$

The constant $\frac{1}{4}$ in both inequalities is sharp. This result improves an earlier result by N. Ujević obtained only for real spaces [21].

In [10], by the use of a different technique, another reverse of Bessel's in-



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 6 of 40

equality has been proven, namely:

$$\begin{aligned}
 (1.16) \quad 0 &\leq \|x\|^2 - \sum_{i \in F} |\langle x, e_i \rangle|^2 \\
 &\leq \frac{1}{4} \cdot \sum_{i \in F} |\phi_i - \varphi_i|^2 - \sum_{i \in F} \left| \frac{\phi_i + \varphi_i}{2} - \langle x, e_i \rangle \right|^2 \\
 &\leq \frac{1}{4} \cdot \sum_{i \in F} |\phi_i - \varphi_i|^2,
 \end{aligned}$$

provided that $(e_i)_{i \in I}$, $(\phi_i)_{i \in I}$, $(\varphi_i)_{i \in I}$, x and F are as above.


Here the constant $\frac{1}{4}$ is sharp in both inequalities.

It has also been shown that the bounds provided by (1.15) and (1.16) for the Bessel's difference $\|x\|^2 - \sum_{i \in F} |\langle x, e_i \rangle|^2$ cannot be compared in general, meaning that there are examples for which one is smaller than the other [10].

Finally, we recall another type of reverse for Bessel inequality that has been obtained in [12]:

$$(1.17) \quad \|x\|^2 \leq \frac{1}{4} \cdot \frac{\sum_{i \in F} (|\phi_i| + |\varphi_i|)^2}{\sum_{i \in F} \operatorname{Re}(\phi_i \overline{\varphi_i})} \sum_{i \in F} |\langle x, e_i \rangle|^2;$$

provided $(\phi_i)_{i \in I}$, $(\varphi_i)_{i \in I}$ satisfy (1.13) (or, equivalently (1.14)) and $\sum_{i \in F} \operatorname{Re}(\phi_i \overline{\varphi_i}) > 0$. Here the constant $\frac{1}{4}$ is also best possible.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 7 of 40


An additive version of (1.17) is

$$(1.18) \quad 0 \leq \|x\|^2 - \sum_{i \in F} |\langle x, e_i \rangle|^2 \\ \leq \frac{1}{4} \cdot \frac{\sum_{i \in F} \{(|\phi_i| - |\varphi_i|)^2 + 4[|\phi_i \varphi_i| - \operatorname{Re}(\phi_i \overline{\varphi_i})]\}}{\sum_{i \in F} \operatorname{Re}(\phi_i \overline{\varphi_i})} \sum_{i \in F} |\langle x, e_i \rangle|^2.$$

The constant $\frac{1}{4}$ is best possible.

It is the main aim of the present paper to point out new reverse inequalities to Schwarz's, triangle and Bessel's inequalities.

Some results related to Grüss' inequality in inner product spaces are also pointed out. Natural applications for integrals are also provided.



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 8 of 40

2. Some Reverses of Schwarz's Inequality

The following result holds.

Theorem 2.1. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} ($\mathbb{K} = \mathbb{R}$, $\mathbb{K} = \mathbb{C}$) and $x, a \in H$, $r > 0$ are such that*

$$(2.1) \quad x \in \overline{B}(a, r) := \{z \in H \mid \|z - a\| \leq r\}.$$

(i) *If $\|a\| > r$, then we have the inequalities*

$$(2.2) \quad 0 \leq \|x\|^2 \|a\|^2 - |\langle x, a \rangle|^2 \leq \|x\|^2 \|a\|^2 - [\operatorname{Re} \langle x, a \rangle]^2 \leq r^2 \|x\|^2.$$

The constant 1 in front of r^2 is best possible in the sense that it cannot be replaced by a smaller one.

(ii) *If $\|a\| = r$, then*

$$(2.3) \quad \|x\|^2 \leq 2 \operatorname{Re} \langle x, a \rangle \leq 2 |\langle x, a \rangle|.$$

The constant 2 is best possible in both inequalities.


(iii) *If $\|a\| < r$, then*

$$(2.4) \quad \|x\|^2 \leq r^2 - \|a\|^2 + 2 \operatorname{Re} \langle x, a \rangle \leq r^2 - \|a\|^2 + 2 |\langle x, a \rangle|.$$

Here the constant 2 is also best possible.

Proof. Since $x \in \overline{B}(a, r)$, then obviously $\|x - a\|^2 \leq r^2$, which is equivalent to

$$(2.5) \quad \|x\|^2 + \|a\|^2 - r^2 \leq 2 \operatorname{Re} \langle x, a \rangle.$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 9 of 40



Title Page

Contents



Go Back

Close

Quit

Page 10 of 40

(i) If $\|a\| > r$, then we may divide (2.5) by $\sqrt{\|a\|^2 - r^2} > 0$ getting

$$(2.6) \quad \frac{\|x\|^2}{\sqrt{\|a\|^2 - r^2}} + \sqrt{\|a\|^2 - r^2} \leq \frac{2 \operatorname{Re} \langle x, a \rangle}{\sqrt{\|a\|^2 - r^2}}.$$

Using the elementary inequality

$$\alpha p + \frac{1}{\alpha} q \geq 2\sqrt{pq}, \quad \alpha > 0, \quad p, q \geq 0,$$

we may state that

$$(2.7) \quad 2\|x\| \leq \frac{\|x\|^2}{\sqrt{\|a\|^2 - r^2}} + \sqrt{\|a\|^2 - r^2}.$$

Making use of (2.6) and (2.7), we deduce


$$(2.8) \quad \|x\| \sqrt{\|a\|^2 - r^2} \leq \operatorname{Re} \langle x, a \rangle.$$

Taking the square in (2.8) and re-arranging the terms, we deduce the third inequality in (2.2). The others are obvious.

To prove the sharpness of the constant, assume, under the hypothesis of the theorem, that, there exists a constant $c > 0$ such that

$$(2.9) \quad \|x\|^2 \|a\|^2 - [\operatorname{Re} \langle x, a \rangle]^2 \leq cr^2 \|x\|^2,$$

provided $x \in \overline{B}(a, r)$ and $\|a\| > r$.



Title Page

Contents



Go Back

Close

Quit

Page 11 of 40

Let $r = \sqrt{\varepsilon} > 0$, $\varepsilon \in (0, 1)$, $a, e \in H$ with $\|a\| = \|e\| = 1$ and $a \perp e$. Put $x = a + \sqrt{\varepsilon}e$. Then obviously $x \in \overline{B}(a, r)$, $\|a\| > r$ and $\|x\|^2 = \|a\|^2 + \varepsilon \|e\|^2 = 1 + \varepsilon$, $\operatorname{Re} \langle x, a \rangle = \|a\|^2 = 1$, and thus $\|x\|^2 \|a\|^2 - [\operatorname{Re} \langle x, a \rangle]^2 = \varepsilon$. Using (2.9), we may write that

$$\varepsilon \leq c\varepsilon(1 + \varepsilon), \quad \varepsilon > 0$$

giving

$$(2.10) \quad c + c\varepsilon \geq 1 \quad \text{for any } \varepsilon > 0.$$

Letting $\varepsilon \rightarrow 0+$, we get from (2.10) that $c \geq 1$, and the sharpness of the constant is proved.

- (ii) The inequality (2.3) is obvious by (2.5) since $\|a\| = r$. The best constant follows in a similar way to the above.
- (iii) The inequality (2.3) is obvious. The best constant may be proved in a similar way to the above. We omit the details.

□

The following reverse of Schwarz's inequality holds.

Theorem 2.2. Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over \mathbb{K} and $x, y \in H$, $\gamma, \Gamma \in \mathbb{K}$ such that either

$$(2.11) \quad \operatorname{Re} \langle \Gamma y - x, x - \gamma y \rangle \geq 0,$$

or, equivalently,

$$(2.12) \quad \left\| x - \frac{\Gamma + \gamma}{2} y \right\| \leq \frac{1}{2} |\Gamma - \gamma| \|y\|,$$

holds.

(i) If $\operatorname{Re}(\Gamma\bar{\gamma}) > 0$, then we have the inequalities

$$(2.13) \quad \begin{aligned} \|x\|^2 \|y\|^2 &\leq \frac{1}{4} \cdot \frac{\{\operatorname{Re}[(\bar{\Gamma} + \bar{\gamma}) \langle x, y \rangle]\}^2}{\operatorname{Re}(\Gamma\bar{\gamma})} \\ &\leq \frac{1}{4} \cdot \frac{|\Gamma + \gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} |\langle x, y \rangle|^2. \end{aligned}$$

The constant $\frac{1}{4}$ is best possible in both inequalities.


(ii) If $\operatorname{Re}(\Gamma\bar{\gamma}) = 0$, then

$$(2.14) \quad \|x\|^2 \leq \operatorname{Re}[(\bar{\Gamma} + \bar{\gamma}) \langle x, y \rangle] \leq |\Gamma + \gamma| |\langle x, y \rangle|.$$

(iii) If $\operatorname{Re}(\Gamma\bar{\gamma}) < 0$, then

$$(2.15) \quad \begin{aligned} \|x\|^2 &\leq -\operatorname{Re}(\Gamma\bar{\gamma}) \|y\|^2 + \operatorname{Re}[(\bar{\Gamma} + \bar{\gamma}) \langle x, y \rangle] \\ &\leq -\operatorname{Re}(\Gamma\bar{\gamma}) \|y\|^2 + |\Gamma + \gamma| |\langle x, y \rangle|. \end{aligned}$$

Proof. The proof of the equivalence between the inequalities (2.11) and (2.12) follows by the fact that in an inner product space $\operatorname{Re} \langle Z - x, x - z \rangle \geq 0$ for $x, z, Z \in H$ is equivalent with $\left\| x - \frac{z+Z}{2} \right\| \leq \frac{1}{2} \|Z - z\|$ (see for example [9]).



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 12 of 40

Consider, for $y \neq 0$, $a = \frac{\gamma+\Gamma}{2}y$ and $r = \frac{1}{2}|\Gamma - \gamma| \|y\|$. Then

$$\|a\|^2 - r^2 = \frac{|\Gamma + \gamma|^2 - |\Gamma - \gamma|^2}{4} \|y\|^2 = \operatorname{Re}(\Gamma\bar{\gamma}) \|y\|^2.$$

- (i) If $\operatorname{Re}(\Gamma\bar{\gamma}) > 0$, then the hypothesis of (i) in Theorem 2.1 is satisfied, and by the second inequality in (2.2) we have

$$\|x\|^2 \frac{|\Gamma + \gamma|^2}{4} \|y\|^2 - \frac{1}{4} \{ \operatorname{Re} [(\bar{\Gamma} + \bar{\gamma}) \langle x, y \rangle] \}^2 \leq \frac{1}{4} |\Gamma - \gamma|^2 \|x\|^2 \|y\|^2$$

from where we derive

$$\frac{|\Gamma + \gamma|^2 - |\Gamma - \gamma|^2}{4} \|x\|^2 \|y\|^2 \leq \frac{1}{4} \{ \operatorname{Re} [(\bar{\Gamma} + \bar{\gamma}) \langle x, y \rangle] \}^2,$$

giving the first inequality in (2.13).

The second inequality is obvious.


To prove the sharpness of the constant $\frac{1}{4}$, assume that the first inequality in (2.13) holds with a constant $c > 0$, i.e.,

$$(2.16) \quad \|x\|^2 \|y\|^2 \leq c \cdot \frac{\{ \operatorname{Re} [(\bar{\Gamma} + \bar{\gamma}) \langle x, y \rangle] \}^2}{\operatorname{Re}(\Gamma\bar{\gamma})},$$

provided $\operatorname{Re}(\Gamma\bar{\gamma}) > 0$ and either (2.11) or (2.12) holds.

Assume that $\Gamma, \gamma > 0$, and let $x = \gamma y$. Then (2.11) holds and by (2.16) we deduce

$$\gamma^2 \|y\|^4 \leq c \cdot \frac{(\Gamma + \gamma)^2 \gamma^2 \|y\|^4}{\Gamma\gamma}$$



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 13 of 40



Title Page

Contents



Go Back

Close

Quit

Page 14 of 40

giving

$$(2.17) \quad \Gamma\gamma \leq c(\Gamma + \gamma)^2 \quad \text{for any } \Gamma, \gamma > 0.$$

Let $\varepsilon \in (0, 1)$ and choose in (2.17), $\Gamma = 1 + \varepsilon$, $\gamma = 1 - \varepsilon > 0$ to get $1 - \varepsilon^2 \leq 4c$ for any $\varepsilon \in (0, 1)$. Letting $\varepsilon \rightarrow 0+$, we deduce $c \geq \frac{1}{4}$, and the sharpness of the constant is proved.

(ii) and (iii) are obvious and we omit the details.

□

Remark 2.1. We observe that the second bound in (2.13) for $\|x\|^2 \|y\|^2$ is better than the second bound provided by (1.5).

The following corollary provides a reverse inequality for the additive version of Schwarz's inequality.

Corollary 2.3. With the assumptions of Theorem 2.2 and if $\operatorname{Re}(\Gamma\bar{\gamma}) > 0$, then we have the inequality:


$$(2.18) \quad 0 \leq \|x\|^2 \|y\|^2 - |\langle x, y \rangle|^2 \leq \frac{1}{4} \cdot \frac{|\Gamma - \gamma|^2}{\operatorname{Re}(\Gamma\bar{\gamma})} |\langle x, y \rangle|^2.$$

The constant $\frac{1}{4}$ is best possible in (2.18).

The proof is obvious from (2.13) on subtracting in both sides the same quantity $|\langle x, y \rangle|^2$. The sharpness of the constant may be proven in a similar manner to the one incorporated in the proof of (i), Theorem 2.2. We omit the details.

Remark 2.2. *It is obvious that the inequality (2.18) is better than (1.6) obtained in [9].*

For some recent results in connection to Schwarz's inequality, see [2], [13] and [15].



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 15 of 40

3. Reverses of the Triangle Inequality

The following reverse of the triangle inequality holds.

Proposition 3.1. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} ($\mathbb{K} = \mathbb{R}, \mathbb{C}$) and $x, a \in H$, $r > 0$ are such that*

$$(3.1) \quad \|x - a\| \leq r < \|a\|.$$

Then we have the inequality


$$(3.2) \quad 0 \leq \|x\| + \|a\| - \|x + a\| \\ \leq \sqrt{2}r \cdot \sqrt{\frac{\operatorname{Re} \langle x, a \rangle}{\sqrt{\|a\|^2 - r^2} (\sqrt{\|a\|^2 - r^2} + \|a\|)}}.$$

Proof. Using the inequality (2.8), we may write that

$$\|x\| \|a\| \leq \frac{\|a\| \operatorname{Re} \langle x, a \rangle}{\sqrt{\|a\|^2 - r^2}},$$

giving

$$(3.3) \quad 0 \leq \|x\| \|a\| - \operatorname{Re} \langle x, a \rangle \\ \leq \frac{\|a\| - \sqrt{\|a\|^2 - r^2}}{\sqrt{\|a\|^2 - r^2}} \operatorname{Re} \langle x, a \rangle$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 16 of 40

$$= \frac{r^2 \operatorname{Re} \langle x, a \rangle}{\sqrt{\|a\|^2 - r^2} \left(\sqrt{\|a\|^2 - r^2} + \|a\| \right)}.$$

Since

$$(\|x\| + \|a\|)^2 - \|x + a\|^2 = 2(\|x\| \|a\| - \operatorname{Re} \langle x, a \rangle),$$

hence, by (3.3), we have

$$\begin{aligned} \|x\| + \|a\| &\leq \sqrt{\|x + a\|^2 + \frac{2r^2 \operatorname{Re} \langle x, a \rangle}{\sqrt{\|a\|^2 - r^2} \left(\sqrt{\|a\|^2 - r^2} + \|a\| \right)}} \\ &\leq \|x + a\| + \sqrt{2}r \cdot \sqrt{\frac{\operatorname{Re} \langle x, a \rangle}{\sqrt{\|a\|^2 - r^2} \left(\sqrt{\|a\|^2 - r^2} + \|a\| \right)}}, \end{aligned}$$

giving the desired inequality (3.2). \square


The following proposition providing a simpler reverse for the triangle inequality also holds.

Proposition 3.2. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over \mathbb{K} and $x, y \in H$, $M \geq m > 0$ such that either*

$$(3.4) \quad \operatorname{Re} \langle My - x, x - my \rangle \geq 0,$$

or, equivalently,

$$(3.5) \quad \left\| x - \frac{M + m}{2} \cdot y \right\| \leq \frac{1}{2} (M - m) \|y\|,$$



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 17 of 40

holds. Then we have the inequality

$$(3.6) \quad 0 \leq \|x\| + \|y\| - \|x + y\| \leq \frac{\sqrt{M} - \sqrt{m}}{\sqrt[4]{mM}} \sqrt{\operatorname{Re} \langle x, y \rangle}.$$

Proof. Choosing in (2.8), $a = \frac{M+m}{2}y$, $r = \frac{1}{2}(M - m)\|y\|$ we get


$$\|x\| \|y\| \sqrt{Mm} \leq \frac{M + m}{2} \operatorname{Re} \langle x, y \rangle$$

giving

$$0 \leq \|x\| \|y\| - \operatorname{Re} \langle x, y \rangle \leq \frac{(\sqrt{M} - \sqrt{m})^2}{2\sqrt{mM}} \operatorname{Re} \langle x, y \rangle.$$

Following the same arguments as in the proof of Proposition 3.1, we deduce the desired inequality (3.6). \square

For some results related to triangle inequality in inner product spaces, see [3], [17], [18] and [19].



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 18 of 40

4. Some Grüss Type Inequalities

We may state the following result.

Theorem 4.1. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over the real or complex number field \mathbb{K} ($\mathbb{K} = \mathbb{R}, \mathbb{K} = \mathbb{C}$) and $x, y, e \in H$ with $\|e\| = 1$. If $r_1, r_2 \in (0, 1)$ and*

$$(4.1) \quad \|x - e\| \leq r_1, \quad \|y - e\| \leq r_2,$$

then we have the inequality

$$(4.2) \quad |\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle| \leq r_1 r_2 \|x\| \|y\|.$$

The inequality (4.2) is sharp in the sense that the constant 1 in front of $r_1 r_2$ cannot be replaced by a smaller quantity.

Proof. Apply Schwarz's inequality in $(H; \langle \cdot, \cdot \rangle)$ for the vectors $x - \langle x, e \rangle e$, $y - \langle y, e \rangle e$, to get (see also [9])


$$(4.3) \quad |\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle|^2 \leq (\|x\|^2 - |\langle x, e \rangle|^2) (\|y\|^2 - |\langle y, e \rangle|^2).$$

Using Theorem 2.1 for $a = e$, we may state that

$$(4.4) \quad \|x\|^2 - |\langle x, e \rangle|^2 \leq r_1^2 \|x\|^2, \quad \|y\|^2 - |\langle y, e \rangle|^2 \leq r_2^2 \|y\|^2.$$

Utilizing (4.3) and (4.4), we deduce

$$(4.5) \quad |\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle|^2 \leq r_1^2 r_2^2 \|x\|^2 \|y\|^2,$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 19 of 40

which is clearly equivalent to the desired inequality (4.2).

The sharpness of the constant follows by the fact that for $x = y$, $r_1 = r_2 = r$, we get from (4.2) that

$$(4.6) \quad \|x\|^2 - |\langle x, e \rangle|^2 \leq r^2 \|x\|^2,$$

provided $\|e\| = 1$ and $\|x - e\| \leq r < 1$. The inequality (4.6) is sharp, as shown in Theorem 2.1, and the proof is completed. \square

Another companion of the Grüss inequality may be stated as well.

Theorem 4.2. *Let $(H; \langle \cdot, \cdot \rangle)$ be an inner product space over \mathbb{K} and $x, y, e \in H$ with $\|e\| = 1$. Suppose also that $a, A, b, B \in \mathbb{K}$ ($\mathbb{K} = \mathbb{R}, \mathbb{C}$) such that $\operatorname{Re}(A\bar{a})$, $\operatorname{Re}(B\bar{b}) > 0$. If either*

$$(4.7) \quad \operatorname{Re} \langle Ae - x, x - ae \rangle \geq 0, \quad \operatorname{Re} \langle Be - y, y - be \rangle \geq 0,$$


or, equivalently,

$$(4.8) \quad \left\| x - \frac{a + A}{2} e \right\| \leq \frac{1}{2} |A - a|, \quad \left\| y - \frac{b + B}{2} e \right\| \leq \frac{1}{2} |B - b|,$$

holds, then we have the inequality

$$(4.9) \quad |\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle| \leq \frac{1}{4} \cdot \frac{|A - a| |B - b|}{\sqrt{\operatorname{Re}(A\bar{a}) \operatorname{Re}(B\bar{b})}} |\langle x, e \rangle \langle e, y \rangle|.$$

The constant $\frac{1}{4}$ is best possible.



Proof. We know, by (4.3), that

$$(4.10) \quad |\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle|^2 \leq (\|x\|^2 - |\langle x, e \rangle|^2) (\|y\|^2 - |\langle y, e \rangle|^2).$$

If we use Corollary 2.3, then we may state that

$$(4.11) \quad \|x\|^2 - |\langle x, e \rangle|^2 \leq \frac{1}{4} \cdot \frac{|A - a|^2}{\operatorname{Re}(A\bar{a})} |\langle x, e \rangle|^2$$

and

$$(4.12) \quad \|y\|^2 - |\langle y, e \rangle|^2 \leq \frac{1}{4} \cdot \frac{|B - b|^2}{\operatorname{Re}(B\bar{b})} |\langle y, e \rangle|^2.$$

Utilizing (4.10) – (4.12), we deduce

$$|\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle|^2 \leq \frac{1}{16} \cdot \frac{|A - a|^2 |B - b|^2}{\operatorname{Re}(A\bar{a}) \operatorname{Re}(B\bar{b})} |\langle x, e \rangle \langle e, y \rangle|^2,$$


which is clearly equivalent to the desired inequality (4.9).

The sharpness of the constant follows from Corollary 2.3, and we omit the details. \square

Remark 4.1. *With the assumptions of Theorem 4.2 and if $\langle x, e \rangle, \langle y, e \rangle \neq 0$ (that is actually the interesting case), then one has the inequality*

$$(4.13) \quad \left| \frac{\langle x, y \rangle}{\langle x, e \rangle \langle e, y \rangle} - 1 \right| \leq \frac{1}{4} \cdot \frac{|A - a| |B - b|}{\sqrt{\operatorname{Re}(A\bar{a}) \operatorname{Re}(B\bar{b})}}.$$

The constant $\frac{1}{4}$ is best possible.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit


Page 21 of 40

Remark 4.2. *The inequality (4.9) provides a better bound for the quantity*

$$|\langle x, y \rangle - \langle x, e \rangle \langle e, y \rangle|$$

than (2.3) of [9].

For some recent results on Grüss type inequalities in inner product spaces, see [4], [6] and [20].



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 22 of 40

5. Reverses of Bessel's Inequality

Let $(H; \langle \cdot, \cdot \rangle)$ be a real or complex infinite dimensional Hilbert space and $(e_i)_{i \in \mathbb{N}}$ an orthonormal family in H , i.e., we recall that $\langle e_i, e_j \rangle = 0$ if $i, j \in \mathbb{N}$, $i \neq j$ and $\|e_i\| = 1$ for $i \in \mathbb{N}$.

It is well known that, if $x \in H$, then the series $\sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2$ is convergent and the following inequality, called *Bessel's inequality*

$$(5.1) \quad \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2 \leq \|x\|^2,$$

holds.

If $\ell^2(\mathbb{K}) := \{ \mathbf{a} = (a_i)_{i \in \mathbb{N}} \subset \mathbb{K} \mid \sum_{i=1}^{\infty} |a_i|^2 < \infty \}$, where $\mathbb{K} = \mathbb{C}$ or $\mathbb{K} = \mathbb{R}$, is the Hilbert space of all complex or real sequences that are 2-summable and $\boldsymbol{\lambda} = (\lambda_i)_{i \in \mathbb{N}} \in \ell^2(\mathbb{K})$, then the series $\sum_{i=1}^{\infty} \lambda_i e_i$ is convergent in H and if $y := \sum_{i=1}^{\infty} \lambda_i e_i \in H$, then $\|y\| = (\sum_{i=1}^{\infty} |\lambda_i|^2)^{\frac{1}{2}}$.


We may state the following result.

Theorem 5.1. *Let $(H; \langle \cdot, \cdot \rangle)$ be an infinite dimensional Hilbert space over the real or complex number field \mathbb{K} , $(e_i)_{i \in \mathbb{N}}$ an orthonormal family in H , $\boldsymbol{\lambda} = (\lambda_i)_{i \in \mathbb{N}} \in \ell^2(\mathbb{K})$ and $r > 0$ with the property that*

$$(5.2) \quad \sum_{i=1}^{\infty} |\lambda_i|^2 > r^2.$$

If $x \in H$ is such that

$$(5.3) \quad \left\| x - \sum_{i=1}^{\infty} \lambda_i e_i \right\| \leq r,$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 23 of 40

then we have the inequality

$$\begin{aligned}
 (5.4) \quad \|x\|^2 &\leq \frac{\left(\sum_{i=1}^{\infty} \operatorname{Re} [\bar{\lambda}_i \langle x, e_i \rangle]\right)^2}{\sum_{i=1}^{\infty} |\lambda_i|^2 - r^2} \\
 &\leq \frac{\left|\sum_{i=1}^{\infty} \bar{\lambda}_i \langle x, e_i \rangle\right|^2}{\sum_{i=1}^{\infty} |\lambda_i|^2 - r^2} \\
 &\leq \frac{\sum_{i=1}^{\infty} |\lambda_i|^2}{\sum_{i=1}^{\infty} |\lambda_i|^2 - r^2} \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2;
 \end{aligned}$$


and

$$(5.5) \quad 0 \leq \|x\|^2 - \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2$$

$$(5.6) \quad \leq \frac{r^2}{\sum_{i=1}^{\infty} |\lambda_i|^2 - r^2} \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2.$$

Proof. Applying the third inequality in (2.2) for $a = \sum_{i=1}^{\infty} \lambda_i e_i \in H$, we have

$$(5.7) \quad \|x\|^2 \left\| \sum_{i=1}^{\infty} \lambda_i e_i \right\|^2 - \left[\operatorname{Re} \left\langle x, \sum_{i=1}^{\infty} \lambda_i e_i \right\rangle \right]^2 \leq r^2 \|x\|^2$$



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 24 of 40

and since

$$\left\| \sum_{i=1}^{\infty} \lambda_i e_i \right\|^2 = \sum_{i=1}^{\infty} |\lambda_i|^2,$$

$$\operatorname{Re} \left\langle x, \sum_{i=1}^{\infty} \lambda_i e_i \right\rangle = \sum_{i=1}^{\infty} \operatorname{Re} [\bar{\lambda}_i \langle x, e_i \rangle],$$

hence, by (5.7) we deduce

$$\|x\|^2 \sum_{i=1}^{\infty} |\lambda_i|^2 - \left[\operatorname{Re} \left\langle x, \sum_{i=1}^{\infty} \lambda_i e_i \right\rangle \right]^2 \leq r^2 \|x\|^2,$$

giving the first inequality in (5.4).


The second inequality is obvious by the modulus property.

The last inequality follows by the Cauchy-Bunyakovsky-Schwarz inequality

$$\left| \sum_{i=1}^{\infty} \bar{\lambda}_i \langle x, e_i \rangle \right|^2 \leq \sum_{i=1}^{\infty} |\lambda_i|^2 \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2.$$

The inequality (5.5) follows by the last inequality in (5.4) on subtracting in both sides the quantity $\sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2 < \infty$. \square

The following result provides a generalization for the reverse of Bessel's inequality obtained in [12].



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 25 of 40

Theorem 5.2. Let $(H; \langle \cdot, \cdot \rangle)$ and $(e_i)_{i \in \mathbb{N}}$ be as in Theorem 5.1. Suppose that $\Gamma = (\Gamma_i)_{i \in \mathbb{N}} \in \ell^2(\mathbb{K})$, $\gamma = (\gamma_i)_{i \in \mathbb{N}} \in \ell^2(\mathbb{K})$ are sequences of real or complex numbers such that

$$(5.8) \quad \sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i) > 0.$$

If $x \in H$ is such that either


$$(5.9) \quad \left\| x - \sum_{i=1}^{\infty} \frac{\Gamma_i + \gamma_i}{2} e_i \right\| \leq \frac{1}{2} \left(\sum_{i=1}^{\infty} |\Gamma_i - \gamma_i|^2 \right)^{\frac{1}{2}}$$

or, equivalently,

$$(5.10) \quad \operatorname{Re} \left\langle \sum_{i=1}^{\infty} \Gamma_i e_i - x, x - \sum_{i=1}^{\infty} \gamma_i e_i \right\rangle \geq 0$$

holds, then we have the inequalities

$$(5.11) \quad \begin{aligned} \|x\|^2 &\leq \frac{1}{4} \cdot \frac{(\sum_{i=1}^{\infty} \operatorname{Re}[(\bar{\Gamma}_i + \bar{\gamma}_i) \langle x, e_i \rangle])^2}{\sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i)} \\ &\leq \frac{1}{4} \cdot \frac{|\sum_{i=1}^{\infty} (\bar{\Gamma}_i + \bar{\gamma}_i) \langle x, e_i \rangle|^2}{\sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i)} \\ &\leq \frac{1}{4} \cdot \frac{\sum_{i=1}^{\infty} |\Gamma_i + \gamma_i|^2}{\sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i)} \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2. \end{aligned}$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 26 of 40

The constant $\frac{1}{4}$ is best possible in all inequalities in (5.11).

We also have the inequalities:

$$(5.12) \quad 0 \leq \|x\|^2 - \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2 \leq \frac{1}{4} \cdot \frac{\sum_{i=1}^{\infty} |\Gamma_i - \gamma_i|^2}{\sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i)} \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2.$$

Here the constant $\frac{1}{4}$ is also best possible.

Proof. Since $\Gamma, \gamma \in \ell^2(\mathbb{K})$, then also $\frac{1}{2}(\Gamma \pm \gamma) \in \ell^2(\mathbb{K})$, showing that the series

$$\sum_{i=1}^{\infty} \left| \frac{\Gamma_i + \gamma_i}{2} \right|^2, \quad \sum_{i=1}^{\infty} \left| \frac{\Gamma_i - \gamma_i}{2} \right|^2 \quad \text{and} \quad \sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i)$$

are convergent. Also, the series


$$\sum_{i=1}^{\infty} \Gamma_i e_i, \quad \sum_{i=1}^{\infty} \gamma_i e_i \quad \text{and} \quad \sum_{i=1}^{\infty} \frac{\gamma_i + \Gamma_i}{2} e_i$$

are convergent in the Hilbert space H .

The equivalence of the conditions (5.9) and (5.10) follows by the fact that in an inner product space we have, for $x, z, Z \in H$, $\operatorname{Re} \langle Z - x, x - z \rangle \geq 0$ is equivalent to $\|x - \frac{z+Z}{2}\| \leq \frac{1}{2} \|Z - z\|$, and we omit the details.

Now, we observe that the inequalities (5.11) and (5.12) follow from Theorem 5.1 on choosing $\lambda_i = \frac{\gamma_i + \Gamma_i}{2}$, $i \in \mathbb{N}$ and $r = \frac{1}{2} \left(\sum_{i=1}^{\infty} |\Gamma_i - \gamma_i|^2 \right)^{\frac{1}{2}}$.

The fact that $\frac{1}{4}$ is the best constant in both (5.11) and (5.12) follows from Theorem 2.2 and Corollary 2.3, and we omit the details. \square



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back


Close

Quit

Page 27 of 40

Remark 5.1. Note that (5.11) improves (1.17) and (5.12) improves (1.18), that have been obtained in [12].

For some recent results related to Bessel inequality, see [1], [5], [14], and [16].



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 28 of 40

6. Some Grüss Type Inequalities for Orthonormal Families

The following result related to Grüss inequality in inner product spaces, holds.

Theorem 6.1. *Let $(H; \langle \cdot, \cdot \rangle)$ be an infinite dimensional Hilbert space over the real or complex number field \mathbb{K} , and $(e_i)_{i \in \mathbb{N}}$ an orthonormal family in H . Assume that $\lambda = (\lambda_i)_{i \in \mathbb{N}}$, $\mu = (\mu_i)_{i \in \mathbb{N}} \in \ell^2(\mathbb{K})$ and $r_1, r_2 > 0$ with the properties that*


$$(6.1) \quad \sum_{i=1}^{\infty} |\lambda_i|^2 > r_1^2, \quad \sum_{i=1}^{\infty} |\mu_i|^2 > r_2^2.$$

If $x, y \in H$ are such that

$$(6.2) \quad \left\| x - \sum_{i=1}^{\infty} \lambda_i e_i \right\| \leq r_1, \quad \left\| y - \sum_{i=1}^{\infty} \mu_i e_i \right\| \leq r_2,$$

then we have the inequalities

$$(6.3) \quad \left| \langle x, y \rangle - \sum_{i=1}^{\infty} \langle x, e_i \rangle \langle e_i, y \rangle \right| \leq \frac{r_1 r_2}{\sqrt{\sum_{i=1}^{\infty} |\lambda_i|^2 - r_1^2} \sqrt{\sum_{i=1}^{\infty} |\mu_i|^2 - r_2^2}} \cdot \sqrt{\sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2 \sum_{i=1}^{\infty} |\langle y, e_i \rangle|^2}$$



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 29 of 40



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 30 of 40

$$\leq \frac{r_1 r_2 \|x\| \|y\|}{\sqrt{\sum_{i=1}^{\infty} |\lambda_i|^2 - r_1^2} \sqrt{\sum_{i=1}^{\infty} |\mu_i|^2 - r_2^2}}.$$

Proof. Applying Schwarz's inequality for the vectors $x - \sum_{i=1}^{\infty} \langle x, e_i \rangle e_i$, $y - \sum_{i=1}^{\infty} \langle y, e_i \rangle e_i$, we have

$$(6.4) \quad \left| \left\langle x - \sum_{i=1}^{\infty} \langle x, e_i \rangle e_i, y - \sum_{i=1}^{\infty} \langle y, e_i \rangle e_i \right\rangle \right|^2 \leq \left\| x - \sum_{i=1}^{\infty} \langle x, e_i \rangle e_i \right\|^2 \left\| y - \sum_{i=1}^{\infty} \langle y, e_i \rangle e_i \right\|^2.$$

Since

$$\left\langle x - \sum_{i=1}^{\infty} \langle x, e_i \rangle e_i, y - \sum_{i=1}^{\infty} \langle y, e_i \rangle e_i \right\rangle = \langle x, y \rangle - \sum_{i=1}^{\infty} \langle x, e_i \rangle \langle e_i, y \rangle$$

and

$$\left\| x - \sum_{i=1}^{\infty} \langle x, e_i \rangle e_i \right\|^2 = \|x\|^2 - \sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2,$$

hence, by (5.5) applied for x and y , and from (6.4), we deduce the first part of (6.3). □

The second part follows by Bessel's inequality.

The following Grüss type inequality may be stated as well.

Theorem 6.2. Let $(H; \langle \cdot, \cdot \rangle)$ be an infinite dimensional Hilbert space and $(e_i)_{i \in \mathbb{N}}$ an orthonormal family in H . Suppose that $(\Gamma_i)_{i \in \mathbb{N}}, (\gamma_i)_{i \in \mathbb{N}}, (\phi_i)_{i \in \mathbb{N}}, (\Phi_i)_{i \in \mathbb{N}} \in \ell^2(\mathbb{K})$ are sequences of real and complex numbers such that

$$(6.5) \quad \sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \overline{\gamma_i}) > 0, \quad \sum_{i=1}^{\infty} \operatorname{Re}(\Phi_i \overline{\phi_i}) > 0.$$

If $x, y \in H$ are such that either


$$(6.6) \quad \left\| x - \sum_{i=1}^{\infty} \frac{\Gamma_i + \gamma_i}{2} \cdot e_i \right\| \leq \frac{1}{2} \left(\sum_{i=1}^{\infty} |\Gamma_i - \gamma_i|^2 \right)^{\frac{1}{2}}$$

$$\left\| y - \sum_{i=1}^{\infty} \frac{\Phi_i + \phi_i}{2} \cdot e_i \right\| \leq \frac{1}{2} \left(\sum_{i=1}^{\infty} |\Phi_i - \phi_i|^2 \right)^{\frac{1}{2}}$$

or, equivalently,

$$(6.7) \quad \operatorname{Re} \left\langle \sum_{i=1}^{\infty} \Gamma_i e_i - x, x - \sum_{i=1}^{\infty} \gamma_i e_i \right\rangle \geq 0,$$

$$\operatorname{Re} \left\langle \sum_{i=1}^{\infty} \Phi_i e_i - y, y - \sum_{i=1}^{\infty} \phi_i e_i \right\rangle \geq 0,$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 31 of 40

holds, then we have the inequality


$$\begin{aligned}
 (6.8) \quad & \left| \langle x, y \rangle - \sum_{i=1}^{\infty} \langle x, e_i \rangle \langle e_i, y \rangle \right| \\
 & \leq \frac{1}{4} \cdot \frac{(\sum_{i=1}^{\infty} |\Gamma_i - \gamma_i|^2)^{\frac{1}{2}} (\sum_{i=1}^{\infty} |\Phi_i - \phi_i|^2)^{\frac{1}{2}}}{(\sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i))^{\frac{1}{2}} (\sum_{i=1}^{\infty} \operatorname{Re}(\Phi_i \bar{\phi}_i))^{\frac{1}{2}}} \\
 & \quad \times \left(\sum_{i=1}^{\infty} |\langle x, e_i \rangle|^2 \right)^{\frac{1}{2}} \left(\sum_{i=1}^{\infty} |\langle y, e_i \rangle|^2 \right)^{\frac{1}{2}} \\
 & \leq \frac{1}{4} \cdot \frac{(\sum_{i=1}^{\infty} |\Gamma_i - \gamma_i|^2)^{\frac{1}{2}} (\sum_{i=1}^{\infty} |\Phi_i - \phi_i|^2)^{\frac{1}{2}}}{[\sum_{i=1}^{\infty} \operatorname{Re}(\Gamma_i \bar{\gamma}_i)]^{\frac{1}{2}} [\sum_{i=1}^{\infty} \operatorname{Re}(\Phi_i \bar{\phi}_i)]^{\frac{1}{2}}} \|x\| \|y\|.
 \end{aligned}$$

The constant $\frac{1}{4}$ is best possible in the first inequality.

Proof. Follows by (5.12) and (6.4).

The best constant follows from Theorem 4.2, and we omit the details. \square

Remark 6.1. We note that the inequality (6.8) is better than the inequality (3.3) in [12]. We omit the details.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 32 of 40

7. Integral Inequalities

Let (Ω, Σ, μ) be a measurable space consisting of a set Ω , a σ -algebra of parts Σ and a countably additive and positive measure μ on Σ with values in $\mathbb{R} \cup \{\infty\}$. Let $\rho \geq 0$ be a Lebesgue measurable function on Ω with $\int_{\Omega} \rho(s) d\mu(s) = 1$. Denote by $L_{\rho}^2(\Omega, \mathbb{K})$ the Hilbert space of all real or complex valued functions defined on Ω and $2 - \rho$ -integrable on Ω , i.e.,

$$(7.1) \quad \int_{\Omega} \rho(s) |f(s)|^2 d\mu(s) < \infty.$$

It is obvious that the following inner product

$$(7.2) \quad \langle f, g \rangle_{\rho} := \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s),$$


generates the norm $\|f\|_{\rho} := \left(\int_{\Omega} \rho(s) |f(s)|^2 d\mu(s) \right)^{\frac{1}{2}}$ of $L_{\rho}^2(\Omega, \mathbb{K})$, and all the above results may be stated for integrals.

It is important to observe that, if

$$(7.3) \quad \operatorname{Re} \left[f(s) \overline{g(s)} \right] \geq 0 \text{ for } \mu - \text{a.e. } s \in \Omega,$$

then, obviously,

$$(7.4) \quad \begin{aligned} \operatorname{Re} \langle f, g \rangle_{\rho} &= \operatorname{Re} \left[\int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right] \\ &= \int_{\Omega} \rho(s) \operatorname{Re} \left[f(s) \overline{g(s)} \right] d\mu(s) \geq 0. \end{aligned}$$



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 33 of 40

The reverse is evidently not true in general.

Moreover, if the space is real, i.e., $\mathbb{K} = \mathbb{R}$, then a sufficient condition for (7.4) to hold is:

$$(7.5) \quad f(s) \geq 0, \quad g(s) \geq 0 \text{ for } \mu - \text{a.e. } s \in \Omega.$$

We provide now, by the use of certain results obtained in Section 2, some integral inequalities that may be used in practical applications.


Proposition 7.1. *Let $f, g \in L^2_\rho(\Omega, \mathbb{K})$ and $r > 0$ with the properties that*

$$(7.6) \quad |f(s) - g(s)| \leq r \leq |g(s)| \text{ for } \mu - \text{a.e. } s \in \Omega.$$

Then we have the inequalities

$$(7.7) \quad \begin{aligned} 0 &\leq \int_{\Omega} \rho(s) |f(s)|^2 d\mu(s) \int_{\Omega} \rho(s) |g(s)|^2 d\mu(s) \\ &\quad - \left| \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right|^2 \\ &\leq \int_{\Omega} \rho(s) |f(s)|^2 d\mu(s) \int_{\Omega} \rho(s) |g(s)|^2 d\mu(s) \\ &\quad - \left[\int_{\Omega} \rho(s) \operatorname{Re} \left(f(s) \overline{g(s)} \right) d\mu(s) \right]^2 \\ &\leq r^2 \int_{\Omega} \rho(s) |g(s)|^2 d\mu(s). \end{aligned}$$

The constant 1 in front of r^2 is best possible.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 34 of 40

The proof follows by Theorem 2.1 and we omit the details.

Proposition 7.2. Let $f, g \in L^2_\rho(\Omega, \mathbb{K})$ and $\gamma, \Gamma \in \mathbb{K}$ such that $\text{Re}(\Gamma\bar{\gamma}) > 0$ and

$$(7.8) \quad \text{Re} \left[(\Gamma g(s) - f(s)) \left(\overline{f(s)} - \bar{\gamma} \overline{g(s)} \right) \right] \geq 0 \text{ for } \mu - \text{a.e. } s \in \Omega.$$

Then we have the inequalities

$$(7.9) \quad \int_{\Omega} \rho(s) |f(s)|^2 d\mu(s) \int_{\Omega} \rho(s) |g(s)|^2 d\mu(s) \leq \frac{1}{4} \cdot \frac{\left\{ \text{Re} \left[(\bar{\Gamma} + \bar{\gamma}) \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right] \right\}^2}{\text{Re}(\Gamma\bar{\gamma})} \leq \frac{1}{4} \cdot \frac{|\Gamma + \gamma|^2}{\text{Re}(\Gamma\bar{\gamma})} \left| \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right|^2.$$


The constant $\frac{1}{4}$ is best possible in both inequalities.

The proof follows by Theorem 2.2 and we omit the details.

Corollary 7.3. With the assumptions of Proposition 7.2, we have the inequality

$$(7.10) \quad 0 \leq \int_{\Omega} \rho(s) |f(s)|^2 d\mu(s) \int_{\Omega} \rho(s) |g(s)|^2 d\mu(s) - \left| \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right|^2 \leq \frac{1}{4} \cdot \frac{|\Gamma - \gamma|^2}{\text{Re}(\Gamma\bar{\gamma})} \left| \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right|^2.$$

The constant $\frac{1}{4}$ is best possible.



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 35 of 40

Remark 7.1. *If the space is real and we assume, for $M \geq m > 0$, that*

$$(7.11) \quad mg(s) \leq f(s) \leq Mg(s), \text{ for } \mu - \text{a.e. } s \in \Omega,$$

then, by (7.9) and (7.10), we deduce the inequalities

$$(7.12) \quad \int_{\Omega} \rho(s) [f(s)]^2 d\mu(s) \int_{\Omega} \rho(s) [g(s)]^2 d\mu(s) \\ \leq \frac{1}{4} \cdot \frac{(M+m)^2}{mM} \left[\int_{\Omega} \rho(s) f(s) g(s) d\mu(s) \right]^2.$$


and

$$(7.13) \quad 0 \leq \int_{\Omega} \rho(s) [f(s)]^2 d\mu(s) \int_{\Omega} \rho(s) [g(s)]^2 d\mu(s) \\ - \left[\int_{\Omega} \rho(s) f(s) g(s) d\mu(s) \right]^2 \\ \leq \frac{1}{4} \cdot \frac{(M-m)^2}{mM} \left[\int_{\Omega} \rho(s) f(s) g(s) d\mu(s) \right]^2.$$

The inequality (7.12) is known in the literature as Cassel's inequality.

The following Grüss type integral inequality for real or complex-valued functions also holds.

Proposition 7.4. *Let $f, g, h \in L^2_{\rho}(\Omega, \mathbb{K})$ with $\int_{\Omega} \rho(s) |h(s)|^2 d\mu(s) = 1$ and*



**Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces**

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 36 of 40

$a, A, b, B \in \mathbb{K}$ such that $\operatorname{Re}(A\bar{a}), \operatorname{Re}(B\bar{b}) > 0$ and

$$\operatorname{Re} \left[(Ah(s) - f(s)) \left(\overline{f(s)} - \overline{ah(s)} \right) \right] \geq 0,$$

$$\operatorname{Re} \left[(Bh(s) - g(s)) \left(\overline{g(s)} - \overline{bh(s)} \right) \right] \geq 0,$$


for μ -a.e. $s \in \Omega$. Then we have the inequalities

$$\begin{aligned} & \left| \int_{\Omega} \rho(s) f(s) \overline{g(s)} d\mu(s) \right. \\ & \quad \left. - \int_{\Omega} \rho(s) f(s) \overline{h(s)} d\mu(s) \int_{\Omega} \rho(s) h(s) \overline{g(s)} d\mu(s) \right| \\ & \leq \frac{1}{4} \cdot \frac{|A - a| |B - b|}{\sqrt{\operatorname{Re}(A\bar{a}) \operatorname{Re}(B\bar{b})}} \\ & \quad \times \left| \int_{\Omega} \rho(s) f(s) \overline{h(s)} d\mu(s) \int_{\Omega} \rho(s) h(s) \overline{g(s)} d\mu(s) \right| \end{aligned}$$

The constant $\frac{1}{4}$ is best possible.

The proof follows by Theorem 4.2.

Remark 7.2. All the other inequalities in Sections 3–6 may be used in a similar way to obtain the corresponding integral inequalities. We omit the details.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back


Close

Quit

Page 37 of 40

References

- [1] X.H. CAO, Bessel sequences in a Hilbert space, *Gongcheng Shuxue Xuebao*, **17**(2) (2000), 92–98.
- [2] A. De ROSSI, A strengthened Cauchy-Schwarz inequality for biorthogonal wavelets, *Math. Inequal. Appl.*, **2**(2) (1999), 263–282.
- [3] J.B. DIAZ AND F.T. METCALF, A complementary triangle inequality in Hilbert and Banach spaces, *Proc. Amer. Math. Soc.*, **17** (1966), 88–97.
- [4] S.S. DRAGOMIR, A generalization of Grüss inequality in inner product spaces and applications. *J. Math. Anal. Appl.*, **237**(1) (1999), 74–82.
- [5] S.S. DRAGOMIR, A note on Bessel's inequality, *Austral. Math. Soc. Gaz.*, **28**(5) (2001), 246–248.
- [6] S.S. DRAGOMIR, Some Grüss type inequalities in inner product spaces, *J. Inequal. Pure & Appl. Math.*, **4**(2) (2003), Article 42, [ONLINE <http://jipam.vu.edu.au/article.php?sid=280>].
- [7] S.S. DRAGOMIR, A counterpart of Schwarz's inequality in inner product spaces, *RGMA Res. Rep. Coll.*, **6**(2003), *Supplement*, Article 18, [ONLINE [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html)].
- [8] S.S. DRAGOMIR, A generalisation of the Cassels and Grueb-Reinboldt inequalities in inner product spaces, Preprint, *Mathematics ArXiv*, math.CA/0307130, [ONLINE <http://front.math.ucdavis.edu/>]. *Nonlinear Analysis Forum*, **8**(2) (2003), 169–178.



Reverses of Schwarz, Triangle and Bessel Inequalities in Inner Product Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 38 of 40

- [9] S.S. DRAGOMIR, Some companions of the Grüss inequality in inner product spaces, *J. Inequal. Pure & Appl. Math.*, **4**(5) (2003), Article 87, [ONLINE <http://jipam.vu.edu.au/article.php?sid=328>].
- [10] S.S. DRAGOMIR, On Bessel and Grüss inequalities for orthonormal families in inner product spaces, *RGMIA Res. Rep. Coll.*, **6**(2003), *Supplement*, Article 12, [ONLINE [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html)]. *Bull. Austral. Math. Soc.*, **69**(2) (2004), 327–340.
- [11] S.S. DRAGOMIR, A counterpart of Bessel’s inequality in inner product spaces and some Grüss type related results, *RGMIA Res. Rep. Coll.*, **6**(2003), *Supplement*, Article 10, [ONLINE [http://rgmia.vu.edu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html)]
- [12] S.S. DRAGOMIR, Some new results related to Bessel and Grüss inequalities for orthonormal families in inner product spaces, *RGMIA Res. Rep. Coll.*, **6**(2003), *Supplement*, Article 13, [ONLINE [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html)].
- [13] S.S. DRAGOMIR AND B. MOND, On the superadditivity and monotonicity of Schwarz’s inequality in inner product spaces, *Makedon. Akad. Nauk. Umet. Oddel. Mat.-Tehn. Nauk. Prilozi*, **15**(2) (1994), 5–22 (1996).
- [14] S.S. DRAGOMIR AND J. SÁNDOR, On Bessel’s and Gram’s inequalities in pre-Hilbertian spaces, *Period. Math. Hungar.*, **29**(3) (1994), 197–205.
- [15] H. GUNAWAN, On n -inner products, n -norms, and the Cauchy-Schwarz inequality, *Sci. Math. Jpn.*, **55**(1) (2002), 53–60.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents




Go Back

Close

Quit

Page 39 of 40

- [16] H. GUNAWAN, A generalization of Bessel's inequality and Parseval's identity, *Period. Math. Hungar.*, **44**(2) (2002), 177–181.
- [17] S.M. KHALEELULLA, On Diaz-Metcalf's complementary triangle inequality, *Kyungpook Math. J.*, **15** (1975), 9–11.
- [18] P.M. MILIČIĆ, On a complementary inequality of the triangle inequality (French), *Mat. Vesnik*, **41**(2) (1989), 83–88.
- [19] D.K. RAO, A triangle inequality for angles in a Hilbert space, *Rev. Colombiana Mat.*, **10**(3) (1976), 95–97.
- [20] P.F. RENAUD, A matrix formulation of Grüss inequality, *Linear Algebra Appl.*, **335** (2001), 95–100.
- [21] N. UJEVIĆ, A new generalisation of Grüss inequality in inner product spaces, *Math. Inequal. & Appl.*, **6**(4) (2003), 617–623.



Reverses of Schwarz, Triangle
and Bessel Inequalities in Inner
Product Spaces

S.S. Dragomir

Title Page

Contents



Go Back

Close

Quit

Page 40 of 40