

SOME PROPERTIES FOR AN INTEGRAL OPERATOR DEFINED BY AL-BOUDI DIFFERENTIAL OPERATOR

SERAP BULUT

Kocaeli University
Civil Aviation College
Arslanbey Campus
41285 İzmit-Kocaeli, TURKEY
EMail: serap.bulut@kocaeli.edu.tr

Received: 08 July, 2008

Accepted: 12 November, 2008

Communicated by: S.S. Dragomir

2000 AMS Sub. Class.: Primary 30C45.

Key words: Analytic functions, Differential operator.

Abstract: In this paper, we investigate some properties for an integral operator defined by Al-Oboudi differential operator.





Some Properties for
an Integral Operator

Serap Bulut

vol. 9, iss. 4, art. 115, 2008

[Title Page](#)

[Contents](#)



Page 1 of 12

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

- | | | |
|---|--|----|
| 1 | Introduction | 3 |
| 2 | Some Properties for I_n on the Class $\mathcal{S}^n(\delta, \alpha)$ | 7 |
| 3 | Some Properties for I_n on the class $\mathcal{M}^n(\delta, \beta)$ | 10 |



Some Properties for
an Integral Operator

Serap Bulut

vol. 9, iss. 4, art. 115, 2008

Title Page

Contents



Page 2 of 12


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 3 of 12

Go Back

Full Screen

Close

1. Introduction

Let \mathcal{A} denote the class of all functions of the form

$$(1.1) \quad f(z) = z + \sum_{j=2}^{\infty} a_j z^j$$

which are analytic in the open unit disk $\mathbb{U} := \{z \in \mathbb{C} : |z| < 1\}$, and $\mathcal{S} := \{f \in \mathcal{A} : f \text{ is univalent in } \mathbb{U}\}$.

For $f \in \mathcal{A}$, Al-Oboudi [2] introduced the following operator:

$$(1.2) \quad D^0 f(z) = f(z),$$

$$(1.3) \quad D^1 f(z) = (1 - \delta)f(z) + \delta z f'(z) = D_\delta f(z), \quad \delta \geq 0,$$

$$(1.4) \quad D^n f(z) = D_\delta(D^{n-1} f(z)), \quad (n \in \mathbb{N} := \{1, 2, 3, \dots\}).$$

If f is given by (1.1), then from (1.3) and (1.4) we see that

$$(1.5) \quad D^n f(z) = z + \sum_{j=2}^{\infty} [1 + (j-1)\delta]^n a_j z^j, \quad (n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}),$$

with $D^n f(0) = 0$.

When $\delta = 1$, we get Sălăgean's differential operator [10].

A function $f \in \mathcal{A}$ is said to be *starlike of order* α if it satisfies the inequality:

$$\operatorname{Re} \left\{ \frac{z f'(z)}{f(z)} \right\} > \alpha \quad (z \in \mathbb{U})$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 12

Go Back

Full Screen

Close

for some $0 \leq \alpha < 1$. We say that f is in the class $\mathcal{S}^*(\alpha)$ for such functions.

A function $f \in \mathcal{A}$ is said to be *convex of order* α if it satisfies the inequality:

$$\operatorname{Re} \left\{ \frac{zf''(z)}{f'(z)} + 1 \right\} > \alpha \quad (z \in \mathbb{U})$$

for some $0 \leq \alpha < 1$. We say that f is in the class $\mathcal{K}(\alpha)$ if it is convex of order α in \mathbb{U} .

We note that $f \in \mathcal{K}(\alpha)$ if and only if $zf' \in \mathcal{S}^*(\alpha)$.

In particular, the classes

$$\mathcal{S}^*(0) := \mathcal{S}^* \quad \text{and} \quad \mathcal{K}(0) := \mathcal{K}$$

are familiar classes of starlike and convex functions in \mathbb{U} , respectively.

Now, we introduce two new classes $\mathcal{S}^n(\delta, \alpha)$ and $\mathcal{M}^n(\delta, \beta)$ as follows:

Let $\mathcal{S}^n(\delta, \alpha)$ denote the class of functions $f \in \mathcal{A}$ which satisfy the condition

$$\operatorname{Re} \left\{ \frac{z(D^n f(z))'}{D^n f(z)} \right\} > \alpha \quad (z \in \mathbb{U})$$

for some $0 \leq \alpha < 1$, $\delta \geq 0$, and $n \in \mathbb{N}_0$.


It is clear that

$$\mathcal{S}^0(\delta, \alpha) \equiv \mathcal{S}^*(\alpha) \equiv \mathcal{S}^n(0, \alpha), \quad \mathcal{S}^n(0, 0) \equiv \mathcal{S}^*.$$

Let $\mathcal{M}^n(\delta, \beta)$ be the subclass of \mathcal{A} , consisting of the functions f , which satisfy the inequality

$$\operatorname{Re} \left\{ \frac{z(D^n f(z))'}{D^n f(z)} \right\} < \beta \quad (z \in \mathbb{U})$$

for some $\beta > 1$, $\delta \geq 0$, and $n \in \mathbb{N}_0$.



Title Page

Contents



Page 5 of 12

Go Back

Full Screen

Close

Also, let $\mathcal{N}(\beta)$ be the subclass of \mathcal{A} , consisting of the functions f , which satisfy the inequality

$$\operatorname{Re} \left\{ \frac{zf''(z)}{f'(z)} + 1 \right\} < \beta \quad (z \in \mathbb{U}).$$

It is obvious that

$$\mathcal{M}^0(\delta, \beta) \equiv \mathcal{M}(\beta) \equiv \mathcal{M}^n(0, \beta).$$

The classes $\mathcal{M}(\beta)$ and $\mathcal{N}(\beta)$ were studied by Uralegaddi et al. in [11], Owa and Srivastava in [9], and Breaz in [4].

Definition 1.1. Let $n, m \in \mathbb{N}_0$ and $k_i > 0$, $1 \leq i \leq m$. We define the integral operator $I_n(f_1, \dots, f_m) : \mathcal{A}^m \rightarrow \mathcal{A}$

$$I_n(f_1, \dots, f_m)(z) := \int_0^z \left(\frac{D^n f_1(t)}{t} \right)^{k_1} \cdots \left(\frac{D^n f_m(t)}{t} \right)^{k_m} dt, \quad (z \in \mathbb{U}),$$

where $f_i \in \mathcal{A}$ and D^n is the Al-Oboudi differential operator.

Remark 1.

(i) For $n = 0$, we have the integral operator

$$I_0(f_1, \dots, f_m)(z) = \int_0^z \left(\frac{f_1(t)}{t} \right)^{k_1} \cdots \left(\frac{f_m(t)}{t} \right)^{k_m} dt$$

introduced in [5]. More details about $I_0(f_1, \dots, f_m)$ can be found in [3] and [4].

(ii) For $n = 0$, $m = 1$, $k_1 = 1$, $k_2 = \dots = k_m = 0$ and $D^0 f_1(z) := D^0 f(z) = f(z) \in \mathcal{A}$, we have the integral operator of Alexander

$$I_0(f)(z) := \int_0^z \frac{f(t)}{t} dt$$

introduced in [1].


- (iii) For $n = 0$, $m = 1$, $k_1 = k \in [0, 1]$, $k_2 = \dots = k_m = 0$ and $D^0 f_1(z) := D^0 f(z) = f(z) \in \mathcal{S}$, we have the integral operator

$$I(f)(z) := \int_0^z \left(\frac{f(t)}{t} \right)^k dt$$

studied in [8].

- (iv) If $k_i \in \mathbb{C}$ for $1 \leq i \leq m$, then we have the integral operator $I_n(f_1, \dots, f_m)$ studied in [7].

In this paper, we investigate some properties for the operators I_n on the classes $\mathcal{S}^n(\delta, \alpha)$ and $\mathcal{M}^n(\delta, \beta)$.





Some Properties for
an Integral Operator

Serap Bulut

vol. 9, iss. 4, art. 115, 2008

Title Page

Contents



Page 6 of 12


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 7 of 12

Go Back

Full Screen

Close

2. Some Properties for I_n on the Class $\mathcal{S}^n(\delta, \alpha)$

Theorem 2.1. Let $f_i \in \mathcal{S}^n(\delta, \alpha_i)$ for $1 \leq i \leq m$ with $0 \leq \alpha_i < 1$, $\delta \geq 0$ and $n \in \mathbb{N}_0$. Also let $k_i > 0$, $1 \leq i \leq m$. If

$$\sum_{i=1}^m k_i(1 - \alpha_i) \leq 1,$$

then $I_n(f_1, \dots, f_m) \in \mathcal{K}(\lambda)$ with $\lambda = 1 + \sum_{i=1}^m k_i(\alpha_i - 1)$.

Proof. By (1.5), for $1 \leq i \leq m$, we have

$$\frac{D^n f_i(z)}{z} = 1 + \sum_{j=2}^{\infty} [1 + (j-1)\delta]^n a_{j,i} z^{j-1}, \quad (n \in \mathbb{N}_0)$$

and

$$\frac{D^n f_i(z)}{z} \neq 0$$

for all $z \in \mathbb{U}$.

On the other hand, we obtain

$$I_n(f_1, \dots, f_m)'(z) = \left(\frac{D^n f_1(z)}{z} \right)^{k_1} \cdots \left(\frac{D^n f_m(z)}{z} \right)^{k_m}$$

for $z \in \mathbb{U}$. This equality implies that

$$\ln I_n(f_1, \dots, f_m)'(z) = k_1 \ln \frac{D^n f_1(z)}{z} + \cdots + k_m \ln \frac{D^n f_m(z)}{z}$$

or equivalently

$$\ln I_n(f_1, \dots, f_m)'(z) = k_1 [\ln D^n f_1(z) - \ln z] + \cdots + k_m [\ln D^n f_m(z) - \ln z].$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 8 of 12

[Go Back](#)

[Full Screen](#)

[Close](#)

By differentiating the above equality, we get

$$\frac{I_n(f_1, \dots, f_m)''(z)}{I_n(f_1, \dots, f_m)'(z)} = \sum_{i=1}^m k_i \left[\frac{(D^n f_i(z))'}{D^n f_i(z)} - \frac{1}{z} \right].$$

Thus, we obtain

$$\frac{z I_n(f_1, \dots, f_m)''(z)}{I_n(f_1, \dots, f_m)'(z)} + 1 = \sum_{i=1}^m k_i \frac{z (D^n f_i(z))'}{D^n f_i(z)} - \sum_{i=1}^m k_i + 1.$$

This relation is equivalent to

$$\operatorname{Re} \left\{ \frac{z I_n(f_1, \dots, f_m)''(z)}{I_n(f_1, \dots, f_m)'(z)} + 1 \right\} = \sum_{i=1}^m k_i \operatorname{Re} \left\{ \frac{z (D^n f_i(z))'}{D^n f_i(z)} \right\} - \sum_{i=1}^m k_i + 1.$$

Since $f_i \in \mathcal{S}^n(\delta, \alpha_i)$, we get


$$\operatorname{Re} \left\{ \frac{z I_n(f_1, \dots, f_m)''(z)}{I_n(f_1, \dots, f_m)'(z)} + 1 \right\} > \sum_{i=1}^m k_i \alpha_i - \sum_{i=1}^m k_i + 1 = 1 + \sum_{i=1}^m k_i (\alpha_i - 1).$$

So, the integral operator $I_n(f_1, \dots, f_m)$ is convex of order λ with $\lambda = 1 + \sum_{i=1}^m k_i (\alpha_i - 1)$. \square

Corollary 2.2. Let $f_i \in \mathcal{S}^n(\delta, \alpha)$ for $1 \leq i \leq m$ with $0 \leq \alpha < 1$, $\delta \geq 0$ and $n \in \mathbb{N}_0$. Also let $k_i > 0$, $1 \leq i \leq m$. If

$$\sum_{i=1}^m k_i \leq \frac{1}{1 - \alpha},$$

then $I_n(f_1, \dots, f_m) \in \mathcal{K}(\rho)$ with $\rho = 1 + (\alpha - 1) \sum_{i=1}^m k_i$.



Title Page

Contents



Page 9 of 12

Go Back

Full Screen

Close

Proof. In Theorem 2.1, we consider $\alpha_1 = \dots = \alpha_m = \alpha$. □

Corollary 2.3. Let $f \in \mathcal{S}^n(\delta, \alpha)$ with $0 \leq \alpha < 1$, $\delta \geq 0$ and $n \in \mathbb{N}_0$. Also let $0 < k \leq 1/(1 - \alpha)$. Then the function

$$I_n(f)(z) = \int_0^z \left(\frac{D^n f(t)}{t} \right)^k dt$$


is in $\mathcal{K}(1 + k(\alpha - 1))$.

Proof. In Corollary 2.2, we consider $m = 1$ and $k_1 = k$. □

Corollary 2.4. Let $f \in \mathcal{S}^n(\delta, \alpha)$. Then the integral operator

$$I_n(f)(z) = \int_0^z (D^n f(t)/t) dt \in \mathcal{K}(\alpha).$$

Proof. In Corollary 2.3, we consider $k = 1$. □



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 10 of 12

Go Back

Full Screen

Close

3. Some Properties for I_n on the class $\mathcal{M}^n(\delta, \beta)$

Theorem 3.1. Let $f_i \in \mathcal{M}^n(\delta, \beta_i)$ for $1 \leq i \leq m$ with $\beta_i > 1$. Then $I_n(f_1, \dots, f_m) \in \mathcal{N}(\lambda)$ with $\lambda = 1 + \sum_{i=1}^m k_i(\beta_i - 1)$ and $k_i > 0$, ($1 \leq i \leq m$).

Proof. Proof is similar to the proof of Theorem 2.1. □

Remark 2. For $n = 0$, we have Theorem 2.1 in [4].

Corollary 3.2. Let $f_i \in \mathcal{M}^n(\delta, \beta)$ for $1 \leq i \leq m$ with $\beta > 1$. Then $I_n(f_1, \dots, f_m) \in \mathcal{N}(\rho)$ with $\rho = 1 + (\beta - 1) \sum_{i=1}^m k_i$ and $k_i > 0$, ($1 \leq i \leq m$).

Corollary 3.3. Let $f \in \mathcal{M}^n(\delta, \beta)$ with $\beta > 1$. Then the integral operator

$$I_n(f)(z) = \int_0^z \left(\frac{D^n f(t)}{t} \right)^k dt \in \mathcal{N}(1 + k(\beta - 1))$$


and $k > 0$.

Corollary 3.4. Let $f \in \mathcal{M}^n(\delta, \beta)$ with $\beta > 1$. Then the integral operator

$$I_n(f)(z) = \int_0^z \frac{D^n f(t)}{t} dt \in \mathcal{N}(\beta).$$

References

- [1] I.W. ALEXANDER, Functions which map the interior of the unit circle upon simple regions, *Ann. of Math.*, **17** (1915), 12–22.
- [2] F.M. AL-BOUDI, On univalent functions defined by a generalized Sălăgean operator, *Int. J. Math. Math. Sci.*, (25-28) 2004, 1429–1436.
- [3] D. BREAZ, A convexity property for an integral operator on the class $S_p(\beta)$, *J. Inequal. Appl.*, (2008), Art. ID 143869.
- [4] D. BREAZ, Certain integral operators on the classes $\mathcal{M}(\beta_i)$ and $\mathcal{N}(\beta_i)$, *J. Inequal. Appl.*, (2008), Art. ID 719354.
- [5] D. BREAZ AND N. BREAZ, Two integral operators, *Studia Univ. Babeş-Bolyai Math.*, **47**(3) (2002), 13–19.
- [6] D. BREAZ, S. OWA AND N. BREAZ, A new integral univalent operator, *Acta Univ. Apulensis Math. Inform.*, **16** (2008), 11–16.
- [7] S. BULUT, Sufficient conditions for univalence of an integral operator defined by Al-Oboudi differential operator, *J. Inequal. Appl.*, (2008), Art. ID 957042.
- [8] S.S. MILLER, P.T. MOCANU AND M.O. READE, Starlike integral operators, *Pacific J. Math.*, **79**(1) (1978), 157–168.
- [9] S. OWA AND H.M. SRIVASTAVA, Some generalized convolution properties associated with certain subclasses of analytic functions, *J. Inequal. Pure Appl. Math.*, **3**(3) (2002), Art. 42. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=194>].



Some Properties for
an Integral Operator

Serap Bulut

vol. 9, iss. 4, art. 115, 2008

Title Page

Contents



Page 11 of 12

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [10] G.Ş. SĂLĂGEAN, Subclasses of univalent functions, *Complex Analysis-Fifth Romanian-Finnish seminar, Part 1* (Bucharest, 1981), Lecture Notes in Math., vol. 1013, Springer, Berlin, 1983, pp. 362–372.
- [11] B.A. URALEGADDI, M.D. GANIGI AND S.M. SARANGI, Univalent functions with positive coefficients, *Tamkang J. Math.*, **25**(3) (1994), 225–230.

Some Properties for
an Integral Operator

Serap Bulut

vol. 9, iss. 4, art. 115, 2008

Title Page

Contents



Page 12 of 12

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756