

Journal of Inequalities in Pure and Applied Mathematics

LOWER BOUNDS FOR THE SPECTRAL NORM

JORMA K. MERIKOSKI AND RAVINDER KUMAR

Department of Mathematics, Statistics and Philosophy
FI-33014 University of Tampere, Finland
EMail: jorma.merikoski@uta.fi

Department of Mathematics
Dayalbagh Educational Institute
Dayalbagh, Agra 282005
Uttar Pradesh, India
EMail: ravinder_dei@yahoo.com

©2000 Victoria University
ISSN (electronic): 1443-5756
249-04



volume 6, issue 4, article 124,
2005.

*Received 20 December, 2004;
accepted 01 September, 2005.*

Communicated by: S. Puntanen

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

Let A be a complex $m \times n$ matrix. We find simple and good lower bounds for its spectral norm $\|A\| = \max\{\|Ax\| \mid x \in \mathbb{C}^n, \|x\| = 1\}$ by choosing x smartly. Here $\|\cdot\|$ applied to a vector denotes the Euclidean norm.

2000 Mathematics Subject Classification: 15A60, 15A18.

Key words: Spectral norm, Singular values.

Contents

1	Introduction	3
2	Simple bounds	4
3	Improved bounds	6
4	Further improvements	8
5	Experiments	11
6	Conclusions	12
7	Remark	13

References



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 2 of 14

1. Introduction

Throughout this paper, \mathbf{A} denotes a complex $m \times n$ matrix ($m, n \geq 2$). We denote by $\|\mathbf{A}\|$ its spectral norm or largest singular value.

The singular values of \mathbf{A} are square roots of the eigenvalues of $\mathbf{A}^* \mathbf{A}$. Since much is known about bounds for eigenvalues of Hermitian matrices, we may apply this knowledge to $\mathbf{A}^* \mathbf{A}$ to find bounds for singular values, but the bounds so obtained are very complicated in general. However, as we will see, we can find simple and good lower bounds for $\|\mathbf{A}\|$ by choosing \mathbf{x} smartly in the variational characterization of the largest eigenvalue of $\mathbf{A}^* \mathbf{A}$,


$$(1.1) \quad \|\mathbf{A}\| = \max \left\{ (\mathbf{x}^* \mathbf{A}^* \mathbf{A} \mathbf{x})^{1/2} \mid \mathbf{x} \in \mathbb{C}^n, \mathbf{x}^* \mathbf{x} = 1 \right\},$$

or, equivalently, in the definition

$$(1.1') \quad \|\mathbf{A}\| = \max \left\{ \|\mathbf{A} \mathbf{x}\| \mid \mathbf{x} \in \mathbb{C}^n, \|\mathbf{x}\| = 1 \right\},$$

where $\|\cdot\|$ applied to a vector denotes the Euclidean norm.

Our earlier papers [3] and [4] are based on somewhat similar ideas to find lower bounds for the spread and numerical radius of \mathbf{A} .



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 3 of 14

2. Simple bounds

Consider the partition $\mathbf{A} = (\mathbf{a}_1 \dots \mathbf{a}_n)$ of \mathbf{A} into columns. For $H \subseteq N = \{1, \dots, n\}$, denote by \mathbf{A}_H the block of the columns \mathbf{a}_h with $h \in H$. We accept also the empty block \mathbf{A}_\emptyset .

Throughout this paper, we let $I (\neq \emptyset)$, K , and L be disjoint subsets of N satisfying $N = I \cup K \cup L$. Since multiplication by permutation matrices does not change singular values, we can reorder the columns, and so we are allowed to assume that

$$\mathbf{A} = \left(\mathbf{A}_I \quad \mathbf{A}_K \quad \mathbf{A}_L \right).$$

Then

$$\mathbf{A}^* \mathbf{A} = \begin{pmatrix} \mathbf{A}_I^* \mathbf{A}_I & \mathbf{A}_I^* \mathbf{A}_K & \mathbf{A}_I^* \mathbf{A}_L \\ \mathbf{A}_K^* \mathbf{A}_I & \mathbf{A}_K^* \mathbf{A}_K & \mathbf{A}_K^* \mathbf{A}_L \\ \mathbf{A}_L^* \mathbf{A}_I & \mathbf{A}_L^* \mathbf{A}_K & \mathbf{A}_L^* \mathbf{A}_L \end{pmatrix}.$$


We denote $\mathbf{e}_H = \sum_{h \in H} \mathbf{e}_h$, where \mathbf{e}_h is the h 'th standard basis vector of \mathbb{C}^n .

We choose $\mathbf{x} = \mathbf{e}_I / \sqrt{|I|}$ in (1.1), where $|\cdot|$ stands for the number of the elements. Then

$$(2.1) \quad \|\mathbf{A}\| \geq \left(\frac{\text{su } \mathbf{A}_I^* \mathbf{A}_I}{|I|} \right)^{\frac{1}{2}},$$

where su denotes the sum of the entries. Hence

$$(2.2) \quad \|\mathbf{A}\| \geq \max_{I \neq \emptyset} \left(\frac{\text{su } \mathbf{A}_I^* \mathbf{A}_I}{|I|} \right)^{\frac{1}{2}} = \max_{I \neq \emptyset} \frac{1}{\sqrt{|I|}} \left\| \sum_{i \in I} \mathbf{a}_i \right\|,$$



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 4 of 14


and, restricting to $I = \{1\}, \dots, \{n\}$,

$$(2.3) \quad \|\mathbf{A}\| \geq \max_i \|\mathbf{a}_i\| = \max_i \left(\sum_j |a_{ji}|^2 \right)^{\frac{1}{2}},$$

and also, restricting to $I = N$,

$$(2.4) \quad \|\mathbf{A}\| \geq \left(\frac{\text{su } \mathbf{A}_I^* \mathbf{A}_I}{n} \right)^{\frac{1}{2}} = \left(\frac{|r_1|^2 + \dots + |r_n|^2}{n} \right)^{\frac{1}{2}},$$

where r_1, \dots, r_n are the row sums of \mathbf{A} .



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 5 of 14

3. Improved bounds

To improve (2.2), choose

$$\mathbf{x} = \frac{\mathbf{e}_I + z\mathbf{e}_K}{\sqrt{|I| + |K|}},$$

where $z \in \mathbb{C}$ satisfies $|z| = 1$. Then

$$\begin{aligned} \|\mathbf{A}\| &\geq \frac{1}{\sqrt{|I| + |K|}} \left[\text{su} \begin{pmatrix} \mathbf{A}_I^* \mathbf{A}_I & z \mathbf{A}_I^* \mathbf{A}_K \\ \bar{z} \mathbf{A}_K^* \mathbf{A}_I & \mathbf{A}_K^* \mathbf{A}_K \end{pmatrix} \right]^{\frac{1}{2}} \\ &= \frac{1}{\sqrt{|I| + |K|}} \left[\left\| \sum_{i \in I} \mathbf{a}_i \right\|^2 + \left\| \sum_{k \in K} \mathbf{a}_k \right\|^2 + 2 \operatorname{Re} \left(z \sum_{i \in I} \sum_{k \in K} \mathbf{a}_i^* \mathbf{a}_k \right) \right]^{\frac{1}{2}}, \end{aligned}$$

which, for $z = \bar{w}/|w|$ if $w = \sum_{i \in I} \sum_{k \in K} \mathbf{a}_i^* \mathbf{a}_k \neq 0$, and z arbitrary if $w = 0$, implies

$$(3.1) \quad \|\mathbf{A}\| \geq \frac{1}{\sqrt{|I| + |K|}} \left(\left\| \sum_{i \in I} \mathbf{a}_i \right\|^2 + \left\| \sum_{k \in K} \mathbf{a}_k \right\|^2 + 2 \left| \sum_{i \in I} \sum_{k \in K} \mathbf{a}_i^* \mathbf{a}_k \right| \right)^{\frac{1}{2}}.$$

Hence

$$(3.2) \quad \|\mathbf{A}\| \geq \max_{I \neq \emptyset, I \cap K = \emptyset} \frac{1}{\sqrt{|I| + |K|}} \left(\left\| \sum_{i \in I} \mathbf{a}_i \right\|^2 + \left\| \sum_{k \in K} \mathbf{a}_k \right\|^2 + 2 \left| \sum_{i \in I} \sum_{k \in K} \mathbf{a}_i^* \mathbf{a}_k \right| \right)^{\frac{1}{2}},$$



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 6 of 14

and, restricting to $I, K = \{1\}, \dots, \{n\}$,


$$(3.3) \quad \|\mathbf{A}\| \geq \frac{1}{\sqrt{2}} \max_{i \neq k} (\|\mathbf{a}_i\|^2 + \|\mathbf{a}_k\|^2 + 2|\mathbf{a}_i^* \mathbf{a}_k|)^{\frac{1}{2}}.$$

It is well-known that the largest eigenvalue of a principal submatrix of a Hermitian matrix is less or equal to that of the original matrix. So, computing the largest eigenvalue of

$$\begin{pmatrix} \mathbf{a}_i^* \mathbf{a}_i & \mathbf{a}_i^* \mathbf{a}_k \\ \mathbf{a}_k^* \mathbf{a}_i & \mathbf{a}_k^* \mathbf{a}_k \end{pmatrix}$$

improves (3.3) to

$$(3.4) \quad \|\mathbf{A}\| \geq \frac{1}{\sqrt{2}} \max_{i \neq k} \left\{ \|\mathbf{a}_i\|^2 + \|\mathbf{a}_k\|^2 + \left[(\|\mathbf{a}_i\|^2 - \|\mathbf{a}_k\|^2)^2 + 4|\mathbf{a}_i^* \mathbf{a}_k|^2 \right]^{\frac{1}{2}} \right\}^{\frac{1}{2}}.$$



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 7 of 14

4. Further improvements

Let \mathbf{B} be a Hermitian $n \times n$ matrix with largest eigenvalue λ . If $\mathbf{0} \neq \mathbf{x} \in \mathbb{C}^n$, then

$$(4.1) \quad \lambda \geq \frac{\mathbf{x}^* \mathbf{B} \mathbf{x}}{\mathbf{x}^* \mathbf{x}}.$$

We replace \mathbf{x} with $\mathbf{B} \mathbf{x}$ (assumed nonzero) and ask whether the bound so obtained,

$$(4.2) \quad \lambda \geq \frac{\mathbf{x}^* \mathbf{B}^3 \mathbf{x}}{\mathbf{x}^* \mathbf{B}^2 \mathbf{x}},$$

is better. In other words, is

$$\frac{\mathbf{x}^* \mathbf{B}^3 \mathbf{x}}{\mathbf{x}^* \mathbf{B}^2 \mathbf{x}} \geq \frac{\mathbf{x}^* \mathbf{B} \mathbf{x}}{\mathbf{x}^* \mathbf{x}}$$


generally valid? The answer is yes if \mathbf{B} is nonnegative definite (and $\mathbf{B} \mathbf{x} \neq \mathbf{0}$). In fact, the function

$$f(p) = \frac{\mathbf{x}^* \mathbf{B}^{p+1} \mathbf{x}}{\mathbf{x}^* \mathbf{B}^p \mathbf{x}} \quad (p \geq 0)$$

is then increasing. We omit the easy proof but note that several interesting questions arise if we instead of nonnegative definiteness assume symmetry and nonnegativity, see [2] and its references.

If $\mathbf{B} = \mathbf{A}^* \mathbf{A}$, then (4.1) implies

$$(4.1') \quad \|\mathbf{A}\| \geq \frac{\|\mathbf{A} \mathbf{x}\|}{\|\mathbf{x}\|},$$



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 8 of 14

and (4.2) implies a better bound

$$(4.2') \quad \|\mathbf{A}\| \geq \frac{\|\mathbf{A}\mathbf{A}^*\mathbf{A}\mathbf{x}\|}{\|\mathbf{A}^*\mathbf{A}\mathbf{x}\|}.$$

Since (2.2) is obtained by choosing in (4.1') $\mathbf{x} = \mathbf{e}_I$ and maximizing over I , we get a better bound by applying (4.2') instead. Because

$$\mathbf{A}^*\mathbf{A}\mathbf{e}_I = \mathbf{A}^* \sum_{i \in I} \mathbf{a}_i = \left(\mathbf{a}_1^* \sum_{i \in I} \mathbf{a}_i \quad \dots \quad \mathbf{a}_n^* \sum_{i \in I} \mathbf{a}_i \right)^T$$

and


$$\mathbf{A}\mathbf{A}^*\mathbf{A}\mathbf{e}_I = \left(\sum_j \mathbf{a}_j \mathbf{a}_j^* \right) \sum_{i \in I} \mathbf{a}_i,$$

we have

$$(4.3) \quad \|\mathbf{A}\| \geq \max \left\{ \frac{\left\| \left(\sum_j \mathbf{a}_j \mathbf{a}_j^* \right) \sum_{i \in I} \mathbf{a}_i \right\|}{\left\| \left(\mathbf{a}_1^* \sum_{i \in I} \mathbf{a}_i \quad \dots \quad \mathbf{a}_n^* \sum_{i \in I} \mathbf{a}_i \right)^T \right\|} \mid I \neq \emptyset, \sum_{i \in I} \mathbf{a}_i \neq \mathbf{0} \right\}.$$

Hence, restricting to $I = \{1\}, \dots, \{n\}$ and assuming all the \mathbf{a}_i 's nonzero,

$$(4.4) \quad \|\mathbf{A}\| \geq \max_i \left\| \left(\sum_j \mathbf{a}_j \mathbf{a}_j^* \right) \mathbf{a}_i \right\| / \left\| \left(\mathbf{a}_1^* \mathbf{a}_i \quad \dots \quad \mathbf{a}_n^* \mathbf{a}_i \right)^T \right\|,$$



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back


Close

Quit

Page 9 of 14

and, restricting to $I = N$ and assuming the row sum vector $\mathbf{r} = (r_1 \dots r_n)^\top$ nonzero,

$$(4.5) \quad \|\mathbf{A}\| \geq \left\| \left(\sum_j \mathbf{a}_j \mathbf{a}_j^* \right) \mathbf{r} \right\| / \left\| (\mathbf{a}_1^* \mathbf{r} \dots \mathbf{a}_n^* \mathbf{r})^\top \right\|.$$



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 10 of 14

5. Experiments


We studied our bounds by random matrices of order 10. In the case of (2.2), (3.2), and (4.3), to avoid big complexities, we did not maximize over sets but studied only $I = \{1, 3, 5, 7, 9\}$, $K = \{2, 4, 6, 8, 10\}$. We considered various types of matrices (positive, normal, etc.). For each type, we performed one hundred experiments and computed means (m) and standard deviations (s) of

$$\frac{\|A\| - \text{bound}}{\text{bound}}.$$

For positive symmetric matrices, (4.5) was by far the best with very surprising success: $m = 0.0000215$, $s = 0.0000316$. For all the remaining types, (4.4) was the best also with surprising success. We mention a few examples.

Type	m	s
Normal	0.0463	0.0227
Positive	0.0020	0.0012
Real	0.0261	0.0151
Complex	0.0429	0.0162

For positive matrices, also the very simple bound (2.4) was surprisingly good: $m = 0.0150$, $s = 0.0067$.



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and Ravinder Kumar

Title Page

Contents



Go Back

Close


Quit

Page 11 of 14

6. Conclusions

Our bounds (2.1), (2.3), (2.4), (3.1), (3.3), (3.4), (4.4), and (4.5) have complexity $O(n^2)$. Also the bounds (2.2), (3.2), and (4.3) have this complexity if we do not include all the subsets of N but only some suitable subsets. Our best $O(n^2)$ bounds seem to be in general better than all the $O(n^2)$ bounds we have found from the literature (e.g., the bound of [1, Theorem 3.7.15]).

One natural way [5, 6] of finding a lower bound for $\|\mathbf{A}\|$ is to compute the Wolkowicz-Styan [7] lower bound for the largest eigenvalue of $\mathbf{A}^* \mathbf{A}$ and to take the square root. The bound so obtained [5, 6] is fairly simple but seems to be in general worse than many of our bounds and has complexity $O(n^3)$.



Lower Bounds for the Spectral
Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back


Close

Quit

Page 12 of 14

7. Remark

As $\|A\| = \|A^T\| = \|A^*\|$, all our results remain valid if we take row vectors instead of column vectors, and column sums instead of row sums. We can also do both and choose the better one.



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 13 of 14

References

- [1] R.A. HORN AND C.R. JOHNSON, *Topics in Matrix Analysis*, Cambridge University Press, Cambridge, 1991.
- [2] H. KANKAANPÄÄ AND J.K. MERIKOSKI, Two inequalities for the sum of elements of a matrix, *Linear and Multilinear Algebra*, **18** (1985), 9–22.
- [3] J.K. MERIKOSKI AND R. KUMAR, Characterizations and lower bounds for the spread of a normal matrix, *Linear Algebra Appl.*, **364** (2003), 13–31.
- [4] J.K. MERIKOSKI AND R. KUMAR, Lower bounds for the numerical radius, *Linear Algebra Appl.*, **410** (2005), 135–142.
- [5] J.K. MERIKOSKI, H. SARRIA AND P. TARAZAGA, Bounds for singular values using traces, *Linear Algebra Appl.*, **210** (1994), 227–254.
- [6] O. ROJO, R. SOTO AND H. ROJO, Bounds for the spectral radius and the largest singular value, *Comput. Math. Appl.*, **36** (1998), 41–50.
- [7] H. WOLKOWICZ AND G.P.H. STYAN, Bounds for eigenvalues using traces, *Linear Algebra Appl.*, **29** (1980), 471–506.



Lower Bounds for the Spectral Norm

Jorma K. Merikoski and
Ravinder Kumar

Title Page

Contents



Go Back

Close

Quit

Page 14 of 14