

REVERSE TRIANGLE INEQUALITY IN HILBERT C^* -MODULES

MARYAM KHOSRAVI, HAKIMEH MAHYAR

Department of Mathematics
Tarbiat Moallem University
Tehran, Iran.

EEmail: {khosravi_m, mahyar}@saba.tmu.ac.ir

MOHAMMAD SAL MOSLEHIAN

Department of Pure Mathematics
Centre of Excellence in Analysis on Algebraic Structures (CEAAS)
Ferdowsi University of Mashhad
P.O. Box 1159, Mashhad 91775, Iran.

EEmail: moslehian@ferdowsi.um.ac.ir

Received: 16 October, 2009

Accepted: 13 November, 2009

Communicated by: S.S. Dragomir



2000 AMS Sub. Class.: Primary 46L08; Secondary 15A39, 26D15, 46L05, 51M16.

Key words: Triangle inequality, Reverse inequality, Hilbert C^* -module, C^* -algebra.

Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian
vol. 10, iss. 4, art. 110, 2009

Title Page

Contents



Page 1 of 21

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Abstract:

We prove several versions of reverse triangle inequality in Hilbert C^* -modules. We show that if e_1, \dots, e_m are vectors in a Hilbert module \mathfrak{X} over a C^* -algebra \mathfrak{A} with unit 1 such that $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $\|e_i\| = 1$ ($1 \leq i \leq m$), and also $r_k, \rho_k \in \mathbb{R}$ ($1 \leq k \leq m$) and $x_1, \dots, x_n \in \mathfrak{X}$ satisfy

$$0 \leq r_k^2 \|x_j\| \leq \operatorname{Re} \langle r_k e_k, x_j \rangle, \quad 0 \leq \rho_k^2 \|x_j\| \leq \operatorname{Im} \langle \rho_k e_k, x_j \rangle,$$

then

$$\left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right]^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|,$$

and the equality holds if and only if


$$\sum_{j=1}^n x_j = \sum_{j=1}^n \|x_j\| \sum_{k=1}^m (r_k + i\rho_k) e_k.$$



Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian
vol. 10, iss. 4, art. 110, 2009

[Title Page](#)

[Contents](#)



Page 2 of 21

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction and Preliminaries	4
2	Multiplicative Reverse of the Triangle Inequality	6
3	Additive Reverse of the Triangle Inequality	16



Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian
vol. 10, iss. 4, art. 110, 2009

Title Page

Contents



Page 3 of 21

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



1. Introduction and Preliminaries

The triangle inequality is one of the most fundamental inequalities in mathematics. Several mathematicians have investigated its generalizations and its reverses.

In 1917, Petrovitch [17] proved that for complex numbers z_1, \dots, z_n ,

$$(1.1) \quad \left| \sum_{j=1}^n z_j \right| \geq \cos \theta \sum_{j=1}^n |z_j|,$$

where $0 < \theta < \frac{\pi}{2}$ and $\alpha - \theta < \arg z_j < \alpha + \theta$ ($1 \leq j \leq n$) for a given real number α .

The first generalization of the reverse triangle inequality in Hilbert spaces was given by Diaz and Metcalf [5]. They proved that for x_1, \dots, x_n in a Hilbert space H , if e is a unit vector of H such that $0 \leq r \leq \frac{\operatorname{Re}\langle x_j, e \rangle}{\|x_j\|}$ for some $r \in \mathbb{R}$ and each $1 \leq j \leq n$, then

$$(1.2) \quad r \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|.$$

Moreover, the equality holds if and only if $\sum_{j=1}^n x_j = r \sum_{j=1}^n \|x_j\| e$.

Recently, a number of mathematicians have presented several refinements of the reverse triangle inequality in Hilbert spaces and normed spaces (see [1, 2, 4, 7, 8, 10, 13, 16]). Recently a discussion of C^* -valued triangle inequalities in Hilbert C^* -modules was given in [3]. Our aim is to generalize some of the results of Dragomir in Hilbert spaces to the framework of Hilbert C^* -modules. For this purpose, we first recall some fundamental definitions in the theory of Hilbert C^* -modules.

Suppose that \mathfrak{A} is a C^* -algebra and \mathfrak{X} is a linear space, which is an algebraic right \mathfrak{A} -module. The space \mathfrak{X} is called a pre-Hilbert \mathfrak{A} -module (or an inner product \mathfrak{A} -module) if there exists an \mathfrak{A} -valued inner product $\langle \cdot, \cdot \rangle : \mathfrak{X} \times \mathfrak{X} \rightarrow \mathfrak{A}$ with the following properties:

Title Page

Contents




Page 4 of 21

Go Back

Full Screen

Close



Title Page

Contents



Page 5 of 21

Go Back

Full Screen

Close

(i) $\langle x, x \rangle \geq 0$ and $\langle x, x \rangle = 0$ if and only if $x = 0$;

(ii) $\langle x, \lambda y + z \rangle = \lambda \langle x, y \rangle + \langle x, z \rangle$;

(iii) $\langle x, ya \rangle = \langle x, y \rangle a$;

(iv) $\langle x, y \rangle^* = \langle y, x \rangle$

for all $x, y, z \in \mathfrak{X}$, $a \in \mathfrak{A}$, $\lambda \in \mathbb{C}$. By (ii) and (iv), $\langle \cdot, \cdot \rangle$ is conjugate linear in the first variable. Using the Cauchy–Schwartz inequality $\langle y, x \rangle \langle x, y \rangle \leq \|\langle x, x \rangle\| \|\langle y, y \rangle\|$ [11, Page 5] (see also [14]), it follows that $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$ is a norm on \mathfrak{X} making it a right normed module. The pre-Hilbert module \mathfrak{X} is called a Hilbert \mathfrak{A} -module if it is complete with respect to this norm. Notice that the inner structure of a C^* -algebra is essentially more complicated than that for complex numbers. For instance, properties such as orthogonality and theorems such as Riesz’ representation in complex Hilbert space theory cannot simply be generalized or transferred to the theory of Hilbert C^* -modules.

One may define an “ \mathfrak{A} -valued norm” $|\cdot|$ by $|x| = \langle x, x \rangle^{1/2}$. Clearly, $\| |x| \| = \|x\|$ for each $x \in \mathfrak{X}$. It is known that $|\cdot|$ does not satisfy the triangle inequality in general. See [11, 12] for more information on Hilbert C^* -modules.

We also use elementary C^* -algebra theory, in particular we utilize the property that if $a \leq b$ then $a^{1/2} \leq b^{1/2}$, where a, b are positive elements of a C^* -algebra \mathfrak{A} . We also repeatedly apply the following known relation:

$$(1.3) \quad \frac{1}{2}(aa^* + a^*a) = (\operatorname{Re} a)^2 + (\operatorname{Im} a)^2,$$

where a is an arbitrary element of \mathfrak{A} . For details on C^* -algebra theory, we refer readers to [15].

Throughout the paper, we assume that \mathfrak{A} is a unital C^* -algebra with unit 1 and for every $\lambda \in \mathbb{C}$, we write λ for $\lambda 1$.



2. Multiplicative Reverse of the Triangle Inequality

Utilizing some C^* -algebraic techniques we present our first result as a generalization of [7, Theorem 2.3].

Theorem 2.1. *Let \mathfrak{A} be a C^* -algebra, let \mathfrak{X} be a Hilbert \mathfrak{A} -module and let $x_1, \dots, x_n \in \mathfrak{X}$. If there exist real numbers $k_1, k_2 \geq 0$ with*

$$0 \leq k_1 \|x_j\| \leq \operatorname{Re}\langle e, x_j \rangle, \quad 0 \leq k_2 \|x_j\| \leq \operatorname{Im}\langle e, x_j \rangle,$$

for some $e \in \mathfrak{X}$ with $|e| \leq 1$ and all $1 \leq j \leq n$, then

$$(2.1) \quad (k_1^2 + k_2^2)^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|.$$

Proof. Applying the Cauchy–Schwarz inequality, we get

$$\left| \left\langle e, \sum_{j=1}^n x_j \right\rangle \right|^2 \leq \|e\|^2 \left| \sum_{j=1}^n x_j \right|^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2,$$

and

$$\left| \left\langle \sum_{j=1}^n x_j, e \right\rangle \right|^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2 |e|^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2,$$

whence

$$\begin{aligned} \left\| \sum_{j=1}^n x_j \right\|^2 &\geq \frac{1}{2} \left(\left| \left\langle e, \sum_{j=1}^n x_j \right\rangle \right|^2 + \left| \left\langle \sum_{j=1}^n x_j, e \right\rangle \right|^2 \right) \\ &= \frac{1}{2} \left(\left\langle e, \sum_{j=1}^n x_j \right\rangle^* \left\langle e, \sum_{j=1}^n x_j \right\rangle + \left\langle \sum_{j=1}^n x_j, e \right\rangle^* \left\langle \sum_{j=1}^n x_j, e \right\rangle \right) \end{aligned}$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 7 of 21

[Go Back](#)

[Full Screen](#)

[Close](#)

$$\begin{aligned}
 &= \left(\operatorname{Re} \left\langle e, \sum_{j=1}^n x_j \right\rangle \right)^2 + \left(\operatorname{Im} \left\langle e, \sum_{j=1}^n x_j \right\rangle \right)^2 && \text{(by (1.3))} \\
 &= \left(\operatorname{Re} \sum_{j=1}^n \langle e, x_j \rangle \right)^2 + \left(\operatorname{Im} \sum_{j=1}^n \langle e, x_j \rangle \right)^2 \\
 &\geq k_1^2 \left(\sum_{j=1}^n \|x_j\| \right)^2 + k_2^2 \left(\sum_{j=1}^n \|x_j\| \right)^2 \\
 &= (k_1^2 + k_2^2) \left(\sum_{j=1}^n \|x_j\| \right)^2 .
 \end{aligned}$$

□

Using the same argument as in the proof of Theorem 2.1, one can obtain the following result, where k_1, k_2 are hermitian elements of \mathfrak{A} .

Theorem 2.2. *If the vectors $x_1, \dots, x_n \in \mathfrak{X}$ satisfy the conditions*

$$0 \leq k_1^2 \|x_j\|^2 \leq (\operatorname{Re} \langle e, x_j \rangle)^2, \quad 0 \leq k_2^2 \|x_j\|^2 \leq (\operatorname{Im} \langle e, x_j \rangle)^2,$$

for some hermitian elements k_1, k_2 in \mathfrak{A} , some $e \in \mathfrak{X}$ with $|e| \leq 1$ and all $1 \leq j \leq n$ then the inequality (2.1) holds.

One may observe an integral version of inequality (2.1) as follows:

Corollary 2.3. *Suppose that \mathfrak{X} is a Hilbert \mathfrak{A} -module and $f : [a, b] \rightarrow \mathfrak{X}$ is strongly measurable such that the Lebesgue integral $\int_a^b \|f(t)\| dt$ exists and is finite. If there exist self-adjoint elements a_1, a_2 in \mathfrak{A} with*

$$a_1^2 \|f(t)\|^2 \leq \operatorname{Re} \langle f(t), e \rangle^2, \quad a_2^2 \|f(t)\|^2 \leq \operatorname{Im} \langle f(t), e \rangle^2 \quad (\text{a.e. } t \in [a, b]),$$



where $e \in \mathfrak{X}$ with $|e| \leq 1$, then

$$(a_1^2 + a_2^2)^{\frac{1}{2}} \int_a^b \|f(t)\| dt \leq \left\| \int_a^b f(t) dt \right\|.$$

Now we prove a useful lemma, which is frequently applied in the next theorems (see also [3]).

Lemma 2.4. *Let \mathfrak{X} be a Hilbert \mathfrak{A} -module and let $x, y \in \mathfrak{X}$. If $|\langle x, y \rangle| = \|x\| \|y\|$, then*

$$y = \frac{x \langle x, y \rangle}{\|x\|^2}.$$

Proof. For $x, y \in \mathfrak{X}$ we have

$$\begin{aligned} 0 &\leq \left| y - \frac{x \langle x, y \rangle}{\|x\|^2} \right|^2 = \left\langle y - \frac{x \langle x, y \rangle}{\|x\|^2}, y - \frac{x \langle x, y \rangle}{\|x\|^2} \right\rangle \\ &= \langle y, y \rangle - \frac{1}{\|x\|^2} \langle y, x \rangle \langle x, y \rangle + \frac{1}{\|x\|^4} \langle y, x \rangle \langle x, x \rangle \langle x, y \rangle - \frac{1}{\|x\|^2} \langle y, x \rangle \langle x, y \rangle \\ &\leq |y|^2 - \frac{1}{\|x\|^2} |\langle x, y \rangle|^2 = |y|^2 - \frac{1}{\|x\|^2} \|x\|^2 \|y\|^2 \\ &= |y|^2 - \|y\|^2 \leq 0, \end{aligned}$$

whence $\left| y - \frac{x \langle x, y \rangle}{\|x\|^2} \right| = 0$. Hence $y = \frac{x \langle x, y \rangle}{\|x\|^2}$. \square

Using the Cauchy–Schwarz inequality, we have the following theorem for Hilbert modules, which is similar to [1, Theorem 2.5].

Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian

vol. 10, iss. 4, art. 110, 2009

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 8 of 21

[Go Back](#)


[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.



Theorem 2.5. Let e_1, \dots, e_m be a family of vectors in a Hilbert module \mathfrak{X} over a C^* -algebra \mathfrak{A} such that $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $\|e_i\| = 1$ ($1 \leq i \leq m$). Suppose that $r_k, \rho_k \in \mathbb{R}$ ($1 \leq k \leq m$) and that the vectors $x_1, \dots, x_n \in \mathfrak{X}$ satisfy

$$0 \leq r_k^2 \|x_j\| \leq \operatorname{Re} \langle r_k e_k, x_j \rangle, \quad 0 \leq \rho_k^2 \|x_j\| \leq \operatorname{Im} \langle \rho_k e_k, x_j \rangle,$$

Then

$$(2.2) \quad \left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right]^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left\| \sum_{j=1}^n x_j \right\|,$$

and the equality holds if and only if


$$(2.3) \quad \sum_{j=1}^n x_j = \sum_{j=1}^n \|x_j\| \sum_{k=1}^m (r_k + i\rho_k) e_k.$$

Proof. There is nothing to prove if $\sum_{k=1}^m (r_k^2 + \rho_k^2) = 0$. Assume that $\sum_{k=1}^m (r_k^2 + \rho_k^2) \neq 0$. From the hypothesis, by $\operatorname{Im}(a) = \operatorname{Re}(ia^*)$, $\operatorname{Re}(a^*) = \operatorname{Re}(a)$ ($a \in \mathfrak{A}$), we have

$$\begin{aligned} & \left(\sum_{k=1}^m (r_k^2 + \rho_k^2) \right)^2 \left(\sum_{j=1}^n \|x_j\| \right)^2 \\ & \leq \left(\operatorname{Re} \left\langle \sum_{k=1}^m r_k e_k, \sum_{j=1}^n x_j \right\rangle + \operatorname{Im} \left\langle \sum_{k=1}^m \rho_k e_k, \sum_{j=1}^n x_j \right\rangle \right)^2 \\ & = \left(\operatorname{Re} \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle \right)^2. \end{aligned}$$

Title Page

Contents



Page 9 of 21

Go Back


Full Screen

Close



Title Page

Contents



Page 10 of 21

Go Back

Full Screen

Close

By (1.3),

$$\begin{aligned}
 & \left(\operatorname{Re} \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle \right)^2 \\
 & \leq \frac{1}{2} \left(\left| \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle \right|^2 + \left| \left\langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{j=1}^n x_j \right\rangle \right|^2 \right) \\
 & \leq \frac{1}{2} \left(\left\| \sum_{j=1}^n x_j \right\|^2 \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2 + \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2 \left\| \sum_{j=1}^n x_j \right\|^2 \right) \\
 & \leq \left\| \sum_{j=1}^n x_j \right\|^2 \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2
 \end{aligned}$$

and since $|a| \leq \|a\|$ ($a \in \mathfrak{A}$),

$$\begin{aligned}
 \left\| \sum_{j=1}^n x_j \right\|^2 \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2 & \leq \left\| \sum_{j=1}^n x_j \right\|^2 \left\| \left\langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle \right\|^2 \\
 & = \left\| \sum_{j=1}^n x_j \right\|^2 \sum_{k=1}^m |r_k + i\rho_k|^2 \|e_k\|^2 \\
 & = \left\| \sum_{j=1}^n x_j \right\|^2 \sum_{k=1}^m (r_k^2 + \rho_k^2).
 \end{aligned}$$



Title Page

Contents



Page 11 of 21

Go Back

Full Screen

Close

Hence

$$\left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right] \left(\sum_{j=1}^n \|x_j\| \right)^2 \leq \left\| \sum_{j=1}^n x_j \right\|^2.$$

By taking square roots the desired result follows.

Clearly we have equality in (2.2) if condition (2.3) holds. To see the converse, first note that if equality holds in (2.2), then all inequalities in the relations above should be equality. Therefore

$$r_k^2 \|x_j\| = \operatorname{Re} \langle r_k e_k, x_j \rangle, \quad \rho_k^2 \|x_j\| = \operatorname{Im} \langle \rho_k e_k, x_j \rangle,$$

$$\operatorname{Re} \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle = \left\langle \sum_{j=1}^n x_j, \sum_{k=1}^m (r_k + i\rho_k) e_k \right\rangle,$$

and

$$\left| \left\langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{j=1}^n x_j \right\rangle \right| = \left\| \sum_{j=1}^n x_j \right\| \left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|.$$

From Lemma 2.4 and the above equalities we have

$$\begin{aligned} \sum_{j=1}^n x_j &= \frac{\sum_{k=1}^m (r_k + i\rho_k) e_k}{\left\| \sum_{k=1}^m (r_k + i\rho_k) e_k \right\|^2} \left\langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{j=1}^n x_j \right\rangle \\ &= \frac{\sum_{k=1}^m (r_k + i\rho_k) e_k}{\sum_{k=1}^m (r_k^2 + \rho_k^2)} \operatorname{Re} \left\langle \sum_{k=1}^m (r_k + i\rho_k) e_k, \sum_{j=1}^n x_j \right\rangle \\ &= \frac{\sum_{k=1}^m (r_k + i\rho_k) e_k}{\sum_{k=1}^m (r_k^2 + \rho_k^2)} \sum_{k=1}^m \sum_{j=1}^n (r_k^2 \|x_j\| + \rho_k^2 \|x_j\|) \end{aligned}$$

$$= \sum_{j=1}^n \|x_j\| \sum_{k=1}^m (r_k + i\rho_k) e_k,$$

which is the desired result. \square

In the next results of this section, we assume that \mathfrak{X} is a right Hilbert \mathfrak{A} -module, which is an algebraic left A -module subject to

$$\langle x, ay \rangle = a \langle x, y \rangle \quad (x, y \in \mathfrak{X}, a \in \mathfrak{A}). \quad (\dagger)$$

For example if \mathfrak{A} is a unital C^* -algebra and \mathfrak{I} is a commutative right ideal of \mathfrak{A} , then \mathfrak{I} is a right Hilbert module over \mathfrak{A} and

$$\langle x, ay \rangle = x^*(ay) = ax^*y = a \langle x, y \rangle \quad (x, y \in \mathfrak{I}, a \in \mathfrak{A}).$$

The next theorem is a refinement of [7, Theorem 2.1]. To prove it we need the following lemma.

Lemma 2.6. *Let \mathfrak{X} be a Hilbert \mathfrak{A} -module and $e_1, \dots, e_n \in \mathfrak{X}$ be a family of vectors such that $\langle e_i, e_j \rangle = 0$ ($i \neq j$) and $\|e_i\| = 1$. If $x \in \mathfrak{X}$, then*

$$|x|^2 \geq \sum_{k=1}^n |\langle e_k, x \rangle|^2 \quad \text{and} \quad |x|^2 \geq \sum_{k=1}^n |\langle x, e_k \rangle|^2.$$

Proof. The first result follows from the following inequality:

$$\begin{aligned} 0 &\leq \left| x - \sum_{k=1}^n e_k \langle e_k, x \rangle \right|^2 = \left\langle x - \sum_{k=1}^n e_k \langle e_k, x \rangle, x - \sum_{j=1}^n e_j \langle e_j, x \rangle \right\rangle \\ &= \langle x, x \rangle + \sum_{k=1}^n \sum_{j=1}^n \langle e_k, x \rangle^* \langle e_k, e_j \rangle \langle e_j, x \rangle - 2 \sum_{k=1}^n |\langle e_k, x \rangle|^2 \end{aligned}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 12 of 21

Go Back

Full Screen

Close



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 21

Go Back

Full Screen

Close

$$\begin{aligned}
 &= \langle x, x \rangle + \sum_{k=1}^n \langle e_k, x \rangle^* \langle e_k, e_k \rangle \langle e_k, x \rangle - 2 \sum_{k=1}^n |\langle e_k, x \rangle|^2 \\
 &\leq |x|^2 + \sum_{k=1}^n \langle e_k, x \rangle^* \langle e_k, x \rangle - 2 \sum_{k=1}^n |\langle e_k, x \rangle|^2 \\
 &= |x|^2 - \sum_{k=1}^n |\langle e_k, x \rangle|^2.
 \end{aligned}$$

By considering $|x - \sum_{k=1}^n \langle e_k, x \rangle e_k|^2$, we similarly obtain the second one. \square

Now we will prove the next theorem without using the Cauchy–Schwarz inequality.

Theorem 2.7. *Let $e_1, \dots, e_m \in \mathfrak{X}$ be a family of vectors with $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $\|e_i\| = 1$ ($1 \leq i \leq m$). If the vectors $x_1, \dots, x_n \in \mathfrak{X}$ satisfy the conditions*

$$(2.4) \quad 0 \leq r_k \|x_j\| \leq \operatorname{Re} \langle e_k, x_j \rangle, \quad 0 \leq \rho_k \|x_j\| \leq \operatorname{Im} \langle e_k, x_j \rangle,$$

for $1 \leq j \leq n, 1 \leq k \leq m$, where $r_k, \rho_k \in [0, \infty)$ ($1 \leq k \leq m$), then

$$(2.5) \quad \left[\sum_{k=1}^m (r_k^2 + \rho_k^2) \right]^{\frac{1}{2}} \sum_{j=1}^n \|x_j\| \leq \left| \sum_{j=1}^n x_j \right|.$$

Proof. Applying the previous lemma for $x = \sum_{j=1}^n x_j$, we obtain

$$\left| \sum_{j=1}^n x_j \right|^2 \geq \frac{1}{2} \left(\sum_{k=1}^m \left| \left\langle e_k, \sum_{j=1}^n x_j \right\rangle \right|^2 + \sum_{k=1}^m \left| \left\langle \sum_{j=1}^n x_j, e_k \right\rangle \right|^2 \right)$$



$$\begin{aligned}
 &= \sum_{k=1}^m \frac{1}{2} \left(\left\langle e_k, \sum_{j=1}^n x_j \right\rangle^* \left\langle e_k, \sum_{j=1}^n x_j \right\rangle + \left\langle \sum_{j=1}^n x_j, e_k \right\rangle^* \left\langle \sum_{j=1}^n x_j, e_k \right\rangle \right) \\
 &= \sum_{k=1}^m \left(\operatorname{Re} \left\langle e_k, \sum_{j=1}^n x_j \right\rangle \right)^2 + \left(\operatorname{Im} \left\langle e_k, \sum_{j=1}^n x_j \right\rangle \right)^2 \quad (\text{by (1.3)}) \\
 &= \sum_{k=1}^m \left(\operatorname{Re} \sum_{j=1}^n \langle e_k, x_j \rangle \right)^2 + \left(\operatorname{Im} \sum_{j=1}^n \langle e_k, x_j \rangle \right)^2 \\
 &\geq \sum_{k=1}^m \left(r_k^2 \left(\sum_{j=1}^n \|x_j\| \right)^2 + \rho_k^2 \left(\sum_{j=1}^n \|x_j\| \right)^2 \right) \quad (\text{by (2.4)}) \\
 &= \sum_{k=1}^m (r_k^2 + \rho_k^2) \left(\sum_{j=1}^n \|x_j\| \right)^2 .
 \end{aligned}$$

□

Proposition 2.8. *In Theorem 2.7, if $\langle e_k, e_k \rangle = 1$, then the equality holds in (2.5) if and only if*

$$(2.6) \quad \sum_{j=1}^n x_j = \left(\sum_{j=1}^n \|x_j\| \right) \sum_{k=1}^m (r_k + i\rho_k) e_k .$$

Proof. If (2.6) holds, then the inequality in (2.5) turns trivially into equality.

Next, assume that equality holds in (2.5). Then the two inequalities in the proof

Title Page

Contents



Page 14 of 21

Go Back

Full Screen

Close

journal of **inequalities**
 in pure and applied
 mathematics

issn: 1443-5756



of Theorem 2.7 should be equalities. Hence

$$\left| \sum_{j=1}^n x_j \right|^2 = \sum_{k=1}^m \left| \left\langle e_k, \sum_{j=1}^n x_j \right\rangle \right|^2 \quad \text{and} \quad \left| \sum_{j=1}^n x_j \right|^2 = \sum_{k=1}^m \left| \left\langle \sum_{j=1}^n x_j, e_k \right\rangle \right|^2,$$

which is equivalent to

$$\sum_{j=1}^n x_j = \sum_{k=1}^m \sum_{j=1}^n e_k \langle e_k, x_j \rangle = \sum_{k=1}^m \sum_{j=1}^n \langle e_k, x_j \rangle e_k,$$

and

$$r_k \|x_j\| = \operatorname{Re} \langle e_k, x_j \rangle, \quad \rho_k \|x_j\| = \operatorname{Im} \langle e_k, x_j \rangle.$$

So

$$\begin{aligned} \sum_{j=1}^n x_j &= \sum_{k=1}^m \sum_{j=1}^n e_k \langle e_k, x_j \rangle \\ &= \sum_{k=1}^m \sum_{j=1}^n e_k (r_k + i\rho_k) \|x_j\| \\ &= \left(\sum_{j=1}^n \|x_j\| \right) \sum_{k=1}^m (r_k + i\rho_k) e_k. \end{aligned}$$

□

Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian

vol. 10, iss. 4, art. 110, 2009

Title Page

Contents



Page 15 of 21


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 16 of 21

[Go Back](#)

[Full Screen](#)

[Close](#)

3. Additive Reverse of the Triangle Inequality

We now present some versions of the additive reverse of the triangle inequality. In [6], Dragomir established the following theorem:

Theorem 3.1. Let $\{e_k\}_{k=1}^m$ be a family of orthonormal vectors in a Hilbert space H and $M_{jk} \geq 0$ ($1 \leq j \leq n, 1 \leq k \leq m$) such that

$$\|x_j\| - \operatorname{Re}\langle e_k, x_j \rangle \leq M_{jk},$$

for each $1 \leq j \leq n$ and $1 \leq k \leq m$. Then

$$\sum_{j=1}^n \|x_j\| \leq \frac{1}{\sqrt{m}} \left\| \sum_{j=1}^n x_j \right\| + \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk};$$

and the equality holds if and only if

$$\sum_{j=1}^n \|x_j\| \geq \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk},$$


and

$$\sum_{j=1}^n x_j = \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \sum_{k=1}^m e_k.$$

We can prove this theorem for Hilbert C^* -modules using some different techniques.



Theorem 3.2. Let $\{e_k\}_{k=1}^m$ be a family of vectors in a Hilbert module \mathfrak{X} over a C^* -algebra \mathfrak{A} with unit 1, $|e_k| \leq 1$ ($1 \leq k \leq m$), $\langle e_i, e_j \rangle = 0$ ($1 \leq i \neq j \leq m$) and $x_j \in \mathfrak{X}$ ($1 \leq j \leq n$). If for some scalars $M_{jk} \geq 0$ ($1 \leq j \leq n, 1 \leq k \leq m$),

$$(3.1) \quad \|x_j\| - \operatorname{Re}\langle e_k, x_j \rangle \leq M_{jk} \quad (1 \leq j \leq n, 1 \leq k \leq m),$$



Title Page

Contents



Page 17 of 21

Go Back

Full Screen

Close

then

$$(3.2) \quad \sum_{j=1}^n \|x_j\| \leq \frac{1}{\sqrt{m}} \left\| \sum_{j=1}^n x_j \right\| + \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk}.$$

Moreover, if $|e_k| = 1$ ($1 \leq k \leq m$), then the equality in (3.2) holds if and only if

$$(3.3) \quad \sum_{j=1}^n \|x_j\| \geq \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk},$$

and

$$(3.4) \quad \sum_{j=1}^n x_j = \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \sum_{k=1}^m e_k.$$

Proof. Taking the summation in (3.1) over j from 1 to n , we obtain

$$\sum_{j=1}^n \|x_j\| \leq \operatorname{Re} \left\langle e_k, \sum_{j=1}^n x_j \right\rangle + \sum_{j=1}^n M_{jk},$$

for each $k \in \{1, \dots, m\}$. Summing these inequalities over k from 1 to m , we deduce

$$(3.5) \quad \sum_{j=1}^n \|x_j\| \leq \frac{1}{m} \operatorname{Re} \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle + \frac{1}{m} \sum_{k=1}^m \sum_{j=1}^n M_{jk}.$$



Title Page

Contents



Page 18 of 21

Go Back

Full Screen

Close

Using the Cauchy–Schwarz we obtain

$$\begin{aligned}
 (3.6) \quad & \left(\operatorname{Re} \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle \right)^2 \\
 & \leq \frac{1}{2} \left(\left| \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle \right|^2 + \left| \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle^* \right|^2 \right) \\
 & \leq \frac{1}{2} \left(\left\| \sum_{k=1}^m e_k \right\|^2 \left\| \sum_{j=1}^n x_j \right\|^2 + \left\| \sum_{k=1}^m e_k \right\|^2 \left\| \sum_{j=1}^n x_j \right\|^2 \right) \\
 & \leq \left\| \sum_{k=1}^m e_k \right\|^2 \left\| \sum_{j=1}^n x_j \right\|^2 \leq m \left\| \sum_{j=1}^n x_j \right\|^2,
 \end{aligned}$$

since

$$\left\| \sum_{k=1}^m e_k \right\|^2 = \left\| \left\langle \sum_{k=1}^m e_k, \sum_{k=1}^m e_k \right\rangle \right\| = \left\| \sum_{k=1}^m \sum_{l=1}^m \langle e_k, e_l \rangle \right\| = \left\| \sum_{k=1}^m |e_k|^2 \right\| \leq m.$$

Using (3.6) in (3.5), we deduce the desired inequality.

If (3.3) and (3.4) hold, then

$$\begin{aligned}
 \frac{1}{\sqrt{m}} \left\| \sum_{j=1}^n x_j \right\| &= \frac{1}{\sqrt{m}} \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \left\| \sum_{k=1}^m e_k \right\| \\
 &= \sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk},
 \end{aligned}$$



and the equality in (3.2) holds true.

Conversely, if the equality holds in (3.2), then obviously (3.3) is valid and we have equalities throughout the proof above. This means that

$$\|x_j\| - \operatorname{Re}\langle e_k, x_j \rangle = M_{jk},$$

$$\operatorname{Re}\left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle = \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle,$$

and

$$\left| \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle \right| = \left\| \sum_{k=1}^m e_k \right\| \left\| \sum_{j=1}^n x_j \right\|.$$

It follows from Lemma 2.4 and the previous relations that


$$\begin{aligned} \sum_{j=1}^n x_j &= \frac{\sum_{k=1}^m e_k}{\left\| \sum_{k=1}^m e_k \right\|^2} \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle \\ &= \frac{\sum_{k=1}^m e_k}{m} \operatorname{Re} \left\langle \sum_{k=1}^m e_k, \sum_{j=1}^n x_j \right\rangle \\ &= \frac{\sum_{k=1}^m e_k}{m} \sum_{k=1}^m \sum_{j=1}^n (\|x_j\| - M_{jk}) \\ &= \left(\sum_{j=1}^n \|x_j\| - \frac{1}{m} \sum_{j=1}^n \sum_{k=1}^m M_{jk} \right) \sum_{k=1}^m e_k. \end{aligned}$$

□

Title Page	
Contents	
◀◀	▶▶
◀	▶
Page 19 of 21	
Go Back	
Full Screen	
Close	

References

- [1] A.H. ANSARI AND M.S. MOSLEHIAN, Refinements of reverse triangle inequality in inner product spaces, *J. Inequal. Pure Appl. Math.*, **6**(3) (2005), Art. 64. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=537>].
- [2] A.H. ANSARI AND M.S. MOSLEHIAN, More on reverse triangle inequality in inner product spaces, *Inter. J. Math. Math. Sci.*, **18** (2005), 2883–2893.
- [3] Lj. ARAMBAČIĆ AND R. RAJIĆ, On the C^* -valued triangle equality and inequality in Hilbert C^* -modules, *Acta Math. Hungar.*, **119**(4) (2008), 373–380.
- [4] I. BRNETIC, S.S. DRAGOMIR, R. HOXHA AND J. PEČARIĆ, A reverse of the triangle inequality in inner product spaces and applications for polynomials, *Aust. J. Math. Anal. Appl.*, **3**(2) (2006), Art. 9.
- [5] J.B. DIAZ AND F.T. METCALF, A complementary triangle inequality in Hilbert and Banach spaces, *Proc. Amer. Math. Soc.*, **17**(1) (1966), 88–97.
- [6] S.S. DRAGOMIR, Reverses of the triangle inequality in inner product spaces, *Linear Algebra Appl.*, **402** (2005), 245–254.
- [7] S.S. DRAGOMIR, Some reverses of the generalized triangle inequality in complex inner product spaces, *RGMI Res. Rep. Coll.*, **7**(E) (2004), Art. 7.
- [8] S.S. DRAGOMIR, Reverse of the continuous triangle inequality for Bochner integral in complex Hilbert spaces, *J. Math. Anal. Appl.*, **329** (2007), 65–76.
- [9] J. KARAMATA, *Teorija i Praksa Stieltjesova Integrala (Sebro-Coratian)*(*Stieltjes Integral, Theory and Practice*), SANU, Posebna izdanja, **154**, Beograd, 1949.




Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian

vol. 10, iss. 4, art. 110, 2009

Title Page

Contents



Page 20 of 21

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



- [10] M. KATO, K.S. SAITO AND T. TAMURA, Sharp triangle inequality and its reverse in Banach spaces, *Math. Inequal. Appl.*, **10** (2007), 451–460.
- [11] E.C. LANCE, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, 210, Cambridge University Press, Cambridge, 1995.
- [12] V.M. MANUILOV AND E.V. TROITSKY, *Hilbert C^* -Modules*, Translations of Mathematical Monographs, 226. American Mathematical Society, Providence, RI, 2005.
- [13] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic, Dordrecht, 1993.
- [14] M.S. MOSLEHIAN AND L.-E. PERSSON, Reverse Cauchy–Schwarz inequalities for positive C^* -valued sesquilinear forms, *Math. Inequal. Appl.* (to appear).
- [15] J.G. MURPHY, *C^* -Algebras and Operator Theory*, Academic Press, Boston, 1990.
- [16] M. NAKAI AND T. TADA, The reverse triangle inequality in normed spaces, *New Zealand J. Math.*, **25**(2) (1996), 181–193.
- [17] M. PETROVICH, Module d'une somme, *L'Enseignement Math.*, **19** (1917), 53–56.

Reverse Triangle Inequality
Maryam Khosravi, Hakimeh Mahyar
and Mohammad Sal Moslehian

vol. 10, iss. 4, art. 110, 2009

Title Page

Contents



Page 21 of 21

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756