



SECOND ORDER DIFFERENTIAL SUBORDINATIONS OF HOLOMORPHIC MAPPINGS ON BOUNDED CONVEX BALANCED DOMAINS IN C^n

YU-CAN ZHU

Department of Mathematics
Fuzhou University,
Fuzhou, 350002 Fujian, P. R. China
EMail: zhuyucan@fzu.edu.cn

MING-SHENG LIU

Department of Mathematics
South China Normal University,
Guangzhou, 510631 Guangdong, P. R. China
EMail: liumsh@sncu.edu.cn

Received: 09 November, 2006

Accepted: 04 November, 2007

Communicated by: **G. Kohr**

2000 AMS Sub. Class.: 32H02, 30C45.

Key words: Differential subordination, biholomorphic convex mapping, convex balanced domain, Minkowski functional.

Abstract: In this paper, we obtain some second order differential subordinations of holomorphic mappings on a bounded convex balanced domain Ω in C^n . These results imply some first order differential subordinations of holomorphic mappings on a bounded convex balanced domain Ω in C^n . When Ω is the unit disc in the complex plane C , these results are just ones of Miller and Mocanu et al. about differential subordinations of analytic functions on the unit disc in the complex plane C .

2nd Order Differential Subordinations
of Holomorphic Mappings

Yu-Can Zhu and Ming-Sheng Liu

vol. 8, iss. 4, art. 104, 2007

[Title Page](#)

[Contents](#)

[«](#) [»](#)

[◀](#) [▶](#)

Page 1 of 30

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics


issn: 1443-5756

© 2007 Victoria University. All rights reserved.

Acknowledgements:

The authors thank the referee for his helpful comments and suggestions to improve our manuscript.

This research is partly supported by the National Natural Science Foundation of China(No.10471048), the Doctoral Foundation of the Education Committee of China(No.20050574002), the Natural Science Foundation of Fujian Province, China (No.Z0511013) and the Education Commission Foundation of Fujian Province, China (No.JB04038).



**2nd Order Differential Subordinations
of Holomorphic Mappings**

Yu-Can Zhu and Ming-Sheng Liu

vol. 8, iss. 4, art. 104, 2007

Title Page

Contents



Page 2 of 30

Go Back

Full Screen


Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	4
2	Main Results and Their Proofs	7



**2nd Order Differential Subordinations
of Holomorphic Mappings**

Yu-Can Zhu and Ming-Sheng Liu

vol. 8, iss. 4, art. 104, 2007

Title Page

Contents



Page 3 of 30


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 30

Go Back

Full Screen

Close

1. Introduction

Let \mathbb{C}^n be the space of n complex variables $z = (z_1, z_2, \dots, z_n)$ with the Euclidian inner product $\langle z, w \rangle = \sum_{j=1}^n z_j \bar{w}_j$ and the norm $\|z\| = \sqrt{\langle z, z \rangle}$. A domain Ω is called a balanced domain in \mathbb{C}^n if $\lambda z \in \Omega$ for all $z \in \Omega$ and $\lambda \in \mathbb{C}$ with $|\lambda| \leq 1$. The Minkowski functional of the balanced domain Ω is

$$\rho(z) = \inf \left\{ t > 0, \frac{z}{t} \in \Omega \right\}, \quad z \in \mathbb{C}^n.$$

Suppose that Ω is a bounded convex balanced domain in \mathbb{C}^n , and $\rho(z)$ is the Minkowski functional of Ω . Then $\rho(\cdot)$ is a norm of \mathbb{C}^n such that

$$\Omega = \{z \in \mathbb{C}^n : \rho(z) < 1\}, \quad \rho(\lambda z) = |\lambda| \rho(z)$$

for $\lambda \in \mathbb{C}$, $z \in \mathbb{C}^n$ (see [20]).

Let $p_j > 1$ ($j = 1, 2, \dots, n$). Then


$$D_p = \left\{ (z_1, z_2, \dots, z_n) \in \mathbb{C}^n : \sum_{j=1}^n |z_j|^{p_j} < 1 \right\}$$

is a bounded convex balanced domain, and the Minkowski functional $\rho(z)$ of D_p satisfies

$$(1.1) \quad \sum_{j=1}^n \left| \frac{z_j}{\rho(z)} \right|^{p_j} = 1.$$

$\rho(z) = \left(\sum_{j=1}^n |z_j|^{p_j} \right)^{1/p}$ is the Minkowski functional of domain

$$B_p = \left\{ z \in \mathbb{C}^n : \sum_{j=1}^n |z_j|^p < 1 \right\}, \quad \text{where } p > 1.$$



Let $Df(z)$ and $D^2f(z)(\cdot, \cdot)$ denote the first Fréchet derivative and the second Fréchet derivative for a holomorphic mapping $f : \Omega \rightarrow \mathbb{C}^n$ respectively. Then they have the matrix representation

$$Df(z) = \left(\frac{\partial f_j(z)}{\partial z_k} \right)_{1 \leq j, k \leq n}, \quad D^2f(z)(b, \cdot) = \left(\sum_{l=1}^n \frac{\partial^2 f_j(z)}{\partial z_k \partial z_l} b_l \right)_{1 \leq j, k \leq n},$$

where $b = (b_1, b_2, \dots, b_n) \in \mathbb{C}^n$. The mapping $f : \Omega \rightarrow \mathbb{C}^n$ is called locally biholomorphic if the matrix $Df(z)$ is nonsingular at each point z in Ω .

The class of all holomorphic mappings $f : \Omega \rightarrow \mathbb{C}^n$ is denoted by $H(\Omega, \mathbb{C}^n)$. Assume $f, g \in H(\Omega, \mathbb{C}^n)$. Then we say that the mapping f is subordinate to g , written $f \prec g$ or $f(z) \prec g(z)$, if there exists a holomorphic mapping $w : \Omega \rightarrow \Omega$ with $w(0) = 0$ such that $f(z) \equiv g(w(z))$ for all $z \in \Omega$. If g is a biholomorphic mapping, then $f(z) \prec g(z)$ if and only if $f(\Omega) \subset g(\Omega)$ and $f(0) = g(0)$.

In classical results of geometric function theory, differential subordinations provide some simple proofs. They play a key role in the study of some integral operators, differential equations, and properties of subclasses of univalent functions, etc. S.S. Miller and P.T. Mocanu et al. have obtained some deep results for differential subordinations [10, 11, 12, 13, 16, 14]. There is an excellent text *Differential Subordinations Theory and Applications*, by S.S. Miller and P.T. Mocanu [15].

The geometric function theory of several complex variables has been studied by many authors. Many important results for biholomorphic convex or starlike mappings in \mathbb{C}^n have been obtained (see [2, 3]). Some differential subordinations of analytic functions in the complex plane are also extended to \mathbb{C}^n [4, 6, 8, 15, 22]. But there are very few results on second order differential subordinations of holomorphic mappings in \mathbb{C}^n .

In this paper, we obtain some second order differential subordinations of holomorphic mappings on a bounded convex balanced domain Ω in \mathbb{C}^n . These results

Title Page

Contents

◀◀ ▶▶

◀ ▶


Page 5 of 30

Go Back

Full Screen

Close

imply some first order differential subordinations of holomorphic mappings on a bounded convex balanced domain Ω in \mathbb{C}^n . When Ω is the unit disc in the complex plane \mathbb{C} , these results are just those of Miller and Mocanu et al. about differential subordinations of analytic functions on the unit disc in the complex plane \mathbb{C} .



**2nd Order Differential Subordinations
of Holomorphic Mappings**

Yu-Can Zhu and Ming-Sheng Liu

vol. 8, iss. 4, art. 104, 2007

Title Page

Contents



Page 6 of 30

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 7 of 30

Go Back

Full Screen

Close

2. Main Results and Their Proofs

In the following, we always assume that the domain Ω is a bounded convex balanced domain in \mathbb{C}^n and $\rho(z)$ is the Minkowski functional of Ω . Then $\rho(\cdot)$ is a norm of \mathbb{C}^n such that

$$\Omega = \{z \in \mathbb{C}^n : \rho(z) < 1\}, \quad \rho(\lambda z) = |\lambda|\rho(z)$$

for $\lambda \in \mathbb{C}, z \in \mathbb{C}^n$.

In order to derive our main results, we need the following lemmas.

Lemma 2.1. *Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$, and let $w \in H(\Omega, \mathbb{C}^n)$ with $w(z) \neq 0$ and $w(0) = 0$. If $z_0 \in \Omega - \{0\}$ satisfies*

$$\rho(w(z_0)) = \max_{\rho(z) \leq \rho(z_0)} \rho(w(z)),$$


then there exists a real number $t \geq 1/2$ such that

$$(2.1) \quad \left\langle Dw(z_0)(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle = t\rho(w_0),$$

and


$$(2.2) \quad \operatorname{Re} \left\langle D^2w(z_0)(z_0, z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \\ \geq \operatorname{Re} \left\{ \sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_l}(w_0) b_j \bar{b}_l - \sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial z_l}(w_0) b_j b_l \right\} - t\rho(w_0),$$

where $w_0 = w(z_0), Dw(z_0)(z_0) = (b_1, b_2, \dots, b_n)$.



Title Page

Contents



Page 8 of 30

Go Back

Full Screen

Close

Proof. Since $\rho(w_0) = \max_{\rho(z) \leq \rho(z_0)} \rho(w(z))$, then we have $w_0 \neq 0$. Otherwise, there is $w(z) \equiv 0$, which contradicts the hypothesis of Lemma 2.1.

Let $w(z) = (w_1(z), w_2(z), \dots, w_n(z))$, $\gamma(t) = w(e^{it}z_0) = (\gamma_1(t), \gamma_2(t), \dots, \gamma_n(t))$. Then we have $\gamma_j(t) = w_j(e^{it}z_0)$ ($j = 1, 2, \dots, n$), $\gamma(0) = w(z_0) = w_0$ and

$$\frac{d\gamma_j(t)}{dt} = ie^{it} \sum_{k=1}^n \frac{\partial w_j(e^{it}z_0)}{\partial z_k} z_k^0, \quad \overline{\frac{d\gamma_j(t)}{dt}} = -ie^{-it} \sum_{k=1}^n \left(\overline{\frac{\partial w_j(e^{it}z_0)}{\partial z_k} z_k^0} \right),$$

where $z_0 = (z_1^0, z_2^0, \dots, z_n^0)$. Set $L(t) = \rho(\gamma(t))$ ($-\pi \leq t \leq \pi$). Some straightforward calculations yield

$$\begin{aligned} L'(t) &= \sum_{j=1}^n \frac{\partial \rho}{\partial z_j}(\gamma(t)) \cdot \frac{d\gamma_j(t)}{dt} + \sum_{j=1}^n \frac{\partial \rho}{\partial \bar{z}_j}(\gamma(t)) \cdot \frac{d\overline{\gamma_j(t)}}{dt} \\ &= -2 \operatorname{Im} \left[e^{it} \sum_{j,k=1}^n \frac{\partial \rho}{\partial z_j}(\gamma(t)) \cdot \frac{\partial w_j(e^{it}z_0)}{\partial z_k} z_k^0 \right] \\ &= -2 \operatorname{Im} \left\langle Dw(e^{it}z_0)(e^{it}z_0), \overline{\frac{\partial \rho}{\partial z}}(\gamma(t)) \right\rangle, \end{aligned}$$

$$\begin{aligned} L''(t) &= -2 \operatorname{Im} \left[ie^{it} \sum_{j,k=1}^n \frac{\partial \rho}{\partial z_j}(\gamma(t)) \cdot \frac{\partial w_j}{\partial z_k} + ie^{2it} \sum_{j,k=1}^n \frac{\partial \rho}{\partial z_j}(\gamma(t)) \sum_{l=1}^n \frac{\partial^2 w_j}{\partial z_k \partial z_l} z_k^0 z_l^0 \right] \\ &\quad - 2 \operatorname{Im} \left[i \sum_{j,k=1}^n \sum_{l,m=1}^n \frac{\partial^2 \rho}{\partial z_j \partial z_l}(\gamma(t)) \cdot \frac{\partial w_l}{\partial z_m} \cdot (e^{it}z_m^0) \frac{\partial w_j}{\partial z_k} \cdot (e^{it}z_k^0) \right] \\ &\quad + 2 \operatorname{Im} \left[i \sum_{j,k=1}^n \sum_{l,m=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_l}(\gamma(t)) \cdot \overline{\left(\frac{\partial w_l}{\partial z_m} \cdot (e^{it}z_m^0) \right)} \frac{\partial w_j}{\partial z_k} \cdot (e^{it}z_k^0) \right]. \end{aligned}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 9 of 30

Go Back

Full Screen

Close

Noting $L(0) = \max_{-\pi \leq t \leq \pi} L(t)$, we have $L'(0) = 0$ and $L''(0) \leq 0$. It follows that

$$(2.3) \quad \text{Im} \left\langle Dw(z_0)(z_0), \frac{\partial \bar{\rho}}{\partial z}(w_0) \right\rangle = 0,$$

and

$$(2.4) \quad \text{Re} \left\langle D^2w(z_0)(z_0, z_0), \frac{\partial \bar{\rho}}{\partial z}(w_0) \right\rangle + \text{Re} \left\langle Dw(z_0)(z_0), \frac{\partial \bar{\rho}}{\partial z}(w_0) \right\rangle \\ + \text{Re} \left[\sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial z_l}(w_0) b_j b_l - \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_l}(w_0) b_j \bar{b}_l \right] \geq 0.$$

On the other hand, by Schwarz's Lemma in \mathbb{C}^n [19], we have

$$\frac{\rho(w(z))}{\rho(z)} \leq \frac{\rho(w_0)}{\rho(z_0)} \quad \text{for } 0 < \rho(z) \leq \rho(z_0).$$

Let

$$\varphi(r) = \frac{\rho(w(rz_0))}{\rho(rz_0)} = \frac{\rho(w(rz_0))}{r\rho(z_0)}.$$

Then $\varphi(1) = \max_{0 < r \leq 1} \varphi(r)$. It follows that

$$\varphi'(1) = \lim_{r \rightarrow 1^-} \frac{\varphi(r) - \varphi(1)}{r - 1} \geq 0.$$

By a simple calculation, we obtain

$$\varphi'(1) = -\frac{\rho(w_0)}{\rho(z_0)} + \frac{2}{\rho(z_0)} \text{Re} \left\langle Dw(z_0)(z_0), \frac{\partial \bar{\rho}}{\partial z}(w_0) \right\rangle \geq 0.$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 10 of 30

[Go Back](#)

[Full Screen](#)

[Close](#)

If we let

$$t = \frac{1}{\rho(w_0)} \operatorname{Re} \left\langle Dw(z_0)(z_0), \frac{\overline{\partial \rho}}{\partial z}(w_0) \right\rangle,$$

then we have $t \geq 1/2$, therefore (2.1) of Lemma 2.1 holds, and (2.2) follows from (2.3) and (2.4). This completes the proof. \square

Remark 1. Since $\rho(tz) = t\rho(z)$ for $t > 0$, then for $z \in \mathbb{C}^n - \{0\}$, we have

$$(2.5) \quad \rho(z) = \frac{d\rho(tz)}{dt} \Big|_{t=1} = \sum_{j=1}^n \frac{\partial \rho}{\partial z_j} z_j + \sum_{j=1}^n \frac{\partial \rho}{\partial \bar{z}_j} \bar{z}_j = 2 \operatorname{Re} \left\langle z, \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle.$$

For any $z \in \mathbb{C}^n - \{0\}$, we have $\rho(\frac{z}{\rho(z)}) = 1$. Letting $w(z) \equiv z$ in (2.1), we obtain that there exists a real number $t \geq \frac{1}{2}$ such that

$$\left\langle z, \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle = t\rho(z) \geq 0, \quad z \in \mathbb{C}^n - \{0\}.$$

Hence it follows from (2.5) that

$$\rho(z) = 2 \left\langle z, \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle, \quad z \in \mathbb{C}^n - \{0\}.$$

Lemma 2.2 ([10]). Let $g(\xi) = a + b_1\xi + b_2\xi^2 + \dots$ be analytic in $|\xi| < 1$ with $g(\xi) \not\equiv 0$. If $\xi_0 = r_0 e^{i\theta_0}$ ($0 < r_0 < 1$) and $\operatorname{Re} g(\xi_0) = \min_{|\xi| \leq r_0} \operatorname{Re} g(\xi)$, then

$$(2.6) \quad \xi_0 g'(\xi_0) \leq -\frac{|a - g(\xi_0)|^2}{2 \operatorname{Re}(a - g(\xi_0))},$$

and

$$(2.7) \quad \operatorname{Re}\{\xi_0^2 g''(\xi_0) + \xi_0 g'(\xi_0)\} \leq 0.$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 11 of 30

[Go Back](#)

[Full Screen](#)

[Close](#)

Lemma 2.3. Suppose that $\rho(z)$ is differentiable in $\Omega - \{0\}$. Let $h : \Omega \rightarrow \mathbb{C}^n$ be a biholomorphic convex mapping with $h(0) = 0$. Then for every $z \in \Omega - \{0\}$, we have

$$\left| 2 \left\langle Dh(z)^{-1}h(z), \overline{\frac{\partial \rho}{\partial z}}(z) \right\rangle - \rho(z) \right| \leq \rho(z).$$

Proof. For each $z \in \Omega - \{0\}$, we let $g(\xi) = \langle Dh(z)^{-1}(h(z) - h(\xi z)), \overline{\frac{\partial \rho}{\partial z}}(z) \rangle$ for $|\xi| \leq 1$. Then $g(\xi)$ is analytic in $|\xi| \leq 1$ and

$$g(\xi) = \left\langle Dh(z)^{-1}h(z), \overline{\frac{\partial \rho}{\partial z}}(z) \right\rangle + b_1\xi + \dots.$$

From the result in [3, 7], we have $\operatorname{Re} g(\xi) > 0$ for all $|\xi| < 1$. Hence we obtain

$$0 = \operatorname{Re} g(1) = \min_{|\xi| \leq 1} \operatorname{Re} g(\xi).$$

By a simple calculation, we may obtain

$$g'(1) = - \left\langle Dh(z)^{-1}Dh(z)(z), \overline{\frac{\partial \rho}{\partial z}}(z) \right\rangle = - \left\langle z, \overline{\frac{\partial \rho}{\partial z}}(z) \right\rangle = -\frac{\rho(z)}{2}.$$

By (2.6), we have

$$-\rho(z) \operatorname{Re} a + |a|^2 \leq 0,$$

where $a = \left\langle (Dh(z)^{-1}h(z), \overline{\frac{\partial \rho}{\partial z}}(z)) \right\rangle$. It follows that

$$\left| 2 \left\langle Dh(z)^{-1}h(z), \overline{\frac{\partial \rho}{\partial z}}(z) \right\rangle - \rho(z) \right| \leq \rho(z).$$

□



Title Page

Contents



Page 12 of 30

Go Back

Full Screen

Close

Lemma 2.4 ([23]). Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$. If $f : \Omega \rightarrow \mathbb{C}^n$ is a biholomorphic convex mapping, then we have

$$(2.8) \quad \operatorname{Re} \left\{ \sum_{l,m=1}^n \frac{\partial^2 \rho}{\partial z_l \partial z_m} b_l b_m + \sum_{l,m=1}^n \frac{\partial^2 \rho}{\partial z_l \partial \bar{z}_m} b_l \bar{b}_m - \left\langle Df(z)^{-1} D^2 f(z)(b, b), \frac{\partial \rho}{\partial z} \right\rangle \right\} \geq 0$$

for every $z = (z_1, z_2, \dots, z_n) \in \Omega - \{0\}$, $b = (b_1, b_2, \dots, b_n) \in \mathbb{C}^n$ with $\operatorname{Re} \left\langle b, \frac{\partial \rho}{\partial z} \right\rangle = 0$.

Lemma 2.5. Assume that $\rho(z)$ is differentiable in $\Omega - \{0\}$. Then

$$(2.9) \quad \rho(z) = 2 \left\langle z, \frac{\partial \rho}{\partial z}(z) \right\rangle, \quad z \in \mathbb{C}^n - \{0\},$$

and

$$(2.10) \quad \left| 2 \left\langle w, \frac{\partial \rho}{\partial z}(z) \right\rangle \right| \leq \rho(w), \quad z \in \mathbb{C}^n - \{0\}, w \in \mathbb{C}^n.$$

Proof. From Remark 1, we only need to prove (2.10). Let $z \in \mathbb{C}^n - \{0\}$ and

$$\Omega_z = \{w \in \mathbb{C}^n : \rho(w) < \rho(z)\}.$$

Then Ω_z is a convex domain in \mathbb{C}^n , and $\frac{\partial \rho}{\partial z}(z)$ is the normal vector of $\partial \Omega_z$ at z . For every $z, w \in \mathbb{C}^n$ with $\rho(z) = 1, \rho(w) = 1$, we have $\operatorname{Re} \left\langle z - w, \frac{\partial \rho}{\partial z}(z) \right\rangle \geq 0$. It follows that

$$(2.11) \quad 2 \operatorname{Re} \left\langle w, \frac{\partial \rho}{\partial z}(z) \right\rangle \leq 2 \operatorname{Re} \left\langle z, \frac{\partial \rho}{\partial z}(z) \right\rangle = \rho(z) = 1.$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 13 of 30

Go Back

Full Screen

Close

When $\left\langle w, \frac{\partial \bar{\rho}}{\partial z}(z) \right\rangle = 0$, it is obvious that (2.10) holds.

When $\left\langle w, \frac{\partial \bar{\rho}}{\partial z}(z) \right\rangle \neq 0$, then $\rho(w) \neq 0$. Using $\frac{z}{\rho(z)}$ to substitute for z and $\frac{w}{\rho(w)}e^{-i\theta}$ to substitute for w in (2.11), we obtain

$$\left| 2 \operatorname{Re} \left\langle w, \frac{\partial \bar{\rho}}{\partial z}(z) \right\rangle \right| \leq \rho(w), \quad z \in \mathbb{C}^n - \{0\}, \quad w \in \mathbb{C}^n,$$

where $\theta = \arg \left\langle w, \frac{\partial \bar{\rho}}{\partial z}(z) \right\rangle$ and $\frac{\partial \rho}{\partial z}(\lambda z) = \frac{\partial \rho}{\partial z}(z)$ for all $\lambda \in (0, +\infty)$ and $z \in \Omega - \{0\}$. This completes the proof. \square

Lemma 2.6. *Suppose that $\rho(z)$ is differentiable in $\Omega - \{0\}$, and let $h : \Omega \rightarrow \mathbb{C}^n$ be a biholomorphic convex mapping with $h(0) = 0$. Then for every $z \in \Omega - \{0\}$ and vector $\xi \in \mathbb{C}^n$, the inequality*

$$\left| 2 \left\langle Dh(z)^{-1}(\xi), \frac{\partial \bar{\rho}}{\partial z}(z) \right\rangle \right| \leq (1 + \rho(z))^2 \rho(Dh(0)^{-1}(\xi))$$

holds.

Proof. Without loss of generality, we may assume that h is a biholomorphic convex mapping on $\bar{\Omega}$. If not, then we can replace $h(z)$ by $h_r(z) = h(rz)$, where $0 < r < 1$.

For any fixed $z \in \Omega - \{0\}$, from the proof of Theorem 2.1 in [5, 9], there exist $\tilde{z} \in \partial\Omega$ and $\mu \in (0, 1)$ such that $h(z) = \mu h(\tilde{z})$ and

$$1 - \mu \geq \frac{1 - \rho(z)}{1 + \rho(z)}.$$

Let $g(w) = h^{-1}[(1 - \mu)h(w) + \mu h(\tilde{z})]$. Since h is a biholomorphic convex mapping on Ω , then $g \in H(\Omega, \mathbb{C}^n)$ with $g(\Omega) \subset \Omega$ and $g(0) = z$. For every $\xi \in \mathbb{C}^n - \{0\}$,



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 14 of 30

[Go Back](#)

[Full Screen](#)

[Close](#)

we set

$$\psi(\lambda) = 2 \left\langle g \left(\lambda \frac{\xi}{\rho(\xi)} \right), \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle,$$

then $\psi(\lambda)$ is an analytic function in $|\lambda| < 1$. By Lemma 2.5, we obtain

$$|\psi(\lambda)| \leq \rho \left(g \left(\lambda \frac{\xi}{\rho(\xi)} \right) \right) < 1$$

for all $|\lambda| < 1$, and

$$\psi(\lambda) = \rho(z) + 2 \left\langle Dg(0) \left(\frac{\xi}{\rho(\xi)} \right), \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle \lambda + \dots$$

From the classical result in [1], we have $|\psi'(0)| \leq 1 - |\psi(0)|^2$. It follows that

$$\left| 2 \left\langle Dg(0)(\xi), \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle \right| \leq (1 - \rho(z)^2) \rho(\xi).$$

Since $Dh(z)Dg(0) = (1 - \mu)Dh(0)$, then


$$\left| 2 \left\langle Dh(z)^{-1}Dh(0)(\xi), \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle \right| \leq \frac{1}{1 - \mu} (1 - \rho(z)^2) \rho(\xi) \leq (1 + \rho(z))^2 \rho(\xi)$$

for all $\xi \in \mathbb{C}^n$, $z \in \Omega - \{0\}$.

Set $\zeta = Dh(0)(\xi)$, then $\xi = Dh(0)^{-1}\zeta$ and

$$\left| 2 \left\langle Dh(z)^{-1}(\zeta), \frac{\overline{\partial \rho}}{\partial z}(z) \right\rangle \right| \leq (1 + \rho(z))^2 \rho(Dh(0)^{-1}(\zeta)),$$

which completes the proof. □



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 15 of 30

Go Back

Full Screen

Close

Theorem 2.7. Let $f, g \in H(\Omega, \mathbb{C}^n)$ with $f(0) = g(0)$, and let g be biholomorphic convex on $\bar{\Omega}$. Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$. If f is not subordinate to g , then there exist points $z_0 \in \Omega - \{0\}$, $w_0 \in \partial\Omega$ with $0 < \rho(z_0) < 1$, $\rho(w_0) = 1$ and there is a real number $t \geq 1/2$ such that

1. $f(z_0) = g(w_0)$,
2. $\left\langle Dg(w_0)^{-1}Df(z_0)(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle = t$, and
3. $\operatorname{Re} \left\langle Dg(w_0)^{-1}D^2f(z_0)(z_0, z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \geq -t$.

Proof. If f is not subordinate to g , then there exist points $z_0 \in \Omega - \{0\}$, $w_0 \in \partial\Omega$ with $0 < \rho(z_0) < 1$, $\rho(w_0) = 1$ such that $f(z_0) = g(w_0)$ and $f(D_r) \subset g(\Omega)$, where $D_r = \{z \in \mathbb{C}^n : \rho(z) < r\}$ and $r = \rho(z_0)$.

Let $w(z) = g^{-1}(f(z))$. Then $w : D_r \rightarrow \Omega$ is a holomorphic mapping with $w(z) \not\equiv 0$ and $w(0) = 0$ satisfying $f(z) = g(w(z))$ for $z \in D_r$. Hence


$$1 = \rho(w_0) = \max_{\rho(z) \leq \rho(z_0)} \rho(w(z)).$$

By a simple calculation, we have

$$\begin{aligned} Dw(z_0)(z_0) &= Dg(w_0)^{-1}Df(z_0)(z_0), \\ Dg(w_0)^{-1}D^2f(z_0)(z_0, z_0) &= Dg(w_0)^{-1}D^2g(w_0)(Dw(z_0)(z_0), Dw(z_0)(z_0)) \\ &\quad + D^2w(z_0)(z_0, z_0). \end{aligned}$$

From (2.1), there is a real number $t \geq 1/2$ such that

$$\left\langle Dw(z_0)(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle = t\rho(w_0) = t.$$



Title Page

Contents



Page 16 of 30

Go Back

Full Screen

Close

So we obtain

$$\left\langle Dg(w_0)^{-1}Df(z_0)(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle = t,$$

and

$$\begin{aligned} & \operatorname{Re} \left\langle Dg(w_0)^{-1}D^2f(z_0)(z_0, z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \\ &= \operatorname{Re} \left\langle Dg(w_0)^{-1}D^2g(w_0)(Dw(z_0)(z_0), Dw(z_0)(z_0)), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \\ & \quad + \operatorname{Re} \left\langle D^2w(z_0)(z_0, z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \\ & \geq \operatorname{Re} \left\langle Dg(w_0)^{-1}D^2g(w_0)(a, a), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \\ & \quad + \operatorname{Re} \left\{ \sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_l}(w_0) a_j \bar{a}_l - \sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial z_l}(w_0) a_j a_l \right\} - t, \end{aligned}$$

where $a = Dw(z_0)(z_0) = (a_1, a_2, \dots, a_n)$. If we let $b = (b_1, b_2, \dots, b_n)$ with $b_j = ia_j$, then we have

$$\operatorname{Re} \left\langle b, \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle = \operatorname{Re} \left\{ i \left\langle Dw(z_0)(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \right\} = \operatorname{Re}\{it\} = 0.$$

From Lemma 2.4, we obtain

$$\operatorname{Re} \left\langle Dg(w_0)^{-1}D^2f(z_0)(z_0, z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle$$

$$\begin{aligned} &\geq -\operatorname{Re} \left\langle Dg(w_0)^{-1} D^2g(w_0)(b, b), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \\ &\quad + \operatorname{Re} \left\{ \sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_l}(w_0) b_j \bar{b}_l + \sum_{j,l=1}^n \frac{\partial^2 \rho}{\partial z_j \partial z_l}(w_0) b_j b_l \right\} - t \\ &\geq -t. \end{aligned}$$

This completes the proof. \square

Remark 2. When $n = 1$, Ω is the unit disc in the complex plane \mathbb{C} and $\rho(z) = |z|$ ($z \in \mathbb{C}$), we may obtain Lemma 1 in [14] from Theorem 2.7. Theorem 2.7 will play a key role in studying some second order differential subordinations of holomorphic mappings on a bounded convex balanced domain Ω in \mathbb{C}^n .


Let Ω_1 be a set of \mathbb{C}^n , and let h be a biholomorphic convex mapping on $\bar{\Omega}$. Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$. We define $\Psi(\Omega_1, h)$ to be the class of maps $\psi : \mathbb{C}^n \times \mathbb{C}^n \times \mathbb{C}^n \times \Omega \rightarrow \mathbb{C}^n$ that satisfy the following conditions:

1. $\psi(h(0), 0, 0, 0) \in \Omega_1$, and
2. $\psi(\alpha, \beta, \gamma, z) \notin \Omega_1$ for $\alpha = h(w)$, $\left\langle Dh(w)^{-1}(\beta), \overline{\frac{\partial \rho}{\partial z}}(w) \right\rangle = t$, $\operatorname{Re} \left\langle Dh(w)^{-1}(\gamma), \overline{\frac{\partial \rho}{\partial z}}(w) \right\rangle \geq -t$, and $z \in \Omega$, where $\rho(w) = 1$ and $t \geq 1/2$.

Theorem 2.8. Let $\psi \in \Psi(\Omega_1, h)$. If $f \in H(\Omega, \mathbb{C}^n)$ with $f(0) = h(0)$ satisfies



$$(2.12) \quad \psi(f(z), Df(z)(z), D^2f(z)(z, z), z) \in \Omega_1$$

for all $z \in \Omega$, then $f(z) \prec h(z)$.



Title Page

Contents




Page 17 of 30

Go Back

Full Screen

Close



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 18 of 30

Go Back

Full Screen

Close

Proof. If f is not subordinate to h , then by Theorem 2.7, there exist points $z_0 \in \Omega - \{0\}$, $w_0 \in \partial\Omega$ with $0 < \rho(z_0) < 1$, $\rho(w_0) = 1$ and there is a real number $t \geq 1/2$ such that

$$f(z_0) = h(w_0), \quad \left\langle Dh(w_0)^{-1}Df(z_0)(z_0), \frac{\partial \bar{\rho}}{\partial z}(w_0) \right\rangle = t,$$

and

$$\operatorname{Re} \left\langle Dh(w_0)^{-1}D^2f(z_0)(z_0, z_0), \frac{\partial \bar{\rho}}{\partial z}(w_0) \right\rangle \geq -t.$$

Set $\alpha = f(z_0)$, $\beta = Df(z_0)(z_0)$, $\gamma = D^2f(z_0)(z_0, z_0)$, then according to the definition of $\Psi(\Omega_1, h)$, we have

$$\psi(f(z_0), Df(z_0)(z_0), D^2f(z_0)(z_0, z_0), z_0) \notin \Omega_1$$

which contradicts (2.12). Hence $f(z) \prec h(z)$, and the proof of Theorem 2.8 is complete. \square


Theorem 2.9. Let $A \geq 0$, $h \in H(\Omega, \mathbb{C}^n)$ be biholomorphic convex with $h(0) = 0$, and let $\psi(z) \in H(\Omega, \mathbb{C}^n)$ with $\psi(0) = 0$. Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$, $k > 4\|Dh(0)^{-1}\|$, and $\varphi, \phi : \Omega_1 \times \Omega \rightarrow \mathbb{C}$ are holomorphic such that

$$\operatorname{Re} \phi(\alpha, z) \geq A + |\varphi(\alpha, z) - 1| - \operatorname{Re}[\varphi(\alpha, z) - 1] + k\rho(\psi(z))$$

for all $(\alpha, z) \in h(\Omega) \times \Omega$, where Ω_1 is a domain of \mathbb{C}^n with $h(\Omega) \subset \Omega_1$ and $\|Dh(0)^{-1}\| = \sup_{\rho(\xi) \leq 1} \rho(Dh(0)^{-1}(\xi))$. If $f \in H(\Omega, \mathbb{C}^n)$ with $f(0) = 0$ satisfies

$$AD^2f(z)(z, z) + \phi(f(z), z)Df(z)(z) + \varphi(f(z), z)f(z) + \psi(z) \prec h(z),$$

then $f(z) \prec h(z)$.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 19 of 30

Go Back

Full Screen

Close

Proof. Without loss of generality, we may assume that f and g satisfy the conditions of Theorem 2.9 on $\bar{\Omega}$. If not, then we can replace $f(z)$ by $f_r(z) = f(rz)$, $\psi(z)$ by $\psi_r(z) = \psi(rz)$, and $h(z)$ by $h_r(z) = h(rz)$, where $0 < r < 1$. We would then prove $f_r(z) \prec h_r(z)$ for all $0 < r < 1$. By letting $r \rightarrow 1^-$, we obtain $f(z) \prec h(z)$.

Let

$$\psi(\alpha, \beta, \gamma, z) = A\gamma + \phi(\alpha, z)\beta + \varphi(\alpha, z)\alpha + \psi(z),$$

and let $\alpha = h(w)$, $\left\langle Dh(w)^{-1}(\beta), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle = t$, $\operatorname{Re} \left\langle Dh(w)^{-1}(\gamma), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle \geq -t$, where $\rho(w) = 1$, $t \geq 1/2$. If we set

$$\psi(\alpha, \beta, \gamma, z) = h(w) + \lambda Dh(w)(w),$$

then we have

$$\begin{aligned} \lambda w &= ADh(w)^{-1}(\gamma) + \phi(\alpha, z)Dh(w)^{-1}(\beta) \\ &\quad + [\varphi(\alpha, z) - 1]Dh(w)^{-1}h(w) + Dh(w)^{-1}(\psi(z)). \end{aligned}$$

Since $2 \left\langle w, \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle = \rho(w) = 1$ from Remark 1, we obtain

$$\begin{aligned} (2.13) \quad \lambda &= 2A \left\langle Dh(w)^{-1}(\gamma), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle + 2\phi(\alpha, z) \left\langle Dh(w)^{-1}(\beta), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle \\ &\quad + 2[\varphi(\alpha, z) - 1] \left\langle Dh(w)^{-1}h(w), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle + 2 \left\langle Dh(w)^{-1}(\psi(z)), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle. \end{aligned}$$

By Lemma 2.3, we have

$$\left| 2 \left\langle Dh(w)^{-1}h(w), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle - 1 \right| \leq 1.$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 20 of 30

Go Back

Full Screen

Close

By Lemma 2.6, we obtain

$$\begin{aligned}
 \operatorname{Re} \lambda &\geq -2At + 2t \operatorname{Re} \phi(\alpha, z) + \operatorname{Re}[\varphi(\alpha, z) - 1] \\
 &\quad - |\varphi(\alpha, z) - 1| - 4\|Dh(0)^{-1}\|\rho(\psi(z)) \\
 &\geq (2t - 1)\{|\varphi(\alpha, z) - 1| - \operatorname{Re}[\varphi(\alpha, z) - 1]\} \\
 &\quad + (k - 4\|Dh(0)^{-1}\|)\rho(\psi(z)) \geq 0.
 \end{aligned}
 \tag{2.14}$$

Now we verify that $\psi(\alpha, \beta, \gamma, z) \notin h(\Omega)$. Suppose not, then there exists $w_1 \in \Omega$ such that $\psi(\alpha, \beta, \gamma, z) = h(w_1)$. From the result in [3, 7, 18, 19], we have

$$-\operatorname{Re} \lambda = 2 \operatorname{Re} \left\langle Dh(w)^{-1}(h(w) - h(w_1)), \frac{\partial \bar{\rho}}{\partial z}(w) \right\rangle > 0,$$

which contradicts (2.14), hence $\psi(\alpha, \beta, \gamma, z) \notin h(\Omega)$. By Theorem 2.8, we obtain $f(z) \prec h(z)$, and the proof is complete. \square

Corollary 2.10. Let $A \geq 0, h \in H(\Omega, \mathbb{C}^n)$ be biholomorphic convex with $h(0) = 0$, and let $\psi(z) \in H(\Omega, \mathbb{C}^n)$ with $\psi(0) = 0$. Suppose that $k > 4\|Dh(0)^{-1}\|, \rho(z)$ is twice differentiable in $\Omega - \{0\}$, and $B(z), C(z) \in H(\Omega, \mathbb{C})$ satisfy

$$\operatorname{Re} B(z) \geq A + |C(z) - 1| - \operatorname{Re}[C(z) - 1] + k\rho(\psi(z))$$

for all $z \in \Omega$, where $\|Dh(0)^{-1}\| = \sup_{\rho(\xi) \leq 1} \rho(Dh(0)^{-1}(\xi))$. If $f \in H(\Omega, \mathbb{C}^n)$ with

$f(0) = 0$ satisfies

$$AD^2 f(z)(z, z) + B(z)Df(z)(z) + C(z)f(z) + \psi(z) \prec h(z),$$

then $f(z) \prec h(z)$.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 21 of 30

Go Back

Full Screen

Close

Corollary 2.11. Let $A \geq 0, h \in H(\Omega, \mathbb{C}^n)$ be biholomorphic convex. Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$, and $B(z) \in H(\Omega, \mathbb{C})$ with $\text{Re } B(z) \geq A$ for all $z \in \Omega$. If $f \in H(\Omega, \mathbb{C}^n)$ with $f(0) = h(0)$ satisfies

$$AD^2 f(z)(z, z) + B(z)Df(z)(z) + f(z) \prec h(z),$$

then $f(z) \prec h(z)$.

Corollary 2.12. Let $h \in H(\Omega, \mathbb{C}^n)$ be biholomorphic convex with $h(0) = 0$. Suppose that $\rho(z)$ is twice differentiable in $\Omega - \{0\}$ and $\phi : \Omega_1 \rightarrow \mathbb{C}$ is holomorphic such that $\text{Re } \phi(h(z)) \geq 0$ for all $z \in \Omega$. If $f \in H(\Omega, \mathbb{C}^n)$ with $f(0) = 0$ satisfies

$$f(z) + \phi(f(z))Df(z)(z) \prec h(z),$$

then $f(z) \prec h(z)$.

Remark 3. When $n = 1$, we have $Df(z)(z) = zf'(z)$ and $D^2 f(z)(z, z) = z^2 f''(z)$. From Corollary 2.10, we may obtain Theorem 2 in [13], Theorem 3.1a in [15], Theorem 1 for case 1 in [12] and Theorem 1 in [14]. From Corollary 2.11, we may obtain Corollary 2.1 in [13].

Example 2.1. Let $\beta > 0$ and $\gamma \in \mathbb{C}$ with $2 \text{Re } \gamma \geq \beta$. The unit ball in \mathbb{C}^n is denoted by $B = \{z \in \mathbb{C}^n : \|z\| < 1\}$. If $u \in \mathbb{C}^n$ with $\|u\| = 1$, then $h(z) = \frac{z}{1 - \langle z, u \rangle}$ is a biholomorphic convex mapping on B (see [17]). By a simple calculation, we have

$$\begin{aligned} (2.15) \quad & \text{Re} \left[\beta \left\langle \frac{z}{1 - \langle z, u \rangle}, u \right\rangle + \gamma \right] \\ &= \frac{\text{Re } \gamma |1 - \langle z, u \rangle|^2 + \beta [\text{Re} \langle z, u \rangle - |\langle z, u \rangle|^2]}{|1 - \langle z, u \rangle|^2} \\ &\geq \frac{\beta |1 - \langle z, u \rangle|^2 + 2\beta [\text{Re} \langle z, u \rangle - |\langle z, u \rangle|^2]}{2|1 - \langle z, u \rangle|^2} \end{aligned}$$

$$= \frac{\beta(1 - |\langle z, u \rangle|^2)}{2|1 - \langle z, u \rangle|^2} > 0$$

for all $z \in B$. If $f \in H(B, \mathbb{C}^n)$ with $f(0) = 0$, then by Corollary 2.12, we have

$$f(z) + \frac{Df(z)(z)}{\beta \langle f(z), u \rangle + \gamma} \prec \frac{z}{1 - \langle z, u \rangle} \implies f(z) \prec \frac{z}{1 - \langle z, u \rangle}.$$

Example 2.2. Let $A \geq 0, \beta \geq 0$ and $\gamma \in \mathbb{C}$ with $\operatorname{Re} \gamma \geq \beta/2 + A$, $u \in \mathbb{C}^n$ with $\|u\| = 1$. If $f \in H(B, \mathbb{C}^n)$ with $f(0) = 0$, then by Theorem 2.9, Corollary 2.11 and (2.15), we have

$$AD^2 f(z)(z, z) + \left[\beta \frac{\langle z, u \rangle}{1 - \langle z, u \rangle} + \gamma \right] Df(z)(z) + f(z) \prec \frac{z}{1 - \langle z, u \rangle} \implies f(z) \prec \frac{z}{1 - \langle z, u \rangle},$$

and

$$AD^2 f(z)(z, z) + [\beta \langle f(z), u \rangle + \gamma] Df(z)(z) + f(z) \prec \frac{z}{1 - \langle z, u \rangle} \implies f(z) \prec \frac{z}{1 - \langle z, u \rangle}.$$



Let $\rho(z)$ be differentiable in $\Omega - \{0\}$. For $M > 0$, we define $\Psi(M)$ to be the class of maps $\psi : \mathbb{C}^n \times \mathbb{C}^n \times \mathbb{C}^n \times \Omega \rightarrow \mathbb{C}^n$ that satisfy the following conditions:

1. $\rho(\psi(0, 0, 0, 0)) < M$ and
2. $\rho(\psi(\alpha, \beta, \gamma, z)) \geq M$ for all $\rho(\alpha) = M$, $2 \left\langle \beta, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle = tM$, $2 \operatorname{Re} \left\langle \gamma, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle \geq (t^2 - t)M$, and $t \geq 1$.



Title Page

Contents



Page 22 of 30

Go Back

Full Screen

Close



Title Page

Contents



Page 23 of 30

Go Back

Full Screen

Close

Theorem 2.13. Let $\psi \in \Psi(M)$. If $w \in H(\Omega, \mathbb{C}^n)$ with $w(0) = 0$ satisfies

$$(2.16) \quad \rho(\psi(w(z), Dw(z)(z), D^2w(z)(z, z), z)) < M$$

for all $z \in \Omega - \{0\}$, then $\rho(w(z)) < M$ for $z \in \Omega$.

Proof. Suppose that the conclusion of Theorem 2.13 is false. Then there exists a point $z_0 \in \Omega - \{0\}$ such that $\rho(w(z_0)) = M$ and $\rho(w(z)) \leq M$ for $\rho(z) \leq \rho(z_0)$. It implies $w_0 = w(z_0) \neq 0$.

Let

$$\varphi(\xi) = 2 \left\langle w \left(\frac{z_0}{\rho(z_0)} \xi \right), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle, \quad \xi \in \mathbb{C}.$$

Then $\varphi(\xi)$ is an analytic function in $|\xi| < 1$. By Lemma 2.5, we have

$$|\varphi(\xi)| \leq \left| 2 \left\langle w \left(\frac{z_0}{\rho(z_0)} \xi \right), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle \right| \leq \rho \left(w \left(\frac{z_0}{\rho(z_0)} \xi \right) \right) \leq \rho(w(z_0))$$

for all $|\xi| \leq \rho(z_0)$, and

$$\varphi(\rho(z_0)) = 2 \left\langle w(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle = \rho(w(z_0)) = \max_{|\xi| \leq \rho(z_0)} |\varphi(\xi)|.$$

By a simple calculation, we have

$$\rho(z_0)\varphi'(\rho(z_0)) = 2 \left\langle Dw(z_0)(z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle,$$

$$\rho(z_0)^2\varphi''(\rho(z_0)) = 2 \left\langle D^2w(z_0)(z_0, z_0), \overline{\frac{\partial \rho}{\partial z}}(w_0) \right\rangle.$$

Using Lemma A in [10] (also see [15, p. 19]), there exists a real number $t \geq 1$ such that

$$2 \left\langle Dw(z_0)(z_0), \frac{\partial \rho}{\partial z}(w_0) \right\rangle = tM,$$

$$2 \operatorname{Re} \left\langle D^2w(z_0)(z_0, z_0), \frac{\partial \rho}{\partial z}(w_0) \right\rangle \geq (t^2 - t)M.$$

By the definition of ψ , we have

$$\rho(\psi(w(z_0), Dw(z_0)(z_0), D^2w(z_0)(z_0, z_0), z_0)) \geq M,$$

which contradicts (2.16). Hence $\rho(w(z)) < M$ for $z \in \Omega$, and the proof is complete. \square

Theorem 2.14. *Let $\rho(z)$ be differentiable in $\Omega - \{0\}$. Suppose that $A(z), B(z), C(z) \in H(\Omega, \mathbb{C})$ with $A(z) \neq 0$ for all $z \in \Omega$ satisfy*

$$(2.17) \quad \operatorname{Re} \frac{B(z)}{A(z)} \geq \max \left\{ -1, \frac{1}{|A(z)|} - \operatorname{Re} \frac{C(z)}{A(z)} + \frac{\rho(\varphi(z))}{|A(z)|} \right\},$$

or


$$(2.18) \quad \operatorname{Re} \frac{C(z)}{A(z)} \geq \frac{1}{|A(z)|} + \frac{\rho(\varphi(z))}{|A(z)|} + 1$$

$$\text{and } 1 - 2\sqrt{\operatorname{Re} \frac{C(z)}{A(z)} - \frac{1}{|A(z)|} - \frac{\rho(\varphi(z))}{|A(z)|}} \leq \operatorname{Re} \frac{B(z)}{A(z)} \leq -1$$

for all $z \in \Omega$. If $w(z) \in H(\Omega, \mathbb{C}^n)$ with $w(0) = 0$ satisfies

$$\rho(A(z)D^2w(z)(z, z) + B(z)Dw(z)(z) + C(z)w(z) + \varphi(z)) < 1$$

for all $z \in \Omega$, then $\rho(w(z)) < 1$ for $z \in \Omega$.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 24 of 30

Go Back

Full Screen

Close



Title Page

Contents



Page 25 of 30

Go Back

Full Screen

Close

Proof. Let

$$\psi(\alpha, \beta, \gamma, z) = A(z)\gamma + B(z)\beta + C(z)\alpha + \varphi(z),$$

where $\rho(\alpha) = 1$, $2 \left\langle \beta, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle = t$, $2 \operatorname{Re} \left\langle \gamma, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle \geq (t^2 - t)$ and $t \geq 1$. From (2.9) and (2.10), we have

$$\begin{aligned} \rho(\psi(\alpha, \beta, \gamma, z)) &\geq \left| 2 \left\langle \psi(\alpha, \beta, \gamma, z) e^{-i\theta}, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle \right| \\ &= \left| |A(z)| 2 \left\langle \gamma, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle + B(z) e^{-i\theta} 2 \left\langle \beta, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle \right. \\ &\quad \left. + C(z) e^{-i\theta} 2 \left\langle \alpha, \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle + 2 e^{-i\theta} \left\langle \varphi(z), \frac{\partial \bar{\rho}}{\partial z}(\alpha) \right\rangle \right| \\ &\geq |A(z)| \left\{ t^2 + t \left[\operatorname{Re} \frac{B(z)}{A(z)} - 1 \right] + \operatorname{Re} \frac{C(z)}{A(z)} - \frac{\rho(\varphi(z))}{|A(z)|} \right\}, \end{aligned}$$

where $\theta = \arg A(z)$. Let


$$L(t) = t^2 + t \left[\operatorname{Re} \frac{B(z)}{A(z)} - 1 \right] + \operatorname{Re} \frac{C(z)}{A(z)} - \frac{\rho(\varphi(z))}{|A(z)|}$$

for $t \geq 1$. Then we have

$$L'(t) = 2t + \operatorname{Re} \frac{B(z)}{A(z)} - 1.$$



If $\operatorname{Re} \frac{B(z)}{A(z)} \geq -1$ for $z \in \Omega$, then $L'(t) \geq \operatorname{Re} \frac{B(z)}{A(z)} + 1 \geq 0$. Hence we obtain

$$\min_{t \geq 1} L(t) = L(1) = \operatorname{Re} \frac{B(z)}{A(z)} + \operatorname{Re} \frac{C(z)}{A(z)} - \frac{\rho(\varphi(z))}{|A(z)|} \geq \frac{1}{|A(z)|}.$$



Title Page

Contents



Page 26 of 30

Go Back

Full Screen

Close

It follows that $\rho(\psi(\alpha, \beta, \gamma, z)) \geq 1$.

If $\operatorname{Re} \frac{B(z)}{A(z)} \leq -1$ for $z \in \Omega$, then

$$\begin{aligned} \min_{t \geq 1} L(t) &= L\left(\frac{1}{2} \left(1 - \operatorname{Re} \frac{B(z)}{A(z)}\right)\right) \\ &= -\frac{1}{4} \left(\operatorname{Re} \frac{B(z)}{A(z)} - 1\right)^2 + \operatorname{Re} \frac{C(z)}{A(z)} - \frac{\rho(\varphi(z))}{|A(z)|} \\ &\geq \frac{1}{|A(z)|}. \end{aligned}$$

It also follows that $\rho(\psi(\alpha, \beta, \gamma, z)) \geq 1$.

Hence we have $\psi \in \Psi(1)$. From Theorem 2.13, we obtain $\rho(w(z)) < 1$ for $z \in \Omega$. \square

Remark 4. Setting $n = 1$, $\varphi(z) \equiv 0$, $A(z) \equiv A$ and $C(z) = 1 - B(z)$ in Theorem 2.14, we get Theorem 4 in [13].


Corollary 2.15. *Suppose that $B(z) \in H(B, \mathbb{C})$ and $A \geq 0$ satisfy $\operatorname{Re} B(z) \geq 0$ for all $z \in B$. If $w(z) \in H(B, \mathbb{C}^m)$ with $w(0) = 0$ satisfy*

$$\|AD^2w(z)(z, z) + B(z)Dw(z)(z) + w(z)\| < 1$$

for all $z \in B$, then $\|w(z)\| < 1$ for $z \in B$.



Example 2.3. Let k and n_1 be positive integers and let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_m) \in \mathbb{C}^m$. We define $\alpha^k = (\alpha_1^k, \alpha_2^k, \dots, \alpha_m^k)$. Suppose $A_1, A_2, \dots, A_{n_1} \in H(B, \mathbb{C})$ ($n_1 \geq 2$) satisfy

$$\operatorname{Re} A_1(z) \geq \sum_{k=2}^{n_1} |A_k(z)|$$



Title Page

Contents



Page 27 of 30

Go Back

Full Screen

Close

for $z \in B$, where B is the unit ball in \mathbb{C}^n . If $w(z) = (w_1(z), w_2(z), \dots, w_m(z)) \in H(B, \mathbb{C}^m)$ with $w(0) = 0$ satisfies

$$\sum_{v=1}^m \left| \sum_{j,l=1}^n \frac{\partial^2 w_v(z)}{\partial z_j \partial z_l} z_j z_l + \sum_{j=1}^n \frac{\partial w_v(z)}{\partial z_j} z_j + \sum_{k=1}^{n_1} A_k(z) [w_v(z)]^k \right|^2 < 1$$

for all $z \in B$, then $\sum_{v=1}^m |w_v(z)|^2 < 1$ for all $z \in B$.

In fact, if we let

$$\psi(\alpha, \beta, \gamma, z) = \gamma + \beta + \sum_{k=1}^{n_1} A_k(z) \alpha^k$$


for $\|\alpha\| = 1$, $\langle \beta, \alpha \rangle = t$, $\operatorname{Re} \langle \gamma, \alpha \rangle \geq t^2 - t$ and $t \geq 1$, then we have

$$\begin{aligned} \|\psi(\alpha, \beta, \gamma, z)\| &\geq \left| \langle \gamma, \alpha \rangle + \langle \beta, \alpha \rangle + A_1(z) + \sum_{k=2}^{n_1} A_k(z) \langle \alpha^k, \alpha \rangle \right| \\ &\geq \operatorname{Re} \{ \langle \gamma, \alpha \rangle + \langle \beta, \alpha \rangle + A_1(z) \} - \sum_{k=2}^{n_1} |A_k(z)| \left(\sum_{j=1}^m |\alpha_j|^{k+1} \right) \\ &\geq t^2 + \operatorname{Re} A_1(z) - \sum_{k=2}^{n_1} |A_k(z)| \geq 1. \end{aligned}$$

Hence $\psi \in \Psi(1)$ for $\rho(z) = \sqrt{\sum_{j=1}^n |z_j|^2}$. According to Theorem 2.13, we have $\sum_{v=1}^m |w_v(z)|^2 < 1$ for all $z \in B$.

References

- [1] J.B. CONWAY, *Functions of One Complex Variable* (Second Edition), Springer-Verlag, New York, 1978.
- [2] I. GRAHAM AND G. KOHR, *Geometric Function Theory in One and Higher Dimensions*, Dekker, New York, 2003.
- [3] S. GONG, *Convex and Starlike Mappings in Several Complex Variables*, Science Press/Kluwer Academic Publishers, 1998.
- [4] S. GONG, A note on partial differential inequalities, *Chin. Ann. of Math.*, **5A**(6) (1984), 771–780 (Chinese).
- [5] S. GONG AND T. LIU, Distortion theorems for biholomorphic convex mappings on bounded convex circular domains, *Chin. Ann. of Math.*, **20B**(3) (1999), 297–304.
- [6] S. GONG AND S.S. MILLER, Partial differential subordinations and inequalities define on complete circular domains, *Comm. Partial Diff. Equation*, **11**(11) (1986), 1243–1255.
- [7] S. GONG AND T. LIU, Criterion for the family of ε starlike mappings, *J. Math. Anal. Appl.*, **274** (2002), 696–704.
- [8] H. HAMADA, G. KOHR AND M. KOHR, First order partial differential subordinations on bounded balanced pseudoconvex domains in \mathbb{C}^n , *Mathematica (Cluj)*, **41(64)** (1999), 161–175.
- [9] T. LIU AND G. REN, The growth theorem of convex mapping on bounded convex circular domains, *Science in China (Series A)*, **41**(2) (1998), 123–130.



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 28 of 30

Go Back

Full Screen

Close



- [10] S.S. MILLER AND P.T. MOCANU, Second order differential inequalities in the complex plane, *J. Math. Anal. Appl.*, **65** (1978), 289–305.
- [11] S.S. MILLER AND P.T. MOCANU, Differential subordinations and univalent functions, *Michigan Math. J.*, **28** (1981), 157–171.
- [12] S.S. MILLER AND P.T. MOCANU, On some classes of first-order differential subordinations, *Michigan Math. J.*, **35** (1985), 185–195.
- [13] S.S. MILLER AND P.T. MOCANU, Differential subordinations and inequalities in the complex plane, *J. Diff. Equation*, **67**(2) (1987), 199–211.
- [14] S.S. MILLER AND P.T. MOCANU, Averaging operator and generalized Robinson differential inequality, *J. Math. Anal. Appl.*, **173** (1993), 459–467.
- [15] S.S. MILLER AND P.T. MOCANU, *Differential Subordinations Theory and Applications*, New York, Marcel Dekker Inc. 2000.
- [16] P.T. MOCANU, Second order averaging operators for analytic functions, *Rev. Roumaine Math. Appl.*, **33**(10) (1988), 875–881.
- [17] K. ROPER AND T. SUFFRIDGE, Convex mappings on the unit ball of \mathbb{C}^n , *J. d'Analyse Math.*, **65** (1995), 333–347.
- [18] T.J. SUFFRIDGE, Starlike and convex maps in Banach spaces, *Pacific J. Math.*, **46**(2) (1973), 475–489.
- [19] T.J. SUFFRIDGE, Starlikeness, convexity and other geometric properties of holomorphic maps in higher dimensions, *Lecture Notes in Math.*, **599** (1976), 146–159.
- [20] A.E. TAYLOR AND D.C. LAY, *Introduction to Functional Analysis*, New York: John Wiley & Sons Inc., 1980, 111–115.

Title Page

Contents



Page 29 of 30

Go Back

Full Screen

Close

- [21] YUCAN ZHU AND HEZENG LIN, First-order differential subordinations, *Acta Sci. Nature Unvi. Sunyatseni*, **30**(4) (1991), 114–118 (Chinese).
- [22] YUCAN ZHU, Differential subordinations of holomorphic mappings in Banach spaces, *J. of Fuzhou University*, **30**(4) (2002), 430–434 (Chinese).
- [23] YUCAN ZHU, Criteria of biholomorphic convex mappings on bounded convex balanced domains, *Acta Math. Sinica*, **46**(6) (2003), 1153–1162 (Chinese).



2nd Order Differential Subordinations
of Holomorphic Mappings

Yu-Can Zhu and Ming-Sheng Liu

vol. 8, iss. 4, art. 104, 2007

Title Page

Contents



Page 30 of 30

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756