

# NOTE ON AN OPEN PROBLEM

**LAZHAR BOUGOFFA**

Al-imam Muhammad Ibn Saud Islamic University  
Faculty of Science, Department of Mathematics  
P.O.Box 84880, Riyadh 11681, Saudi Arabia  
EMail: [bougoffa@hotmail.com](mailto:bougoffa@hotmail.com)

*Received:* 17 December, 2006

*Accepted:* 1 April, 2007

*Communicated by:* **P.S. Bullen**

*2000 AMS Sub. Class.:* 26D15.

*Key words:* Integral inequality.

*Abstract:* The aim of this short note is to establish an integral inequality and its reverse which give an affirmative answer to an open problem posed by QUÔC ANH NGÔ, DU DUC THANG, TRANT TAT DAT, and DANG ANH TUAN, in the paper [*Notes on an integral inequality*, J. Ineq. Pure and Appl. Math., 7(4)(2006), Art. 120.]



---

Note on an Open Problem



Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

---

[Title Page](#)

[Contents](#)



Page 1 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

# Contents

1	<a href="#">Introduction</a>	3
2	<a href="#">Main Results</a>	5



---

## Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

---

[Title Page](#)

[Contents](#)



Page 2 of 10

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

Title Page

Contents



Page 3 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

## 1. Introduction

Very recently, in the paper [1] the authors studied some integral inequalities and proposed the following open problem:

**Problem 1.1.** Let  $f$  be a continuous function on  $[0, 1]$  satisfying

$$(1.1) \quad \int_x^1 f(t)dt \geq \int_x^1 tdt, \quad \forall x \in [0, 1].$$

Under what conditions does the inequality

$$(1.2) \quad \int_0^1 f^{\alpha+\beta}(x)dx \geq \int_0^1 x^\alpha f^\beta(x)dx$$

hold for  $\alpha$  and  $\beta$ ?

This type of integral inequality is a complement, variant and continuation of Qi's inequality [2]. Before giving an affirmative answer to Problem 1.1 and its reverse, we establish the following essential lemma:

**Lemma 1.2.** Let  $f(x)$  be nonnegative function, continuous on  $[a, b]$  and differentiable on  $(a, b)$ .

If  $\int_x^b f(t)dt \leq \int_x^b (t - a)dt$ ,  $\forall x \in [a, b]$ , then

$$(1.3) \quad f(x) \leq x - a.$$

If  $\int_x^b f(t)dt \geq \int_x^b (t - a)dt$ ,  $\forall x \in [a, b]$ , then

$$(1.4) \quad f(x) \geq x - a.$$



*Proof.* In order to prove (1.3), set

$$G(x) = \int_x^b [f(t) - (t - a)]dt \leq 0, \quad \forall x \in [a, b],$$

we have

$$G'(x) = x - a - f(x), \quad \forall x \in [a, b].$$

We shall give an indirect proof, we suppose  $f(x) \geq x - a$ , then  $G'(x) \leq 0$ ,  $G(x)$  decreases, and  $G(x) \geq 0$ , because of  $G(b) = 0$ . This contradiction establishes (1.3).

The proof of (1.4) is the same as the proof of (1.3).  $\square$

Now, our results can be stated as follows:

---

**Note on an Open Problem**

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

---

Title Page

Contents



Page 4 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

Title Page

Contents



Page 5 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

## 2. Main Results

**Theorem 2.1.** Let  $f(x)$  be a nonnegative function, continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and let  $\alpha$  and  $\beta$  be positive numbers.

If  $\int_x^b f(t)dt \leq \int_x^b (t-a)dt$ ,  $\forall x \in [a, b]$ , then

$$(2.1) \quad \int_a^b f^{\alpha+\beta}(x)dx \leq \int_a^b (x-a)^\alpha f^\beta(x)dx.$$

If  $\int_x^b f(t)dt \geq \int_x^b (t-a)dt$ ,  $\forall x \in [a, b]$ , then

$$(2.2) \quad \int_a^b f^{\alpha+\beta}(x)dx \geq \int_a^b (x-a)^\alpha f^\beta(x)dx.$$

*Proof.* Set

$$F(x) = \int_a^x [f^{\alpha+\beta}(t) - (t-a)^\alpha f^\beta(t)dt], \quad \forall x \in [a, b].$$

We can see that

$$F'(x) = f^{\alpha+\beta}(x) - (x-a)^\alpha f^\beta(x),$$

so that

$$F'(x) = [f^\alpha(x) - (x-a)^\alpha] f^\beta(x).$$

If

$$\int_x^b f(t)dt \leq \int_x^b (t-a)dt, \quad \forall x \in [a, b],$$

and from (1.3) of Lemma 1.2, we have,  $f(x) \leq (x-a)$ , so that  $f^\alpha(x) \leq (x-a)^\alpha$ . Thus  $F'(x) \leq 0$ , and  $F(x)$  is decreasing on  $[a, b]$ . Since  $F(a) = 0$ , we have  $F(x) \leq 0$ ,  $\forall x \in [a, b]$ , which gives the inequality (2.1).



Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

Title Page

Contents



Page 6 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-575b

© 2007 Victoria University. All rights reserved.

When

$$\int_x^b f(t)dt \geq \int_x^b (t-a)dt, \quad \forall x \in [a, b],$$

we have from (1.4)  $f^\alpha(x) \geq (x-a)^\alpha$ , and the rest of the proof is the same as that of (2.1).  $\square$

*Remark 1.* If  $f(x) = 0$  or  $f(x) = x - a$ , the equality in (2.1) holds.

Now we establish new integral inequalities similar to (2.1) and (2.2) involving  $n$  functions:  $f_i(x)$ ,  $i = 1, \dots, n$ .

**Theorem 2.2.** Let  $f_i(x)$ ,  $i = 1, 2, \dots, n$  be nonnegative functions, continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and let  $\alpha_i, \beta_i$ ,  $i = 1, \dots, n$  be positive numbers.

If

$$\int_x^b f_i(t)dt \leq \int_x^b (t-a)dt, \quad \forall x \in [a, b],$$

then

$$(2.3) \quad \int_a^b \prod_{i=1}^n f_i^{\alpha_i + \beta_i}(x) dx \leq \int_a^b (x-a)^{\sum_{i=1}^n \alpha_i} \prod_{i=1}^n f_i^{\beta_i}(x) dx.$$

If

$$\int_x^b f_i(t)dt \geq \int_x^b (t-a)dt, \quad \forall x \in [a, b],$$

then

$$(2.4) \quad \int_a^b \prod_{i=1}^n f_i^{\alpha_i + \beta_i}(x) dx \geq \int_a^b (x-a)^{\sum_{i=1}^n \alpha_i} \prod_{i=1}^n f_i^{\beta_i}(x) dx.$$



Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

Title Page

Contents



Page 7 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

*Proof.* As in proof of (2.1) and (2.2), we let

$$F(x) = \int_a^x \left[ \prod_{i=1}^n f_i^{\alpha_i + \beta_i}(t) - (t-a)^{\sum_{i=1}^n \alpha_i} \prod_{i=1}^n f_i^{\beta_i}(t) \right] dt,$$

thus

$$F'(x) = \left[ \prod_{i=1}^n f_i^{\alpha_i}(x) - (x-a)^{\sum_{i=1}^n \alpha_i} \prod_{i=1}^n f_i^{\beta_i}(x) \right],$$

when

$$\int_x^b f_i(t) dt \leq \int_x^b (t-a) dt \quad \left( \text{resp. } \int_x^b f_i(t) dt \geq \int_x^b (t-a) dt \right), \quad \forall x \in [a, b],$$

using Lemma 1.2, we have

$$f_i(x) \leq x-a \quad \left( \text{resp. } f_i(x) \geq x-a \right), \quad i = 1, 2, \dots, n,$$

and


$$f_i^{\alpha_i}(x) \leq (x-a)^{\alpha_i} \quad \left( \text{resp. } f_i^{\alpha_i}(x) \geq (x-a)^{\alpha_i} \right), \quad i = 1, 2, \dots, n,$$

thus

$$\prod_{i=1}^n f_i^{\alpha_i}(x) \leq (x-a)^{\sum_{i=1}^n \alpha_i} \quad \left( \text{resp. } \prod_{i=1}^n f_i^{\alpha_i}(x) \geq (x-a)^{\sum_{i=1}^n \alpha_i} \right).$$

The rest of the proof is the same as in Theorem 2.1, and we omit (2.3) and (2.4).  $\square$

In order to illustrate a possible practical use of this result, we shall give in the following a simple example in which we can apply Theorem 2.1.



Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

Title Page

Contents



Page 8 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

Example 2.1. For  $\alpha = \beta = 1$  :

i) Let  $f(t) = \cos t + t$  on  $[0, \frac{\pi}{2}]$ , we see that all the conditions of Theorem 2.1 are fulfilled. Indeed,

$$\begin{aligned} \int_x^{\frac{\pi}{2}} (\cos t + t) dt &= 1 - \sin x + \frac{1}{2} \left( \frac{\pi^2}{4} - x^2 \right) \\ &\geq \int_x^{\frac{\pi}{2}} t dt = \frac{1}{2} \left( \frac{\pi^2}{4} - x^2 \right), \quad \forall x \in \left[ 0, \frac{\pi}{2} \right], \end{aligned}$$

and straightforward computation yields

$$\begin{aligned} \int_0^{\frac{\pi}{2}} (\cos x + x)^2 dx &= \frac{5\pi}{4} + \frac{\pi^3}{24} - 2 \\ &> \int_0^{\frac{\pi}{2}} x (\cos x + x) dx \\ &= \frac{\pi}{2} + \frac{\pi^3}{24} - 1. \end{aligned}$$

That is,

$$\int_0^{\frac{\pi}{2}} f^2(x) dx > \int_0^{\frac{\pi}{2}} x f(x) dx.$$

ii) Let  $f(t) = t - \frac{\pi}{2} + \cos t$  on  $[\frac{\pi}{2}, \pi]$ , all the conditions of Theorem 2.1 be satisfied. We see that

$$\int_x^{\pi} \left( t - \frac{\pi}{2} + \cos t \right) dt \leq \int_x^{\pi} \left( t - \frac{\pi}{2} \right) dt, \quad \forall x \in \left[ \frac{\pi}{2}, \pi \right],$$





which is equivalent to

$$\int_x^\pi \cos t dt = -\sin x \leq 0, \quad x \in \left[\frac{\pi}{2}, \pi\right],$$

and direct computation yields

$$\begin{aligned} \int_{\frac{\pi}{2}}^\pi \left(x - \frac{\pi}{2} + \cos x\right)^2 dx &= \frac{\pi}{4} + \frac{\pi^3}{24} - 2 \\ &< \int_{\frac{\pi}{2}}^\pi \left(x - \frac{\pi}{2}\right) f(x) dx \\ &= \frac{\pi^3}{24} - 1. \end{aligned}$$

That is

$$\int_{\frac{\pi}{2}}^\pi f^2(x) dx < \int_{\frac{\pi}{2}}^\pi \left(x - \frac{\pi}{2}\right) f(x) dx.$$

---

**Note on an Open Problem**

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

---

Title Page

Contents



Page 9 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

## References

- [1] QUỐC ANH NGÔ, DU DUC THANG, TRAN TAT DAT, AND DANG ANH TUAN, Notes on an integral inequality, *J. Ineq. Pure and Appl. Math.*, 7(4) (2006), Art. 120. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=737>].
- [2] FENG QI, Several integral inequalities, *J. Ineq. Pure and Appl. Math.*, 1(2) (2000), Art. 19. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=113>].



---

Note on an Open Problem

Lazhar Bougoffa

vol. 8, iss. 2, art. 58, 2007

---

Title Page

Contents



Page 10 of 10

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756