



# Another Proof of Nathanson's Theorems

Quan-Hui Yang  
School of Mathematical Sciences  
Nanjing Normal University  
Nanjing 210046  
P. R. CHINA  
[yangquanhui01@163.com](mailto:yangquanhui01@163.com)

## Abstract

In this paper, without using generating functions, we give new combinatorial proofs of several theorems by Nathanson on the representation functions, and we also obtain generalizations of these theorems.

## 1 Introduction

Let  $\mathcal{A}$  be a set of nonnegative integers. Let  $r_h^{\mathcal{A}}(n)$  denote the number of representations of  $n$  as a sum of  $h$  elements of  $\mathcal{A}$  and  $r^{\mathcal{A}}(n)$  denote the number of representations of  $n$  as a sum of an arbitrary number of elements of  $\mathcal{A}$ , where representations differing only in the arrangement of their summands are counted separately. We notice that if  $0 \notin \mathcal{A}$ , then  $r^{\mathcal{A}}(n) = \sum_{h=1}^{\infty} r_h^{\mathcal{A}}(n)$  is finite for all  $n$ . Representation functions have been extensively studied by many authors [1, 2, 3, 5, 6] and are still a fruitful area of research in additive number theory. Using generating functions, Nathanson [4] proved the following results.

**Theorem 1.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be sets of nonnegative integers, and let  $r_h^{\mathcal{A}}(n)$  and  $r_h^{\mathcal{B}}(n)$  denote the number of representations of  $n$  as a sum of  $h$  elements of  $\mathcal{A}$  and  $\mathcal{B}$ , respectively. If  $r_h^{\mathcal{A}}(n) = r_h^{\mathcal{B}}(n)$  for all  $n \geq 0$ , then  $\mathcal{A} = \mathcal{B}$ .*

**Theorem 2.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be sets of positive integers, and let  $r^{\mathcal{A}}(n)$  and  $r^{\mathcal{B}}(n)$  denote the number of representations of  $n$  as a sum of an arbitrary number of elements of  $\mathcal{A}$  and  $\mathcal{B}$ , respectively. If  $r^{\mathcal{A}}(n) = r^{\mathcal{B}}(n)$  for all  $n \geq 1$ , then  $\mathcal{A} = \mathcal{B}$ .*

**Theorem 3.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be sets of positive integers, and let  $p^{\mathcal{A}}(n)$  and  $p^{\mathcal{B}}(n)$  denote the number of representations of  $n$  as a sum of an arbitrary number of elements of  $\mathcal{A}$  and  $\mathcal{B}$ , respectively, where representations differing only in the arrangement of their summands are not counted separately. If  $p^{\mathcal{A}}(n) = p^{\mathcal{B}}(n)$  for all  $n \geq 1$ , then  $\mathcal{A} = \mathcal{B}$ .*

In this paper, we give new proofs of theorems above. Indeed, we shall prove slightly more. We first introduce some notation. If  $\mathcal{A}$  is a strictly increasing sequence of integers, then  $a_n$  denotes the  $n$ th element of  $\mathcal{A}$ . Let  $\mathcal{A}$  be a set of nonnegative integers and  $\mathcal{H}$  be a set of positive integers. If  $|\mathcal{H}|$  is finite, then  $r_{\mathcal{H}}^{\mathcal{A}}(n)$  denotes the number of representations of  $n$  as a sum of  $h_1$  or  $h_2$  or  $\dots$  elements of  $\mathcal{A}$ ; if  $|\mathcal{H}|$  is infinite, then  $r_{\mathcal{H}}^{\mathcal{A}}(n)$  denotes the number of representations of  $n$  as a sum of  $h_1$  or  $h_2$  or  $\dots$  elements of  $\mathcal{A} \setminus \{0\}$ .

**Theorem 4.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be nonempty sets of nonnegative integers. Let  $\mathcal{H}$  be a nonempty set of positive integers, and let  $S = \{\min\{a_i, b_i\} : i = 1, 2, \dots\}$ . Write  $t = (\min(\mathcal{H}) - 1) \min\{a_1, b_1\}$ . If  $r_{\mathcal{H}}^{\mathcal{A}}(n) = r_{\mathcal{H}}^{\mathcal{B}}(n)$  for all  $n \in t + S$ , then  $\mathcal{A} = \mathcal{B}$ .*

Let  $\mathcal{H} = \{h\}$ . Since  $t + S \subseteq \{0, 1, 2, \dots\}$ , Theorem 4 is a generalization of Theorem 1. Let  $\mathcal{H} = \{1, 2, 3, \dots\}$ . If  $\mathcal{A}$  and  $\mathcal{B}$  are sets of positive integers, then  $t + S \subseteq \{1, 2, \dots\}$ . Hence, Theorem 4 is also a generalization of Theorem 2.

**Theorem 5.** *Let  $\mathcal{A}$ ,  $\mathcal{B}$  and  $\mathcal{H}$  be nonempty sets of positive integers. Let  $S = \{\min\{a_i, b_i\} : i = 1, 2, \dots\}$ , and let  $p_{\mathcal{H}}^{\mathcal{A}}(n)$  denote the number of representations of  $n$  as a sum of  $h_1$  or  $h_2$  or  $\dots$  elements of  $\mathcal{A}$ , where representations differing only in the arrangement of their summands are not counted separately. Write  $t = (\min(\mathcal{H}) - 1) \min\{a_1, b_1\}$ . If  $p_{\mathcal{H}}^{\mathcal{A}}(n) = p_{\mathcal{H}}^{\mathcal{B}}(n)$  for all  $n \in t + S$ , then  $\mathcal{A} = \mathcal{B}$ .*

Let  $\mathcal{H} = \{1, 2, 3, \dots\}$ . Since  $t + S \subseteq \{1, 2, \dots\}$ , Theorem 5 is a generalization of Theorem 3.

Let  $A$ ,  $B$ , and  $T$  be finite sets of integers. If each residue class modulo  $m$  contains exactly the same number of elements of  $A$  as elements of  $B$ , then we write  $A \equiv B \pmod{m}$ . If the number of solutions of the congruence  $a + t \equiv n \pmod{m}$  with  $a \in A$ ,  $t \in T$ , equals the number of solutions of the congruence  $b + t \equiv n \pmod{m}$  with  $b \in B$ ,  $t \in T$ , for each residue class  $n$  modulo  $m$ , then we write  $A + T \equiv B + T \pmod{m}$ . Nathanson [4] also proved the following theorem.

**Theorem 6.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be distinct nonempty sets of nonnegative integers such that  $r_2^{\mathcal{A}}(n) = r_2^{\mathcal{B}}(n)$  for all sufficiently large  $n$ . Then there exist finite sets  $A, B$ , and  $T$  with  $A \cup B \subset \{0, 1, \dots, N\}$  and  $T \subset \{0, 1, \dots, m - 1\}$  such that  $A + T \equiv B + T \pmod{m}$ , and  $\mathcal{A} = A \cup \mathcal{C}$  and  $\mathcal{B} = B \cup \mathcal{C}$ , where  $\mathcal{C} = \{c > N \mid c \equiv t \pmod{m} \text{ for some } t \in T\}$ .*

In this paper, we prove theorems above without using generating functions. We notice that for a prime number  $p$ , if  $\mathcal{A}$  and  $\mathcal{B}$  are sets of nonnegative integers such that  $r_p^{\mathcal{A}}(n) = r_p^{\mathcal{B}}(n)$  for all sufficiently large  $n$ , then  $\mathcal{A}$  and  $\mathcal{B}$  eventually coincide. Now, I pose the following problem.

**Problem 7.** *Let  $p \geq 3$  be a prime number and  $\mathcal{A}$  be a set of nonnegative integers. Does there exist a set of nonnegative integers  $\mathcal{B}$  with  $\mathcal{B} \neq \mathcal{A}$  such that  $r_p^{\mathcal{A}}(n) = r_p^{\mathcal{B}}(n)$  for all sufficiently large  $n$ ?*

## 2 Proof of Theorems 4 and 5

Suppose that  $\mathcal{A} \neq \mathcal{B}$ . Let  $h = \min(\mathcal{H})$  and  $j_0$  be the smallest index such that  $a_{j_0} \neq b_{j_0}$ . Without loss of generality, we can assume that  $a_{j_0} < b_{j_0}$ . Let  $C = \{a_j : j < j_0\}$ . Since  $a_j = b_j$  for all  $j < j_0$  and  $t = (h-1)a_1$ , we have  $(h-1)a_1 + a_{j_0} \in t + S$  and

$$\begin{aligned} r_{\mathcal{H}}^{\mathcal{A}}((h-1)a_1 + a_{j_0}) &= r_{\mathcal{H}}^C((h-1)a_1 + a_{j_0}) + 1 \\ &= r_{\mathcal{H}}^{\mathcal{B}}((h-1)a_1 + a_{j_0}) + 1, \end{aligned}$$

which is a contradiction. Hence, we have  $\mathcal{A} = \mathcal{B}$ . This completes the proof of Theorem 4.

The proof of Theorem 5 is very similar to the proof of Theorem 4, and we omit it here.

## 3 Proof of Theorem 6

Clearly,  $r_2^{\mathcal{A}}(2n)$  is odd if and only if  $n \in \mathcal{A}$ . Similarly,  $n \in \mathcal{B}$  if and only if  $r_2^{\mathcal{B}}(2n)$  is odd. If  $r_2^{\mathcal{A}}(n) = r_2^{\mathcal{B}}(n)$  for all  $n > N_0$ , then for all  $n > N_0$  we have  $n \in \mathcal{A}$  if and only if  $n \in \mathcal{B}$ . Let

$$\mathcal{D} = \mathcal{A} \cap [N_0 + 1, \infty) = \mathcal{B} \cap [N_0 + 1, \infty)$$

and write

$$\eta(n) = \begin{cases} 1, & \text{if } n \in \mathcal{D}; \\ 0, & \text{otherwise.} \end{cases}$$

Then for  $n > 2N_0$ , we have

$$\begin{aligned} r_2^{\mathcal{A}}(n) &= 2\#\{(a, d) : a \in \mathcal{A} \setminus \mathcal{D}, d \in \mathcal{D}, a + d = n\} \\ &\quad + \#\{(d', d'') : d', d'' \in \mathcal{D}, d' + d'' = n\} \\ &= 2 \sum_{a \in \mathcal{A} \setminus \mathcal{D}} \eta(n - a) + \#\{(d', d'') : d', d'' \in \mathcal{D}, d' + d'' = n\}. \end{aligned} \tag{1}$$

Similarly, we have

$$r_2^{\mathcal{B}}(n) = 2 \sum_{b \in \mathcal{B} \setminus \mathcal{D}} \eta(n - b) + \#\{(d', d'') : d', d'' \in \mathcal{D}, d' + d'' = n\}. \tag{2}$$

Since  $r_2^{\mathcal{A}}(n) = r_2^{\mathcal{B}}(n)$  for all  $n > N_0$ , by (1) and (2), we have

$$\sum_{a \in \mathcal{A} \setminus \mathcal{D}} \eta(n - a) = \sum_{b \in \mathcal{B} \setminus \mathcal{D}} \eta(n - b) \tag{3}$$

for all  $n > 2N_0$ . Let  $i_0$  be the smallest index such that  $a_{i_0} \neq b_{i_0}$ . Without loss of generality, we may assume that  $a_{i_0} < b_{i_0}$ .

Let

$$t = n - a_{i_0}$$

and

$$\mathcal{D}' = \{a : a < a_{i_0}, a \in \mathcal{A}\}.$$

Then by (3), we have

$$\eta(t) = \sum_{b \in \mathcal{B} \setminus (\mathcal{D} \cup \mathcal{D}')} \eta(t + a_{i_0} - b) - \sum_{a \in \mathcal{A} \setminus (\mathcal{D} \cup \mathcal{D}' \cup a_{i_0})} \eta(t + a_{i_0} - a).$$

Since  $\eta(t)$  defined by a linear recurrence on a finite set  $\{0, 1\}$ , we have that it must be eventually periodic. Hence, for some  $N > N_0$ ,  $\mathcal{D} \cap [N + 1, \infty)$  is periodic. We denote such a period by  $m$ . Let  $T = \{t : t \equiv d \pmod{m} \text{ for some } d \in \mathcal{D} \cap [N + 1, \infty) \text{ and } 0 \leq t < m\}$ . Then we have  $n \in \mathcal{A} \cap \mathcal{B} \cap [N + 1, \infty)$  if and only if  $n \equiv t \pmod{m}$  for some  $t \in T$ .

The remainder of the proof is the same as that of the proof by Nathanson. To make this paper self-contained, we formulate it here. Let

$$A = \{a \leq N : a \in \mathcal{A}\}, \quad B = \{b \leq N : b \in \mathcal{B}\},$$

and

$$\mathcal{C} = \{c > N : c \in \mathcal{A} \cap \mathcal{B}\} = \{c > N : c \equiv t \pmod{m} \text{ for some } t \in T\}.$$

Then  $\mathcal{A} = A \cup \mathcal{C}$  and  $\mathcal{B} = B \cup \mathcal{C}$ . Next we prove that  $A + T \equiv B + T \pmod{m}$ .

For  $n > 2N$ , we have

$$\begin{aligned} r_2^{\mathcal{A}}(n) &= r_2^{\mathcal{C}}(n) + 2\#\{(a, c) : a \in A, c \in \mathcal{C}, a + c = n\} \\ &= r_2^{\mathcal{C}}(n) + 2\#\{(a, t) : a \in A, t \in T, a + t \equiv n \pmod{m}\}. \end{aligned} \tag{4}$$

Similarly,

$$r_2^{\mathcal{B}}(n) = r_2^{\mathcal{C}}(n) + 2\#\{(b, t) : b \in B, t \in T, b + t \equiv n \pmod{m}\}. \tag{5}$$

Since  $r_2^{\mathcal{A}}(n) = r_2^{\mathcal{B}}(n)$  for  $n > 2N$ , by (4) and (5), we have that  $A + T \equiv B + T \pmod{m}$ . This completes the proof of Theorem 6.

## 4 Acknowledgement

I sincerely thank my supervisor Professor Yong-Gao Chen for his valuable suggestions and useful discussions. I am also grateful to the referee for his/her valuable comments.

## References

- [1] Y. G. Chen, A. Sárkőzy, V. T. Sós, and M. Tang, On the monotonicity properties of additive representation functions, *Bull. Aust. Math. Soc.* **72** (2005), 129–138.
- [2] P. Erdős, A. Sárkőzy, and V. T. Sós, Problems and results on additive properties of general sequences, V, *Monatsh. Math.* **102** (1986), 183–197.

- [3] P. Erdős, A. Sárközy, and V. T. Sós, On additive properties of general sequences, *Discrete Math.* **136** (1994), 75–99.
- [4] M. B. Nathanson, Representation functions of sequences in additive number theory, *Proc. Amer. Math. Soc.* **72** (1978), 16–20.
- [5] M. B. Nathanson, Every function is the representation function of an additive basis for the integers, *Port. Math.* **62** (2005), 55–72.
- [6] M. B. Nathanson, Representation functions of bases for binary linear forms, *Funct. Approx. Comment. Math.* **37** (2007), 341–350.

---

2010 *Mathematics Subject Classification*: Primary 11B34; Secondary 11D85.

*Keywords*: Nathanson’s theorems; representation functions; generating functions.

---

Received March 13 2011; revised version received August 13 2011. Published in *Journal of Integer Sequences*, September 25 2011.

---

Return to [Journal of Integer Sequences home page](#).