



Enumerating Minimal Length Lattice Paths

Jackson Evoniuk, Steven Klee, and Van Magnan

Department of Mathematics

Seattle University

901 12th Avenue

Seattle, WA 98122

USA

evoniukj@seattleu.edu

klee@seattleu.edu

magnanv@seattleu.edu

Abstract

Given a finite set of integer vectors, S , we consider the set of all lattice walks comprised as ordered sequences of steps whose directions come from S . We further restrict our attention to walks of minimal length, meaning they cannot be shortened through some linear combination of allowable steps from S . We consider the problem of counting the number of such minimal walks terminating at a fixed point (a, b) for various choices of the set S .

1 Introduction

Let S be a finite set of vectors in \mathbb{Z}^2 . An S -walk is an ordered sequence $\mathbf{s} = s_1, s_2, \dots, s_k$ of steps with $s_i \in S$ for all i . We may visualize an S -walk as a path beginning at the origin and terminating at the point whose coordinates are given by $s_1 + s_2 + \dots + s_k$. We say the number of steps in a path is its *length*, and we refer to the elements of S as *allowable steps*.

The problem of enumerating the walks terminating at a fixed point (a, b) with $a, b \in \mathbb{N}$ is classical in combinatorics. For example, when $S = \{(1, 0), (0, 1)\}$, the number of such walks is $\binom{a+b}{a}$. When S is an arbitrary set of allowable vectors, there may be several paths of different lengths that terminate at a fixed point (a, b) . For example, if $S = \{(1, 0), (0, 1), (1, 1)\}$, then

the path $\mathbf{s} = (1, 0), (1, 0), (0, 1), (1, 1), (0, 1), (1, 0), (1, 1)$ is a path of length 7 terminating at the point $(5, 4)$, and $\mathbf{s}' = (1, 1), (1, 0), (1, 1), (1, 1), (1, 1)$ is a path of length 5 terminating at the point $(5, 4)$. These walks are illustrated in Figure 1.




Figure 1: A non-minimal S -walk (left) and a minimal S -walk (right) terminating at the point $(a, b) = (5, 4)$ when $S = \{(1, 0), (0, 1), (1, 1)\}$.

Our goal in this paper is to enumerate the *minimal S -walks* to a point (a, b) — among all S -walks terminating at (a, b) , we consider only those of minimal length. We will write $d(a, b; S)$ to denote the S -distance of the point (a, b) from the origin, which counts the number of steps in a minimal S -walk to (a, b) . In the previous example, any S -walk terminating at $(5, 4)$ must utilize at least 5 steps among $\{(1, 0), (1, 1)\}$, so $d(5, 4; S) \geq 5$. Therefore, \mathbf{s}' is minimal because it is an S -walk of length 5. In contrast, \mathbf{s} is not minimal. In general, we write $\mathcal{W}(a, b; S)$ to denote the set of minimal S -walks terminating at the point (a, b) . Our goal in this paper is to examine this problem in several different contexts, exhibiting either explicit closed formulas or generating functions to determine $|\mathcal{W}(a, b; S)|$.

The rest of the paper is structured as follows. In Section 2, we warm up with the case that $S = \{(1, 0), (0, 1), (1, 1)\}$. In Section 3, we consider the sets

$$Q_n := \{(1, 0), (0, 1)\} \cup \{(i, n - i) : 0 \leq i \leq n\},$$

consisting of the standard basis vectors along with all nonnegative integer vectors whose coordinate sum equals n for $n \geq 2$. We include the standard basis vectors to ensure that every point (a, b) with $a, b \in \mathbb{N}$ can be reached by a Q_n -walk. Next, we turn our attention to the case that $S = \{(1, 0), (0, 1), (u, v)\}$ for arbitrary $u, v \in \mathbb{N}$ in Section 4. In Section 5, we explore the case that $S = \{(1, 0), (0, 1), (2, 1), (1, 2)\}$. We conclude in Section 6 with some open problems.

2 Minimal walks for $S = \{(1, 0), (0, 1), (1, 1)\}$

We begin by examining the set of allowable steps $S = \{(1, 0), (0, 1), (1, 1)\}$. In this and subsequent sections, we have implemented a simple greedy search in Sage [3] to determine

the number of minimal S -walks terminating at a given point (a, b) for small values of a and b . This data is depicted visually in Figure 2. For example, the circled 20 indicates that there are 20 minimal S -paths terminating at the point $(6, 3)$.

10		1	10	45	120	210	252	210	120	45	10	1
9		1	9	36	84	126	126	84	36	9	1	10
8		1	8	28	56	70	56	28	8	1	9	45
7		1	7	21	35	35	21	7	1	8	36	120
6		1	6	15	20	15	6	1	7	28	84	210
5		1	5	10	10	5	1	6	21	56	126	252
4		1	4	6	4	1	5	15	35	70	126	210
3		1	3	3	1	4	10	20	35	56	84	120
2		1	2	1	3	6	10	15	21	28	36	45
1		1	1	2	3	4	5	6	7	8	9	10
0		1	1	1	1	1	1	1	1	1	1	1
$\frac{b}{a}$		0	1	2	3	4	5	6	7	8	9	10

Figure 2: The number of minimal S -walks terminating at each point (a, b) for $0 \leq a, b \leq 10$ and $S = \{(1, 0), (0, 1), (1, 1)\}$.

In that image, we notice that there appear to be two copies of Pascal’s triangle (OEIS sequence [A007318](#)) glued together along the line $y = x$. Our first result proves that this pattern continues.

Theorem 1. *Let $S = \{(1, 0), (0, 1), (1, 1)\}$ and let (a, b) be a point with $a, b \in \mathbb{N}$. Then $|\mathcal{W}(a, b; S)| = \binom{\max(a, b)}{\min(a, b)}$.*

Proof. By symmetry, we may assume without loss of generality that $a \geq b$. Note that $d(a, b; S) \geq a$ since an allowable step in S increases the x -coordinate by at most 1. Conversely, $d(a, b; S) \leq a$ since (a, b) can be reached by taking b steps in the $(1, 1)$ -direction, followed by $a - b$ steps in the $(1, 0)$ -direction. Thus $d(a, b; S) = a$.

In particular, it follows that every step in a minimal S -walk terminating at (a, b) must increase the x -coordinate, and hence such a walk does not use any $(0, 1)$ steps. Since a $(1, 1)$ step is the only remaining step that can increase the y -coordinate, a minimal S -walk is comprised of a total steps, b of which are $(1, 1)$ steps and the remaining $a - b$ of which are $(1, 0)$ steps. There are $\binom{a}{b}$ such paths. □

3 Minimal walks for steps of fixed length

Our next goal is to consider the set of allowable steps

$$Q_n := \{(1, 0), (0, 1)\} \cup \{(i, n - i) : 0 \leq i \leq n\},$$

for $n \geq 2$. For example, $Q_3 = \{(1, 0), (0, 1), (0, 3), (1, 2), (2, 1), (3, 0)\}$, and Figure 3 shows data for the number of minimal Q_3 -walks. This array did not previously appear in OEIS, but we have added it as sequence [A292435](#).

10	4	50	10	150	1215	101	1416	11046	546	7882	63056
9	1	16	130	20	255	1830	135	1740	12600	580	7882
8	6	3	36	250	31	355	2325	155	1860	12600	546
7	3	24	6	64	380	40	420	2520	155	1740	11046
6	1	9	48	10	88	460	44	420	2325	135	1416
5	3	2	15	72	12	96	460	40	355	1830	101
4	2	9	3	21	84	12	88	380	31	255	1215
3	1	4	12	4	21	72	10	64	250	20	150
2	1	1	4	12	3	15	48	6	36	130	10
1	1	2	1	4	9	2	9	24	3	16	50
0	1	1	1	1	2	3	1	3	6	1	4
$\begin{array}{l} b \\ \hline a \end{array}$	0	1	2	3	4	5	6	7	8	9	10

Figure 3: The number of minimal Q_3 -walks terminating at each point (a, b) for $0 \leq a, b \leq 10$.

In Figure 3, we observe an interesting phenomenon. If we fix a value $m = 3q + r$ with $0 \leq r < 3$, and consider all points (a, b) with $a + b = m$, then the number of minimal Q_3 -walks terminating at (a, b) is a multiple of $\binom{q+r}{r}$. For example, when $m = 7 = 3 \cdot 2 + 1$, the entries along the diagonal $(3, 9, 15, 21, 21, 15, 9, 3)$ are all divisible by $\binom{2+1}{1} = 3$.

In order to justify this phenomenon in general, we introduce an additional piece of terminology. For Q_n , we will call the steps $(1, 0)$ and $(0, 1)$ *short* steps and the remaining steps of the form $(i, n - i)$ *long* steps.

Lemma 2. *Let $a, b \in \mathbb{N}$ and write $a + b = n \cdot q + r$ with $0 \leq r < n$. Then any minimal Q_n -walk terminating at (a, b) uses exactly q long steps and r short steps. Consequently, $d(a, b; Q_n) = q + r$.*

Proof. Let \mathbf{s} be a (not necessarily minimal) Q_n -walk terminating at (a, b) , and suppose \mathbf{s} uses q' long steps and r' short steps. If $r' \geq n$, then we can replace n of those short steps with one long step, which would result in shorter path. Here, it is worth noting that we do not require these n short steps to appear consecutively in \mathbf{s} . We can simply remove them from \mathbf{s} and append their vector sum, which is a long step, to the end of the resulting walk. This creates a new walk terminating at (a, b) with fewer steps.

Therefore, if \mathbf{s} is a minimal Q_n -walk, then $r' < n$. By taking the vector sum of every step in \mathbf{s} , we see that $a + b = n \cdot q' + r'$. Since the quotient and remainder in the division algorithm are unique, we must have $q' = q$ and $r' = r$, meaning \mathbf{s} uses q long steps and r short steps. \square

Let $(a, b) \in \mathbb{N}^2$ and write $a + b = q \cdot n + r$ with $0 \leq r < n$. We can now use Lemma 2 to see why $|\mathcal{W}(a, b; Q_n)|$ is divisible by $\binom{q+r}{r}$. We can partition $\mathcal{W}(a, b; Q_n)$ into equivalence classes by declaring $\mathbf{s} \sim \mathbf{s}'$ if (1) \mathbf{s} and \mathbf{s}' use the same number of each step from Q_n and (2) the relative order of the long steps and the relative order of the short steps in \mathbf{s} is the same as that in \mathbf{s}' . For example, in Q_3 , the paths $(3, 0), (2, 1), (1, 0), (0, 1), (1, 2), (2, 1)$ and $(1, 0), (3, 0), (0, 1), (2, 1), (1, 2), (2, 1)$ are equivalent. The paths equivalent to \mathbf{s} are determined by choosing r positions out of $q + r$ total steps in which we will place the (ordered list of) short steps.

At this point, however, the combinatorics of enumerating minimal Q_n -walks to a fixed point (a, b) is somewhat complicated because the linear algebra problem of determining all ways to write (a, b) as a sum of q long steps and r short steps is difficult to do in generality. Instead, it is easier to exhibit a generating function that will enumerate all such walks.

Theorem 3. *For all $n \geq 2$, the number of minimal Q_n -walks can be computed by the generating function*

$$\sum_{(a,b) \in \mathbb{N}^2} |\mathcal{W}(a, b; Q_n)| x^a y^b = \sum_{q=0}^{\infty} \sum_{r=0}^{n-1} \binom{q+r}{r} \left(\sum_{i=0}^n x^i y^{n-i} \right)^q (x+y)^r. \quad (1)$$

Proof. For fixed q and r , let $\sigma = n \cdot q + r$. Expanding $(\sum_{i=0}^n x^i y^{n-i})^q$ encodes all possible ways to make an ordered list of q long steps, and expanding $(x+y)^r$ encodes all possible ways to make an ordered list of r short steps. Hence, multiplying these quantities together encodes all possible ways to make an ordered list of q long steps followed by r short steps. Multiplying by $\binom{q+r}{r}$ accounts for all possible ways to shuffle these steps together. Therefore, the summand of the generating function for fixed q and r covers all equivalence classes (as described above) of minimal Q_n -walks terminating along the diagonal where $a + b = \sigma$. \square

4 Minimal walks for $S = \{(1, 0), (0, 1), (u, v)\}$

In this section, we consider an asymmetric set of allowable steps in $S = \{(1, 0), (0, 1), (u, v)\}$ for arbitrary integers $u, v \geq 1$.

Given a point $(a, b) \in \mathbb{N}^2$, let $m = m(a, b) = \min(\lfloor \frac{a}{u} \rfloor, \lfloor \frac{b}{v} \rfloor)$. Concretely, m is the largest integer such that $m \cdot u \leq a$ and $m \cdot v \leq b$; or in other words, m is the largest number of steps one can take in the (u, v) -direction without exceeding the x - or y -coordinate of (a, b) .

Theorem 4. *Let $S = \{(1, 0), (0, 1), (u, v)\}$ with $u, v \geq 1$, and let $(a, b) \in \mathbb{N}^2$. A minimal S -walk to the point (a, b) uses exactly m steps in the (u, v) -direction. Consequently,*

$$d(a, b; S) = m + a - m \cdot u + b - m \cdot v,$$

and

$$|\mathcal{W}(a, b; S)| = \binom{m + a - m \cdot u + b - m \cdot v}{m, a - m \cdot u, b - m \cdot v}.$$

Proof. First, we will prove that a minimal S -walk uses exactly m steps in the (u, v) -direction. As noted above, any S -walk terminating at (a, b) uses at most m steps in the (u, v) -direction. We claim an S -walk using fewer than m steps in the (u, v) -direction is non-minimal. Indeed, consider an S -walk using m' steps in the (u, v) -direction, x steps in the $(1, 0)$ -direction, and y steps in the $(0, 1)$ -direction, and assume $m' < m$.

Since

$$a = m' \cdot u + x \leq (m - 1) \cdot u + x = m \cdot u + x - u \leq a + x - u,$$

it follows that $x \geq u$. Similarly, $y \geq v$. Since $x \geq u$ and $y \geq v$, we can replace u steps in the $(1, 0)$ -direction and v steps in the $(0, 1)$ -direction with one step in the (u, v) -direction, resulting in a shorter path. Thus, a walk using m' steps in the (u, v) -direction is non-minimal, and hence a minimal S -walk uses at least m steps in the (u, v) -direction.

Finally, since a minimal S -walk uses m steps in the (u, v) -direction, then it must use $a - m \cdot u$ steps in the $(1, 0)$ -direction and $b - m \cdot v$ steps in the $(0, 1)$ -direction. This immediately implies the stated formulas for the distance and number of minimal S -walks. \square

Remark 5. In the case that $(u, v) = (1, 1)$, note that $m = \min(a, b)$, that $m + a - mu + b - mv = a + b - m = \max(a, b)$, and that one of $a - mu$ and $b - mv$ is equal to 0. Thus Theorem 4 generalizes the results in Section 2.

5 Minimal walks for $\overline{Q}_3 = \{(1, 0), (0, 1), (2, 1), (1, 2)\}$

Recall that in Section 3, we considered the set of allowable steps Q_3 . By removing the vectors $(3, 0)$ and $(0, 3)$ from this set, we obtain the set $\overline{Q}_3 = \{(1, 0), (0, 1), (2, 1), (1, 2)\}$. The combinatorial data coming from this seemingly simple example did not appear in OEIS [1] and seems interesting in its own right. We have since added this data to OEIS as sequence [A292436](#).

Figure 4 shows the number of minimal walks for \overline{Q}_3 . Here, we have included the lines spanned by the vectors $(2, 1)$ and $(1, 2)$ for reference. We observe that these lines divide the nonnegative quadrant into three regions. In the regions weakly above the line $y = 2x$ and

weakly below the line $2y = x$, we observe that the number of minimal \overline{Q}_3 -walks appears to be a binomial coefficient, while the behavior between these two lines appears to be different. Our goal in this section is to explain several patterns in data collected in Figure 4.

10	1	9	28	35	15	1	36	441	15	245	1960
9	1	8	21	20	5	90	5	90	735	20	245
8	1	7	15	10	1	25	225	10	120	735	15
7	1	6	10	4	50	4	50	300	10	90	441
6	1	5	6	1	16	100	6	50	225	5	36
5	1	4	3	24	3	24	100	4	25	90	1
4	1	3	1	9	36	3	16	50	1	5	15
3	1	2	9	2	9	24	1	4	10	20	35
2	1	1	4	9	1	3	6	10	15	21	28
1	1	2	1	2	3	4	5	6	7	8	9
0	1	1	1	1	1	1	1	1	1	1	1
$\frac{b}{a}$	0	1	2	3	4	5	6	7	8	9	10

Figure 4: The number of minimal \overline{Q}_3 -walks terminating at each point (a, b) for $0 \leq a, b \leq 10$.

We begin by establishing some notation that will be used throughout the proofs in this section. Given a point $(a, b) \in \mathbb{N}^2$, consider a minimal \overline{Q}_3 -walk, \mathbf{s} , terminating at (a, b) . Let x, y, z , and w respectively denote the number of $(1, 0)$ -, $(0, 1)$ -, $(2, 1)$ -, and $(1, 2)$ -steps in \mathbf{s} . Thus

$$a = x + 2z + w \quad \text{and} \quad b = y + z + 2w. \quad (2)$$

As in Section 3, we will refer to the steps $(2, 1)$ and $(1, 2)$ in \overline{Q}_3 as *long* steps and the steps $(1, 0)$ and $(0, 1)$ as *short* steps.

Our first goal is to explain the combinatorial data observed in the outer regions where $a \geq 2b$ or $b \geq 2a$.

Theorem 6. *Let $(a, b) \in \mathbb{N}^2$ with $a \geq 2b$. A minimal \overline{Q}_3 -walk terminating at (a, b) does not use any steps in the $(0, 1)$ -direction or in the $(1, 2)$ -direction. Consequently,*

$$d(a, b; \overline{Q}_3) = a - b$$

and

$$|\mathcal{W}(a, b; \overline{Q}_3)| = \binom{a-b}{b}.$$

Proof. Consider a minimal \overline{Q}_3 -walk, \mathbf{s} , terminating at the point (a, b) . Let x, y, z , and w be as defined above. Our first goal is to show $y = 0$ and $w = 0$.

Since $a \geq 2b$, it follows from Eq. (2) that $x \geq 2y + 3w$. Therefore, if $y \geq 1$, then $x \geq 2$. So if \mathbf{s} uses a $(0, 1)$ -step, then it uses (at least) two $(1, 0)$ -steps. But a $(0, 1)$ -step and two $(1, 0)$ -steps can be replaced with a $(2, 1)$ -step, which would give a shorter path. Thus $y = 0$.

Similarly, if $w \geq 1$, then $x \geq 3$. So if \mathbf{s} uses a $(1, 2)$ -step, it uses at least three $(1, 0)$ -steps. But $(1, 2) + 3(1, 0) = (4, 2) = 2(2, 1)$, so these four steps can be replaced with two $(2, 1)$ -steps, which would give a shorter path. Thus $w = 0$ as well.

Since $y = 0$ and $w = 0$, Eq. (2) reduces to

$$a = x + 2z \quad \text{and} \quad b = z,$$

which is equivalent to $x = a - 2b$ (which is nonnegative since $a \geq 2b$) and $z = b$. Since $x + z$ is the total number of steps in \mathbf{s} , it follows that $d(a, b; \overline{Q}_3) = a - b$. Finally, to determine a path to the point (a, b) , we must use $a - b$ total steps, b of which go in the direction $(2, 1)$ and $a - 2b$ of which go in the direction $(1, 0)$. There are $\binom{a-b}{b}$ ways to determine such a path. This completes the proof. \square

Since \overline{Q}_3 is symmetric, the following corollary immediately handles the case that $b \geq 2a$.

Corollary 7. *Let $(a, b) \in \mathbb{N}^2$ with $b \geq 2a$. A minimal \overline{Q}_3 -walk terminating at (a, b) does not use any steps in the $(1, 0)$ -direction or in the $(2, 1)$ -direction. Consequently, $d(a, b; \overline{Q}_3) = b - a$ and $|\mathcal{W}(a, b; \overline{Q}_3)| = \binom{b-a}{a}$.*

Now we turn our attention to the case that (a, b) lies in the central region of Figure 4.

Lemma 8. *Let $(a, b) \in \mathbb{N}^2$ with $\frac{1}{2}b \leq a \leq 2b$. A minimal \overline{Q}_3 -walk terminating at (a, b) uses at most two short steps.*

Proof. As before, consider a minimal \overline{Q}_3 -walk, \mathbf{s} , terminating at the point (a, b) . Let x, y, z , and w be as defined above. Assume by way of contradiction that \mathbf{s} uses at least three short steps, or equivalently that $x + y \geq 3$.

If $x \geq 2$ and $y \geq 1$ (or if $x \geq 1$ and $y \geq 2$), then \mathbf{s} can be shortened by replacing three short steps with a $(2, 1)$ -step (respectively, with a $(1, 2)$ -step). So we need only consider the case that $x \geq 3$ and $y = 0$. By symmetry, this will cover the case that $x = 0$ and $y \geq 3$.

Since $a \leq 2b$, it follows from Eq. (2) that $x \leq 2y + 3w$. Since $y = 0$ and $x \geq 3$, it follows that $w \geq 1$. Thus \mathbf{s} uses at least three $(1, 0)$ -steps and one $(1, 2)$ -step, which can be shortened by replacing $3(1, 0) + (1, 2) = (4, 2)$ with two $(2, 1)$ -steps. This contradicts our assumption that \mathbf{s} is minimal. \square

Theorem 9. Let $(a, b) \in \mathbb{N}^2$ with $\frac{1}{2}b \leq a \leq 2b$, and write $a + b = 3q + r$ with $0 \leq r \leq 2$. Then

$$d(a, b; \overline{Q}_3) = q + r$$

and

$$|\mathcal{W}(a, b; \overline{Q}_3)| = \binom{q+r}{r} \binom{q+r}{a-q}.$$

Proof. As before, consider a minimal \overline{Q}_3 -walk, \mathbf{s} , terminating at the point (a, b) . Let x, y, z , and w be as defined above. By Eq. (2), $a + b = x + y + 3(z + w)$. By Lemma 8, $x + y < 3$, and hence by uniqueness of the quotient and remainder in the division algorithm, it must be the case that $x + y = r$ and $z + w = q$. But $x + y + z + w$ is the total number of steps used in \mathbf{s} , and hence $d(a, b; \overline{Q}_3) = q + r$. In particular, r counts the number of short steps used in a minimal path.

Now we turn our attention to counting the number of minimal \overline{Q}_3 -walks terminating at (a, b) . Let \mathbf{s} be such a walk. We know \mathbf{s} uses r short steps. Define a new \overline{Q}_3 -walk $\hat{\mathbf{s}}$ as follows: replace each $(1, 0)$ -step in \mathbf{s} with a $(2, 1)$ -step and replace each $(0, 1)$ -step in \mathbf{s} with a $(1, 2)$ -step. Note that $\hat{\mathbf{s}}$ is a \overline{Q}_3 -walk that only uses long steps. Moreover, $\hat{\mathbf{s}}$ terminates at the point $(a + r, b + r)$ as each replacement increases the vector sum by $(1, 1)$.

Now let \hat{w} and \hat{z} respectively denote the number of $(2, 1)$ - and $(1, 2)$ -steps in $\hat{\mathbf{s}}$. Applying Eq. (2) to $\hat{\mathbf{s}}$ gives rise to the system of equations

$$\begin{aligned} 2\hat{z} + \hat{w} &= a + r \\ \hat{z} + 2\hat{w} &= b + r. \end{aligned}$$

The coefficient matrix $\begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$ is invertible, so this system has a unique solution. We can easily verify that

$$\begin{aligned} \hat{z} &= a - q \\ \hat{w} &= b - q \end{aligned}$$

is a solution, which can also be derived by inverting the coefficient matrix and making use of the fact that $a + b = 3q + r$.

Moreover, $a - q$ and $b - q$ are both nonnegative because $\frac{1}{2}b \leq a \leq 2b$. Indeed, $3q + r = a + b \leq a + 2a$, and hence $3(a - q) \geq r \geq 0$. The same logic shows $3(b - q) \geq r \geq 0$.

Therefore, the \overline{Q}_3 -minimal walks terminating at (a, b) can be constructed as follows. Start with a \overline{Q}_3 -walk using $a - q$ steps in the direction $(2, 1)$ and $b - q$ steps in the direction $(1, 2)$. There are $\binom{a-q+b-q}{a-q} = \binom{q+r}{a-q}$ such walks. Next, choose r of those steps to transform into short steps. If the chosen step is a $(2, 1)$ -step, turn it into a $(1, 0)$ step; if it is a $(1, 2)$ -step, turn it into a $(0, 1)$ -step. There are $\binom{q+r}{r}$ ways to make these choices. The resulting path terminates at the point (a, b) and has length $q + r = d(a, b; \overline{Q}_3)$. Thus $|\mathcal{W}(a, b; \overline{Q}_3)| = \binom{q+r}{r} \binom{q+r}{a-q}$, as desired. \square

6 Open problems

6.1 Minimal walks for general $S = \{(1, 0), (0, 1), (u, v), (v, u)\}$.

Based on the results in Section 5, it seems natural to explore the more general case that our allowable steps are $S = \{(1, 0), (0, 1), (u, v), (v, u)\}$ for arbitrary u and v such that $u > v \geq 1$.

In this case, the distances seem more complicated than they were in any of our previous examples. The reason for this is that a point's distance from the origin is inherently dependent on its position relative to the \mathbb{N} -span of $\{(u, v), (v, u)\}$. In particular, that distance to point (a, b) can be determined greedily by finding a nearest point in the \mathbb{N} -span of $\{(u, v), (v, u)\}$ that lies weakly south and west of (a, b) and filling in the rest of the walk




Figure 5: The distance $d(a, b; \{(1, 0), (0, 1), (3, 5), (5, 3)\})$ for $0 \leq a, b \leq 15$.

with short steps. This means the distance changes, often dramatically, depending on a point's position in a fundamental parallelogram spanned by (u, v) and (v, u) . For example, consider the case that $S = \{(1, 0), (0, 1), (3, 5), (5, 3)\}$. The distances from the origin to nearby points (a, b) are displayed in Figure 5.

Problem 10. Let $S = \{(1, 0), (0, 1), (u, v), (v, u)\}$ with $u > v \geq 1$ arbitrary. Determine $d(a, b; S)$ and $|\mathcal{W}(a, b; S)|$.

6.2 Catalan generalizations

Based on the wealth of beautiful combinatorics arising from Catalan and Motzkin paths, the following question is very natural.

Problem 11. Let $a \geq b$ and let S be a set of allowable steps. How many minimal S -walks terminating at (a, b) stay weakly below the line $y = x$?

For brevity, let us say that an S -Catalan walk is a minimal S -walk that stays weakly below the line $y = x$. For integers a, b with $a \geq b$, we will write $C(a, b; S)$ to denote the set of all S -Catalan walks terminating at (a, b) .

Consider the set $S_n = \{(i, n - i) : 0 \leq i \leq n\}$; i.e., the nonnegative integer vectors with coordinate sum n . We can explore $C(a, b; S_n)$ for several values of n .

For $S_1 = \{(1, 0), (0, 1)\}$ the number of S_1 -Catalan walks terminating at (a, a) is the a -th Catalan number, which appear in OEIS as sequence [A000108](#). More generally, the number of walks terminating at (a, b) is $\frac{a-b+1}{a+b+1} \binom{a+b+1}{a+1}$, which is OEIS sequence [A009766](#). Krattenthaler [2, Corollary 10.3.2] gives a proof of this fact.

9																	120											
8																0	0	184										
7															0	52	0	0	234									
6														13	0	0	68	0	0	212								
5													0	0	18	0	0	64	0	0	158							
4												0	6	0	0	21	0	0	50	0	0	99						
3											2	0	0	7	0	0	16	0	0	30	0	0	50					
2										0	0	2	0	0	5	0	0	9	0	0	14	0	0	20				
1									0	1	0	0	2	0	0	3	0	0	4	0	0	5	0	0	6			
0	1	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1	0	0	1
b/a	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18									

Figure 6: Number of S_3 -Catalan paths terminating at points (a, b) with $0 \leq a + b \leq 18$.

For $S_2 = \{(2, 0), (1, 1), (0, 2)\}$, the number of S_2 -Catalan walks terminating at (a, a) is the a -th Motzkin number, which is found in sequence [A001006](#). Indeed, the map sending $(2, 0) \mapsto (1, 1)$, $(1, 1) \mapsto (1, 0)$, and $(0, 2) \mapsto (1, -1)$ gives a bijection to classical Motzkin paths comprised of steps $\{(1, 1), (1, 0), (1, -1)\}$ that start at $(0, 0)$, terminate at $(a, 0)$, and stay above the line $y = 0$.

As a next step we can consider $S_3 = \{(3, 0), (2, 1), (1, 2), (0, 3)\}$. Figure 6 shows the number of S_3 -Catalan paths terminating at points (a, b) with $a \geq b$ and $a + b \leq 18$.

The sequence of nonzero numbers along the line $y = x$, which continues as 1, 2, 13, 120, 1288, 15046, . . . , did not appear in OEIS. We have added it as sequence [A292437](#).

Problem 12. Determine the generating function for

$$\sum_{a \geq b \geq 0} |C(a, b; S_3)| x^a y^b \quad \text{or} \quad \sum_{a=0}^{\infty} |C(a, a; S_3)| t^a.$$

7 Acknowledgments

We gratefully acknowledge support from NSF grant DMS-1600048. We also thank Naomi Cameron, Tom Edgar, Lara Pudwell, and Jennifer Quinn for providing helpful insights and suggestions over the course of this project.

References

- [1] OEIS Foundation Inc. (2016). The On-Line Encyclopedia of Integer Sequences, <http://oeis.org>.
- [2] Christian Krattenthaler, Lattice path enumeration. In *Handbook of Enumerative Combinatorics*, Discrete Math. Appl. (Boca Raton), pp. 589–678. CRC Press, 2015.
- [3] W. A. Stein et al., *Sage Mathematics Software (Version 8.0)*. The Sage Development Team, 2017. <http://www.sagemath.org>.

2010 *Mathematics Subject Classification*: Primary 05A15; Secondary 05A19.

Keywords: lattice path enumeration, generating function, Catalan number, Motzkin number.

(Concerned with sequences [A000108](#), [A001006](#), [A007318](#), [A009766](#), [A292435](#), [A292436](#), and [A292437](#).)

Received December 9 2017; revised version received March 27 2018. Published in *Journal of Integer Sequences*, March 28 2018.

Return to [Journal of Integer Sequences home page](#).