

Stein Extensions of Riemann Symmetric Spaces and some Generalization

Dedicated to Professor Keisaku Kumahara on his 60th birthday

Toshihiko Matsuki

Communicated by S. Gindikin

Abstract. It was proved by Huckleberry that the Akhiezer-Gindikin domain is included in the “Iwasawa domain” using complex analysis. We show here that no complex analysis is necessary for a proof of this result. Indeed we generalize the notions of the Akhiezer-Gindikin domain and the Iwasawa domain for two associated symmetric subgroups in real Lie groups and prove the inclusion. Moreover, by the symmetry of two associated symmetric subgroups, we also give a direct proof of the known fact that the Akhiezer-Gindikin domain is included in all cycle spaces.

1. Introduction

1.1. Akhiezer-Gindikin domain and Iwasawa domain

Let $G_{\mathbb{C}}$ be a connected complex semisimple Lie group and $G_{\mathbb{R}}$ a connected real form of $G_{\mathbb{C}}$. Let $K_{\mathbb{C}}$ be the complexification in $G_{\mathbb{C}}$ of a maximal compact subgroup K of $G_{\mathbb{R}}$. Let $\mathfrak{g}_{\mathbb{R}} = \mathfrak{k} \oplus \mathfrak{m}$ denote the Cartan decomposition of $\mathfrak{g}_{\mathbb{R}} = \text{Lie}(G_{\mathbb{R}})$ with respect to K . Let \mathfrak{t} be a maximal abelian subspace in \mathfrak{m} . Put

$$\mathfrak{t}^+ = \{Y \in \mathfrak{t} \mid |\alpha(Y)| < \frac{\pi}{2} \text{ for all } \alpha \in \Sigma\}$$

where Σ is the restricted root system of $\mathfrak{g}_{\mathbb{C}}$ with respect to \mathfrak{t} . Then the “Akhiezer-Gindikin domain” D in $G_{\mathbb{C}}$ is defined in [AG] by

$$D = G_{\mathbb{R}}(\exp \mathfrak{t}^+)K_{\mathbb{C}}.$$

Let B be a Borel subgroup of $G_{\mathbb{C}}$ such that $G_{\mathbb{R}}B$ is closed in $G_{\mathbb{C}}$. Then $K_{\mathbb{C}}B$ is the unique open dense $K_{\mathbb{C}}-B$ double coset in $G_{\mathbb{C}}$ ([M2]). Define an open subset

$$\Omega = \{x \in G_{\mathbb{C}} \mid x^{-1}G_{\mathbb{R}}B \subset K_{\mathbb{C}}B\}$$

in $G_{\mathbb{C}}$. Clearly, Ω is left $G_{\mathbb{R}}$ -invariant and right $K_{\mathbb{C}}$ -invariant. The connected component Ω_0 of Ω containing the identity is often called the “Iwasawa domain”.

Remark 1. Let S_j ($j \in J$) be the $K_{\mathbb{C}}$ - B double cosets in $G_{\mathbb{C}}$ of (complex) codimension one and T_j the closure of S_j . Then the complement of $K_{\mathbb{C}}B$ in $G_{\mathbb{C}}$ is $\bigcup_{j \in J} T_j$. So we can write

$$\begin{aligned}\Omega &= \{x \in G_{\mathbb{C}} \mid x^{-1}G_{\mathbb{R}} \cap T_j = \phi \text{ for all } j \in J\} \\ &= \{x \in G_{\mathbb{C}} \mid x \notin gT_j^{-1} \text{ for all } j \in J \text{ and } g \in G_{\mathbb{R}}\}.\end{aligned}$$

Thus Ω_0 is Stein because Ω is the complement of an infinite family $\{gT_j^{-1} \mid j \in J, g \in G_{\mathbb{R}}\}$ of complex hypersurfaces.

The equality

$$D = \Omega_0 \tag{1.1}$$

was proved in [GM] when $G_{\mathbb{R}}$ is classical type or exceptional Hermitian type. Independently, Krötz and Stanton proved $D \subset \Omega_0$ for classical cases in [KS]. We should note that the proofs in these two papers are based on elementary linear algebraic computations. (Remark that [FH] did not refer [GM] in their historical reference on (1.1).) On the other hand, Barchini proved the inclusion $D \supset \Omega_0$ by a general but elementary argument in [B]. Recently, Huckleberry ([H], Proposition 2.0.2 in [FH]) gave a proof of the opposite inclusion

$$D \subset \Omega_0 \tag{1.2}$$

by using strictly plurisubharmonicity of a function ρ proved in [BHH] (revised version).

But we can see that we need no complex analysis to prove (1.2). In this paper, we will generalize the notions of the Akhiezer-Gindikin domain and the Iwasawa domain for real Lie groups as in the next subsection and prove the inclusion (1.2).

1.2. Generalization to real Lie groups

Let G be a connected real semisimple Lie group and σ an involution of G . Take a Cartan involution θ such that $\sigma\theta = \theta\sigma$. Put $H = G_0^\sigma = \{g \in G \mid \sigma(g) = g\}_0$ and $H' = G_0^{\sigma\theta}$. Then the symmetric subgroup H' is called “associated” to the symmetric subgroup H (c.f. [M1]). The structure of the double coset decomposition $H' \backslash G / H$ is precisely studied in [M3] in a general setting for arbitrary two involutions.

Remark 2. (i) For example, if $G = G_{\mathbb{C}}$, then $K_{\mathbb{C}}$ and $G_{\mathbb{R}}$ are associated.

(ii) We should remark here that all the results on Jordan decompositions and elliptic elements etc. for the decomposition $G_{\mathbb{R}} \backslash G_{\mathbb{C}} / K_{\mathbb{C}}$ in Section 3 of [FH] were already proved in [M3]. In [FH], they are not referred at all.

Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m} = \mathfrak{h} \oplus \mathfrak{q}$ be the $+1, -1$ -eigenspace decompositions of $\mathfrak{g} = \text{Lie}(G)$ with respect to θ and σ , respectively. Then the Lie algebra \mathfrak{h}' of H' is written as $\mathfrak{h}' = (\mathfrak{k} \cap \mathfrak{h}) \oplus (\mathfrak{m} \cap \mathfrak{q})$. Let \mathfrak{t} be a maximal abelian subspace of $\mathfrak{k} \cap \mathfrak{q}$. Then we can define the root space

$$\mathfrak{g}_{\mathbb{C}}(\mathfrak{t}, \alpha) = \{X \in \mathfrak{g}_{\mathbb{C}} \mid [Y, X] = \alpha(Y)X \text{ for all } Y \in \mathfrak{t}\}$$

for any linear form $\alpha : \mathfrak{t} \rightarrow i\mathbb{R}$. Here $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \oplus i\mathfrak{g}$ is the complexification of \mathfrak{g} . Put

$$\Sigma = \Sigma(\mathfrak{g}_{\mathbb{C}}, \mathfrak{t}) = \{\alpha \in i\mathfrak{t}^* - \{0\} \mid \mathfrak{g}_{\mathbb{C}}(\mathfrak{t}, \alpha) \neq \{0\}\}.$$

Then Σ satisfies the axiom of the root system ([R]). Since $\theta(Y) = Y$ for all $Y \in \mathfrak{t}$, we can decompose $\mathfrak{g}_{\mathbb{C}}(\mathfrak{t}, \alpha)$ into $+1, -1$ -eigenspaces for θ as

$$\mathfrak{g}_{\mathbb{C}}(\mathfrak{t}, \alpha) = \mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha) \oplus \mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \alpha). \tag{1.3}$$

Define a subset $\Sigma(\mathfrak{m}_{\mathbb{C}}, \mathfrak{t}) = \{\alpha \in i\mathfrak{t}^* - \{0\} \mid \mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \alpha) \neq \{0\}\}$ of Σ and put

$$\mathfrak{t}^+ = \{Y \in \mathfrak{t} \mid |\alpha(Y)| < \frac{\pi}{2} \text{ for all } \alpha \in \Sigma(\mathfrak{m}_{\mathbb{C}}, \mathfrak{t})\}.$$

Then we define a generalization of the Akhiezer-Gindikin domain D in G by

$$D = H'T^+H.$$

where $T^+ = \exp \mathfrak{t}^+$. (We will show in Proposition 1 that D is open in G .)

Let P be an arbitrary parabolic subgroup of G such that $H'P$ is closed in G . Then HP is open in G by [M2]. Define an open subset

$$\Omega = \{x \in G \mid x^{-1}H'P \subset HP\} = \{x \in G \mid H'x \subset PH\}$$

in G . Then we may call the connected component Ω_0 a “generalized Iwasawa domain”. As a complete generalization of (1.2), we can prove the following theorem.

Theorem. $D \subset \Omega_0$.

Here we should explain the construction of this paper to prove this theorem. In Section 2, we prove properties on the generalized Akhiezer-Gindikin domain D . Lemma 2 is the most important basic technical lemma. It implies $(\mathfrak{h}' \cap \text{Ad}(a)\mathfrak{h}) \oplus (\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q}) \subset \mathfrak{k}$ for $a \in T^+$ where $\mathfrak{q}' = (\mathfrak{k} \cap \mathfrak{q}) \oplus (\mathfrak{m} \cap \mathfrak{h})$. Note that $H' \cap aHa^{-1}$ is the isotropy subgroup of the action of H' at the point $aH \in G/H$. So the inclusion $\mathfrak{h}' \cap \text{Ad}(a)\mathfrak{h} \subset \mathfrak{k}$ is a generalization of Proposition 2 in [AG]. But we do not find statements in [AG] corresponding to the inclusion $\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q} \subset \mathfrak{k}$. This inclusion is the key of this paper.

In Proposition 1, we show that D is open in G . This is a generalization of Proposition 4 in [AG]. We also give a precise orbit structure $H' \backslash D/H \cong T^+/W$ of D where $W = N_{K \cap H}(\mathfrak{t})/Z_{K \cap H}(\mathfrak{t})$ in Proposition 2. This is a generalization of Proposition 8 in [AG]. In Section 3, we construct a left H' -invariant right H -invariant real analytic function ρ on D and prepare the key lemma (Lemma 3) which follows from Lemma 2. In Section 4, we prove Theorem. Basic formulation is the same as Proposition 2.0.2 in [FH]. But we do not need complex analysis.

1.3. Application to cycle spaces

Note that we can exchange the roles of H and H' . We can apply this to the pair of $K_{\mathbb{C}}$ and $G_{\mathbb{R}}$ in $G_{\mathbb{C}}$. Let P be a parabolic subgroup of $G_{\mathbb{C}}$ such that $S = K_{\mathbb{C}}P$ is closed in $G_{\mathbb{C}}$. Then $S' = G_{\mathbb{R}}P$ is open in $G_{\mathbb{C}}$. Put

$$\Omega(S) = \{x \in G_{\mathbb{C}} \mid x^{-1}K_{\mathbb{C}}P \subset G_{\mathbb{R}}P\}.$$

Then by Theorem, we have $D^{-1} = K_{\mathbb{C}}T^+G_{\mathbb{R}} \subset \Omega(S)_0$ by the notation in Section 1.1 and therefore $D \subset \Omega(S)_0^{-1}$. Since $\Omega(S)^{-1} = \{x \in G_{\mathbb{C}} \mid xK_{\mathbb{C}}P \subset G_{\mathbb{R}}P\} = \{x \in G_{\mathbb{C}} \mid xS \subset S'\}$, the domain $\Omega(S)_0^{-1}$ is usually called the “cycle space” for S' . Hence we have given a direct proof of the known fact:

Corollary. Akhizer-Gindikin domain D is included in all cycle spaces.

(Remark: This fact was known by combining (1.2) and Proposition 8.3 in [GM] because Proposition 8.3 implies that the Iwasawa domain Ω_0 is included in all the cycle spaces $\Omega(S)_0^{-1}$.)

Acknowledgement: The author would like to express his hearty thanks to S. Gindikin for his encouragement and for many useful discussions.

2. A generalization of Akhiezer-Gindikin domain

We will use the notations in Section 1.2 (not in Section 1.1). In this section, we will prepare basic results on a generalization of the Akhiezer-Gindikin domain by extending elementary arguments in [M3] Section 3.

First we should note that we have only to consider the problem on each minimal σ -stable ideals of \mathfrak{g} . So we may assume that \mathfrak{g} has no proper σ -stable ideals. We may also assume that \mathfrak{g} is noncompact, otherwise we have $P = G$ and the problem is trivial.

Let $[\mathfrak{m}, \mathfrak{m}]$ denote the linear subspace of \mathfrak{k} spanned by $[Y, Z]$ for $Y, Z \in \mathfrak{m}$. Then $\mathfrak{g}' = [\mathfrak{m}, \mathfrak{m}] \oplus \mathfrak{m}$ becomes a σ -stable ideal of \mathfrak{g} because \mathfrak{g}' is σ -stable and $[\mathfrak{k}, [\mathfrak{m}, \mathfrak{m}]] \subset [[\mathfrak{k}, \mathfrak{m}], \mathfrak{m}] + [\mathfrak{m}, [\mathfrak{k}, \mathfrak{m}]] \subset [\mathfrak{m}, \mathfrak{m}]$. Since $\mathfrak{g}' = \mathfrak{g}$, we have $\mathfrak{k} = [\mathfrak{m}, \mathfrak{m}]$ and therefore

$$\mathfrak{k}_{\mathbb{C}} = [\mathfrak{m}_{\mathbb{C}}, \mathfrak{m}_{\mathbb{C}}]. \tag{2.1}$$

Lemma 1. (i) Every root $\alpha \in \Sigma$ such that $\mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha) \neq \{0\}$ is written as a sum of two elements in $\Sigma(\mathfrak{m}_{\mathbb{C}}, \mathfrak{t}) \cup \{0\}$.

(ii) If $Y \in \mathfrak{t}^+$, then $|\alpha(Y)| < \pi$ for all $\alpha \in \Sigma$.

(iii) If $Y \in \mathfrak{t}$ satisfies $\alpha(Y) = 0$ for all $\alpha \in \Sigma$, then $Y = 0$.

(iv) If $Y, Z \in \mathfrak{t}^+$ satisfy $\exp Y \in (\exp Z)(T \cap H)$, then $Y = Z$. Especially, \exp is injective on \mathfrak{t}^+ .

Proof. (i) Let X be a nonzero element in $\mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha)$. Then by (2.1), we can write $X = \sum_j [Y_j, Z_j]$ with some nonzero elements $Y_j \in \mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \beta_j)$ and $Z_j \in \mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \gamma_j)$ such that $\beta_j + \gamma_j = \alpha$.

(ii) is clear by (i) and the definition of \mathfrak{t}^+ .

(iii) If $\alpha(Y) = 0$ for all $\alpha \in \Sigma$, then Y commutes with $\mathfrak{g}_{\mathbb{C}}(\mathfrak{t}, \alpha)$ for all $\alpha \in \Sigma \cup \{0\}$. Hence Y commutes with \mathfrak{g} and therefore $Y = 0$ because \mathfrak{g} is semisimple.

(iv) Suppose $\exp Y = (\exp Z)h$ with some $Y, Z \in \mathfrak{t}^+$ and $h \in T \cap H$. Applying σ , we have $\exp(-Y) = (\exp(-Z))h$. So we have $\exp 2Y = \exp 2Z$. Put $a = \exp 2Y = \exp 2Z$ and apply $\text{Ad}(a)$ to $\mathfrak{g}_{\mathbb{C}}(\mathfrak{t}, \alpha)$ for $\alpha \in \Sigma(\mathfrak{m}_{\mathbb{C}}, \mathfrak{t})$. Then we have $e^{2\alpha(Y)} = e^{2\alpha(Z)}$ and therefore $\alpha(Y) - \alpha(Z) \in \pi i\mathbb{Z}$. Since $|\alpha(Y)| < \pi/2$

and $|\alpha(Z)| < \pi/2$ by the definition of \mathfrak{t}^+ , we have $\alpha(Y - Z) = \alpha(Y) - \alpha(Z) = 0$ for all $\alpha \in \Sigma(\mathfrak{m}_{\mathbb{C}}, \mathfrak{t})$. This implies $\alpha(Y - Z) = 0$ for all $\alpha \in \Sigma$ by (i) and therefore $Y - Z = 0$ by (iii). ■

Let Y be an element of \mathfrak{t} and put $a = \exp Y$. Consider the conjugate $\sigma_a = \text{Ad}(a)\sigma\text{Ad}(a)^{-1} = \sigma\text{Ad}(a)^{-2}$ of σ by $\text{Ad}(a)$. Then σ_a is an involution of \mathfrak{g} such that $\mathfrak{g}^{\sigma_a} = \text{Ad}(a)\mathfrak{h}$ and $G_0^{\sigma_a} = aHa^{-1}$. Put $\tau = \sigma\theta = \theta\sigma$. The key idea of [M3] was to consider the automorphism $\tau\sigma_a$ (which is not involutive in general) of \mathfrak{g} . Since $\tau\sigma_a = \tau\sigma\text{Ad}(a)^{-2} = \theta\text{Ad}(a)^{-2}$ and since θ and $\text{Ad}(a)$ commutes, the automorphism $\tau\sigma_a$ is semisimple. Hence by Lemma 1 in [M3], we have a direct sum decomposition

$$\mathfrak{g} = (\mathfrak{h}' + \text{Ad}(a)\mathfrak{h}) \oplus (\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q}) \tag{2.2}$$

where $\mathfrak{q}' = (\mathfrak{k} \cap \mathfrak{q}) \oplus (\mathfrak{m} \cap \mathfrak{h})$. On the other hand, we have the $+1, -1$ -eigenspace decomposition of $\mathfrak{g}^{\tau\sigma_a}$ for τ by $\mathfrak{g}^{\tau\sigma_a} = (\mathfrak{g}^{\tau\sigma_a} \cap \mathfrak{h}') \oplus (\mathfrak{g}^{\tau\sigma_a} \cap \mathfrak{q}') = (\mathfrak{h}' \cap \text{Ad}(a)\mathfrak{h}) \oplus (\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q})$ as in [M3] Section 3.

Lemma 2. *If $Y \in \mathfrak{t}^+$, then $\mathfrak{g}^{\tau\sigma_a} = \mathfrak{z}_{\mathfrak{k}}(Y) = \{X \in \mathfrak{k} \mid [Y, X] = 0\}$.*

Proof. By (1.3), we have a direct sum decomposition

$$\mathfrak{g}_{\mathbb{C}} = \bigoplus_{\alpha \in \Sigma \cup \{0\}} (\mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha) \oplus \mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \alpha)). \tag{2.3}$$

If $X \in \mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha)$, then we have $\tau\sigma_a X = \theta\text{Ad}(a)^{-2}X = e^{-2\alpha(Y)}X$. On the other hand, if $X \in \mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \alpha)$, then we have $\tau\sigma_a X = \theta\text{Ad}(a)^{-2}X = -e^{-2\alpha(Y)}X$. Hence the decomposition (2.3) is the eigenspace decomposition of $\mathfrak{g}_{\mathbb{C}}$ for $\tau\sigma_a$. We have only to verify whether every direct summand in (2.3) is contained in $\mathfrak{g}_{\mathbb{C}}^{\tau\sigma_a}$ or not.

Note that $\alpha(Y)$ is pure imaginary. Let X be a nonzero element in $\mathfrak{m}_{\mathbb{C}}(\mathfrak{t}, \alpha)$. Since $|\alpha(Y)| < \pi/2$ by the assumption, we have $-e^{-2\alpha(Y)} \neq 1$. Hence $X \notin \mathfrak{g}_{\mathbb{C}}^{\tau\sigma_a}$. On the other hand, let X be a nonzero element in $\mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha)$. Since $|\alpha(Y)| < \pi$ by Lemma 1 (ii), we have $e^{-2\alpha(Y)} = 1 \iff \alpha(Y) = 0$. Hence $X \in \mathfrak{g}_{\mathbb{C}}^{\tau\sigma_a} \iff \alpha(Y) = 0$. Thus we have proved

$$\mathfrak{g}_{\mathbb{C}}^{\tau\sigma_a} = \bigoplus_{\alpha \in \Sigma \cup \{0\}, \alpha(Y)=0} \mathfrak{k}_{\mathbb{C}}(\mathfrak{t}, \alpha) = \mathfrak{z}_{\mathfrak{k}_{\mathbb{C}}}(Y)$$

and therefore $\mathfrak{g}^{\tau\sigma_a} = \mathfrak{z}_{\mathfrak{k}}(Y)$. ■

Since $\mathfrak{g}^{\tau\sigma_a} = \mathfrak{z}_{\mathfrak{k}}(Y)$ is a compact Lie algebra and since \mathfrak{t} is maximal abelian in $\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q} = \mathfrak{z}_{\mathfrak{h} \cap \mathfrak{q}}(Y)$, we have $\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q} = \text{Ad}(H' \cap aHa^{-1})_0\mathfrak{t}$. Moreover if U is a neighborhood of the origin 0 in \mathfrak{t} , then

$$V = \text{Ad}(H' \cap aHa^{-1})_0U \text{ is a neighborhood of } 0 \text{ in } \mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q}. \tag{2.4}$$

Proposition 1. *D is open in G .*

Proof. By the left H' -action and the right H -action on D , we have only to show that a neighborhood of $a = \exp Y$ for $Y \in \mathfrak{t}^+$ is contained in D . Take a neighborhood $U = \{Z - Y \mid Z \in \mathfrak{t}^+\}$ of 0 in \mathfrak{t} . Then we have $T^+a^{-1} = \exp U$. Hence

$$\begin{aligned} e \in H'T^+Ha^{-1} &= H'T^+a^{-1}aHa^{-1} = H'(\exp U)aHa^{-1} \\ &= H' \exp(\text{Ad}(H' \cap aHa^{-1})_0U)aHa^{-1} = H'(\exp V)aHa^{-1}. \end{aligned} \tag{2.5}$$

Since V is a neighborhood of 0 in $\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q}$ by (2.4), it follows from (2.2) that $H'T^+Ha^{-1}$ contains a neighborhood of e . Hence $H'T^+H$ contains a neighborhood of a . ■

Proposition 2. *Let a and b be elements of T^+ . Then $b = \ell ah^{-1}$ for some $\ell \in H'$ and $h \in H \iff b = waw^{-1}$ for some $w \in N_{K \cap H}(\mathfrak{t})$. Here $N_{K \cap H}(\mathfrak{t})$ is the normalizer of \mathfrak{t} in $K \cap H$. (Write $K = G^\theta$ as usual).*

Proof. Since the implication \Leftarrow is clear, we have only to prove \Rightarrow . Suppose $b = \ell ah^{-1}$ for some $\ell \in H'$ and $h \in H$. Put $\mathfrak{t}' = \text{Ad}(\ell)\mathfrak{t} = \text{Ad}(\ell a)\mathfrak{t} = \text{Ad}(bh)\mathfrak{t}$. Then \mathfrak{t}' is a maximal abelian subspace of $\mathfrak{q}' \cap \text{Ad}(b)\mathfrak{q}$. Since $(\mathfrak{h}' \cap \text{Ad}(b)\mathfrak{h}, \mathfrak{q}' \cap \text{Ad}(b)\mathfrak{q})$ is a compact symmetric pair by Lemma 2, there is an $x \in H' \cap bHb^{-1}$ such that $\text{Ad}(x)\mathfrak{t}' = \mathfrak{t}$. Put $\ell' = x\ell \in H'$ and $h' = b^{-1}xbh \in H$. Then we have $\ell'ah'^{-1} = x\ell ah^{-1}b^{-1}x^{-1}b = xbh h^{-1}b^{-1}x^{-1}b = b$ and

$$\begin{aligned} \ell'Th'^{-1} &= x\ell Th^{-1}b^{-1}x^{-1}b = x\ell aTh^{-1}b^{-1}x^{-1}b \\ &= xbhTh^{-1}b^{-1}x^{-1}b = xT'x^{-1}b = Tb = T \end{aligned} \tag{2.6}$$

where $T = \exp \mathfrak{t}$ and $T' = \exp \mathfrak{t}'$.

Since $\ell'h'^{-1} = \ell'eh'^{-1} \in T$ by (2.6) and since $\ell'T\ell'^{-1}\ell'h'^{-1} = T$ also by (2.6), we have $\ell'T\ell'^{-1} = T$. Write $\ell' = k \exp X$ with $k \in K \cap H$ and $X \in \mathfrak{m} \cap \mathfrak{q}$. Then it is well-known that $[X, \mathfrak{t}] = \{0\}$ by a standard argument. (Suppose $\text{Ad}(k \exp X)Y = Y'$ for some $Y, Y' \in \mathfrak{k}$. Then applying θ , we have $\text{Ad}(k \exp(-X))Y = Y'$ and therefore $\text{Ad}(\exp 2X)Y = Y$. Since $\text{ad}X : \mathfrak{g} \rightarrow \mathfrak{g}$ is expressed by a real symmetric matrix, we have $[X, Y] = 0$.)

Hence we have $kTk^{-1} = T$ and $b = \ell'al'^{-1}\ell'h'^{-1} = kak^{-1}\ell'h'^{-1}$. We have only to prove $c = \ell'h'^{-1} = e$. Write $h' = k' \exp X'$ with $k' \in K \cap H$ and $X' \in \mathfrak{m} \cap \mathfrak{h}$. Then it follows from $\ell' = ch'$ that $k \exp X = ck' \exp X'$. Since $c \in T \subset K$, we have $k = ck'$ and $X = X' = 0$. Hence $c \in T \cap H$. Since $kak^{-1}, b \in \exp \mathfrak{t}^+$, we have $c = e$ by Lemma 1 (iv). ■

3. Construction of a function ρ

Write $W = N_{K \cap H}(\mathfrak{t})/Z_{K \cap H}(\mathfrak{t})$. Let ρ_0 be a W -invariant real analytic function on \mathfrak{t}^+ which has no critical points except the origin. For the sake of later use, we should also assume $\rho_0(Z)$ tends to $+\infty$ when Z goes to the boundary of \mathfrak{t}^+ . For example, we may put

$$\rho_0(Z) = \sum_{\alpha \in \Sigma(\mathfrak{m}_{\mathbb{C}}, \mathfrak{t})} \frac{1}{\pi - 2i\alpha(Z)}. \tag{3.1}$$

This function (3.1) is clearly convex and therefore it has no critical points except the origin. By Proposition 2 and Lemma 1 (iv), we can define a function ρ on D by

$$\rho(\ell(\exp Z)h) = \rho_0(Z) \quad \text{for } \ell \in H', h \in H \text{ and } Z \in \mathfrak{t}^+.$$

Proposition 3. ρ is real analytic on D .

Proof. By the left H' -action and the right H -action on D , we have only to show ρ is real analytic at every $a = \exp Y \in T^+$. Consider the right a -translate $\rho_a(x) = \rho(xa)$ for $x \in Da^{-1}$ of ρ . Since ρ_a is left H' -invariant and right aHa^{-1} -invariant, we have only to show that the function $\rho'_a(X) = \rho_a(\exp X)$ for $X \in V$ is real analytic at 0 by (2.2) and (2.5) where $U \subset \mathfrak{t}$ and $V = \text{Ad}(H' \cap aHa^{-1})_0 U \subset \mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q}$ are as in the proof of Proposition 1. Since $\rho_a(gxg^{-1}) = \rho_a(x)$ for $g \in H' \cap aHa^{-1}$ and $x \in \exp V$, ρ'_a is $\text{Ad}(H' \cap aHa^{-1})$ -invariant. Note that $(H' \cap aHa^{-1})_0 = (Z_{K \cap H}(Y))_0$ by Lemma 2 and that $\rho'_a(Z) = \rho((\exp Z)a) = \rho_0(Z+Y)$ for $Z \in U$ is invariant under the action of $w \in W$ such that $w(Y) = Y$. Since we can easily extend the well-known Chevalley's restriction theorem to real analytic functions at 0, the function ρ'_a on $V = \text{Ad}(H' \cap aHa^{-1})_0 U$ is real analytic at 0. ■

Remark 3. (i) The function ρ on D has no critical points outside $H'H = H'eH$ by the assumption on ρ_0 .

(ii) If ρ_0 is a W -invariant smooth function on \mathfrak{t}^+ . Then we can show that ρ is a smooth function on D by using [S].

The tangent space $T_a(G)$ of G at a is identified with $\mathfrak{g} = T_e(G)$ by the right a -action. In other words, we identify $T_a(G)$ with the left infinitesimal action of \mathfrak{g} at a . Now we have the following key lemma.

Lemma 3. Let $a = \exp Y$ with $Y \in \mathfrak{t}^+ - \{0\}$. Then the hyperplane in $T_a(G)$ defined by $d\rho = 0$ is orthogonal, with respect to the Killing form on \mathfrak{g} , to a nonzero vector Z in \mathfrak{k} .

Proof. Taking the right a -translate ρ_a of ρ as in the proof of Proposition 3, we have only to consider the hyperplane in the tangent space $T_e(G) \cong \mathfrak{g}$ defined by $d\rho_a = 0$.

Since ρ_a is left H' -invariant and right aHa^{-1} -invariant, the differential $d\rho_a$ vanishes on $\mathfrak{h}' + \text{Ad}(a)\mathfrak{h}$. Hence the normal vector Z is contained in the orthogonal complement $\mathfrak{q}' \cap \text{Ad}(a)\mathfrak{q} (\subset \mathfrak{k}$ by Lemma 2) of $\mathfrak{h}' + \text{Ad}(a)\mathfrak{h}$. ■

Proof of Theorem. The basic formulation is the same as Proposition 2.0.2 in [FH]. Suppose that $D \not\subset PH$. We will deduce a contradiction.

Let PxH be a P - H double coset with the least dimension among the P - H double cosets intersecting D . So the intersection $PxH \cap D$ is relatively closed in D . Since $H'P = (K \cap H)P$ by [M2], we have $H' = (K \cap H)(P \cap H') = (P \cap H')(K \cap H)$. Hence PxH intersects $(K \cap H)T^+$ and the image of $\rho|_{PxH \cap D}$ is equal to the image of $\rho|_{PxH \cap (K \cap H)T^+}$. The set $\{x \in (K \cap H)T^+ \mid \rho(x) \leq m\}$ is compact for any $m \in \mathbb{R}$ because we carefully assumed that ρ_0 is $+\infty$ on the boundary of \mathfrak{t}^+ . Hence the

function $\rho|_{PxH \cap D}$ attains its minimum on some point ka with $k \in K \cap H$ and $a \in T^+$. Replacing P by the k -conjugate $k^{-1}Pk$, we may assume $k = e$. Since $a \in PxH$ and $PxH \cap PH = \phi$, we have $a \neq e$.

By Lemma 3, there is a nonzero element Z in \mathfrak{k} such that Z is orthogonal to $\mathfrak{p} = \text{Lie}(P)$. But this leads a contradiction because $Z \in \mathfrak{k}$ is also orthogonal to $\theta\mathfrak{p}$ and therefore Z is orthogonal to $\mathfrak{p} + \theta\mathfrak{p} = \mathfrak{g}$ which cannot happen since the Killing form is nondegenerate on \mathfrak{g} . ■

References

- [AG] Akhiezer, D. N., and S. G. Gindikin, *On Stein extensions of real symmetric spaces*, Math. Ann. **286** (1990), 1–12.
- [B] Barchini, L., *Stein extensions of real symmetric spaces and the geometry of the flag manifold*, preprint.
- [BHH] Burns, D., S. Halverscheid, and R. Hind, *The geometry of Grauert tubes and complexification of symmetric spaces*, preprint (CV/0109186).
- [FH] Fels, G., and A. Huckleberry, *Characterization of cycle domains via Kobayashi hyperbolicity*, preprint (AG/0204341).
- [GM] Gindikin, S., and T. Matsuki, *Stein extensions of Riemann symmetric spaces and dualities of orbits on flag manifolds*, preprint (MSRI-Preprint 2001-028).
- [H] Huckleberry, A., *On certain domains in cycle spaces of flag manifolds*, Math. Annalen, to appear.
- [KS] Krötz, B., and R. Stanton, *Holomorphic extension of a representation I: automorphic functions*, preprint.
- [M1] Matsuki, T., *The orbits of affine symmetric spaces under the action of minimal parabolic subgroups*, J. Math. Soc. Japan **31** (1979), 331–357.
- [M2] —, *Closure relations for orbits on affine symmetric spaces under the action of parabolic subgroups. Intersections of associated orbits*, Hiroshima Math. J. **18** (1988), 59–67.
- [M3] —, *Double coset decompositions of reductive Lie groups arising from two involutions*, J. of Algebra **197** (1997), 49–91.
- [R] Rossmann, W., *The structure of semisimple symmetric spaces*, Canad. J. Math. **31** (1979), 157–180.
- [S] Schwarz, G., *Smooth functions invariant under the action of a compact Lie group*, Topology **14** (1975), 63–68.

Toshihiko Matsuki
 Faculty of Integrated Human Studies
 Kyoto University
 Kyoto 606-8501, Japan
 matsuki@math.h.kyoto-u.ac.jp

Received August 28, 2002
 and in final form September 6, 2002