

On sum-sets and product-sets of complex numbers

par JÓZSEF SOLYMOSI

RÉSUMÉ. On donne une preuve simple que pour tout ensemble fini de nombres complexes A , la taille de l'ensemble de sommes $A + A$ ou celle de l'ensemble de produits $A \cdot A$ est toujours grande.

ABSTRACT. We give a simple argument that for any finite set of complex numbers A , the size of the the sum-set, $A + A$, or the product-set, $A \cdot A$, is always large.

1. Introduction

Let A be a finite subset of complex numbers. The *sum-set* of A is $A + A = \{a + b : a, b \in A\}$, and the *product-set* is given by $A \cdot A = \{a \cdot b : a, b \in A\}$. Erdős conjectured that for any n -element set the sum-set or the product-set should be close to n^2 . For integers, Erdős and Szemerédi [7] proved the lower bound $n^{1+\varepsilon}$.

$$\max(|A + A|, |A \cdot A|) \geq |A|^{1+\varepsilon}.$$

Nathanson [9] proved the bound with $\varepsilon = 1/31$, Ford [8] improved it to $\varepsilon = 1/15$, and the best bound is obtained by Elekes [6] who showed $\varepsilon = 1/4$ if A is a set of real numbers. Very recently Chang [3] proved $\varepsilon = 1/54$ to finite sets of complex numbers. For further results and related problems we refer to [4, 5] and [1, 2].

In this note we prove Elekes' bound for complex numbers.

Theorem 1.1. *There is a positive absolute constant c , such that, for any finite sets of complex numbers A, B , and Q ,*

$$c|A|^{3/2}|B|^{1/2}|Q|^{1/2} \leq |A + B| \cdot |A \cdot Q|,$$

whence $c|A|^{5/4} \leq \max\{|A + A|, |A \cdot A|\}$.

Manuscrit reçu le 26 aout 2003.

This research was supported by NSERC and OTKA grants.

2. Proof

For the proof we need some simple observations and definitions. For each $a \in A$ let us find "the closest" element, an $a' \in A$ so that $a' \neq a$ and for any $a'' \in A$ if $|a - a'| > |a - a''|$ then $a = a''$. If there are more than one closest elements, then let us select any of them. This way we have $|A|$ ordered pairs, let us call them *neighboring pairs*.

Definition. We say that a quadruple (a, a', b, q) is *good* if (a, a') is a neighboring pair, $b \in B$ and $q \in Q$, moreover

$$|\{u \in A + B : |a + b - u| \leq |a - a'|\}| \leq \frac{28|A + B|}{|A|}$$

and

$$|\{v \in A \cdot Q : |aq - v| \leq |aq - a'q|\}| \leq \frac{28|A \cdot Q|}{|A|}.$$

When a quadruple (a, a', b, q) is good, then it means that the neighborhoods of $a + b$ and aq are not very dense in $A + B$ and in $A \cdot Q$.

Lemma 2.1. *For any $b \in B$ and $q \in Q$ the number of good quadruples (a, a', b, q) is at least $|A|/2$.*

Proof. Let us consider the set of disks around the elements of A with radius $|a - a'|$ (i.e. for every $a \in A$ we take the largest disk with center a , which contains no other elements of A in its interior). A simple geometric observation shows that no complex number is covered by more than 7 disks. Therefore

$$\sum_{a \in A} |\{u \in A + B : |a + b - u| \leq |a - a'|\}| \leq 7|A + B|$$

and

$$\sum_{a \in A} |\{v \in A \cdot Q : |aq - v| \leq |aq - a'q|\}| \leq 7|A \cdot Q|$$

providing that at least half of the neighboring pairs form good quadruples with b and q . Indeed, if we had more than a quarter of the neighboring pairs so that, say,

$$|\{v \in A \cdot Q : |aq - v| \leq |aq - a'q|\}| > \frac{28|A \cdot Q|}{|A|}$$

then it would imply

$$7|A \cdot Q| \geq \frac{|A|}{4} |\{v \in A \cdot Q : |aq - v| \leq |aq - a'q|\}| > 7|A \cdot Q|.$$

□

Proof of Theorem 1 To prove the theorem, we count the good quadruples (a, a', b, q) twice. For the sake of simplicity let us suppose that $0 \notin Q$. Such a quadruple is uniquely determined by the quadruple $(a + b, a' + b, aq, a'q)$. Now observe that there are $|A + B|$ possibilities for the first element, and given the value of $a + b$, the second element $a' + b$ must be one of the $28|A + B|/|A|$ nearest element of the sum-set $A + B$. We make the same argument for the third and fourth component to find that the number of such quadruples is at most

$$|A + B| \frac{28|A + B|}{|A|} |A \cdot Q| \frac{28|A \cdot Q|}{|A|}.$$

On the other hand, by Lemma 1 the number of such quadruples is at least

$$\frac{|A|}{2} |B| |Q|$$

that proves the theorem.

A similar argument works for quaternions and for other hypercomplex numbers. In general, if T and Q are sets of similarity transformations and A is a set of points in space such that from any quadruple $(t(p_1), t(p_2), q(p_1), q(p_2))$ the elements $t \in T$, $q \in Q$, and $p_1 \neq p_2 \in A$ are uniquely determined, then

$$c|A|^{3/2}|T|^{1/2}|Q|^{1/2} \leq |T(A)| \cdot |Q(A)|,$$

where c depends on the dimension of the space only.

References

- [1] J. BOURGAIN, S. KONJAGIN, *Estimates for the number of sums and products and for exponential sums over subgroups in finite fields of prime order*. C. R. Acad. Sci. Paris **337** (2003), no. 2, 75–80.
- [2] J. BOURGAIN, N. KATZ, T. TAO, *A sum-product estimate in finite fields, and applications*. Geometric And Functional Analysis GAFA **14** (2004), no. 1, 27–57.
- [3] M. CHANG, *A sum-product estimate in algebraic division algebras over \mathbb{R}* . Israel Journal of Mathematics (to appear).
- [4] M. CHANG, *Factorization in generalized arithmetic progressions and applications to the Erdős-Szemerédi sum-product problems*. Geometric And Functional Analysis GAFA **13** (2003), no. 4, 720–736.
- [5] M. CHANG, *Erdős-Szemerédi sum-product problem*. Annals of Math. **157** (2003), 939–957.
- [6] GY. ELEKES, *On the number of sums and products*. Acta Arithmetica **81** (1997), 365–367.
- [7] P. ERDŐS, E. SZEMERÉDI, *On sums and products of integers*. In: Studies in Pure Mathematics; To the memory of Paul Turán. P. Erdős, L. Alpár, and G. Halász, editors. Akadémiai Kiadó – Birkhauser Verlag, Budapest – Basel-Boston, Mass. 1983, 213–218.
- [8] K. FORD, *Sums and products from a finite set of real numbers*. Ramanujan Journal, **2** (1998), (1-2), 59–66.
- [9] M. B. NATHANSON, *On sums and products of integers*. Proc. Am. Math. Soc. **125** (1997), (1-2), 9–16.

József SOLYMOSI
Department of Mathematics,
University of British Columbia
1984 Mathematics Road, Vancouver,
Colombie-Britannique, Canada V6T 1Z2
E-mail : solymosi@math.ubc.ca