

RING-LIKE STRUCTURES DERIVED FROM λ -LATTICES
WITH ANTITONE INVOLUTIONS

IVAN CHAJDA, Olomouc

(Received December 12, 2005)

Abstract. Using the concept of the λ -lattice introduced recently by V. Snášel we define λ -lattices with antitone involutions. For them we establish a correspondence to ring-like structures similarly as it was done for ortholattices and pseudorings, for Boolean algebras and Boolean rings or for lattices with an antitone involution and the so-called Boolean quasirings.

Keywords: λ -lattice, λ -semilattice, ortholattice, λ -ortholattice, antitone involution, Boolean quasiring

MSC 2000: 06C15, 16Y99, 06B99, 81P10, 06A12

The well-known correspondence between Boolean algebras and Boolean rings (see e.g. [1]) was extended to orthomodular lattices by H. Länger [12] and to ortholattices by the author in [3]. A general setting was described by G. Eigenthaler, H. Länger and the author in [4] and [5]. It was generalized to lattices with antitone involution by D. Dorninger, H. Länger and M. Maćczyński [6], [7] and to generalized orthomodular lattices, see [5]. It was motivated by the use of these ring-like structures in certain logics of quantum mechanics, see e.g. [6], [7] for the description. However, in quantum mechanics it can happen that we cannot distinguish between two possibilities. It leads us to study more general structures than lattices where still ring-like structures can be induced.

Suitable tools for our investigation thus can be the so-called λ -lattices introduced by V. Snášel [13] and treated in [11], [13] and the so-called λ -semilattices known also under the name commutative directoids in [9]. On the other hand, this level of generality can make problems in computing formulas because some well-known

This work is supported by the Czech Government via the research project MSM 6198959214.

results for lattices fail for λ -lattices. For example, if $x \mapsto x'$ is an antitone involution on a lattice $\mathcal{L} = (L; \vee, \wedge)$ then the De Morgan laws

$$(DM) \quad (x \vee y)' = x' \wedge y' \quad \text{and} \quad (x \wedge y)' = x' \vee y'$$

hold in \mathcal{L} , see e.g. [1], [8]. However, this is not the case for λ -lattices. Hence we must often require some additional properties.

1. λ -LATTICES WITH ANTITONE INVOLUTIONS

First, we recall some well-known concepts. Let $(A; \leq)$ be an ordered set. For $a, b \in A$ denote

$$\begin{aligned} U(a, b) &= \{x \in A; a \leq x \text{ and } b \leq x\}, \\ L(a, b) &= \{x \in A; x \leq a \text{ and } x \leq b\}. \end{aligned}$$

An ordered set $(A; \leq)$ is called *up-directed* (*down-directed*) if $U(a, b) \neq \emptyset$ (or $L(a, b) \neq \emptyset$, respectively) for each $a, b \in A$. Further, $(A; \leq)$ is *directed* if it is both an up- and down-directed set.

Let $(A; \leq)$ be a down-directed set. Denote by $\text{Exp } A$ the power set of A . Let λ be a mapping $\lambda: \text{Exp } A \rightarrow A$ such that

- (i) $\lambda(L(a, b)) \in L(a, b)$,
- (ii) if $a \leq b$ then $\lambda(L(a, b)) = a$.

Define a binary operation \wedge on A as follows:

$$a \wedge b = \lambda(L(a, b)).$$

The groupoid $(A; \wedge)$ will be called a λ -*semilattice* (or a commutative directoid in [9]).

It is easy to verify that $(A; \wedge)$ satisfies the identities

- (I) $x \wedge x = x$ (idempotency),
- (C) $x \wedge y = y \wedge x$ (commutativity),
- (SA) $x \wedge ((x \wedge y) \wedge z) = (x \wedge y) \wedge z$ (skew associativity).

Also conversely, if $(A; \wedge)$ is a groupoid satisfying (I), (C), (SA) and \leq is defined by the rule

$$a \leq b \quad \text{if and only if} \quad a \wedge b = a$$

then $(A; \leq)$ is a down-directed set and $\lambda(L(a, b)) = a \wedge b$ satisfies (i) and (ii) mentioned above.

Moreover, if $(A; \leq)$ is a directed set and $\lambda: \text{Exp } A \rightarrow A$ satisfies also

- (iii) $\lambda(U(a, b)) \in U(a, b)$,
- (iv) if $a \leq b$ then $\lambda(U(a, b)) = b$

then we can introduce another operation \vee by setting

$$a \vee b = \lambda(U(a, b))$$

and easily verify the following identities:

$$(I') \quad x \vee x = x,$$

$$(C') \quad x \vee y = y \vee x,$$

$$(SA') \quad x \vee ((x \vee y) \vee z) = (x \vee y) \vee z,$$

$$(Ab) \quad x \vee (x \wedge y) = x, \quad x \wedge (x \vee y) = x \text{ (absorption)}.$$

Then the algebra $(A; \vee, \wedge)$ is called a λ -lattice (see [11], [13]). Also conversely, if $(A; \vee, \wedge)$ is an algebra of type $(2, 2)$ satisfying the identities (I), (I'), (C), (C'), (SA), (SA'), (Ab) then the relation defined by

$$a \leq b \quad \text{if and only if} \quad a \vee b = b$$

coincides with the already introduced induced order on $(A; \wedge)$. If we put

$$\lambda(U(a, b)) = a \vee b$$

then (iii) and (iv) are satisfied.

Hence, λ -semilattices and λ -lattices can be viewed either as algebras satisfying certain identities or as ordered sets with constraints on upper and lower bounds. Contrary to the case of lattices, the choice of $\lambda(L(a, b))$ or $\lambda(U(a, b))$ need not be unique. Consider e.g. the directed sets drawn in Fig. 1 and Fig. 2.




Fig. 1




Fig. 2

In the first case, we have three choices for $a \vee b$, namely c or d or 1 . Analogously, $c \wedge d$ can be a or b or 0 (for comparable elements the choice is unique due to the conditions (ii) and (iv)). In all 9 possible cases, the resulting algebra will be a λ -lattice. Analogously, in the second case (which is a lattice) we can pick up e.g. $a \vee b = c$ or $a \vee b = 1$. In the latter case, the resulting algebra will be a λ -lattice which is not a lattice.

Let $(A; \leq)$ be an ordered set. A mapping $x \mapsto x'$ on A is called an *antitone involution* if $x'' = x$ and $x \leq y$ implies $y' \leq x'$. It is well-known that if $(L; \vee, \wedge)$ is

a lattice and $x \mapsto x'$ is an antitone involution on $(L; \leq)$ where \leq is the induced order then the De Morgan laws (DM) hold.

Consider our λ -lattice $\mathcal{L} = (A; \vee, \wedge)$ visualized in Fig. 1. Define $x \mapsto x'$ on $A = \{0, a, b, c, d, 1\}$ by the table

$$\frac{x \quad \parallel \quad 0 \quad a \quad b \quad c \quad d \quad 1}{x' \quad \parallel \quad 1 \quad d \quad c \quad b \quad a \quad 0}.$$

Then clearly it is an antitone involution on A . Let us pick up e.g. $a \vee b = c$ (for other elements the operation \vee is determined as supremum). Then $(A; \vee)$ is a λ -semilattice (w.r.t. \vee). Now, if e.g. $c \wedge d = a$ (for other elements it is determined as infimum) then $\mathcal{L} = (A; \vee, \wedge)$ is a λ -lattice with an antitone involution but

$$b' \wedge a' = c \wedge d = a \neq b = c' = (b \vee a)',$$

thus in \mathcal{L} the De Morgan laws do not hold. On the contrary, when choosing $c \wedge d = b$, De Morgan laws hold in \mathcal{L} . Hence, validity of (DM) depends not only on the induced order but also on our choice of operations.

2. λ -BOOLEAN QUASIRINGS

By a *Boolean quasiring* (see [4], [6], [7]) we mean an algebra $\mathcal{R} = (R; +, \cdot, 0, 1)$ of type $(2, 2, 0, 0)$ satisfying the identities

- (R1) $x + y = y + x,$
- (R2) $x + 0 = x,$
- (R3) $(x \cdot y) \cdot z = x \cdot (y \cdot z),$
- (R4) $x \cdot y = y \cdot x,$
- (R5) $x \cdot x = x,$
- (R6) $x \cdot 0 = 0,$
- (R7) $x \cdot 1 = x,$
- (R8) $1 + (1 + x \cdot y) \cdot (1 + y) = y.$

For our purposes, we modify this definition as follows. An algebra $\mathcal{R} = (R; +, \cdot, 0, 1)$ of type $(2, 2, 0, 0)$ is called a *λ -Boolean quasiring* if it satisfies the identities (R1), (R2), (R4)–(R8) and

$$(R3^*) \quad x \cdot ((x \cdot y) \cdot z) = (x \cdot y) \cdot z.$$

One can immediately verify that every Boolean quasiring is a λ -Boolean quasiring since (R3*) follows easily by (R3), (R4) and (R5). The mutual correspondence between Boolean quasirings and lattices with an antitone involution was established in [6]. We are going to extend this correspondence to λ -Boolean quasirings.

Theorem 1. *Let $\mathcal{L} = (L; \vee, \wedge, ', 0, 1)$ be a bounded λ -lattice with an antitone involution. Define*

$$x + y = (x \vee y) \wedge (x \wedge y)' \quad \text{and} \quad x \cdot y = x \wedge y.$$

Then $\mathcal{R}(L) = (L; +, \cdot, 0, 1)$ is a λ -Boolean quasiring. If, moreover, the De Morgan laws hold in \mathcal{L} then $\mathcal{R}(L)$ satisfies the correspondence identity

$$\text{(Cor)} \quad (1 + (1 + x) \cdot (1 + y)) \cdot (1 + x \cdot y) = x + y.$$

Proof. Since $(L; \wedge)$ is a bounded λ -semilattice, the identities (R3*), (R4)–(R7) are immediate consequences of (I), (C), (SA) and the properties of the induced order. The identity (R1) is a trivial consequence of the definition and (R2) is evident. It remains to prove (R8) and (Cor).

For (R8) we use (Ab) to compute

$$1 + (1 + x \cdot y) \cdot (1 + y) = ((x \wedge y)' \wedge y')' = (y')' = y'' = y.$$

For (Cor), we apply De Morgan laws to derive

$$(1 + (1 + x) \cdot (1 + y)) \cdot (1 + x \cdot y) = (x' \wedge y')' \wedge (x \wedge y)' = (x \vee y) \wedge (x \wedge y)' = x + y.$$

□

We can prove also the converse.

Theorem 2. *Let $\mathcal{R} = (R; +, \cdot, 0, 1)$ be a λ -Boolean quasiring. Define*

$$x \wedge y = x \cdot y, \quad x' = 1 + x \quad \text{and} \quad x \vee y = 1 + (1 + x) \cdot (1 + y).$$

Then $\mathcal{L}(R) = (R; \vee, \wedge, ', 0, 1)$ is a bounded λ -lattice with an antitone involution in which the De Morgan laws hold.

Proof. By (R3*), (R4)–(R7), $(R; \wedge)$ is a bounded λ -semilattice. If we put $y = x$ in (R8) and apply (R5), we obtain the identity

$$(*) \quad 1 + (1 + x) = x$$

proving that the unary operation $x' = 1 + x$ is an involution on R . Suppose $x \leq y$. Then $x \wedge y = x \cdot y = x$ and, by (R8),

$$1 + x' \cdot y' = 1 + (1 + x) \cdot (1 + y) = 1 + (1 + x \cdot y) \cdot (1 + y) = y.$$

Thus, applying (*), we arrive at

$$x' \wedge y' = x' \cdot y' = 1 + (1 + x' \cdot y') = 1 + y = y'$$

whence $y' \leq x'$, i.e. this operation is an antitone involution on R .

Further, using (*), we obtain

$$x' \vee y' = 1 + (1 + x') \cdot (1 + y') = 1 + x \cdot y = (x \wedge y)'$$

and

$$x' \wedge y' = (1 + x) \cdot (1 + y) = 1 + (1 + (1 + x) \cdot (1 + y)) = (x \vee y)'$$

thus the De Morgan laws hold in $\mathcal{L}(R)$. Due to this fact, $(R; \vee)$ is also a λ -semilattice and, by (R8),

$$(x \wedge y) \vee y = ((x \cdot y)' \cdot y')' = 1 + (1 + x \cdot y) \cdot (1 + y) = y.$$

The dual absorption law can be established by the De Morgan laws and involutorness. Hence, $\mathcal{L}(R) = (R; \vee, \wedge, ', 0, 1)$ is a bounded λ -lattice with an antitone involution. \square

Theorem 3. *Let $\mathcal{R} = (R; +, \cdot, 0, 1)$ be a λ -Boolean quasiring satisfying the correspondence identity (Cor). Then $\mathcal{R}(\mathcal{L}(R)) = \mathcal{R}$.*

Let $\mathcal{L} = (L; \vee, \wedge, ', 0, 1)$ be a bounded λ -lattice with an antitone involution in which De Morgan laws hold. Then $\mathcal{L}(\mathcal{R}(L)) = \mathcal{L}$.

Proof. Evidently, the multiplicative operations coincide in $\mathcal{R}(\mathcal{L}(R))$ and \mathcal{R} . To prove $\mathcal{R}(\mathcal{L}(R)) = \mathcal{R}$ we need only to show that also $\oplus = +$ where \oplus is the additive operation in $\mathcal{R}(\mathcal{L}(R))$. Applying (Cor) we compute

$$x \oplus y = (x \vee y) \wedge (x \wedge y)' = (1 + (1 + x) \cdot (1 + y)) \cdot (1 + x \cdot y) = x + y.$$

Analogously, the operation meet clearly coincides in $\mathcal{L}(\mathcal{R}(L))$ and \mathcal{L} . Hence, it remains to prove $\sqcup = \vee$ and $x^* = x'$ where \sqcup is the join and $*$ is the antitone involution in $\mathcal{L}(\mathcal{R}(L))$. We have

$$x^* = 1 + x = (1 \vee x) \wedge (1 \wedge x)' = 1 \wedge x' = x'$$

and

$$x \sqcup y = 1 + (1 + x) \cdot (1 + y) = (x' \wedge y')' = x \vee y$$

due to the De Morgan laws. \square

3. λ -ORTHOLATTICES

By an *ortholattice* (see e.g. [2], [10]) we mean a bounded lattice $\mathcal{L} = (L; \vee, \wedge, ', 0, 1)$ with an antitone involution which is a complementation, i.e. $x \wedge x' = 0$ and $x \vee x' = 1$ for each $x \in L$. Of course, De Morgan laws hold in \mathcal{L} and hence $x \wedge x' = 0$ is equivalent to $x \vee x' = 1$.

We can extend this concept to λ -lattices.

By a λ -*ortholattice* we mean a bounded λ -lattice $\mathcal{L} = (L; \vee, \wedge, ', 0, 1)$ with an antitone involution such that $x \wedge x' = 0$ and $x \vee x' = 1$.

Formally, the definition is the same as for ortholattices, but we must be careful. For example, the λ -lattice depicted in Fig. 3 is an λ -ortholattice.




Fig. 3

Since it is a λ -lattice, we must specify joins and meets of non-comparable elements. If e.g. $a \vee b = c$ and $a' \wedge b' = d'$ then $a' \wedge b' = d' \neq c' = (a \vee b)'$ and hence the De Morgan laws do not hold. It means that for $a \vee b = c$ we must set $a' \wedge b' = c'$ etc.

The mutual correspondence between ortholattices and the so-called pseudosemirings was settled in [3]. However, it is easy to establish such a correspondence for λ -ortholattices and certain λ -Boolean quasirings.

Theorem 4. *Let $\mathcal{L} = (L; \vee, \wedge, ', 0, 1)$ be a λ -ortholattice in which the De Morgan laws hold. Define*

$$x + y = (x \vee y) \wedge (x \wedge y)' \quad \text{and} \quad x \cdot y = x \wedge y.$$

Then $\mathcal{R}(L) = (L; +, \cdot, 0, 1)$ is a λ -Boolean quasiring of characteristic 2 (i.e. satisfying the identity $x + x = 0$) satisfying the correspondence identity (Cor).

Let $\mathcal{R} = (R; +, \cdot, 0, 1)$ be a λ -Boolean quasiring of characteristic 2. Define

$$x \vee y = 1 + (1 + x) \cdot (1 + y), \quad x \wedge y = x \cdot y, \quad x' = 1 + x.$$

Then $\mathcal{L}(R) = (R; \vee, \wedge, ', 0, 1)$ is a λ -ortholattice in which the De Morgan laws hold.

Moreover, $\mathcal{L}(\mathcal{R}(L)) = \mathcal{L}$ and if \mathcal{R} satisfies (Cor) then also $\mathcal{R}(\mathcal{L}(R)) = \mathcal{R}$.

Proof. Of course, if $x \wedge x' = 0$ then $x + x = x \wedge x' = 0$, thus the induced λ -Boolean quasiring $\mathcal{R}(L)$ is of characteristic 2 for each λ -ortholattice \mathcal{L} . The rest of the proof follows by Theorem 1.

Conversely, if \mathcal{R} satisfies $x + x = 0$ then $x \wedge x' = (x \vee x) \wedge (x \wedge x)' = x + x = 0$ and, due to the De Morgan laws, also $x \vee x' = 1$ and thus $\mathcal{L}(R)$ is really a λ -ortholattice.

The remaining statements follow directly by Theorems 2 and 3. \square

In what follows we concentrate on a special case of a λ -ortholattice, the so-called *orthomodular λ -lattice* which satisfies the *orthomodular law*

$$(OML) \quad x \leq y \Rightarrow x \vee (y \wedge x') = y$$

which is equivalent to the identity

$$(**) \quad (x \wedge y) \vee (x \wedge (x \wedge y)') = x.$$

An example of an orthomodular λ -lattice can be constructed from the lattice drawn in Fig. 4.




Fig. 4

One can easily verify that \mathcal{L} is an orthomodular lattice. To change it into a λ -lattice which is not a lattice, we define $a \vee b = 1$ and $a' \wedge b' = 0$ and leave all other joins and meets as they were in \mathcal{L} . Of course, the new algebra is a λ -lattice but not a lattice, but the orthomodular law remains valid.

We are wonder what a λ -Boolean quasiring corresponds to an orthomodular λ -lattice.

Theorem 5. Let $\mathcal{L} = (L; \vee, \wedge, ', 0, 1)$ an orthomodular λ -lattice in which the De Morgan laws hold. Define

$$x + y = (x \vee y) \wedge (x \wedge y)' \quad \text{and} \quad x \cdot y = x \wedge y.$$

Then $\mathcal{R}(L)$ is a λ -Boolean quasiring of characteristic 2 satisfying the identity

$$(OM) \quad (1 + x \cdot y) + (x + x \cdot y) = 1 + x.$$

Let $\mathcal{R} = (R; +, \cdot, 0, 1)$ be a λ -Boolean quasiring of characteristic 2 satisfying the identity (OM). Define

$$x \vee y = 1 + (1 + x) \cdot (1 + y), \quad x \wedge y = x \cdot y \quad \text{and} \quad x' = 1 + x.$$

Then $\mathcal{L}(R) = (R; \vee, \wedge, ', 0, 1)$ is an orthomodular λ -lattice in which the De Morgan laws hold.

Moreover, $\mathcal{L}(\mathcal{R}(L)) = \mathcal{L}$ and if \mathcal{R} satisfies also (Cor) then $\mathcal{R}(\mathcal{L}(R)) = \mathcal{R}$.

Proof. In virtue of Theorem 4, we need only to verify the orthomodular law for $\mathcal{L}(R)$ and the identity (OM) for $\mathcal{R}(L)$.

Let \mathcal{L} be an orthomodular λ -lattice in which the De Morgan laws hold. We can easily derive $x' \vee (x' \vee y') = x' \vee y'$ since $x' \leq x' \vee y'$ and thus

$$\begin{aligned} x \cdot (1 + x \cdot y) &= x \wedge (x \wedge y)' = x \wedge (x' \vee y') = x \wedge (x' \vee (x' \vee y')) \\ &= x \wedge (x \wedge (x \wedge y))' = (x \vee (x \wedge y)) \wedge (x \wedge (x \wedge y))' = x + x \cdot y. \end{aligned}$$

Since $1 + x \cdot y = (x \wedge y)' \geq x \wedge (x \wedge y)' = x \cdot (1 + x \cdot y)$ and for $a \geq b$ we have

$$a + b = (a \vee b) \wedge (a \wedge b)' = a \wedge b' = a \cdot b',$$

then also $(1 + x \cdot y) + x \cdot (1 + x \cdot y) = (1 + x \cdot y) \cdot (1 + x \cdot (1 + x \cdot y))$. Now we apply (***) to compute

$$\begin{aligned} (1 + x \cdot y) + (x + x \cdot y) &= (1 + x \cdot y) + x \cdot (1 + x \cdot y) = (1 + x \cdot y) \cdot (1 + x \cdot (1 + x \cdot y)) \\ &= [(x \wedge y) \vee (x \wedge (x \wedge y))]' = x' = 1 + x. \end{aligned}$$

Thus $\mathcal{R}(L)$ satisfies (OM).

Conversely, let \mathcal{R} be a λ -Boolean quasiring of characteristic 2 satisfying (OM). Let $x \leq y$ in $\mathcal{L}(R)$. Then

$$\begin{aligned} x \vee (y \wedge x') &= 1 + (1 + x) \cdot (1 + y \cdot (1 + x)) \\ &= 1 + ((1 + x) + y \cdot (1 + x)) = 1 + ((1 + x \cdot y) + y \cdot (1 + x \cdot y)) \\ &= 1 + ((1 + x \cdot y) + (y + x \cdot y)) = 1 + (1 + y) = y \end{aligned}$$

(by (*) of the proof of Theorem 2). Hence, $\mathcal{L}(R)$ is an orthomodular λ -lattice. \square

References

- [1] *G. Birkhof*: Lattice Theory. (Third edition.) Amer. Math. Soc. Colloq. Publ. 25, Providence, RI, 1979. [Zbl 0505.06001](#)
- [2] *L. Beran*: Orthomodular Lattices. Reidel Publ., Dordrecht, 1985. [Zbl 0558.06008](#)
- [3] *I. Chajda*: Pseudorings induced by ortholattices. Czechoslovak Math. J. 46 (1996), 405–411. [Zbl 0879.06003](#)
- [4] *I. Chajda, G. Eigenthaler*: A note on orthopseudorings and Boolean quasirings. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II 207 (1998), 83–94. [Zbl 1040.06003](#)
- [5] *I. Chajda, H. Länger, M. Maćczyński*: Ring-like structures corresponding to generalized orthomodular lattices. Math. Slovaca 54 (2004), 143–150. [Zbl 1068.06008](#)
- [6] *D. Dorminger, H. Länger, M. Maćczyński*: The logic induced by a system of homomorphisms and its algebraic characterizations. Demonstratio Math. 30 (1997), 215–232. [Zbl 0879.06005](#)
- [7] *D. Dorminger, H. Länger, M. Maćczyński*: On ring-like structures occurring in axiomatic quantum mechanics. Österr. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II 206 (1997), 279–289. [Zbl 0945.03095](#)
- [8] *G. Grätzer*: General Lattice Theory. Birkhäuser, Basel, 1978. [Zbl 0385.06015](#)
- [9] *J. Ježek, R. Quackenbush*: Directoids: algebraic models of up-directed sets. Algebra Univers. 27 (1990), 49–69. [Zbl 0699.08002](#)
- [10] *G. Kalmbach*: Orthomodular Lattices. Academic Press, London, 1983. [Zbl 0512.06011](#)
- [11] *J. Karásek*: Rotations of λ -lattices. Math. Bohem. 121 (1996), 293–300. [Zbl 0879.06001](#)
- [12] *H. Länger*: Generalizations of the correspondence between Boolean algebras and Boolean rings to orthomodular lattices. Tatra Mt. Math. Publ. 15 (1998), 97–105. [Zbl 0939.03075](#)
- [13] *V. Snášel*: λ -Lattices. Math. Bohem. 122 (1997), 267–272. [Zbl 0897.06003](#)

Author's address: Ivan Chajda, Department of Algebra and Geometry, Palacký University Olomouc, Tomkova 40, 779 00 Olomouc, Czech Republic, e-mail: chajda@inf.upol.cz.