

FIXED-POINT MAPPINGS ON COMPACT METRIC SPACES

Ljubomir Ćirić

(Communicated August 9, 1978)

Let (M, d) be a metric space and T a selfmapping of M into itself. If

$$(1) \quad d(Tx, Ty) < d(x, y)$$

holds for every x, y in M with $x \neq y$, then T is called a contractive mapping. On complete metric spaces contractive mappings may be without fixed-point. However, if M is compact, then every contractive selfmapping on M has a unique fixed point.

D. Bailey in [1] has proved that if M is compact and T is continuous and such that for every $x, y \in M$ with $x \neq y$ there exists a positive integer $n(x, y)$ such that

$$(2) \quad d(T^{n(x,y)}x, T^{n(x,y)}y) < d(x, y),$$

then T has a unique fixed point in M .

In the following we will extend the result of Bailey to mappings which satisfy a contractive condition which is weaker than (2).

We now prove the following theorem.

THEOREM 1. *Let T be a continuous mapping on the compact metric space M into itself satisfying the inequality*

$$(3) \quad d(T^n x, T^n y) < \max \left\{ d(x, y), d(x, Tx), d(y, Ty), \frac{1}{2} [d(x, Ty) + d(y, Tx)] \right\}$$

for all x, y in M with $x \neq y$ where $n = n(x, y)$ is a positive integer. Then T has a unique fixed point in M .

Proof. Define on M a real-valued function F by $F(x) = d(x, Tx)$. Since T is continuous, it follows that F is continuous too. Therefore, F on M attains its maximum and minimum. Let u in M be such that

$$(4) \quad F(u) = \min\{F(x) : x \in M\}.$$

We will show that u is a fixed-point of T . If we assume that $F(u) = d(u, Tu) > 0$, then by (3), for $n = n(u, Tu)$, we have

$$\begin{aligned} F(T^n u) &= d(T^n u, TT^n u) = d(T^n u, T^n Tu) \\ &< \max\left\{d(u, Tu), d(u, Tu), d(Tu, TTu), \frac{1}{2}[d(u, T^2 u) + 0]\right\} \\ &\leq \max\left\{F(u), F(Tu), \frac{1}{2}[F(u) + F(Tu)]\right\}. \end{aligned}$$

Since by (4) $\max\{F(u), F(Tu), \frac{1}{2}[F(u) + F(Tu)]\} = F(u)$, we have $F(T^n u) < F(u)$, which is a contradiction with (4). Therefore, u is a fixed-point of T . The uniqueness of u follows easily from (3). This completes the proof of the theorem.

Theorem 1 holds if some of the conditions are relaxed. So we have

THEOREM 2. *Let T be an orbitally continuous mapping on the compact metric space M into itself satisfying (3). Then T has a unique fixed-point in M .*

The following example shows that the continuity conditions of T in the theorems 1. and 2. cannot be removed.

Example. If M is the closed interval $[0, 1]$ and $T : M \rightarrow M$ is defined by $T(x) = \frac{x}{2}$ if $x \neq 0$ and $T(0) = 1$, then T satisfies (3) with $n(x, y) = 2$, as $d(T^2 0, T^2 x) = \frac{1}{2} - \frac{x}{4} < \frac{1}{2} \cdot 1 = \frac{1}{2}d(0, T0)$ and $d(T^2 y, T^2 x) \frac{1}{4}d(y, x)$ for $y \neq 0$. However, T has not fixed points.

Now we will show that the condition (3) may be much more weakened.

THEOREM 3. *Let T be a continuous mapping on the compact metric space M into itself satisfying the inequality*

$$(5) \quad d(T^n x, T^n y) < \max_{0 \leq p, q, r, s, t \leq n} \left\{ d(T^p x, T^p y), d(T^q x, T^{q+1} x), \right. \\ \left. d(T^r y, T^{r+1} y), \frac{1}{2}[d(T^s x, T^{s+1} y) + d(T^t y, T^{t+1} x)] \right\}$$

for some positive integer $n = n(x, y)$ and x, y in M for which the right-hand side of inequality is positive. Then T has a unique fixed-point in M .

Proof. We may assume that the right-hand side of inequality (5) is positive for each x, y in M . For if it is not positive, then $x = y = Tx$, which means that T has a fixed-point. Let the mapping F and the point u be defined as in the proof of the theorem 1. and let $n = n(u, Tu)$. Then by (5) it follows

$$\begin{aligned} F(T^n u) &= d(T^n u, T^n Tu) < \max_{0 \leq p, q, r, s, t \leq n} \left\{ d(T^p u, TT^p u), d(T^q u, TT^q u), \right. \\ &\quad \left. d(T^{r+1} u, TT^{r+1} u), \frac{1}{2}[d(T^s u, T^2 T^s u) + 0] \right\}. \end{aligned}$$

Using the triangle inequality and (4) we obtain $F(T^n) < F(u)$, which is a contradiction with (4). Therefore, the right-hand side of (5) is zero for $x = u$ and $y = Tu$. Hence $Tu = u$. The uniqueness of the fixed-point follows easily. This completes the proof of the theorem.

REFERENCES

- [1] D. Bailey, *Some theorems on contractive mappings*, J. London Math. Soc., **41** (1966), 101–106.
- [2] Lj. Ćirić, *Generalized contractions and fixed-point theorems*, Publ. Inst. Math., **12(26)**, 1971, 19–26.
- [3] T. Pal, M. Maiti, I. Achari, *Extensions of Ćirić's generalized contradictions*, Mat. vesnik, **13 (28)**, 1976, 313–317.

Mašinski fakultet
11000 Beograd