

ON REGULAR RINGS AND SELF-INJECTIVE RINGS, IV

Roger Yu Chi Ming

Abstract. This paper is essentially concerned with f -injectivity (a generalization of injectivity) and an analogous concept which generalizes projectivity, called F -projectivity.

Introduction. Throughout, A represents an associative ring with identity and A -modules are unital. J and Z denote respectively the Jacobson radical and the left singular ideal of A . ${}_A M$ is called f -injective (resp. p -injective) if, for any finitely generated (resp. principal) left ideal I of A , every left A -homomorphism of I into M extends to A . Then A is von Neumann regular iff every left A -module is flat iff every left A -module is f -injective (p -injective). Right f -injective (resp. p -injective) modules are similarly defined. Flatness and f -injectivity are distinct concepts. However, if I is a left ideal of A , then ${}_A I$ f -injective implies ${}_A A/I$ flat. A is called left (resp. right) f -injective if ${}_A A$ (resp. A_A) is f -injective.

Injective and projective modules, extensively studied in recent years, are fundamental concepts in ring theory (cf. [2, 3, 4, 8]). Our first result gives a condition for a finitely generated left ideal of a semi-prime left f -injective ring to be generated by a central idempotent. Self-injective regular rings, biregular rings are next considered. Quasi-Frobeniusean rings are characterised in terms of F -projectivity and f -injectivity. If A has a classical left quotient ring Q such that every divisible torsionfree quasi-injective left A -module is an F -projective left Q -module, then Q is semi-simple Artinian. A sufficient condition is given for a classical quotient ring to be Noetherian. Left duo rings whose divisible left modules are f -injective are characterized.

As usual, an ideal of A means a two-sided ideal and A is called left duo (after E. H. Feller) if every left ideal of A is an ideal. A left (right) ideal is called reduced if it contains no non-zero nilpotent element. The concepts of f -injectivity and p -injectivity have been studied by various authors (cf. for example, [1, 11, 12, 13, 14, 15]) and in connection with semigroup and torsion theories, consult [5, 6, 10, 16]. In semigroup theory, f -injectivity (p -injectivity) is also called weak

f -injectivity (weak p -injectivity) (cf. [5,6]). Note that if A is left f -injective, then $J = Z$ (cf. [17]).

The rings considered in the following proposition need not be von Neumann regular (cf. for example, [11]).

PROPOSITION 1. *Let A be a semi-prime left f -injective ring. The following conditions are then equivalent for a finitely generated left ideal F :*

- (1) F is generated by a central idempotent;
- (2) $F = l(T)$, where T is a finitely generated left ideal.

Proof. If $F = Ae$ where e is a central idempotent, then $F = l(u)$ where $u = l - e$ is central and hence $F = l(AuA)$, where $AuA = Au$. This shows that (1) implies (2).

Assume (2). Then $F = l(T)$, where T is an ideal of A , ${}_A T$ is finitely generated. Therefore F is an ideal of A . By [7, Theorem 1], $r(F \cap T) = r(F) + r(T)$ and since A is semi-prime, $A = r(o) = r(l(T) \cap T) = r(F \cap T) = r(F) + r(T)$. Now $F = l(T) = r(T)$ and therefore $A = F + r(F)$. Since $F \cap r(F) = F \cap l(F) = 0$, then $A = F \oplus r(F)$. It follows from the semi-primeness of A that F is generated by a central idempotent.

COROLLARY 1.1. *The following conditions are equivalent for a left- f -injective ring A :*

- (1) A is biregular;
- (2) A is semi-prime such that for each $a \in A$, AaA is the principal left annihilator of AbA for some $b \in A$.

The proof of Proposition 1 yields.

PROPOSITION 2. *Let A be a semi prime left self-injective ring. If T is an ideal of A which is an annihilator, then T is generated by a central idempotent. Consequently, any finitely generated right ideal which is an ideal of A is generated by a central idempotent.*

Combining the results above with [19, Theorem (DL)], we get

PROPOSITION 3. *If A is left self-injective, the following conditions are equivalent:*

- (1) A is regular and biregular;
- (2) A is semi-prime such that for any $a \in A$, AaA is the principal left annihilator of AbA for some $b \in A$;
- (3) A is semi-prime such that for any $a \in A$, AaA is a principal right ideal of A ;
- (4) If $a, b \in A$ such that $AaA + AbA \neq A$, then $AaA + AbA$ is the left annihilator of a non-nilpotent central element of A .

Proposition 3 and Corollary 1.1. yield

PROPOSITION 4. *The following conditions are equivalent for a left self-injective ring A :*

- (1) *A is regular such that every ideal is generated by a central idempotent;*
- (2) *A is right non-singular such that every ideal is a right annihilator ideal;*
- (3) *A is semi-prime such that every ideal is a finitely generated right ideal of A .*

Question. If A is semi-prime, T an ideal of A which is a left annihilator, is T a complement left ideal of A ? (In case A is prime, the answer is positive.)

As an analog of f -injectivity, we introduce F -projectivity.

Definition. A left A -module P is called F -projective if, given any finitely generated left A -modules M, N with an epimorphism $g : M \rightarrow N$ and any left A -homomorphism $f : P \rightarrow N$, there exists a left A -homomorphism $h : P \rightarrow M$ such that $gh = f$.

It may be noted that a direct sum of left A -modules is F -projective if, and only if, each direct summand is F -projective.

PROPOSITION 5. *If P is a finitely generated F -projective left A -module, then ${}_A P$ is projective.*

Proof. Let $P = \sum_{i=1}^n Au_i$ be a F -projective left A -module. If M, N are left A -modules, $p : {}_A M \rightarrow {}_A N$ an epimorphism, $f : P \rightarrow N$ a left A -homomorphism, set $f(u_i) = v_i$ for each i , $1 \leq i \leq n$. Since p is an epimorphism, there exists $w_i \in M$ such that $p(w_i) = v_i$, $1 \leq i \leq n$. If $M' = \sum_{i=1}^n Aw_i$, $N' = \sum_{i=1}^n Av_i$, then p' , the restriction of p to M' , is an epimorphism of M' onto N' . Since ${}_A P$ is F -projective, there exists a left A -homomorphism $h' : P \rightarrow M'$ such that $p'h' = f$. If j is the inclusion map $M' \rightarrow M$, $h = jh'$, for any $y \in P$, $y = \sum_{i=1}^n a_i u_i + i$, $ph(y) = p(\sum_{i=1}^n a_i h(u_i)) = p(\sum_{i=1}^n a_i h'(u_i)) = p'(\sum_{i=1}^n a_i h'(u_i)) = \sum_{i=1}^n a_i p'h'(u_i) = \sum_{i=1}^n a_i f(u_i) = f(y)$, which proves that $ph = f$. The proposition then follows.

COROLLARY 5.1. *If A is commutative, then A is quasi-Frobeniusean iff A is Artinian such that every injective A -module is F -projective.*

Proof. Assume that A is Artinian and every injective A -module is F -projective. If M is injective, then M is a direct sum of finitely generated A -modules M_i . Since M is F -projective, then each M_i is F -projective which implies that M_i is projective, whence M is a projective A -module. The corollary then follows from [3, Theorem 24.20].

If A is left perfect, left f -injective, then every simple left A -module is a homomorphic image of an injective left A -module (cf. [2, P. 481]). If \times is right f -injective satisfying the maximum condition on left annihilator ideals, then A is quasi-Frobeniusean [18]. Recall that (1) A is a left Kasch ring if every maximal left ideal of A is a left annihilator; (2) A is right pseudo-coherent iff the right annihilator of every finitely generated left ideal is a finitely generated right ideal; (3) A left

A -module C is a cogenerator if, for any M in the category of left A -modules, there exists a monomorphism of M into a direct product of copies of C . A is left pseudo-Frobeniusean iff ${}_A A$ is an injective cogenerator iff A is a left self-injective left Kasch ring. Quasi-Frobeniusean rings are both left and right pseudo-Frobeniusean.

A is called left (resp. right) hypercyclic if the injective hull of every cyclic left (resp. right) A -module is cyclic.

Remark 1. The following conditions are equivalent: (1) Every factor ring of A is quasi-Frobeniusean; (2) A is a left and right hypercyclic principal left ideal ring such that every injective one-sided A -module is F -projective. (cf. [3, Proposition 25.4. 6 B]).

If A is von Neumann regular, a theorem of I. Kaplansky asserts that for any projective left A -module P , every finitely generated left submodule is a direct summand. Consequently, the following remark holds.

Remark 2. If A is von Neumann regular, then every F -projective left A -module is projective.

For results on coherent rings, consult [11].

THEOREM 6. *The following conditions are equivalent:*

- (1) A is quasi-Frobeniusean;
- (2) A is coherent left f -injective left cogenerating such that every flat left A -module is F -projective;
- (3) A is left cogenerating such that every injective left A -module is F -projective;
- (4) A is right f -injective left perfect right pseudo-coherent;
- (5) A is left f -injective left perfect right pseudo-coherent.

Proof. Obviously, (1) implies (2).

Assume (2). Since A is left cogenerating, then every left ideal is a left annihilator. Since A is left f -injective, then every finitely generated right ideal is a right annihilator. By [11, Theorem 2.1], every injective left A -module is flat. Therefore (2) implies (3).

Assume (3). Since ${}_A A$ is a cogenerator, for any non-zero left A -module M , any $o \neq y \in M$, there exist a left A -homomorphism $g : M \rightarrow A$ such that $g(y) \neq o$. Now if ${}_A P$ is F -projective, for any left A -module M with an epimorphism $p : M \rightarrow P$, since there exists a non-zero left A -homomorphism $g : M \rightarrow A$, we can show that p splits. But then, we get ${}_A P$ projective. Consequently, every injective left A -module is projective and by [3, Theorem 24.20], A is quasi-Frobeniusean. Therefore (3) implies (4).

Assume (4). Let $I_1 \subseteq I_2 \subseteq \dots \subseteq I_n \subseteq \dots$ be an ascending chain of finitely generated left ideals of A . If $R_i = r(I_i)$ for each i , since A is right pseudo-coherent, then R_i is a finitely generated right ideal for each i . Since A is left perfect,

then $R_1 \supseteq R_2 \supseteq R_3 \dots \supseteq R_n \supseteq \dots$ yields $R_m = R_s$ for some positive integer m and all $s \geq m$ [8, P. 303]. Now A right f -injective implies that every finitely generated left ideal is a left annihilator, whence $I_m = 1(r(I_m)) = 1(R_m) = 1(R_s) = 1(r(I_s)) = I_s$ for all $s \geq m$. This proves that A is left Noetherian. Then (4) implies (5) by [18, Theorem 2].

Assume (5). Let R be a right annihilator ideal, U a subset of A such that $R = r(U)$. Since A is right pseudo-coherent, the right annihilator for any finitely generated left ideal is finitely generated. Since A is left perfect, there exists a finite subset F of U such that $R = r(F)$ which implies that R must be a finitely generated right ideal. Therefore A satisfies the minimum condition on right annihilators which implies that A satisfies the maximum condition on left annihilators. Then Z is nilpotent and since A is left f -injective, $Z = J$ [17, Proposition 3] which implies that A is semi-primary. Therefore A is a left f -injective semi-primary ring which implies that A is left Kash. Now let F be a finitely generated left ideal of A . Suppose that $F \subset 1(r(F))$. If $y \in 1(r(F))$, $y \notin F$, $G = F + Ay$, the set E of left ideals I of A such that $F \subseteq I \subset G$ is an inductive set and by Zorn Lemma, E has a maximal member K . Then G/K is a simple left A -module and since A is left Kash, $G/K = Au$, where Au is a minimal left ideal of A . There exists a left A -homomorphism $G \rightarrow Au$ which yields a non-zero left A -homomorphism $f : G \rightarrow A$ such that $f(K) = o$. Since G is a finitely generated left ideal of A , there exists $c \in A$ such that $f(b) = bc$ for all $b \in G$ (A being left f -injective). Then $f(F) = o$ which implies $c \in r(F)$, whence $f(y) = yc = o$. Thus $f(G) = o$ which contradicts f non-zero. This proves that F must be a left annihilator. Then A satisfies the ascending chain condition on finitely generated left ideals which implies that A is left Noetherian. Therefore A is left self-injective, left Noetherian and hence (5) implies (1).

We now consider classical quotient rings in terms of F -projectivity and f -injectivity. For the definition and properties of classical quotient rings, consult, for example, [4] and [9]. For divisibility and torsionfreeness, see [9].

PROPOSITION 7. *Let A have a classical left quotient ring Q . The following conditions are then equivalent:*

- (1) Q is semi-simple Artinian;
- (2) Every divisible torsionfree quasi-injective left A -module is an F -projective left Q -module.

Proof. Since any divisible torsionfree left A -module is left Q -module [9, p. 140], (1) implies (2).

Assume (2). Let P be a quasi-injective left Q -module. Since any left Q -module is a divisible torsionfree left A -module, then ${}_A P$ is divisible torsionfree. If ${}_A N$ is a submodule of ${}_A P$, $f : N \rightarrow P$ a left A -homomorphism, since ${}_A N$ is torsionfree, we may define a left Q -homomorphism $g : {}_Q N \rightarrow P$ by $g(c^{-1}w) = c^{-1}f(w)$, for any non-zero divisor c of A , $w \in N$. Since ${}_Q P$ is quasi-injective, g extends to an endomorphism of ${}_Q P$ and so f extends to an endomorphism of ${}_A P$. This shows that ${}_A P$ is quasi-injective and by hypothesis, ${}_Q P$ is F -projective.

Therefore every simple left Q -module, being quasi-injective, is F -projective and by Proposition 5, every simple left Q -module is projective. This proves that (2) implies (1).

PROPOSITION 8. *Let A have a classical left quotient ring Q . If every divisible torsionfree f -injective left A -module is injective, then Q is left Noetherian.*

Proof. Let M be an f -injective left Q -module, $F = \sum_{i=1}^n Au_i$ a finitely generated left ideal of A , $g : {}_A F \rightarrow {}_A M$ a left A -homomorphism. Define $G : \sum_{i=1}^n Qu_i \rightarrow M$ by $G(\sum_{i=1}^n q_i u_i) = \sum_{i=1}^n q_i (g(u_i))$ $q_i \in Q$, $1 \leq i \leq n$. Then G is a well-defined left Q -homomorphism and so there exists $y \in M$ such that $G(u_i) = u_i y$ for each i , $1 \leq i \leq n$. Consequently, for any $v \in F$, $v = \sum_{i=1}^n a_i u_i$, $a_i \in A$, $g(v) = \sum_{i=1}^n a_i g(u_i) = \sum_{i=1}^n a_i G(u_i) = \sum_{i=1}^n a_i u_i y = (\sum_{i=1}^n a_i u_i) y = vy$ which proves that ${}_A M$ is f -injective. Since ${}_A M$ is divisible torsionfree, then ${}_A M$ is injective. If $j : {}_Q P \rightarrow {}_Q N$ is a monomorphism, $f : {}_Q P \rightarrow {}_Q M$ a left Q -homomorphism, \hat{j}, \hat{f} the restrictions of j, f respectively to ${}_A P$, since ${}_A M$ is injective; there exists a left A -homomorphism $\hat{h} : {}_A N \rightarrow {}_A M$ such that $\hat{h}\hat{j} = \hat{f}$. If $h : {}_Q N \rightarrow {}_Q M$ is defined by $h(qu) = gh(u)$ for all $q \in Q$, $u \in N$, then $hj = f$ which proves that ${}_Q M$ is injective. Since any direct sum of f -injective left Q -modules is f -injective, then any direct sum of injective left Q -modules is injective which implies that Q is left Noetherian [3, Theorem 20.1].

We now turn to rings whose divisible modules are f -injective.

PROPOSITION 9. *If every divisible left A -module is f -injective, then A is left semi-hereditary.*

Proof. Since any injective left A -module is divisible and any quotient module of a divisible module is divisible, then the quotient of any injective left A -module is f -injective. Let I be a finitely generated left ideal of A . Given any quotient module M/N of a left A -module M , $k : M \rightarrow M/N$ the natural projection, if $f : I \rightarrow M/N$ is a left A -homomorphism, E the injective hull of ${}_A M$, $j : M/N \rightarrow E/N$ the inclusion map, then with $F = jf$, we have $F : I \rightarrow E/N$. Therefore ${}_A E/N$ is f -injective which implies the existence of $\bar{z} = z + N$, $z \in E$, such that $F(b) = b\bar{z}$ for all $b \in I$. Define $g : I \rightarrow E$ by $g(b) = bz$ for all $b \in I$. If $K : E \rightarrow E/N$ is the natural projection, for all $b \in I$, $Kg(b) = K(bz) = bK(z) = b\bar{z} = F(b)$, whence $F = Kg$. As $f(I) \subseteq M/N$, then $F(I) \subseteq M/N$, $g(I) \subseteq M$ and if define $h : I \rightarrow M$ by $h(b) = g(b)$ for all $b \in I$, we get $kh = f$. This proves that ${}_A I$ is projective which yields the proposition.

The next result gives a characterization of commutative rings whose divisible modules are f -injective. It is clear that over von Neumann regular rings, p -injectivity coincides with f -injectivity.

THEOREM 10. *The following conditions are equivalent for a ring A whose complement left ideals are ideals:*

- (1) *Every divisible left A -module is f -injective;*
- (2) *A is left semi-hereditary and every p -injective left A -module is f -injective;*

(3) Every p -injective left A -module is f -injective and for every $a \in A$, there exist an idempotent $e \in 1(a)$ and n elements b_1, \dots, b_n in Aa , n non zero-divisors c_1, \dots, c_n in A such that $a = \sum_{i=1}^n a_i b_i$, $a_i \in A$, $(1-e)a_i c_i = a$ for each i , $1 \leq i \leq n$.

Proof. Since any p -injective left A -module is divisible, then (1) implies (2) by Proposition 9.

Assume (2). Let D be a divisible left A -module. For any $a \in A$, since ${}_A Aa$ is projective, $1(a) = Ae$, $e = e^2 \in A$. Since A is left non-singular and every complement left ideal is an ideal of A , then A is reduced which implies that e is a central idempotent. With $c = a + e$, since $Aa \cap 1(a) = o$, it is easy to see that c is a non zero-divisor of A . If $f : Aa \rightarrow D$ is a left A -homomorphism, set $f(a) = d \in D$. Then $d = cv$ for some $v \in D$ (inasmuch as ${}_A D$ is divisible) and $a = (1-e)a$ implies $f(a) = f((1-e)a) = (1-e)f(a) = (1-e)cv = (1-e)(a+e)v = av$ which shows that ${}_A D$ is p -injective. By hypothesis, ${}_A D$ is f -injective. Thus (2) implies (3) (it is sufficient to take $n = 1$, $b_1 = a$, $c_1 = c$, $a_1 = 1$).

Assume (3). Let M be a divisible left A -module, $a \in A$, $f : Aa \rightarrow M$ a left A -homomorphism. By hypothesis, there exist a positive integer n , $b_1, \dots, b_n \in Aa$, an idempotent e such that $ea = o$, non zero-divisors $c_1, \dots, c_n \in A$ such that $a = \sum_{i=1}^n a_i b_i$, $a_i \in A$, $(1-e)a_i c_i = a$ for each i , $1 \leq i \leq n$. Since ${}_A M$ is divisible, for each i , $1 \leq i \leq n$, $f(b_i) = c_i m_i$, $m_i \in M$ and $f(a) = f((1-e)a) = (1-e)f(a) = (1-e)f(\sum_{i=1}^n a_i b_i) = (1-e)(\sum_{i=1}^n a_i f(b_i)) = (1-e)(\sum_{i=1}^n a_i c_i m_i) = a(\sum_{i=1}^n m_i)$ which proves that ${}_A M$ is p -injective. By hypothesis, every divisible left A -module is f -injective and hence (3) implies (1).

COROLLARY 10.1. *If every complement left ideal of A is an ideal, the following are then equivalent: (1) Every divisible left A -module is injective; (2) A is a left hereditary, left Noetherian ring whose p -injective left modules are f -injective.*

COROLLARY 10.2. *A left duo ring whose divisible left modules are injective admits a classical left quotient ring which is a finite direct sum of division rings.*

The next remark completes nicely Theorem 4 of [20].

Remark 3. The following conditions are equivalent: (a) A is left self-injective regular with non-zero socle; (b) A is left f -injective with an injective non-singular maximal left ideal; (c) A has an injective non-singular maximal left ideal and the left socle of A is f -injective; (d) A is semi-prime with an injective non-singular, maximal left ideal such that every maximal right ideal is f -injective.

Remark 4. A is strongly regular iff A is a left f -injective ring with a non-singular maximal left ideal such that every maximal left ideal is an ideal of A .

Remark 5. Let A be left f -injective. If every finitely generated faithful left A -module is F -projective, then A is left self-injective and consequently, every finitely generated faithful left A -module is injective.

Remark 6. (1) A is von Neumann regular iff every divisible left A -module is flat; (2) the following conditions are equivalent: (a) A is semi-simple Artinian; (b) every divisible left A -module is projective; (c) every left A -module is F -projective.

REFERENCES

- [1] G. Baccela, *Generalized V -rings and von Neumann regular rings*, Rend. Sem. Mat. Univ. Padova **72** (1984), 117–133.
- [2] H. Bass, *Finitistic dimension and a homological generalization of semi-primary rings*, Trans. Amer. Math. Soc. **95** (1960), 466–488.
- [3] C. Faith, *Algebra II: Ring Theory*, Springer-Verlag, **191** (1976).
- [4] K. R. Goddard, *Ring Theory, Nonsingular rings and modules*, Pure Appl. Math. **33** (1976).
- [5] V. Gould, *The characterization of monoids by properties of their S -systems*, Semigroup Forum **32** (1985), 251–265.
- [6] V. Gould, *Divisible S -systems and R -modules*, Proc. Edinburg Math. Soc. **30** (1987), 187–200.
- [7] M. Ikeda and T. Nakayama, *On some characteristic properties of quasi-Frobenius and regular rings*, Proc. Amer. Math. Soc. **5** (1954), 15–19.
- [8] F. Kasch, *Modules and Rings* London Math. Soc., 1982.
- [9] L. Levy, *Torsion free and divisible modules over non-integral domains*, Canad. J. Math. **15** (1963), 132–151.
- [10] J. K. Luedeman, F. R. McMorris, and S. K. Sim, *Semi groups for which every totally irreducible S -system is injective*, Comment. Math. Univ. Carolinae **19** (1978), 27–35.
- [11] J. L. G. Pardo and N. R. Gonzales, *On some properties of IF rings*, Quaestiones Math. **5** (1983), 395–405.
- [12] Sii-Joo Kim, *A note on generalizations of PCI rings*, Honam Math. **5** (1983), 205–214.
- [13] A. K. Tiwary, S. A. Paramhans and B. M. Pandey, *Generalizations of quasi-injectivity*, Progress Math. **13** (1979), 31–40.
- [14] H. Tominaga, *On s -unital rings*, Maths. J. Okayama Univ. **18** (1976), 117–134.
- [15] H. Tominaga, *On s -unital rings, II*, Math. J. Okayama Univ. **19** (1977), 171–182.
- [16] K. Varadarajan and K. Wehrhahn, *P -injectivity of simple pre-torsion modules*, Glasgow Math. J. **28** (1986), 223–225.
- [17] R. Yue Chi Ming, *On von Neumann regular rings, VIII*, J. Korean Math. Soc. **19** (1983), 97–104.
- [18] R. Yue Chi Ming, *On quasi-Frobeniusean and Artinian rings*, Publ. Inst. Math. (Beograd) **33** (47) (1983), 239–245.
- [19] R. Yue Chi Ming, *On regular rings and self-injective rings, II*, Glasnik Mat. **18** (38) (1983), 221–229.
- [20] R. Yue Chi Ming, *On regular rings and self-injective rings, III*, Tamkang J. Math. **17** (1986), n°3, 59–67.

Universite Paris VII
 U. F. R. de Mathematiques, URA 212 CNRS
 2, Place Jussieu
 75251 Paris Cedex 05
 France

(Received 07 09 1987)