

**THE CATEGORY
OF COMPACT METRIC SPACES
AND ITS FUNCTIONAL ANALYTIC DUALS**

Branka Pavlović

ABSTRACT. A Lipschitz algebra $\text{Lip}(X, d_X)$ over a compact metric space (X, d_X) consists of all complex valued continuous functions on (X, d_X) which are Lipschitz with respect to d_X and the standard metric on the complex plane \mathbb{C} (absolute value). The norm on $\text{Lip}(X, d_X)$ is given by $\|f\| = \sup\{|f(x)| : x \in X\} + \sup\{|f(x) - f(y)|/d_X(x, y) : x, y \in X \text{ \& } x \neq y\}$. We show that the category CLip in which objects are Lipschitz algebras and morphisms are algebra homomorphisms is dual to the category CMet in which objects are compact metric spaces and morphisms are Lipschitz maps. Let (X, d) be any metric space, and let $Y = \{(x, y) \in X \times X : x \neq y\}$. De Leeuw derivation defined by the metric d is the operator $D : C_b(X) \rightarrow C_b(Y)$ be defined by $(Df)(x, y) = (f(y) - f(x))/d(x, y)$ for $(x, y) \in Y$. We consider the category CDer in which objects are pairs $(C(X), D_X)$, where (X, d_X) is a compact metric space and D_X is the corresponding de Leeuw derivation, and morphisms are all homomorphisms $\nu : C(X) \rightarrow C(Y)$ for which $f \in \text{Domain}(D_X)$ implies $\nu f \in \text{Domain}(D_Y)$. We show that CDer is equivalent to CLip , and that CDer is dual to CMet .

1. Introduction and definitions

It is well known that the following two categories are dual: (1) the category in which objects are compact Hausdorff spaces and morphisms are continuous maps; (2) the category in which objects are commutative unital C^* -algebras and morphisms are homomorphisms. Recent work in noncommutative geometry ([2, 7]) has prompted a search for functional analytic representation of metric spaces and

2000 *Mathematics Subject Classification*. Primary 18B99; Secondary 18B30, 46J10, 46L89, 46M15.

Key words and phrases. Lipschitz algebras; de Leeuw derivations; dual and equivalent categories.

Work on this paper began while the author was a Post-Doctoral Fellow at the University of Victoria, Canada, with Professors John Phillips and Ian Putnam, where the research was supported by their NSERC grants.

This research was supported in part by the Ministry of Science and Technology of Serbia, through the Institute of Mathematics of the Serbian Academy of Sciences and Arts, project # 04M03.

for a functional analytic characterization of their geometric properties. The goal of this paper is to show that similar to how topological spaces have their functional analytic counterpart in C^* -algebras, metric spaces have their functional analytic counterpart in Lipschitz algebras. We also want to show that just as there is a connection between a metric and the underlying topological space (if X is a compact Hausdorff space, metric is a continuous function on $X \times X$ satisfying certain conditions), there likewise exists a connection between a Lipschitz algebra and the underlying C^* -algebra A via the de Leeuw derivation on A .

DEFINITION 1.1. A map $f : X \rightarrow Y$ from a metric space (X, d_X) to a metric space (Y, d_Y) is said to be *Lipschitz* if there exists a constant M such that for all x, y in X

$$d_Y(f(x), f(y)) \leq M d_X(x, y).$$

The smallest such constant is called the *Lipschitz constant* of f .

The Lipschitz constant of f , $p(f)$, can be expressed explicitly as

$$\begin{aligned} p(f) &= \inf\{M > 0 : d_Y(f(x), f(y)) \leq M d_X(x, y) \quad \forall x, y \in X\} \\ &= \sup\{d_Y(f(x), f(y))/d_X(x, y) : x, y \in X \text{ \& } x \neq y\}. \end{aligned}$$

When Y is a normed space $p(f)$ is also called the *Lipschitz norm* of f (it is in fact a semi-norm).

If (X, d_X) , (Y, d_Y) , and (Z, d_Z) are metric spaces, and $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are Lipschitz maps, then $g \circ f$ is Lipschitz with $p(g \circ f) \leq p(f)p(g)$. Since the composition of two Lipschitz maps is Lipschitz, and the identity map on X is the identity morphism in the categorical sense, it follows that compact metric spaces (as objects) and Lipschitz maps (as morphisms) form a category. We denote it by CMet.

Let (X, d_X) be a metric space. We denote by $\text{Lip}(X, d_X)$ the set of all bounded complex valued continuous functions on (X, d_X) which are Lipschitz with respect to d_X and the standard metric on the complex plane \mathbb{C} (absolute value). Let $\|f\|_\infty = \sup\{|f(x)| : x \in X\}$. Define a norm on $\text{Lip}(X, d_X)$ by $\|f\| = \|f\|_\infty + p(f)$. With respect to pointwise operations $\text{Lip}(X, d_X)$ is a self-adjoint Banach $*$ -algebra over X ($\|a^*\| = \|a\|$, $a \in$ the algebra, and the $*$ -operation is complex conjugation).

DEFINITION 1.2. A commutative Banach $*$ -algebra A is called *Lipschitz* if there exists a metric space (X, d_X) such that $A = \text{Lip}(X, d_X)$.

If (X, d_X) is compact, then $\text{Lip}(X, d_X) = \{f : f \in C(X) \text{ and } p(f) < \infty\}$, and it is a unital, natural, regular, self-adjoint Banach function algebra over X [1, 6, 8]. Clearly, Lipschitz algebras over compact metric spaces (as objects) and unital $*$ -homomorphisms (as morphisms) form a category, which we denote by CLip.

Let (X, d) be any metric space. Let $Y = \{(x, y) \in X \times X : x \neq y\}$, and let $C_b(Y)$ denote the space of all bounded continuous complex valued functions on Y . For $f \in C_b(X)$ and $b \in C_b(Y)$ let $f.b$ and $b.f$ be defined by $(f.b)(x, y) = f(x)b(x, y)$ and $(b.f)(x, y) = f(y)b(x, y)$, $(x, y) \in Y$. Then $C_b(Y)$ is a $C_b(X)$ -bimodule (for this and the subsequent definition see [3] and [7]).

DEFINITION 1.3. Let (X, d) be any metric space, and let $Y = \{(x, y) \in X \times X : x \neq y\}$. Let $D : C_b(X) \rightarrow C_b(Y)$ be defined by

$$(Df)(x, y) = \frac{f(y) - f(x)}{d(x, y)}$$

for $(x, y) \in Y$. We say that D is the *de Leeuw derivation* defined by metric d .

It is easy to see that D is indeed a derivation, and that $\text{Dom}(D) = \text{Lip}(X, d) = \{f \in C_b(X) : \|Df\| < \infty\}$ (see [7]). Clearly $\|Df\| = p(f)$, for $f \in \text{Dom}(D)$.

DEFINITION 1.4. Let (X, d_X) and (Y, d_Y) be compact metric spaces, and let $(C(X), D_X)$ and $(C(Y), D_Y)$ be the corresponding algebras of continuous functions with de Leeuw derivations defined by the corresponding metrics. We say that a homomorphism $\nu : C(X) \rightarrow C(Y)$ is *Lipschitz* if the following condition holds:

$$(1.1) \quad f \in \text{Dom}(D_X) \text{ implies } \nu f \in \text{Dom}(D_Y).$$

Since the composition of two Lipschitz homomorphisms is Lipschitz, and the identity homomorphism on $C(X)$ is identity morphism in categorical sense, it follows that commutative unital C^* -algebras with de Leeuw derivations as objects, and Lipschitz homomorphisms as morphisms form a category. We denote it by CDer .

NOTE 1.1. All homomorphisms of algebras which we encounter here are automatically continuous, since each of them is from a Banach algebra into another commutative semisimple Banach algebra.

The main goal of this paper is to prove the following result, which is a direct consequence of Theorem 2.2 and Theorem 3.1.

THEOREM 1.1. *The category CLip is equivalent to CDer , and CLip and CDer are the duals of CMet .*

This result shows that Lipschitz algebras are indeed a functional analytic counterpart of compact metric spaces. The general direction in which this research aims is to find a reasonable noncommutative analog of a metric space. This result shows that we need a good definition of a noncommutative Lipschitz algebra. So we need a functional analytic characterization of (commutative) Lipschitz algebras in the C^* -algebra setting. This characterization should be similar to the characterization of $C(X)$, the algebras of all continuous functions over compact Hausdorff spaces, as the unital commutative C^* -algebras. This is where the importance of the category CDer comes from. Suppose we can find a characterization of de Leeuw derivations in C^* -algebraic terms, i.e., as derivations from commutative C^* -algebras satisfying certain conditions. Lipschitz algebras would then be characterized as domains of such derivations. Then we can define the noncommutative de Leeuw derivation as a derivation from any C^* -algebra (not necessarily commutative) satisfying the same conditions, and we can simply define the noncommutative Lipschitz algebra as the domain of such a de Leeuw derivation. At the moment we do not know of such a desired characterization of de Leeuw derivations. However, it is clear that such a characterization would have to rely on the results of [7], which state the conditions

which any operator from a C^* -algebra, which defines a metric on the state space of that algebra, has to satisfy.

2. The categories CMet and CLip

We will use the following relation between two metrics on a space.

DEFINITION 2.1. Two metrics d_1 and d_2 on a space X are said to be *boundedly equivalent metrics* if there exist constants $C_1 > 0$ and $C_2 > 0$ such that

$$C_1 d_1(x, y) \leq d_2(x, y) \leq C_2 d_1(x, y)$$

for any $x, y \in X$.

Note that this is the same as saying that the identity map on X , I_X , is Lipschitz both as a map from (X, d_1) to (X, d_2) and as a map from (X, d_2) to (X, d_1) . It is easily seen that the topology on X induced by d_2 is the same as the one induced by d_1 . The Lipschitz algebras over (X, d_1) and (X, d_2) are also similarly related.

PROPOSITION 2.1. [8, Corollary 3.5] *Let d_1 and d_2 be bounded metrics on X . Then $\text{Lip}(X, d_1)$ and $\text{Lip}(X, d_2)$ have the same elements if and only if d_1 and d_2 are boundedly equivalent.*

Similar relationship holds for their norms.

PROPOSITION 2.2. *Let d_1 and d_2 be bounded metrics on X . The norms on $\text{Lip}(X, d_1)$ and on $\text{Lip}(X, d_2)$ are equivalent if and only if d_1 and d_2 are boundedly equivalent.*

PROOF. Suppose that d_1 and d_2 are boundedly equivalent. Then for $f \in \text{Lip}(X, d_1)$

$$\begin{aligned} p_1(f) &= \sup \left\{ \frac{|f(x) - f(y)|}{d_1(x, y)} : x, y \in X \text{ \& } x \neq y \right\} \\ &= \sup \left\{ \frac{|f(x) - f(y)|}{d_2(x, y)} \frac{d_2(x, y)}{d_1(x, y)} : x, y \in X \text{ \& } x \neq y \right\} \\ &\leq \sup \left\{ \frac{|f(x) - f(y)|}{d_2(x, y)} : x, y \in X \text{ \& } x \neq y \right\} \times \\ &\quad \sup \left\{ \frac{d_2(x, y)}{d_1(x, y)} : x, y \in X \text{ \& } x \neq y \right\} = p_2(f) p_{12}(I_X) \end{aligned}$$

where $p_{12}(I_X)$ is the Lipschitz constant of I_X as a map from (X, d_1) to (X, d_2) . Similarly, $p_2(f) \leq p_1(f) p_{21}(I_X)$, where $p_{21}(I_X)$ is the Lipschitz constant of I_X as a map from (X, d_2) to (X, d_1) . Hence

$$\begin{aligned} \|f\|_1 &= \|f\|_\infty + p_1(f) \leq \|f\|_\infty + p_1(f) p_{12}(I_X) \\ &\leq \max\{1, p_{12}(I_X)\} (\|f\|_\infty + p_2(f)) = \max\{1, p_{12}(I_X)\} \|f\|_2. \end{aligned}$$

Similarly, $\|f\|_2 \leq \max\{1, p_{21}(I_X)\} \|f\|_1$.

Conversely, suppose that the norms are equivalent, that is, there are constants C and K such that for any $f \in \text{Lip}(X, d_1)$ we have $p_1(f) \leq C p_2(f)$ and $p_2(f) \leq K p_1(f)$. Let $x, y \in X$. Let the function f_1 be defined by $f_1(z) = d_1(z, y)$, and

let the function f_2 be defined by $f_2(z) = d_2(z, y)$. Then $p_1(f_1) = 1$, $p_2(f_1) \leq K$, $p_2(f_2) = 1$ and $p_1(f_2) \leq C$. We obtain

$$\frac{d_1(x, y)}{d_2(x, y)} = \frac{|f_1(x) - f_1(y)|}{d_2(x, y)} \leq p_2(f_1) \leq K$$

and

$$\frac{d_2(x, y)}{d_1(x, y)} = \frac{|f_2(x) - f_2(y)|}{d_1(x, y)} \leq p_1(f_2) \leq C,$$

which shows that the two metrics are boundedly equivalent. \square

In order to show that CMet is dual to CLip we also need the following result which we quote from [8].

THEOREM 2.1. [8, Theorem 5.1] *Let $A_i = \text{Lip}(X_i, d_i)$ where (X_i, d_i) is compact, $i = 1, 2$. Then every homomorphism $\nu : A_1 \rightarrow A_2$ is of the form*

$$(2.1) \quad (\nu f)(x) = f(F(x)) \quad f \in A_1, x \in X_2$$

where $F : X_2 \rightarrow X_1$ satisfies

$$(2.2) \quad d_1(F(x), F(y)) \leq K d_2(x, y) \quad x, y \in X_2$$

for some positive constant K . Conversely, if ν is defined on A_1 by the equation (2.1) where $F : X_2 \rightarrow X_1$ satisfies the condition (2.2), then ν is a homomorphism of A_1 into A_2 . ν is one-to-one if and only if $F(X_2) = X_1$. ν takes A_1 onto A_2 if and only if F satisfies the additional condition

$$K' d_2(x, y) \leq d_1(F(x), F(y)) \quad x, y \in X$$

for some positive constant K' .

We single out the following easy fact which we need later.

PROPOSITION 2.3. *Let (X, d_X) be a metric space and let Y be a topological space homeomorphic to X via $\tau : X \rightarrow Y$. Then d_Y defined by $d_Y(y_1, y_2) = d_X(\tau^{-1}(y_1), \tau^{-1}(y_2))$ is a metric on Y such that the metric and the original topology on Y coincide. Furthermore, both τ and τ^{-1} are Lipschitz maps between (X, d_X) and (Y, d_Y) of Lipschitz constants $p(\tau) = p(\tau^{-1}) = 1$.*

PROOF. Let U be any open subset of Y . Then $\tau^{-1}(U)$ is open in X and there exists an open ball $B(x_0, r) = \{x \in X : d_X(x_0, x) < r\} \subset \tau^{-1}(U)$. Moreover, $\tau(B(x_0, r)) = \{\tau(x) : x \in B(x_0, r)\} = \{\tau(x) : x \in X \text{ \& } d_X(x_0, x) < r\} = \{y \in Y : d_Y(y, \tau(x_0)) < r\} = B(\tau(x_0), r)$ is an open ball which is contained in U .

Conversely, by the same argument as above, any open ball in (Y, d_Y) is the image of an open ball in (X, d_X) which is an open set in X , and thus so is its τ -image in Y . \square

We denote by I_A the identity map (morphism) on an object A . For example: I_{CMet} is the identity functor on the category CMet; I_X is the identity map on the set X , i.e., $I_X(x) = x$ for all $x \in X$; and I_A is the identity automorphism of an algebra A , i.e., $I_A f = f$ for all $f \in A$.

THEOREM 2.2. *Let (X, d_X) and (Y, d_Y) be compact metric spaces, i.e., objects in CMet , and let $F : (X, d_X) \rightarrow (Y, d_Y)$ be a Lipschitz map, that is a morphism in CMet , i.e., $F \in \text{Hom}((X, d_X), (Y, d_Y))$. Let $T : \text{CMet} \rightarrow \text{CLip}$ be defined by $T(X, d_X) = \text{Lip}(X, d_X)$, and let $TF : \text{Lip}(Y, d_Y) \rightarrow \text{Lip}(X, d_X)$ be defined by $(TF)g = g \circ F$ for $g \in \text{Lip}(Y, d_Y)$.*

Let $A = \text{Lip}(X, d_X)$ and $B = \text{Lip}(Y, d_Y)$ be objects in CLip , and let $\nu : B \rightarrow A$ be a homomorphism, i.e., $\nu \in \text{Hom}(B, A)$. Let $S : \text{CLip} \rightarrow \text{CMet}$ be defined by $SA = (X_{SA}, d_{SA})$, where X_{SA} is the character space of A , and d_{SA} is a metric on X_{SA} defined by

$$(2.3) \quad d_{SA}(\xi, \eta) = \sup\{|\xi f - \eta f| : f \in A \text{ \& } \|f\| \leq 1\},$$

for $\xi, \eta \in X_{SA}$; let $S\nu : SA \rightarrow SB$ be defined by $(S\nu)(\xi)g = \xi(\nu g)$ for $\xi \in X_{SA}$ and $g \in B = \text{Lip}(Y, d_Y)$. Then:

- (a) *T is a contravariant functor from CMet to CLip ;*
- (b) *S is a contravariant functor from CLip to CMet ;*
- (c) *$S \circ T$ is naturally isomorphic to I_{CMet} and $T \circ S$ is naturally isomorphic to I_{CLip} .*

We conclude that CMet and CLip are dual categories.

PROOF. (a) By the quoted Theorem 2.1, $TF \in \text{Hom}(\text{Lip}(Y, d_Y), \text{Lip}(X, d_X))$ for every $F \in \text{Hom}((X, d_X), (Y, d_Y))$. Clearly $T I_X = I_{TX}$, since $(T I_X)f = f \circ I_X = f$, for all $f \in \text{Lip}(X, d_X)$. Let $G \in \text{Hom}((Y, d_Y), (Z, d_Z))$. Then for $h \in \text{Lip}(Z, d_Z)$ we have

$$\begin{aligned} (T(G \circ F))(h) &= h \circ (G \circ F) = (h \circ G) \circ F \\ &= TG(h) \circ F = TF(TG(h)) = (TF \circ TG)(h), \end{aligned}$$

which means that $T(G \circ F) = TF \circ TG$. So T is indeed a contravariant functor.

(b) Since $A = \text{Lip}(X, d_X) \in \text{CLip}$ is a unital natural commutative Banach algebra, by Gelfand theory X_{SA} is a compact Hausdorff space homeomorphic to X via $\tau_X : X \rightarrow X_{SA}$ defined by $(\tau_X(x))f = f(x)$ for $x \in X$ and $f \in A$. Let $\xi, \eta \in X_{SA}$ be such that $\xi = \tau_X(x)$ and $\eta = \tau_X(y)$ for some $x, y \in X$. Then $d_{SA}(\xi, \eta) = d_{SA}(\tau_X(x), \tau_X(y))$ and so

$$\begin{aligned} (2.4) \quad d_{SA}(\xi, \eta) &= \sup\{|\xi f - \eta f| : f \in A \text{ \& } \|f\| \leq 1\} \\ &= \sup\{|(\tau_X(x))f - (\tau_X(y))f| : f \in A \text{ \& } \|f\| \leq 1\} \\ &= \sup\{|f(x) - f(y)| : f \in A \text{ \& } \|f\| \leq 1\} = \frac{d_X(x, y)}{1 + d_X(x, y)}. \end{aligned}$$

To check the validity of last equality in the above equation, note that the value $d_X(x, y)/(1 + d_X(x, y))$ is achieved by the function f defined by $f(z) = \min\{d(x, z), d(x, y)\}/(1 + d_X(x, y))$. On the other hand, if $f(x) = 0$ and $f(y) > d_X(x, y)/(1 + d_X(x, y))$, then $p(f) > 1/(1 + d_X(x, y))$ and $\|f\|_\infty > d_X(x, y)/(1 + d_X(x, y))$, so that $\|f\| > 1$, which should not happen. The important thing is that the metric d_{1X} on X defined by $d_{1X}(x, y) = d_{SA}(\tau(x), \tau(y)) = d_X(x, y)/(1 + d_X(x, y))$ is

boundedly equivalent to d_X . In fact, we have

$$(2.5) \quad \frac{d_X(x, y)}{1 + \text{Diam}(X, d_X)} \leq d_{1X}(x, y) = \frac{d_X(x, y)}{1 + d_X(x, y)} \leq d_X(x, y)$$

for all $x, y \in X$, where $\text{Diam}(X, d_X) = \sup\{d_X(x, y) : x, y \in X\}$. By Proposition 2.3 the metric d_{SA} defined by (2.3) defines a topology on X_{SA} agreeing with the Gelfand topology. Thus $(X_{SA}, d_{SA}) \in \text{CMet}$. To prove that $S\nu$ is a morphism in CMet , we need to show that it is a Lipschitz map. For $\xi, \eta \in X_{SA}$ we have

$$\begin{aligned} d_{SB}(S\nu\xi, S\nu\eta) &= \sup\{|(S\nu\xi)g - (S\nu\eta)g| : g \in B \text{ \& } \|g\| \leq 1\} \\ &= \sup\{|\xi(\nu g) - \eta(\nu g)| : g \in B \text{ \& } \|g\| \leq 1\} \\ &\leq \sup\{|\xi f - \eta f| : f \in \nu B \text{ \& } \|f\| \leq \|\nu\|\} \\ &\leq \|\nu\| \sup\{|\xi f - \eta f| : f \in A \text{ \& } \|f\| \leq 1\} = \|\nu\| d_{SA}(\xi, \eta). \end{aligned}$$

This shows that $p(S\nu) \leq \|\nu\|$. Clearly $S I_A = I_{X_{SA}}$, since for $\xi \in X_{SA}$ $(S I_A(\xi))f = \xi(I_A f) = \xi f$ for all $f \in A$, so $S I_A(\xi) = \xi$. For $\nu \in \text{Hom}(B, A)$, $\mu \in \text{Hom}(C, B)$, $\xi \in X_{SA}$ and $h \in C$ we have

$$\begin{aligned} (S(\nu \circ \mu)(\xi))h &= \xi((\nu \circ \mu)h) = \xi(\nu(\mu h)) \\ &= (S\nu(\xi))(\mu h) = (S\mu(S\nu(\xi)))h = (S\mu \circ S\nu)(\xi)h, \end{aligned}$$

which means that $S(\nu \circ \mu) = S\mu \circ S\nu$. So S is indeed a contravariant functor.

(c) To see that $S \circ T$ is naturally isomorphic to ICMet , let $(X, d_X) \in \text{CMet}$, $A = T(X, d_X) = \text{Lip}(X, d_X)$ and $(S \circ T)(X, d_X) = TA = (X_{SA}, d_{SA})$, and let $\tau_X : X \rightarrow X_{SA}$ be as in (b). We show that τ_X is a natural isomorphism, i.e., that τ_X is an invertible morphism in CMet . Since by Gelfand theory, as pointed out in (b), τ_X is a homeomorphism, we only need to show that τ_X and τ_X^{-1} are both Lipschitz. But, that clearly follows from (2.4) and (2.5), and in fact $p(\tau_X) = \sup\{d_{1X}(x, y)/d_X(x, y) : x, y \in X \text{ \& } x \neq y\} = \sup\{1/(1 + d_X(x, y)) : x, y \in X \text{ \& } x \neq y\} = 1$ and $p(\tau_X^{-1}) = \sup\{d_X(x, y)/d_{1X}(x, y) : x, y \in X \text{ \& } x \neq y\} \leq 1 + \text{Diam}(X, d_X)$. Thus τ_X is an invertible morphism in CMet , therefore ICMet is naturally isomorphic to $S \circ T$.

To see that $T \circ S$ is naturally isomorphic to ICLip , let $A = \text{Lip}(X, d_X) \in \text{CLip}$. Define a map $\sigma_A : A \rightarrow (T \circ S)A = \text{Lip}(X_{SA}, d_{SA})$ by $\sigma_A f = f \circ \tau_X^{-1}$, for $f \in A$, i.e., $(\sigma_A f)(\xi) = f(\tau_X^{-1}(\xi))$ for $\xi \in X_{SA}$. Clearly, $g = \sigma_A f \in C(X_{SA})$ and $\|\sigma_A f\|_\infty = \|f\|_\infty$. Since by Proposition 2.3 τ_X^{-1} is Lipschitz with $p(\tau_X^{-1}) = 1 + \text{Diam}(X, d_X)$, we have that $g = f \circ \tau_X^{-1}$ is Lipschitz, and $p(g) = p(f \circ \tau_X^{-1}) \leq p(f) p(\tau_X^{-1}) = p(f) (1 + \text{Diam}(X, d_X))$. Clearly, σ_A is one-to-one. It is also onto, since for any $g \in \text{Lip}(X_{SA}, d_{SA})$, by the same arguments as above $f = g \circ \tau_X \in \text{Lip}(X, d_X)$, and $\sigma_A f = g$. Note that the situation is similar to the one in Proposition 2.2. Thus σ_A is an invertible morphism in CLip , which means that σ_A is a natural isomorphism. Therefore $T \circ S$ is naturally isomorphic to ICLip . \square

Some remarks concerning the functors T and S are in order.

REMARK 2.1. Since every Lipschitz algebra A is $\text{Lip}(X, d_X)$ for some (X, d_X) , T is onto as a map of objects. Let $F, G \in \text{Hom}((X, d_X), (Y, d_Y))$, and suppose that

$TF = TG$. That means that for all $g \in \text{Lip}(Y, d_Y)$, $TF(g) = TG(g)$, i.e., that $g \circ F = g \circ G$. If $F \neq G$, then there exists $x \in X$ with $F(x) \neq G(x)$. But then there exists $g \in \text{Lip}(Y, d_Y)$ with $g(F(x)) = 1$ and $g(G(x)) = 0$, so that we get $g \circ F \neq g \circ G$. Contradiction. Thus T is one-to-one on morphisms. It is also onto on morphisms, since by the quoted Theorem 2.1 every homomorphism of Lipschitz algebras arises from some Lipschitz map. However, we cannot still claim that CMet and CLip are in fact contravariantly isomorphic categories, since we do not have an explicit inverse for T . Note that the functor S (the would-be inverse of T) does not map $\text{Lip}(X, d_X)$ to (X, d_X) but to (X_{SA}, d_{SA}) .

REMARK 2.2. There are other choices for defining metric d_{SA} than the one given by (2.3). For example, one can define d_{SA} by

$$(2.6) \quad d_{SA}(\xi, \eta) = \sup\{|\xi f - \eta f| : f \in A \text{ \& } p(f) \leq 1\},$$

for $\xi, \eta \in X_{SA}$. The difference between (2.3) and (2.6) is that in (2.6) we are taking $p(f) \leq 1$ instead of $\|f\| \leq 1$ as in (2.3). However, we have to be sure that we can compute $p(f)$ from the given data. This can be done using the following two steps: (1) obtain $\|f\|_\infty$ by $\|f\|_\infty = \sup\{|\xi f| : \xi \in X_{SA}\}$; (2) take $p(f) = \|f\| - \|f\|_\infty$. This approach has the advantage that when the metric d_{1X} on X is defined by $d_{1X}(x, y) = d_{SA}(\tau(x), \tau(y))$, we obtain that $d_{1X} = d_X$, that is, we get back our original metric. This also means that we do not need to use the results concerning boundedly equivalent metrics. We leave it to the taste of the reader to decide which of these two choices we have given is the more suitable one.

3. Category CDer and its relation to CLip and CMet

We now turn our attention to the category CDer .

THEOREM 3.1. *Let $(X, d_X), (Y, d_Y) \in \text{CMet}$, and $F \in \text{Hom}((X, d_X), (Y, d_Y))$. Let $R : \text{CMet} \rightarrow \text{CDer}$ be defined by $R(X, d_X) = (A_{RX}, D_{RX})$, where $A_{RX} = C(X)$ and D_{RX} is the de Leeuw derivation defined by d_X , i.e., $(D_{RX}f)(x, y) = (f(y) - f(x))/d_X(x, y)$ for $x, y \in X$, $x \neq y$, and $f \in \text{Dom}(D_{RX}) = \{f \in C(X) : \|D_{RX}f\|_\infty < \infty\}$. Let $RF : A_{RY} = C(Y) \rightarrow A_{RX} = C(X)$ be defined by $(RF)g = g \circ F$ for $g \in C(Y)$.*

Let $(A, D_A), (B, D_B) \in \text{CDer}$, and let $\nu \in \text{Hom}((A, D_A), (B, D_B))$. Let $Q : \text{CDer} \rightarrow \text{CLip}$ be defined by $QA = \text{Dom}(D_A)$ equipped with the norm $\|f\|_{QA} = \|f\| + \|D_A f\|$ and let $Q\nu : QA \rightarrow QB$ be defined by $Q\nu = \nu|_{QA}$. Then:

- (a) R is a contravariant functor from CMet to CDer ;
- (b) Q is a covariant functor from CDer to CLip ;
- (c) $P = R \circ S : \text{CLip} \rightarrow \text{CDer}$ is a covariant functor;
- (d) $P \circ Q$ is naturally isomorphic to I_{CDer} and $Q \circ P$ is naturally isomorphic to I_{CLip} .

We conclude that CDer is equivalent to CLip and that it is dual to CMet .

PROOF. (a) Clearly, (A_{RX}, D_{RX}) is an object in CDer . Since $\text{Dom}(D_{RX}) = \text{Lip}(X, d_X)$, by (a) of Theorem 2.2, RF is a morphism in CDer , and everything else is as for the functor T .

(b) Clear, since if $A = C(X)$ and $D_A = D_{d_X}$, the derivation defined by d_X , $QA = \text{Lip}(X, d_X)$, and by the condition (1.1) in Definition 1.4, $(Q\nu)f \in \text{Lip}(Y, d_Y)$, so that $Q\nu$ is a homomorphism from $\text{Lip}(X, d_X)$ into $\text{Lip}(Y, d_Y)$.

(c) Clear.

(d) To see that $P \circ Q = R \circ S \circ Q$ is naturally isomorphic to ICDer , let $(A, D_A) \in \text{CDer}$ where $A = C(X)$ and $D_A = D_{d_X}$. Then $QA = \text{Lip}(X, d_X)$, $(S \circ Q)A = (X_{SA}, d_{SA})$, and $(R \circ S \circ Q)A = (C(X_{SA}), D_{d_{SA}})$. Define $\theta_A : A \rightarrow C(X_{SA})$ by $\theta_A f = f \circ \tau_X^{-1}$, i.e., $(\theta_A f)(\xi) = f(\tau_X^{-1}(\xi))$ for $\xi \in X_{SA}$. Note that θ_A is defined similarly as σ_A from the part (c) of the proof of Theorem 2.2. Clearly, $g = \theta_A f \in C(X_{SA})$ and $\|\theta_A f\|_\infty = \|f\|_\infty$, and θ_A is an isomorphism of A and $C(X_{SA})$. We need to show that θ_A and θ_A^{-1} are Lipschitz homomorphisms, that is, that they satisfy the condition (1.1) in Definition 1.4. But, $\text{Dom}(D_A) = \text{Lip}(X, d_X)$ and $\text{Dom}(D_{d_{SA}}) = \text{Lip}(X_{SA}, d_{SA})$, so by the proof of (c) of Theorem 2.2, this is satisfied. Hence θ_A is an invertible morphism in CDer , which means that θ_A is a natural isomorphism. Therefore ICDer is naturally isomorphic to $P \circ Q$.

To see that $Q \circ P = Q \circ R \circ S$ is naturally isomorphic to ICLip , it is enough to observe that $Q \circ R = T$. Thus $Q \circ P = T \circ S$, which we already know from (c) of Theorem 2.2 to be naturally isomorphic to ICLip . \square

REMARK 3.1. A similar remark holds concerning the functors Q , P and R as for the functors T and S as in Remark 2.1. The functor Q is almost an isomorphism of CDer and CLip , but we do not have its explicit inverse, as the functor P is not the one. Likewise, the functor R is almost a contravariant isomorphism from CMet to CDer , but the functor $S \circ Q$ is not its inverse.

Here is an easy, but interesting consequence of the above considerations.

PROPOSITION 3.1. *Let $A = \text{Lip}(X, d_X)$ and $B = \text{Lip}(Y, d_Y)$ be Lipschitz algebras over compact metric spaces, and let $\nu : B \rightarrow A$ be a homomorphism. Then ν is continuous as a map from $C(Y)$ to $C(X)$ and there exists a unique homomorphism $\nu_1 : C(Y) \rightarrow C(X)$ such that $\nu = \nu_1|_B$.*

PROOF. Let $F : X \rightarrow Y$ be defined by $F = \tau_Y^{-1} \circ S\nu \circ \tau_X$, where τ_X , τ_Y and S are defined in Theorem 2.2. Clearly, F is a continuous map, and so $\nu_1 : C(Y) \rightarrow C(X)$ defined by $\nu_1 g = g \circ F$ for $g \in C(Y)$ is a homomorphism. We need to show that $\nu_1|_B = \nu$. So let $g \in B$. Then

$$\begin{aligned} (\nu_1 g)(x) &= g(F(x)) = g(\tau_Y^{-1} \circ S\nu \circ \tau_X(x)) \\ &= (S\nu \circ \tau_X(x))g = \tau_X(x)(\nu g) = (\nu g)(x), \end{aligned}$$

since by definition $g(\tau_Y^{-1}(\eta)) = \eta g$ for $\eta \in X_{SB}$ and $\tau_X(x)f = f(x)$.

To prove the uniqueness of the extended homomorphism $\nu_1 : C(Y) \rightarrow C(X)$, note first that ν_1 is automatically continuous. Furthermore, $B = \text{Lip}(Y, d_Y)$ is dense in $C(Y)$, since it is a point-separating, self-adjoint subalgebra of $C(Y)$. Thus, if there existed another extension $\nu_2 : C(Y) \rightarrow C(X)$ of ν , it would also be continuous and it would coincide with ν_1 on the dense subalgebra $B = \text{Lip}(Y, d_Y)$. Obviously, it then has to be equal to ν_1 . \square

In other words, every homomorphism of Lipschitz algebras is a restriction of some Lipschitz homomorphism of the underlying algebras of continuous functions. To prove the existence of the extension ν_1 of ν without using the approach taken here, one would have to show that the homomorphism ν is continuous as a map from continuous functions on Y to continuous functions on X . Then, one could get the extended homomorphism ν_1 by continuity, using the fact that $\text{Lip}(Y, d_Y)$ is dense in $C(Y)$.

References

- [1] William G. Bade, P. C. Curtis, Jr., and H. G. Dales, *Amenability and weak amenability for Beurling and Lipschitz algebras*, Proc. London Math. Soc. **55** (1987), 359–377.
- [2] Alain Connes, *Compact metric spaces, Fredholm modules, and hyperfiniteness*, Ergodic Theory Dynam. Systems **9** (1989), 207–220.
- [3] Karl De Leeuw, *Banach spaces of Lipschitz functions*, Studia Math. **21** (1961), 55–66.
- [4] Jerry A. Johnson, *Banach spaces of Lipschitz functions and vector-valued Lipschitz functions*, Trans. Amer. Math. Soc. **148** (1970), 147–169.
- [5] Saunders Mac Lane, *Categories for the Working Mathematician*, Springer-Verlag, 1971.
- [6] Branka Pavlović, *Automatic continuity of Lipschitz algebras*, J. Funct. Anal. **131** (1995), 115–144.
- [7] Branka Pavlović, *Defining metric spaces via operators from unital C^* -algebras*, Pacific J. Math. **186** (1998), 285–313.
- [8] Donald R. Sherbert, *Banach algebras of Lipschitz functions*, Pacific J. Math. **13** (1963), 1387–1399.

Matematički institut
Kneza Mihaila 35
11001 Beograd, p.p. 367
Serbia and Montenegro

(Received 13 03 2002)

(Revised 09 02 2003)

and
School of Science
Griffith University
Nathan Qld. 4111, Brisbane
Australia
and
Department of Mathematics
The University of Queensland
Brisbane Qld. 4072
Australia

Current address: Department of Mathematics, The University of Queensland, St. Lucia Qld. 4072, Brisbane, Australia

pavlovic@maths.uq.edu.au