

**EXISTENCE OF SOLUTIONS FOR SOME
NONLINEAR BEAM EQUATIONS ***

P. AMSTER and P.P. CÁRDENAS ALZATE

Recommended by Luís Sanchez

Abstract: We study the existence of solutions for some nonlinear ordinary differential equations under a nonlinear boundary condition which arise on beam theory. Assuming suitable conditions we prove the existence of at least one solution applying topological methods.

1 – Introduction

This work is devoted to the study of the existence of solutions for some nonlinear ordinary differential equations under a nonlinear boundary condition. In 1995 Rebelo and Sanchez [9] have considered the second order problem

$$(1) \quad \begin{cases} u'' + g(t, u) = 0 & 0 < t < T \\ u'(0) = -f(u(0)) \\ u'(T) = f(u(T)) \end{cases}$$

with $g : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ for g satisfying a sign condition or either nondecreasing with respect to u , and $f \in C(\mathbb{R}, \mathbb{R})$ continuous and strictly nondecreasing. This equation may be regarded as a mathematical model for the axial deformation of a nonlinear elastic beam, with two nonlinear elastic springs acting at the extremities according to the law $u'(0) = -f(u(0))$, $u'(T) = f(u(T))$, and the total force exerted by the nonlinear spring undergoing the displacement u given by $g(t, u)$.

Received: February 28, 2005; *Revised:* June 8, 2005.

* This work was partially supported by Fundación Antorchas.

On the other hand, the following fourth order problem for the deflection of a beam resting on elastic bearings was considered, among other authors, by Grossinho and Ma (see [3], [6], and also [4] for asymmetric boundary conditions):

$$(2) \quad \begin{cases} u^{(4)} + g(t, u) = 0 & 0 < t < T \\ u''(0) = u''(T) = 0 \\ u'''(0) = -f(u(0)) \\ u'''(T) = f(u(T)) . \end{cases}$$

In section 2 we study (1) for $g = g(t, u, u')$. We remark that in this more general situation the problem is no longer variational; for this reason we shall apply instead topological methods. On the other hand, in order to find a priori bounds for the derivative we shall assume as in [2] the following Nagumo type condition:

$$(3) \quad |g(t, u, v)| \leq \psi(|v|) \quad \forall (t, u, v) \in \mathcal{E} .$$

Here \mathcal{E} is a subset of $[0, T] \times \mathbb{R}^2$ to be specified, and $\psi : [0, +\infty) \rightarrow (0, +\infty)$ is a continuous function satisfying the inequality

$$\int_r^M \frac{1}{\psi(s)} ds > T$$

for some constants M and r to be specified. Under these assumptions we shall prove the existence of solutions by the method of upper and lower solutions.

Moreover, in section 3 we obtain an existence result under Landesman–Lazer type conditions (see e.g. [8]) applying topological degree methods [7].

Finally, in section 4 we consider the fourth order problem (2) for $g = g(t, u, u', u'', u''')$. More precisely, we prove the existence of symmetric solutions, i.e. such that $u(t) = u(T - t)$, under appropriate Landesman–Lazer and Nagumo type conditions.

2 – The second order case. Upper and lower solutions

In this section we prove an existence result for the following second order problem:

$$(4) \quad \begin{cases} u'' + g(t, u, u') = 0 & 0 < t < T \\ u'(0) = -f(u(0)) \\ u'(T) = f(u(T)) . \end{cases}$$

We shall assume the existence of an ordered couple of a lower and an upper solution. Namely, we shall assume there exist $\alpha, \beta : [0, T] \rightarrow \mathbb{R}$ such that $\alpha(t) \leq \beta(t)$,

$$(5) \quad \alpha''(t) + g(t, \alpha, \alpha') \geq 0 ,$$

$$(6) \quad \beta''(t) + g(t, \beta, \beta') \leq 0 ,$$

and

$$(7) \quad \begin{cases} \alpha'(0) \geq -f(\alpha(0)), & \alpha'(T) \leq f(\alpha(T)) \\ \beta'(0) \leq -f(\beta(0)), & \beta'(T) \geq f(\beta(T)) . \end{cases}$$

In this context, set

$$r = \min \left\{ \max \left\{ \frac{|\alpha(0) - \beta(T)|}{T}, \frac{|\alpha(T) - \beta(0)|}{T} \right\}, \max_{\alpha(0), \alpha(T) \leq s \leq \beta(0), \beta(T)} |f(s)| \right\} ,$$

fix a constant $M > r$ such that

$$M \geq \max \left\{ \|\alpha'\|_{C([0, T])}, \|\beta'\|_{C([0, T])} \right\}$$

and define

$$\mathcal{E} = \left\{ (t, u, v) \in [0, T] \times \mathbb{R}^2 : \alpha(t) \leq u \leq \beta(t), |v| \leq M \right\} .$$

Theorem 2.1. *With the previous notations, assume there exists an ordered couple of a lower and an upper solution of (4). Furthermore, assume that g satisfies the Nagumo condition (3). Then the boundary value problem (4) admits at least one solution u , with*

$$\alpha(t) \leq u(t) \leq \beta(t) , \quad |u'(t)| < M \quad \forall t \in [0, T] .$$

Proof: Set $\lambda > 0$ and consider the functions $P : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$, $Q : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$P(t, x) = \begin{cases} x & \alpha(t) \leq x \leq \beta(t) \\ \beta(t) & x > \beta(t) \\ \alpha(t) & x < \alpha(t) , \end{cases}$$

$$Q(x) = \begin{cases} x & -M \leq x \leq M \\ M & x > M \\ -M & x < -M . \end{cases}$$

We define a compact fixed point operator $\phi : C^1([0, T]) \rightarrow C^1([0, T])$ in the following way: for each $v \in C^1([0, T])$, let $u = \phi(v)$ be the unique solution of the linear Neumann problem

$$\begin{aligned} u'' - \lambda u &= g(t, P(t, v), Q(v')) - \lambda P(t, v) , \\ u'(0) &= -f(P(0, v(0))) , \quad u'(T) = f(P(T, v(T))) . \end{aligned}$$

By standard results, ϕ is well defined and compact. Moreover, multiplying the previous equation by u it follows that

$$- \int_0^T (u'' - \lambda u) u \leq C \|u\|_{L^2}$$

for some constant C . Hence

$$\|u'\|_{L^2}^2 + \lambda \|u\|_{L^2}^2 \leq C \|u\|_{L^2} + f(P(T, v(T))) u(T) + f(P(0, v(0))) u(0) ,$$

and it follows that $\|u\|_{H^1} \leq C$ for some constant C . We conclude that $\|u\|_{C^1} \leq C$ for some constant C , and by a straightforward application of Schauder Theorem it follows that ϕ has a fixed point u . We claim that

$$\alpha(t) \leq u(t) \leq \beta(t) , \quad |u'(t)| < M \quad \forall t \in [0, T] ,$$

and hence u is a solution of the problem. Indeed, if for example $(u - \beta)(t_0) > 0$ for some $t_0 \in (0, T)$ maximum, then $P(t_0, u(t_0)) = \beta(t_0)$, $u'(t_0) = \beta'(t_0)$, and

$$\begin{aligned} (u - \beta)''(t_0) - \lambda(u - \beta)(t_0) &\geq g(t_0, P(t_0, u(t_0)), Q(u'(t_0))) - \lambda P(t_0, u(t_0)) \\ &\quad - \left[g(t_0, \beta(t_0), \beta'(t_0)) - \lambda \beta(t_0) \right] = 0 , \end{aligned}$$

a contradiction. Now, if $u - \beta$ attains an absolute positive maximum for example at $t = 0$, then $(u - \beta)'(0) \leq 0$. Moreover, as $P(0, u(0)) = \beta(0)$ we deduce that $(u - \beta)'(0) = -f(P(0, u(0))) - \beta'(0) \geq 0$, and hence $(u - \beta)'(0) = 0$. On the other hand, in a neighborhood of 0 we have that $u(t) > \beta(t)$ and then

$$\begin{aligned} (u - \beta)'' - \lambda(u - \beta) &\geq g(t, P(t, u), Q(u')) - \lambda P(t, u) - [g(t, \beta, \beta') - \lambda \beta] \\ &= g(t, \beta, Q(u')) - g(t, \beta, \beta') . \end{aligned}$$

As $u'(0) = \beta'(0) \in [-M, M]$, the right-hand term vanishes at $t = 0$, meanwhile $u(0) - \beta(0) > 0$. It follows that $(u - \beta)'' \geq \lambda(u - \beta) + g(t, \beta, Q(u')) - g(t, \beta, \beta') > 0$ in $(0, \delta)$ for some $\delta > 0$, which contradicts the fact that 0 is an absolute maximum

of $u - \beta$. In the same way, it follows that $u - \beta$ cannot attain a positive absolute maximum at T . We deduce in a similar way that $u(t) \geq \alpha(t)$ for every $t \in [0, T]$.

Next, assume for example that $u'(t_0) = M$ for some t_0 .

If $r = \max_{\alpha(0), \alpha(T) \leq s \leq \beta(0), \beta(T)} |f(s)|$, then $|u'(0)|, |u'(T)| \leq r$; otherwise there exists \tilde{t} such that

$$u'(\tilde{t}) = \frac{u(T) - u(0)}{T} \leq \frac{\beta(T) - \alpha(0)}{T} \leq r .$$

In both cases, we deduce the existence of t_1 such that $u'(t_1) = r$. We may assume that $r < u'(t) < M$ for any t between t_1 and t_0 , and hence

$$T < \int_r^M \frac{1}{\psi(s)} ds = \int_{t_1}^{t_0} \frac{u''(t)}{\psi(u'(t))} dt \leq \left| \int_{t_1}^{t_0} \frac{g(t, u, u')}{\psi(u'(t))} dt \right| \leq |t_0 - t_1| ,$$

a contradiction. The proof is analogous if $u'(t_0) = -M$. ■

Remark 2.2. In particular, the conditions of the previous theorem hold if there exist two constants $\alpha < \beta$ such that

$$g(t, \alpha, 0) \geq 0 \geq g(t, \beta, 0)$$

and

$$f(\alpha) \geq 0 \geq f(\beta)$$

provided that g satisfies $|g(t, u, v)| \leq \psi(|v|)$ for $\alpha \leq u \leq \beta$, $|v| < M$ and $\int_0^M \frac{1}{\psi(s)} ds > T$. □

Remark 2.3. When f is nondecreasing, a more general result is proved in [1]. □

3 – Landesman–Lazer type conditions

In this section we prove the existence of solutions of (4) under Landesman–Lazer type conditions. We shall assume that f is one-side globally bounded, i.e. $f \leq r$ or $f \geq -r$ for some positive constant r , and that g satisfies the Nagumo condition (3) over the set

$$\mathcal{E} = \left\{ (t, u, v) \in [0, T] \times \mathbb{R}^2 : |v| \leq M \right\}$$

for some $M > r$.

Moreover, we shall assume that the limits

$$\limsup_{u \rightarrow \pm\infty} g(t, u, v) := g_s^\pm(t)$$

and

$$\liminf_{u \rightarrow \pm\infty} g(t, u, v) := g_i^\pm(t)$$

exist, and that they are uniform for $|v| < M$. We also define the (possibly infinite) quantities

$$\limsup_{u \rightarrow \pm\infty} f(u) := f_s^\pm$$

and

$$\liminf_{u \rightarrow \pm\infty} f(u) := f_i^\pm .$$

Then we have:

Theorem 3.1. *Under the previous assumptions, problem (4) admits at least one solution, provided that one of the following conditions holds:*

$$(8) \quad 2f_s^+ + \int_0^T g_s^+(t) dt < 0 < 2f_i^- + \int_0^T g_i^-(t) dt$$

$$(9) \quad 2f_s^- + \int_0^T g_s^-(t) dt < 0 < 2f_i^+ + \int_0^T g_i^+(t) dt .$$

Remark 3.2. Conditions of this kind are known in the literature as Landesman–Lazer type conditions after the pioneering paper of E. Landesman and A. Lazer [5]. In particular, taking $f = 0$ in Theorem 3.1 we obtain standard Landesman–Lazer conditions for the Neumann problem. \square

For the sake of completeness, we summarize the main aspects of coincidence degree theory. Let \mathbb{V} and \mathbb{W} be real normed spaces, $L : \text{Dom}(L) \subset \mathbb{V} \rightarrow \mathbb{W}$ a linear Fredholm mapping of index 0, and $N : \mathbb{V} \rightarrow \mathbb{W}$ continuous. Moreover, set two continuous projectors $\pi_{\mathbb{V}} : \mathbb{V} \rightarrow \mathbb{V}$ and $\pi_{\mathbb{W}} : \mathbb{W} \rightarrow \mathbb{W}$ such that $\text{R}(\pi_{\mathbb{V}}) = \text{Ker}(L)$ and $\text{Ker}(\pi_{\mathbb{W}}) = \text{R}(L)$, and an isomorphism $J : \text{R}(\pi_{\mathbb{W}}) \rightarrow \text{Ker}(L)$. It is readily seen that

$$L_{\pi_{\mathbb{V}}} := L|_{\text{Dom}(L) \cap \text{Ker}(\pi_{\mathbb{V}})} : \text{Dom}(L) \cap \text{Ker}(\pi_{\mathbb{V}}) \rightarrow \text{R}(L)$$

is one-to-one; denote its inverse by $K_{\pi_{\mathbb{V}}}$. If Ω is a bounded open subset of \mathbb{V} , N is called L -compact on Ω if $\pi_{\mathbb{W}}N(\Omega)$ is bounded and $K_{\pi_{\mathbb{V}}}(I - \pi_{\mathbb{W}})N : \Omega \rightarrow \mathbb{V}$ is compact.

The following continuation theorem is due to Mawhin [7]:

Theorem 3.3. *Let L be a Fredholm mapping of index zero and N be L -compact on a bounded domain $\Omega \subset \mathbb{V}$. Suppose that:*

1. $Lx \neq \lambda Nx$ for each $\lambda \in (0, 1]$ and each $x \in \partial\Omega$.
2. $\pi_{\mathbb{W}}Nx \neq 0$ for each $x \in \text{Ker}(L) \cap \partial\Omega$.
3. $d(J\pi_{\mathbb{W}}N, \Omega \cap \text{Ker}(L), 0) \neq 0$, where d denotes the Brouwer degree.

Then the equation $Lx = Nx$ has at least one solution in $\text{Dom}(L) \cap \Omega$. ■

Proof of Theorem 3.1: Set $\mathbb{V} = C^1([0, T])$, $\mathbb{W} = L^2(0, T) \times \mathbb{R}^2$, and the operators $L : H^2(0, T) \rightarrow \mathbb{W}$, $N : \mathbb{V} \rightarrow \mathbb{W}$ given by

$$Lu = (u'', u'(0), u'(T)) , \quad Nu = (-g(\cdot, u, u'), -f(u(0)), f(u(T))) .$$

It is easy to verify that

$$\text{Ker}(L) = \mathbb{R} , \quad \text{R}(L) = \left\{ (\varphi, A, B) \in \mathbb{W} : \bar{\varphi} = \frac{B-A}{T} \right\} ,$$

where $\bar{\varphi}$ denotes the usual average given by $\bar{\varphi} = \frac{1}{T} \int_0^T \varphi(t) dt$. Then, we may define $\pi_{\mathbb{V}}(X) = \bar{u}$, $\pi_{\mathbb{W}}(\varphi, A, B) = (\bar{\varphi} - \frac{B-A}{T}, 0, 0)$, and $J : \text{R}(\pi_{\mathbb{W}}) \rightarrow \mathbb{R}$ given by $J(C, 0, 0) = C$. In this case, for $(\varphi, A, B) \in \text{R}(L)$, the function $U = K_{\pi_{\mathbb{V}}}(\varphi, A, B)$ is defined as the unique solution of the problem

$$U'' = \varphi , \quad U'(0) = A, \quad U'(T) = B$$

that satisfies $\bar{U} = 0$. Writing $U'(t) = A + \int_0^t \varphi$ and using Wirtinger inequality, L -compactness of N follows.

We claim there exists a constant R such that if $Lu = \lambda Nu$ with $0 < \lambda \leq 1$ then $\|u\|_{C^1} \leq R$. Indeed, suppose by contradiction that $Lu_n = \lambda_n Nu_n$, with $0 < \lambda_n \leq 1$ and $\|u_n\|_{C^1} \rightarrow \infty$. As $u_n'' = -\lambda_n g(t, u_n, u_n')$ and $u_n'(0) = -\lambda_n f(u_n(0))$, $u_n'(T) = \lambda_n f(u_n(T))$, by the Nagumo condition and using the fact that

$$\min \{ u_n'(0), u_n'(T) \} \leq r \quad \text{and} \quad \max \{ u_n'(0), u_n'(T) \} \geq -r ,$$

it follows as in the previous section that $\|u_n'\|_{C([0, T])} < M$ for every n . Hence $\|u_n\|_{C([0, T])} \rightarrow \infty$, and $\|u_n - \bar{u}_n\|_{C([0, T])} \leq C$ for some constant C . Taking

a subsequence, assume for example that $\bar{u}_n \rightarrow +\infty$ and that (8) holds; then integrating the equation we obtain the equality

$$f(u_n(T)) + f(u_n(0)) = - \int_0^T g(t, u_n, u'_n) dt ,$$

and thus

$$0 \leq \limsup_{n \rightarrow \infty} f(u_n(T)) + \limsup_{n \rightarrow \infty} f(u_n(0)) + \int_0^T g_s^+(t) dt < 0$$

a contradiction. The proof is similar for the other cases; hence, taking $\Omega = B_R(0)$ for R large enough, the first condition in Theorem 3.3 is fulfilled.

Further, the function $J\pi_{\mathbb{W}}N|_{\bar{\Omega} \cap \text{Ker}(L)} = [-R, R]$ is given by

$$J\pi_{\mathbb{W}}N(s) = -\frac{1}{T} \left(\int_0^T g(t, s, 0) dt + 2f(s) \right) ,$$

and in the same way as before it follows that for R large enough

$$J\pi_{\mathbb{W}}N(R) J\pi_{\mathbb{W}}N(-R) < 0 .$$

Thus, $\deg(J\pi_{\mathbb{W}}N, \Omega \cap \text{Ker}(L), 0) = \pm 1$, and the proof is complete. ■

4 – Symmetric solutions for the general fourth order case

In this section we study the existence of symmetric solutions for the problem

$$(10) \quad \begin{cases} u^{(4)} + g(t, u, u', u'', u''') = 0 & 0 < t < T \\ u''(0) = u''(T) = 0 \\ u'''(0) = -f(u(0)) \\ u'''(T) = f(u(T)) . \end{cases}$$

We shall assume that g is symmetric with respect to t , namely:

$$(11) \quad g(t, u, v, w, x) = g(T - t, u, v, w, x) .$$

Our Nagumo condition for this problem reads:

$$(12) \quad |g(t, u, v, w, x)| \leq \psi(|x|) \quad \forall (t, u, v, w, x) \in \mathcal{E}$$

with $\mathcal{E} = [0, T] \times \mathbb{R}^3 \times [-M, M]$, and $\psi : [0, +\infty) \rightarrow (0, +\infty)$ continuous, with

$$\int_0^M \frac{1}{\psi(s)} ds > T .$$

Moreover, assume that the limits

$$\limsup_{s \rightarrow \pm\infty} g(t, s, v, w, x) := g_s^\pm(t)$$

and

$$\liminf_{s \rightarrow \pm\infty} g(t, s, v, w, x) := g_i^\pm(t)$$

exist, and that they are uniform over the set

$$\mathcal{C} = \left\{ (v, w, x) \in \mathbb{R}^3 : |v| < \frac{T^2}{4}M, |w| < \frac{T}{2}M \text{ and } |x| < M \right\} .$$

The quantities f_s^\pm and f_i^\pm are defined as before. Then we have:

Theorem 4.1. *Under the previous assumptions, problem (10) admits at least one symmetric solution, provided that one of the conditions (8) or (9) holds.*

Proof: We proceed as in the proof of Theorem 3.1. Let

$$\mathbb{V} = \left\{ u \in C^3([0, T]) : u(t) = u(T - t), u''(0) = 0 \right\} ,$$

$$\mathbb{W} = \left\{ u \in L^2(0, T) : u(t) = u(T - t) \right\} \times \mathbb{R}$$

and define the operators $L : H^4(0, T) \cap \mathbb{V} \rightarrow \mathbb{W}$, $N : \mathbb{V} \rightarrow \mathbb{W}$ by

$$Lu = (u^{(4)}, u'''(0)) , \quad Nu = -(g(\cdot, u, u', u'', u'''), f(u(0))) .$$

Again, it is easy to verify that

$$\text{Ker}(L) = \mathbb{R} , \quad \text{R}(L) = \left\{ (\varphi, c) \in \mathbb{W} : \int_0^T \varphi(t) dt + 2c = 0 \right\} .$$

Then, we may define $\pi_{\mathbb{V}}(u) = \bar{u}$, $\pi_{\mathbb{W}}(\varphi, c) = (\bar{\varphi} + 2c, 0)$, and $J : \text{R}(\pi_{\mathbb{W}}) \rightarrow \mathbb{R}$ given by $J(C, 0) = C$. For $(\varphi, c) \in \text{R}(L)$, the function $U = K_{\pi_{\mathbb{V}}}(\varphi, c)$ is defined as the unique solution of the problem

$$\begin{cases} U^{(4)} = \varphi \\ U''(0) = 0, \quad U'''(0) = c \\ U(t) = U(T - t) \\ \bar{U} = 0 . \end{cases}$$

As before, it is easy to prove that N is L -compact. Next, if $Lu_n = \lambda_n Nu_n$, with $0 < \lambda_n \leq 1$ and $\|u_n\|_{C^3} \rightarrow \infty$, by the Nagumo condition and using the fact that $u_n'''(\frac{T}{2}) = 0$, it follows that $\|u_n'''\|_{C([0,T])} < M$ for every n . Moreover, for $t \leq \frac{T}{2}$ we have:

$$|u_n''| \leq \int_0^t |u_n''| < \frac{T}{2} M$$

and

$$|u_n'| \leq \left| \int_t^{\frac{T}{2}} u_n'' \right| < \frac{T^2}{4} M .$$

As u_n is symmetric, we conclude that $(u_n'(t), u_n''(t), u_n'''(t)) \in \mathcal{C}$ for every $t \in [0, T]$. Then $\|u_n\|_{C([0,T])} \rightarrow \infty$, and $\|u_n - \bar{u}_n\|_{C^3([0,T])} \leq C$ for some constant C . The rest of the proof follows as in the second order case. ■

Some examples and remarks

Example 4.2. As an example of Theorem 4.1 we may consider a symmetric function g such that

$$g(t, u, v, w, x) = g_0(t, u) + \gamma(u) g_1(t, u, v, w, x) ,$$

where g_0 is bounded, $|g_1(t, u, v, w, x)| \leq A + B|x|$ and $\gamma(u) \rightarrow 0$ as $|u| \rightarrow \infty$.

Then $|g(t, u, v, w, x)| \leq C + D|x|$ for some positive constants C and D and the Nagumo condition is satisfied taking $\psi(x) = C + Dx$ and M large enough. Moreover,

$$\limsup_{u \rightarrow \pm\infty} g_0(t, u) = g_s^\pm(t) , \quad \liminf_{u \rightarrow \pm\infty} g_0(t, u) = g_i^\pm(t) ,$$

and the assumptions of Theorem 4.1 are fulfilled if (8) or (9) holds. For example, it suffices to assume that

$$\lim_{|u| \rightarrow \infty} f(u) \operatorname{sgn}(u) = +\infty \quad \text{or} \quad \lim_{|u| \rightarrow \infty} f(u) \operatorname{sgn}(u) = -\infty . \quad \square$$

Remark 4.3. In the situation of Theorem 4.1, if $g_s^\pm = g_i^\pm := g^\pm$ and $f_s^\pm = f_i^\pm := f^\pm$, integrating the equation it follows that if for example

$$g^+(t) \leq g \leq g^-(t) \quad \text{and} \quad f^+ < f < f^-$$

or

$$g^-(t) \leq g \leq g^+(t) \quad \text{and} \quad f^- < f < f^+$$

then the respective conditions (8) and (9) are also necessary. □

Remark 4.4. The Nagumo condition (12) can be dropped if we assume that g has a linear growth of the type

$$|g(t, u, v, w, x)| \leq A + B|u| + C|v| + D|w| + E|x|$$

(with B, C, D and E small enough), and that the limits g_t^\pm and g_s^\pm are uniform on \mathbb{R}^3 . Indeed, in this case if $Lu_n = \lambda_n Nu_n$, with $0 < \lambda_n \leq 1$, then using the fact that $u_n''' = \lambda_n \int_{\frac{t}{2}}^t g(s, u_n, u_n', u_n'', u_n''') ds$, we deduce:

$$\left(1 - \frac{TE}{2}\right) \|u_n'''\|_{C([0, T])} \leq \frac{T}{2} \left(A + B\|u_n\|_{C([0, T])} + C\|u_n'\|_{C([0, T])} + D\|u_n''\|_{C([0, T])} \right).$$

Integrating twice, as E, D and C are small, we obtain:

$$\|u_n'\|_{C([0, T])} \leq \delta(A + B\|u_n\|_{C([0, T])})$$

for some constant δ . By the mean value theorem, for $B < \delta$ we conclude that if for example $\bar{u}_n \rightarrow +\infty$ then $\inf_{t \in [0, T]} u_n(t) \rightarrow +\infty$, and the rest of the proof follows as before. In particular, for $g = g(t, u)$ it suffices to take $B < \frac{16}{T^4}$. \square

Remark 4.5. In [3], Theorem 2, it is proved by variational methods that if $g = g(t, u)$ is symmetric on t , and $f, g(t, \cdot)$ are nondecreasing, then problem (10) admits a symmetric solution if and only if

$$2f(a) + \int_0^T g(t, a) dt = 0 \quad \text{for some } a \in \mathbb{R}.$$

By monotonicity, this condition is equivalent to (9), unless $f(u) \equiv f(a)$ and $g(t, u) \equiv g(t, a)$ for all $u \geq a$ or for all $u \leq a$. Note that, in this last case, existence of solutions can be easily proved; thus, taking into account the previous remarks 4.3 and 4.4, when $|g(t, u)| \leq A + B|u|$ (with $B < \frac{16}{T^4}$) we may conclude that Theorem 4.1 is essentially equivalent to Theorem 2 in [3].

Moreover, without the monotonicity condition the authors prove (see [3], Theorem 5) the existence of a symmetric solution of (10) for g and f sublinear, i.e.

$$\frac{g(t, u)}{u} \rightarrow 0 \quad \text{as } |u| \rightarrow \infty$$

uniformly in t , and

$$\frac{f(u)}{u} \rightarrow 0 \quad \text{as } |u| \rightarrow \infty,$$

assuming a growth condition for f and g , and that one of the following hypotheses holds:

- i) $g(t, u) \rightarrow \pm\infty$ as $u \rightarrow \pm\infty$ uniformly in t and f bounded by below.
- ii) $f(u) \rightarrow \pm\infty$ as $u \rightarrow \pm\infty$ and g bounded by below.

It is clear that the sublinearity condition implies that $|g(t, u)| \leq A + B|u|$ for some $B < \frac{16}{7^4}$ and some A , and that if i) or ii) holds then the second inequality in condition (9) is fulfilled. Thus, some cases of Theorem 5 in [3] are covered by Theorem 4.1; in particular, if f is bounded by above for $u < 0$ in i) or if g is bounded by above for $u < 0$ in ii).

However, the first inequality in (9) does not necessarily hold under assumptions i) or ii): one may consider for instance the (sublinear) functions $f(u) = |u|^{1/2}$ and $g(t, u) = u^{1/3}$. \square

ACKNOWLEDGEMENTS – The authors would like to thank the anonymous referee for the careful reading of the original manuscript and his/her fruitful corrections and remarks.

REFERENCES

- [1] FABRY, CH. and HABETS, P. – Upper and lower solutions for second-order boundary value problems with nonlinear boundary conditions, *Nonlinear Analysis*, 10 (1986), 985–1007.
- [2] FRANCO, D. and O’REGAN, D. – *Existence of solutions to second order problems with nonlinear boundary conditions*, Proc. of the Fourth Int. Conf. on Dynamical Systems and Diff. Equations, Discrete and Continuous Dynamical Systems 2003, 273–280.
- [3] GROSSINHO, M. and MA, T.F. – Symmetric equilibria for a beam with a nonlinear elastic foundation, *Portugaliae Mathematica*, 51 (1994), 375–393.
- [4] GROSSINHO, M. and TERSIAN, S. – The dual variational principle and equilibria for a beam resting on a discontinuous nonlinear elastic foundation, *Nonlinear Analysis, Theory, Methods, and Applications*, 41 (2000), 417–431.
- [5] LANDESMAN, E. and LAZER, A. – Nonlinear perturbations of linear elliptic boundary value problems at resonance, *J. Math. Mech.*, 19 (1970), 609–623.
- [6] MA, T.F. – Existence results for a model of nonlinear beam on elastic bearings, *Applied Mathematical Letters*, 13 (2000), 11–15.
- [7] MAWHIN, J. – *Topological degree methods in nonlinear boundary value problems*, NSF-CBMS Regional Conference in Mathematics no. 40, American Mathematical Society, Providence, RI, 1979.

- [8] MAWHIN, J. – Landesman–Lazer conditions for boundary value problems: A nonlinear version of resonance, *Bol. de la Sociedad Española de Mat. Aplicada*, 16 (2000), 45–65.
- [9] REBELO, C. and SANCHEZ, L. – Existence and multiplicity for an O.D.E. with nonlinear boundary conditions, *Differential Equations and Dynamical Systems*, 3(4) (1995), 383–396.

Pablo Amster,
FCEyN, Departamento de Matemática, Universidad de Buenos Aires,
Ciudad Universitaria, Pabellón I, (1428) Buenos Aires – ARGENTINA
E-mail: pamster@dm.uba.ar
and
CONICET

and

Pedro Pablo Cárdenas Alzate,
Departamento de Matemáticas, Universidad Tecnológica de Pereira,
Pereira – COLOMBIA
E-mail: ppablo@utp.edu.co