

Doran–Harder–Thompson Conjecture via SYZ Mirror Symmetry: Elliptic Curves

Atsushi KANAZAWA

Department of Mathematics, Kyoto University,
Kitashirakawa-Oiwake, Sakyo, Kyoto, 606-8502, Japan
E-mail: akanazawa@math.kyoto-u.ac.jp

Received December 20, 2016, in final form April 06, 2017; Published online April 11, 2017

<https://doi.org/10.3842/SIGMA.2017.024>

Abstract. We prove the Doran–Harder–Thompson conjecture in the case of elliptic curves by using ideas from SYZ mirror symmetry. The conjecture claims that when a Calabi–Yau manifold X degenerates to a union of two quasi-Fano manifolds (Tyurin degeneration), a mirror Calabi–Yau manifold of X can be constructed by gluing the two mirror Landau–Ginzburg models of the quasi-Fano manifolds. The two crucial ideas in our proof are to obtain a complex structure by gluing the underlying affine manifolds and to construct the theta functions from the Landau–Ginzburg superpotentials.

Key words: Calabi–Yau manifolds; Fano manifolds; SYZ mirror symmetry; Landau–Ginzburg models; Tyurin degeneration; affine geometry

2010 Mathematics Subject Classification: 53D37; 14J33; 14J32; 14J45; 14D06

1 Introduction

The aim of this short article is to prove the Doran–Harder–Thompson conjecture in the case of elliptic curves by using ideas from SYZ mirror symmetry.

Given a Tyurin degeneration of a Calabi–Yau manifold X to a union of two quasi-Fano manifolds $X_1 \cup_Z X_2$ intersecting along a common smooth anti-canonical divisor $Z \in |-K_{X_i}|$ for $i = 1, 2$, it is natural to investigate a potential relationship between geometry of the Calabi–Yau manifold X and that of the quasi-Fano manifolds X_1 and X_2 . Motivated by the works of Dolgachev [5], Tyurin [21] and Auroux [3], recently Doran–Harder–Thompson proposed a remarkable conjecture (Conjecture 2.3), which builds a bridge between mirror symmetry for the Calabi–Yau manifold X and that for the quasi-Fano manifolds X_1 and X_2 [6]. It claims that we should be able to glue together the mirror Landau–Ginzburg models $W_i: Y_i \rightarrow \mathbb{C}$ of the pair (X_i, Z) for $i = 1, 2$ to construct a mirror Calabi–Yau manifold Y of X equipped with a fibration $W: Y \rightarrow \mathbb{P}^1$. They provided supporting evidence for the conjecture in various different settings. For instance it was shown that under suitable assumptions we can glue together the Landau–Ginzburg models $W_i: Y_i \rightarrow \mathbb{C}$ for $i = 1, 2$ to obtain a C^∞ -manifold Y with the expected topological Euler number $\chi(Y) = (-1)^{\dim X} \chi(X)$. Thus the topological version of the conjecture is essentially proven. However, the real difficulty of this conjecture lies in constructing Y as a complex manifold, which should be mirror to the symplectic manifold X (or vice versa).

In this article, we will prove the conjecture in the case of elliptic curves, beginning with a symplectic elliptic curve X and constructing the mirror complex elliptic curve Y . In order to obtain the correct complex manifold Y by gluing the mirror Landau–Ginzburg models $W_i: Y_i \rightarrow \mathbb{C}$ for $i = 1, 2$, we need to keep track of the subtle complex structures of the (not necessarily algebraic) Kähler manifold Y_i for $i = 1, 2$. To this end, we find ideas of SYZ mirror symmetry very useful.

This paper is a contribution to the Special Issue on Modular Forms and String Theory in honor of Noriko Yui. The full collection is available at <http://www.emis.de/journals/SIGMA/modular-forms.html>

The key idea in our proof is twofold. The first is to obtain the correct complex structure by gluing the underlying affine base manifolds of X_1 and X_2 in SYZ mirror symmetry. This is based on the philosophy that a Tyurin degeneration of a Calabi–Yau manifold X can be thought to be fibred over a Heegaard splitting of the base B of a special Lagrangian torus fibration $\phi: X \rightarrow B$. The second is to construct theta functions out of the Landau–Ginzburg superpotentials. As a corollary, we observe that the product formulae of the theta functions are the manifestation of quantum corrections appearing in SYZ mirror symmetry.

It is worth mentioning that a variant of the conjecture was discussed in the work of Auroux [3]. He studied a $\mathbb{Z}/2\mathbb{Z}$ -invariant version of mirror symmetry (Calabi–Yau double covers) and investigated elliptic curves from a different perspective from ours [3, Example 3.2]. Advantages of the present work are firstly to work with not necessarily identical affine manifolds (doubling) and secondly to construct theta functions out of the Landau–Ginzburg superpotentials in an interesting way based on the geometry of the conjecture.

Structure of article

We will provide a self-contained description of the subjects for completeness. Section 2 builds basic setup and formulates the main conjecture. Section 3 reviews basics of SYZ mirror symmetry both in the Calabi–Yau and quasi-Fano settings. Section 4 is the main part of this article and proves the conjecture in the case of elliptic curves. Section 5 comments on further research directions.

2 Doran–Harder–Thompson conjecture

In this section, we will provide background materials and review the Doran–Harder–Thompson conjecture, following the original article [6]. In general mirror symmetry is a conjecture about a Calabi–Yau manifold near a large complex structure limit, which is thought to be a maximal degeneration, in the complex moduli space. However in this article we will be interested in another class of loci in the complex moduli space, where a Calabi–Yau manifold degenerates to a union of two quasi-Fano manifolds.

2.1 Tyurin degeneration

A Calabi–Yau manifold X is a compact Kähler manifold such that the canonical bundle is trivial $K_X = 0$ and $H^i(X, \mathcal{O}_X) = 0$ for $0 < i < \dim X$. A quasi-Fano manifold X is a smooth variety X such that $|-K_X|$ contains a smooth Calabi–Yau member and $H^i(X, \mathcal{O}_X) = 0$ for $0 < i$. A Tyurin degeneration is a degeneration $\mathcal{X} \rightarrow \Delta$ of Calabi–Yau manifolds over the unit disc $\Delta = \{|z| < 1\} \subset \mathbb{C}$, such that the total space \mathcal{X} is smooth and the central fibre $\mathcal{X}_0 = X_1 \cup_Z X_2$ is a union of two quasi-Fano manifolds X_1 and X_2 intersecting normally along a common anti-canonical divisor $Z \in |-K_{X_i}|$ for $i = 1, 2$. Conversely, we have the following result of Kawamata–Namikawa [15], which is slightly modified for our setting.

Theorem 2.1 ([15, Theorem 4.2]). *Let X_1 and X_2 be quasi-Fano manifolds and $Z \in |-K_{X_i}|$ a common smooth anti-canonical divisor for $i = 1, 2$. Assume that there exist ample divisors $D_i \in \text{Pic}(X_i)$ for $i = 1, 2$ which restrict to an ample divisor $D_1|_Z = D_2|_Z$ on Z . Then the union $X_1 \cup_Z X_2$ of X_1 and X_2 intersecting normally along Z is smoothable to a Calabi–Yau manifold X if and only if $N_{Z/X_1} \cong N_{Z/X_2}^{-1}$ (d -semistability). Moreover the resulting Calabi–Yau manifold X is unique up to deformation.*

A Tyurin degeneration is thought to be a complex analogue of a Heegaard splitting of a compact oriented real 3-fold without boundary. Based on this analogy, in his posthumous article [21]

Tyurin proposed to study geometry of a Calabi–Yau 3-fold by using that of quasi-Fano 3-folds when they are related by a Tyurin degeneration.

2.2 Mirror symmetry for quasi-Fano manifolds

We consider a pair (X, Z) consisting of a quasi-Fano manifold X and an anti-canonical divisor $Z \in |-K_X|$. The complement $X \setminus Z$ can be thought of as a log Calabi–Yau manifold as there exists a nowhere vanishing volume form Ω on $X \setminus Z$ with poles along Z .

Example 2.2. Let X be a toric Fano n -fold and Z the toric boundary, which is the complement of the dense torus $(\mathbb{C}^\times)^n \subset X$. Then $X \setminus Z = (\mathbb{C}^\times)^n$ carries a standard holomorphic volume form $\Omega = \wedge_{i=1}^n \sqrt{-1} d \log z_i$, where (z_i) are the coordinates of $(\mathbb{C}^\times)^n$.

Definition 2.3. A Landau–Ginzburg model is a pair (Y, W) of a Kähler manifold Y and a holomorphic function $W: Y \rightarrow \mathbb{C}$, which is called a superpotential.

It is classically known that there is a version of mirror symmetry for Fano manifolds together with an anti-canonical divisor. We expect that such mirror symmetry should hold also for quasi-Fano manifolds (or even for varieties with effective anti-canonical divisors [2]). Here we formulate a mirror conjecture for quasi-Fano manifolds (see for example Katzarkov–Kontsevich–Pantev [14], Harder [9]).

Conjecture 2.4. *For a pair (X, Z) of a quasi-Fano n -fold X and a smooth anti-canonical divisor $Z \in |-K_X|$, there exists a Landau–Ginzburg model (Y, W) such that*

- 1) $\sum_j h^{n-i+j, j}(X) = h^i(Y, W^{-1}(s))$ for a regular value $s \in \mathbb{C}$ of W ,
- 2) *the generic fibres of W and the generic anti-canonical hypersurfaces in X are mirror families of compact Calabi–Yau $(n-1)$ -folds,*

where $h^i(Y, W^{-1}(s))$ is the rank of the relative cohomology group $H^i(Y, W^{-1}(s))$. The pair (Y, W) is called a mirror Landau–Ginzburg model of (X, Z) .

The anti-canonical divisor Z can be thought of as an obstruction for the quasi-Fano manifold X to be a Calabi–Yau manifold and W is an obstruction (or potential function) for the Floer homology of a Lagrangian torus in X to be defined in the sense of Fukaya–Oh–Ohta–Ono [4, 7] as we will see in the next section.

Example 2.5. Let $X = \mathbb{P}^1$ and $Z = \{0, \infty\}$ equipped with a toric Kähler form ω . Then the mirror Landau–Ginzburg model of (X, Z) is given by $(\mathbb{C}^\times, W = z + \frac{q}{z})$, where $q = \exp(-\int_{\mathbb{P}^1} \omega)$. One justification of this mirror duality is given by the ring isomorphism

$$\mathrm{QH}(\mathbb{P}^1) = \mathbb{C}[H]/(H^2 - q) \cong \mathbb{C}[z^{\pm 1}]/(z^2 - q) = \mathrm{Jac}(W).$$

Here $\mathrm{QH}(\mathbb{P}^1)$ is the quantum cohomology ring of \mathbb{P}^1 and $\mathrm{Jac}(W)$ is the Jacobian ring of the superpotential W .

Conjecture 2.4 can be generalized to the case when Z is mildly singular, but the superpotential W will no longer be proper. In fact, it is anticipated that Z is smooth if and only if W is proper.

Example 2.6. For $X = \mathbb{P}^2$ with a toric Kähler form ω , we define Z_0 to be the toric boundary, Z_1 the union of a smooth conic and a line intersecting 2 points, and Z_2 a nodal cubic curve, and Z_3 a smooth cubic curve. The mirror Landau–Ginzburg model of (X, Z_0) is given by

$$\left(Y_0 = (\mathbb{C}^\times)^2, W_0 = x + y + \frac{q}{xy} \right),$$

where $q = \exp(-\int_H \omega)$ for the line class H . A generic fiber of W_0 is an elliptic curve with 3 punctures. On the other hand, the mirror Landau–Ginzburg model (Y_i, W_i) of the pair (X, Z_i) is a fiberwise partial compactification of W_0 such that a generic fiber of W_i is an elliptic curve with $3 - i$ punctures for $1 \leq i \leq 3$.

2.3 Doran–Harder–Thompson conjecture

Mirror symmetry for Calabi–Yau manifolds and that for (quasi-)Fano manifolds have been studied for a long time, but somewhat independently. A natural question to ask is, *how are mirror symmetry for these manifolds related to each other?* Motivated by works of Dolgachev [5], Tyurin [21], and Auroux [3], Doran–Harder–Thompson proposed the following remarkable conjecture, which we call the DHT conjecture for short.

Conjecture 2.7 (Doran–Harder–Thompson [6]). *Given a Tyurin degeneration of a Calabi–Yau manifold X to the union $X_1 \cup_Z X_2$ of quasi-Fano manifolds intersecting along their common smooth anti-canonical divisor Z , then the mirror Landau–Ginzburg models $W_i: Y_i \rightarrow \mathbb{C}$ of (X_i, Z) for $i = 1, 2$ can be glued together to be a Calabi–Yau manifold Y equipped with a Calabi–Yau fibration $W: Y \rightarrow \mathbb{P}^1$. Moreover, Y is mirror symmetric to X .*

The above gluing process can be understood as follows. We denote by n the dimension of X and by Z_i^\vee a fiber of the superpotential W_i mirror to a Calabi–Yau $(n - 1)$ -fold Z .

1. Firstly, we assume that all the important information about the Landau–Ginzburg model $W_i: Y_i \rightarrow \mathbb{C}$ is contained in the critical locus of the superpotential W_i . Therefore, without much loss of information, we may replace it with a new Landau–Ginzburg model $W_i: Y_i \rightarrow \mathbb{D}_i$ for a sufficiently large disc \mathbb{D}_i which contains all the critical values.
2. Secondly, the Calabi–Yau manifolds Z_1^\vee and Z_2^\vee are both mirror symmetric to Z , and thus we expect that they are topologically identified¹.
3. Thirdly, Theorem 2.1 implies that we have $N_{Z/X_1} \cong N_{Z/X_2}^{-1}$ because $X_1 \cup_Z X_2$ is smoothable to a Calabi–Yau manifold X . According to Kontsevich’s homological mirror symmetry [16], we have an equivalence of triangulated categories

$$D^b\text{Coh}(Z) \cong D^b\text{Fuk}(Z_i^\vee).$$

Then the monodromy symplectomorphism on Z_i^\vee associated to the anti-clockwise loop $\partial\mathbb{D}_i$ can be identified with the autoequivalence $(-) \otimes \omega_{X_i}[n]|_Z \cong (-) \otimes N_{Z/X_i}^{-1}[n]$ on $D^b\text{Coh}(Z)$ (see [14, 19] for details). Therefore $N_{Z/X_1} \cong N_{Z/X_2}^{-1}$ implies, under mirror symmetry, that the monodromy action on Z_1^\vee along the anti-clockwise loop $\partial\mathbb{D}_1$ and that on Z_2^\vee along the clockwise loop $-\partial\mathbb{D}_2$ can be identified.

Therefore, assuming various mirror symmetry statements, we are able to glue the fibrations $W_i: Y_i \rightarrow \mathbb{D}_i$ for $i = 1, 2$ along open neighborhoods of the boundaries $\partial\mathbb{D}_1$ and $\partial\mathbb{D}_2$ to construct a C^∞ -manifold Y equipped with a fibration $W: Y \rightarrow S^2$ (Fig. 1). Note that the smoothness of Z implies the compactness of Y , which follows from the properness of the superpotentials W_i for $i = 1, 2$ (cf. Example 2.6). The highly non-trivial part of the conjecture is that there exist a Calabi–Yau structure on Y and a complex structure on S^2 in such a way that $W: Y \rightarrow \mathbb{P}^1$ is holomorphic and Y is mirror symmetric to the Calabi–Yau manifold X . Moreover, a fiber of W is mirror to Z by the above construction.

¹Two Calabi–Yau manifolds may be topologically different even if they share the same mirror manifold. There is no problem if $\dim X = 1, 2$ or 3 for example.




Figure 1. Gluing Landau–Ginzburg models.

In [6] the authors provide supporting evidence for this conjecture in various different settings, including Batyrev–Borisov mirror symmetry and Dolgachev–Nikulin mirror symmetry. For example, in the 3-fold case, the Hodge number mirror symmetry is shown to be equivalent to a version of Dolgachev–Nikulin mirror symmetry for K3 surfaces, provided that Y admits a Calabi–Yau structure. Another important result is that, under reasonable assumptions, the resulting C^∞ -manifold Y has the expected Euler number $\chi(Y) = (-1)^{\dim X} \chi(X)$. Thus the conjecture is essentially proven at the topological level. However, the real difficulty of the conjecture lies in constructing Y as a complex manifold, which should be mirror to the symplectic manifold X (or vice versa). The goal of this article is to overcome this difficulty in the 1-dimensional case.

One warning is in order. The DHT conjecture is likely to be false unless we impose a condition on how the Tyurin degeneration of a Calabi–Yau manifold X occurs in the complex moduli space. For example, if the complex moduli space of X is 1-dimensional, the Kähler moduli space of a mirror Calabi–Yau manifold Y is also 1-dimensional and thus Y cannot have a fibration structure (unless it is 1-dimensional). The conjecture should be modified so that the Tyurin degeneration occurs in a locus which contains a large complex structure limit. In such a case, the Calabi–Yau manifold Y should be the mirror corresponding to the large complex structure limit. Otherwise what we could expect is that there exists a *homological mirror* W of X equipped with a *non-commutative Calabi–Yau fibration* $D^b\text{Coh}(\mathbb{P}^1) \rightarrow D^b(W)$. This can be thought of as homological mirror to the Tyurin degeneration (see [6, Section 6] for more details).

3 SYZ mirror symmetry

The Strominger–Yau–Zaslow (SYZ) mirror symmetry conjecture [20] provides a foundational geometric understanding of mirror symmetry for Calabi–Yau manifolds. It claims that a mirror pair of Calabi–Yau manifolds should admit dual special Lagrangian torus fibrations. It is Hitchin [10] who first observed that the base of the fibration, which is locally the moduli space of the special Lagrangian fibers [18], carries two natural integral affine structures. These integral affine structures are essential in SYZ mirror symmetry and appear to be more fundamental than symplectic and complex geometry [8]. One of the integral affine structures will play a vital role in our proof of the DHT conjecture.

3.1 SYZ mirror symmetry for Calabi–Yau manifolds

Let X be a Calabi–Yau n -fold equipped with a Kähler form ω and holomorphic volume form Ω . An n -dimensional real submanifold $L \subset X$ is called special Lagrangian if $\omega|_L = 0$ and $\text{Im}(\Omega)|_L = 0$ (after suitable change of the phase of Ω). The celebrated SYZ mirror symmetry conjecture [20] asserts that, for a mirror pair of Calabi–Yau n -folds X and Y , there exist special Lagrangian

T^n -fibrations ϕ and ϕ^\vee

$$\begin{array}{ccc} X & & Y \\ & \searrow \phi & \swarrow \phi^\vee \\ & B & \end{array}$$

over the common base B , which are fiberwisely dual to each other away from singular fibers. The treatment of singular fibers constitutes the essential part where the quantum corrections come into the play. The SYZ conjecture not only provides a powerful way to construct a mirror manifold Y out of X as a fiberwise dual, but also explains why mirror symmetry should hold via Fourier–Mukai type transformations [17]. It is worth noting that a mirror manifold Y depends on a choice of a special Lagrangian fibration on X , and conjecturally this is equivalent to a choice of a large complex structure limit, where the Gromov–Hausdorff limit of the Calabi–Yau manifold X with its Ricci-flat metric normalized to have a fixed diameter is identified with the base B .

3.2 Integral affine structures on B

Let $\phi: X \rightarrow B$ be a special Lagrangian T^n -fibration of a Calabi–Yau n -fold X . We denote by L_b the fiber of ϕ at $b \in B$. The complement $B^\circ \subset B$ of the discriminant locus carries two natural integral affine structures², which we call symplectic and complex. They are defined by ω and $\text{Im}(\Omega)$ respectively as follows.

Symplectic integral affine structure. Let $U \subset B^\circ$ be a small open neighborhood of $b \in B^\circ$. For a basis $\{\alpha_i\}$ of $H_1(L_b, \mathbb{Z})$, the symplectic integral affine coordinates at $b' \in U$ are defined by $x_i := \int_{A_i} \omega$, where $A_i \in H_2(X, L_b \cup L_{b'})$ is the 2-dimensional cylinder traced out by α_i . This is classically known in the theory of action-angle variables.

Complex integral affine structure. The construction is parallel to the above. Let $U \subset B^\circ$ be a small open neighborhood of $b \in B^\circ$. For a basis $\{\beta_i\}$ of $H_{n-1}(L_b, \mathbb{Z})$, the complex integral affine coordinates at $b' \in U$ are defined by $x_i^\vee := \int_{B_i} \text{Im}(\Omega)$, where $B_i \in H_n(X, L_b \cup L_{b'})$ is the n -dimensional cylinder traced out by β_i .

We can easily check that the above coordinates are well-defined because a generic fiber L is a special Lagrangian and both ω and Ω are closed. These coordinates depend on a choice of a base point $b \in B^\circ$ and a choice of a basis of $H_1(L_b, \mathbb{Z})$ or $H_{n-1}(L_b, \mathbb{Z})$. Different base points and different bases of the homology group give rise to changes of coordinates by elements in $\text{GL}_n(\mathbb{Z}) \ltimes \mathbb{R}^n$.

Example 3.1. For $a, b \in \mathbb{R}_+$, let $X = \mathbb{C}/(\mathbb{Z} + \sqrt{-1}a\mathbb{Z})$ be the elliptic curve equipped with a holomorphic volume form $\Omega = dz$ and a Kähler form $\omega = \frac{b}{a} dx \wedge dy$. Then $X \rightarrow B \simeq T^1$, $z \mapsto \text{Im}(z)$ defines a smooth special Lagrangian T^1 -fibration. The symplectic and complex affine lengths of the base B are given by $b = \int_E \omega$ and $a = \int_0^a \text{Im}(\Omega)$ respectively. We observe that the mirror elliptic curve $Y = \mathbb{C}/(\mathbb{Z} + \sqrt{-1}b\mathbb{Z})$ is obtained by switching the two affine structures on the base B . We can complexify this picture by introducing B -fields.

On the other hand, given an integral affine manifold B of dimension n , we have smooth dual T^n -fibrations:

$$\begin{array}{ccc} TB/\Lambda & & T^*B/\Lambda^* \\ & \searrow \phi & \swarrow \phi^\vee \\ & B & \end{array}$$

²An integral affine manifold is a manifold whose local coordinate changes are given by elements in the integral affine transformation group $\text{GL}_n(\mathbb{Z}) \ltimes \mathbb{R}^n$.

Here $\Lambda = \mathbb{Z}\langle \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \rangle$ is a fiberwise lattice in the tangent bundle TB generated by integral affine coordinates $\{x_i\}$ of B , and Λ^* is the dual lattice in the cotangent bundle T^*B . In a natural way, TB/Λ and T^*B/Λ^* are complex and symplectic manifolds respectively. In order to make them (possibly non-compact) Calabi–Yau manifolds, we need a dual integral affine structure on B . More precisely, we need a potential $f: B \rightarrow \mathbb{R}$ satisfying the real Monge–Ampère equation $\det(\frac{\partial^2 f}{\partial x_i \partial x_j}) = C$ for a constant $C \in \mathbb{R}$. Then the dual integral affine structure is given by the Legendre transformation of the original one. For example, the symplectic and complex affine structures discussed above are Legendre dual to each other. This is called semi-flat mirror symmetry and serves as a local model for SYZ mirror symmetry without quantum correction [17]. In general it is a very hard problem to extend this picture when singular fibers are present.

In this article we will begin with a symplectic manifold X and construct a complex manifold Y . So let us take a close look at this case. Given a special Lagrangian T^n -fibration $\phi: X \rightarrow B$ of a Calabi–Yau manifold X , we endow B^o with the symplectic integral affine structure. We may think of the semi-flat mirror Y^o of $X^o := \phi^{-1}(B^o)$ as the space of pairs (b, ∇) where $b \in B^o$ and ∇ is a flat $U(1)$ -connection on the trivial complex line bundle over L_b up to gauge. There is a natural map $\phi^\vee: Y^o \rightarrow B^o$ given by forgetting the second coordinate. With the same notation as before, the complex structure of Y^o is given by the following semi-flat complex coordinates

$$z_i(b, \nabla) := \exp\left(-2\pi \int_{A_i} \omega\right) \text{Hol}_\nabla(\gamma_i),$$

where $\text{Hol}_\nabla(\gamma_i)$ denotes the holonomy of ∇ along the path γ_i . Then we observe that the dual fibration ϕ^\vee is locally given by the tropicalization map $(z_i) \mapsto \left(-\frac{\log |z_i|}{2\pi}\right)_i$.

3.3 SYZ mirror symmetry for quasi-Fano manifolds

Let us now consider a quasi-Fano n -fold X with an anti-canonical divisor $Z \in |-K_X|$. Observing that the complement $X \setminus Z$ carries a holomorphic n -form with poles along Z , we think of $X \setminus Z$ as a log Calabi–Yau manifold, to which the SYZ construction can be applied. More generally, in the framework of SYZ mirror symmetry for a manifold with an effective anti-canonical divisor [2, 4], the superpotential W of a mirror Landau–Ginzburg model is obtained as the weighted count of holomorphic discs of Maslov index $\mu = 2$ with boundary in a smooth fiber L of a given special Lagrangian torus fibration $\phi: X \rightarrow B$. To be more explicit, the superpotential W is a function on the semi-flat mirror Y^o given by

$$W(b, \nabla) := \sum_{\substack{\beta \in \pi_2(X, L_b) \\ \mu(\beta) = 2}} n_\beta z_\beta(b, \nabla),$$

where z_β is defined to be

$$z_\beta(b, \nabla) := \exp\left(-2\pi \int_\beta \omega\right) \text{Hol}_\nabla(\partial\beta)$$

and n_β denotes the one-point open Gromov–Witten invariant of class $\beta \in \pi_2(X, L)$ defined by the machinery of Fukaya–Oh–Ohta–Ono [7]. It is not difficult to check that W is locally a holomorphic function on Y^o .

In the following, we shall focus on the toric Fano case. Namely, we consider a toric Fano n -fold X equipped with a toric Kähler form ω and a meromorphic volume form $\Omega = \wedge_{i=1}^n \sqrt{-1} d \log z_i$, where $(z_i)_i$ are the standard coordinates of the open dense torus $(\mathbb{C}^\times)^n \subset X$. Let $Z \subset X$ be the toric boundary (Example 2.2). Then the toric moment map $\phi: X \rightarrow \mathbb{R}^n$ gives a smooth

special Lagrangian T^n -fibration $\phi: X \setminus Z \rightarrow B^\circ$, where $B := \phi(X) \subset \mathbb{R}^n$ is the moment polytope and B° is its interior³. By construction of the semi-flat mirror, it is straightforward to check the following assertion.

Proposition 3.2. *Let $\text{Trop}: (\mathbb{C}^\times)^n \rightarrow \mathbb{R}^n$, $(z_i)_i \mapsto \left(-\frac{\log|z_i|}{2\pi}\right)_i$ be the tropicalization map. Then the semi-flat mirror Y° of the complement $X \setminus Z \cong (\mathbb{C}^\times)^n$ is given by the polyannulus $\text{Trop}^{-1}(B^\circ)$. Moreover the dual fibration ϕ^\vee is identified with the restriction $\phi^\vee = \text{Trop}|_{Y^\circ}: Y^\circ \rightarrow B^\circ$.*

In the toric Fano case, we do not modify Y° further, so henceforth we simply write $Y = Y^\circ$. In general, there is a discriminant locus in the interior of B and then the semi-flat mirror Y° needs quantum corrections by the wall-crossing formulae of the superpotential W .

Let us take a close look at the projective line \mathbb{P}^1 . We have a special Lagrangian T^1 -fibration $\phi: \mathbb{P}^1 \rightarrow B = [0, \text{Im}(\tau)]$ given by the moment map, where $\tau := \sqrt{-1} \int_{\mathbb{P}^1} \omega$. By Proposition 3.2, the mirror Y of $\mathbb{P}^1 \setminus \{0, \infty\} \cong \mathbb{C}^\times$ is given by the annulus

$$Y = A_{(q,1)} := \{q < |z| < 1\} \subset \mathbb{C},$$

where $q := e^{2\pi\sqrt{-1}\tau}$. Each special Lagrangian fiber separates \mathbb{P}^1 into two discs, one containing 0 and the other containing ∞ . The classes β_1 and β_2 representing these disc classes satisfy $\beta_1 + \beta_2 = [\mathbb{P}^1]$, and hence the coordinates on Y should satisfy $z_{\beta_1} z_{\beta_2} = q$. Moreover we can easily check that these are the only holomorphic discs of Maslov index 2 and $n_{\beta_1} = n_{\beta_2} = 1$. Using $z = z_{\beta_1}$ as a new coordinate on the mirror $Y = A_{(q,1)}$, we obtain the Landau–Ginzburg superpotential

$$W = z_{\beta_1} + z_{\beta_2} = z + \frac{q}{z}.$$

So far, we discuss only the real Kähler structure for simplicity, but we can easily complexify it in the above discussion. We will do this in the next section.

Remark 3.3. The moment map for the toric $(S^1)^n$ -action is defined only up to addition of a constant in the range \mathbb{R}^n . In other words, the only intrinsic property of the base space B is its affine structure and an affine embedding $B \subset \mathbb{R}^n$ is a choice. For example, we may take another moment map $\phi': \mathbb{P}^1 \rightarrow B' = \left[-\frac{\text{Im}(\tau)}{2}, \frac{\text{Im}(\tau)}{2}\right]$, and then the mirror Landau–Ginzburg model becomes

$$W': Y' = A_{(q^{\frac{1}{2}}, q^{-\frac{1}{2}})} \rightarrow \mathbb{C}, \quad z \mapsto z + \frac{q}{z},$$

where

$$A_{(a,b)} := \{z \in \mathbb{C} \mid a < |z| < b\}$$

for positive real numbers $a < b$. Note that we have a biholomorphism $A_{(q,1)} \cong A_{(q^{\frac{1}{2}}, q^{-\frac{1}{2}})}$, which is induced by the translation of the underlying affine manifolds $B \cong B'$ in \mathbb{R}^n . Moreover, near the large volume limit, where $\int_{\beta} \omega \rightarrow \infty$, we may identify Y' with $\mathbb{C}^\times = A_{(0,\infty)}$. In this way, we understand that the SYZ mirror coincides with the Hori–Vafa mirror $(\mathbb{C}^\times, z + \frac{q}{z})$, after suitably renormalizing the superpotential (Example 2.5). We refer the reader to [11] for a physical derivation of the mirror Landau–Ginzburg model.

³By abuse of notation, we use the same ϕ for the restriction of ϕ to $X \setminus Z$.

It is also worth noting that the superpotential W is in general not known to converge if X is not a toric Fano manifold. Moreover, if a fiber Lagrangian L bounds a holomorphic disc of class $\beta \in \pi_2(X, L)$ of Maslov index 0, then the one-point open Gromov–Witten invariant n_β depends on the fiber L as well as the point $p \in L$ which the holomorphic discs are required to pass through. On the other hand, we often want to take a smooth anti-canonical divisor Z instead of the toric boundary so that the mirror Landau–Ginzburg superpotential W is proper. In that case there appears to be a discriminant locus in the interior of the base B and we need quantum corrections in the above toric SYZ construction [2].

4 Gluing Landau–Ginzburg models: elliptic curves

Finally we are in the position to confirm the DHT conjecture in the case of elliptic curves. In the language of SYZ mirror symmetry, we will *glue the Landau–Ginzburg models* by essentially gluing the affine base manifolds of special Lagrangian fibrations. The inspiration comes from the fact that a Tyurin degeneration is a complex analogue of a Heegaard splitting. Another key ingredient of the proof is to construct theta functions out of the Landau–Ginzburg superpotentials based on the geometry of the conjecture.

Let us consider a Tyurin degeneration of an elliptic curve $\mathcal{X} \rightarrow \Delta$, where a generic fiber is a smooth elliptic curve and the central fiber $\mathcal{X}_0 = X_1 \cup_Z X_2$ is the union of two rational curves X_1 and X_2 glued at two points Z . We complexify the Kähler structure $B + \sqrt{-1}\omega$ of \mathcal{X} by introducing the B -field $B \in H^2(\mathcal{X}, \mathbb{R})/2\pi H^2(\mathcal{X}, \mathbb{Z})$, and define

$$\tau_i := \int_{X_i} (B + \sqrt{-1}\omega), \quad q_i := e^{2\pi\sqrt{-1}\tau_i}, \quad i = 1, 2.$$

Then the complexified Kähler structure of a generic elliptic fiber X of the family $\mathcal{X} \rightarrow \Delta$ is given by

$$\tau := \tau_1 + \tau_2 = \int_{\mathcal{X}_0} (B + \sqrt{-1}\omega)$$

so that $q := e^{2\pi\sqrt{-1}\tau} = q_1 q_2$.

Let us consider the moment maps $\phi_i: X_i \rightarrow B_i$ for $i = 1, 2$, where the base affine manifolds are $B_1 = [0, \text{Im}(\tau_1)]$ and $B_2 = [-\text{Im}(\tau_2), 0]$. Then the mirror Landau–Ginzburgs of (X_1, Z) and (X_2, Z) are respectively given by

$$\begin{aligned} W_1: \quad Y_1 &= A_{(|q_1|, 1)} \rightarrow \mathbb{C}, & z_1 &\mapsto z_1 + \frac{q_1}{z_1}, \\ W_2: \quad Y_2 &= A_{(1, |q_2|^{-1})} \rightarrow \mathbb{C}, & z_2 &\mapsto z_2 + \frac{q_2}{z_2}, \end{aligned}$$

where $q_1 z_2 = z_1$ (Remark 3.3). We observe that the boundary of the closure $\overline{Y_1 \cup Y_2} \subset \mathbb{C}^\times$ can be glued by the multiplication map $z \mapsto qz$ to form an elliptic curve Y , which is identified with the mirror elliptic curve $\mathbb{C}^\times/q^{\mathbb{Z}} := \mathbb{C}^\times/(z \sim qz)$ of X . This construction corresponds to the gluing of the boundary of the affine manifold $B_1 \cup B_2 = [-\text{Im}(\tau_2), \text{Im}(\tau_1)]$ by the shift $\text{Im}(\tau)$ (twisted by the B -field upstairs).

In order to confirm the DHT conjecture, we moreover want a map $\mathbb{C}^\times \rightarrow (\mathbb{C} \times \mathbb{C}) \setminus (0, 0)$ which descends to a double covering $W: \mathbb{C}^\times/q^{\mathbb{Z}} \rightarrow \mathbb{P}^1$ that locally looks like the superpotential W_1 (resp. W_2) over the upper (resp. lower) semisphere of the base \mathbb{P}^1 . Unfortunately, the naive analytic continuation of (W_1, W_2) over \mathbb{C}^\times does not work. The correct answer is given by

considering all the Landau–Ginzburg models of the above sort, namely for $i \in \mathbb{Z}$ the Landau–Ginzburg models

$$\begin{aligned} W_{2i+1}: \quad Y_{2i+1} &= A_{(|q^{-i}q_1|, |q^{-i}|)} \rightarrow \mathbb{C}, & z_{2i+1} &\mapsto z_{2i+1} + \frac{q_1}{z_{2i+1}}, \\ W_{2i}: \quad Y_{2i} &= A_{(|q^{1-i}|, |q^{1-i}q_2^{-1}|)} \rightarrow \mathbb{C}, & z_{2i} &\mapsto z_{2i} + \frac{q_2}{z_{2i}}, \end{aligned}$$

where the variable z_i is defined inductively by $qz_{i+2} = z_i$. This means to eliminate the above arbitrary choice of a fundamental domain of the \mathbb{Z} -action on \mathbb{C}^\times to construct the mirror elliptic curve $\mathbb{C}^\times/q^\mathbb{Z}$. A crucial observation is that if we consider all the even or odd superpotential W_i 's at once (in the sense of Remark 4.1 below), they descend to the elliptic curve $\mathbb{C}^\times/q^\mathbb{Z}$ as sections of an ample line bundle as follows. Let us first consider the infinite product

$$\begin{aligned} W'_1(z) &:= \prod_{i=1}^{\infty} \left(1 + \frac{q_1}{z_{2i-1}^2}\right) \left(1 + \frac{z_{-2i+1}^2}{q_1}\right) \\ &= \prod_{i=1}^{\infty} (1 + q^{2i-1}(q_2z^2)^{-1})(1 + q^{2i-1}q_2z^2) \\ &= \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \sum_{l \in \mathbb{Z}} q^{l^2} (q_2z^2)^l \\ &= \frac{e^{\frac{\pi\sqrt{-1}\tau}{6}}}{\eta(2\tau)} \vartheta_{\frac{1}{2}, 0}(2\zeta - \tau_1, 2\tau), \end{aligned}$$

where we set $z = z_1 = e^{2\pi\sqrt{-1}\zeta}$ and

$$\eta(\tau) := e^{\frac{\pi\sqrt{-1}\tau}{12}} \prod_{m=1}^{\infty} (1 - e^{2\pi\sqrt{-1}\tau m}), \quad \vartheta_{a,b}(\zeta, \tau) := \sum_{n \in \mathbb{Z}} e^{\pi\sqrt{-1}(n+a)^2\tau} e^{2\pi\sqrt{-1}(n+a)(\zeta+b)}$$

are the Dedekind eta function and theta function with characteristic $(a, b) \in \mathbb{R}^2$ respectively.

Remark 4.1. We can think of W'_1 as the product of all the Landau–Ginzburg superpotential W_i 's for odd $i \in \mathbb{Z}$ because of the formula

$$\left(z_j + \frac{q_k}{z_j}\right) \left(z_{-j} + \frac{q_k}{z_{-j}}\right) = q_k q^j \left(1 + \frac{q_k}{z_j^2}\right) \left(1 + \frac{z_{-j}^2}{q_k}\right),$$

for all $j \in \mathbb{Z}$ and $k = 1, 2$.

In a similar manner, we next consider

$$\begin{aligned} W'_2(z) &:= \prod_{i=1}^{\infty} \left(1 + \frac{q_2}{z_{2i}^2}\right) \left(1 + \frac{z_{-2i+2}^2}{q_2}\right) \\ &= \prod_{i=1}^{\infty} \left(1 + q^{2i-1} \frac{q_1}{z^2}\right) \left(1 + q^{2i-1} \frac{z^2}{q_1}\right) \\ &= \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \sum_{l \in \mathbb{Z}} q^{l^2} \left(\frac{z^2}{q_1}\right)^l \\ &= \frac{e^{\frac{\pi\sqrt{-1}\tau}{6}}}{\eta(2\tau)} \vartheta_{0,0}(2\zeta - \tau_1, 2\tau), \end{aligned}$$

which can be thought of as the product of all the Landau–Ginzburg superpotential W_i 's for even $i \in \mathbb{Z}$. It is a classical fact that the theta functions with characteristics

$$\vartheta_{\frac{1}{2},0}(2\zeta - \tau_1, 2\tau), \quad \vartheta_{0,0}(2\zeta - \tau_1, 2\tau)$$

form a basis of the (2)-polarization of the elliptic curve $Y = \mathbb{C}^\times/q^{\mathbb{Z}}$. Therefore we obtain a double covering

$$W: Y = \mathbb{C}^\times/q^{\mathbb{Z}} \rightarrow \mathbb{P}^1, \quad z \mapsto [W'_1(z) : W'_2(z)].$$

Observing that W locally looks like the superpotential W_i on each piece Y_i , we confirm that it is precisely the gluing of the two Landau–Ginzburg models argued in the DHT conjecture. This completes a proof of the conjecture in the case of elliptic curves.

Remark 4.2. It is crucial in the above proof not to take the large volume limit but to keep track of the complex structures on the mirror annuli. In this way, we are able to naturally glue the Landau–Ginzburg models (without the heuristic cutting process discussed in the conjecture). It is also interesting to observe that the product expressions of the theta functions are the manifestation of quantum corrections, which are encoded in the Landau–Ginzburg superpotentials, in SYZ mirror symmetry.

The elliptic curves are somewhat special and our construction readily generalizes to a degeneration of an elliptic curve to a nodal union of n rational curves forming a cycle. The superpotential of each rational curve corresponds to a theta function with an appropriate characteristic, and they span a basis of the (n) -polarization of the mirror elliptic curve. Note that the same result is obtained in [1, Section 8.4] and [13, Section 4] from different perspectives. This is due to the accidental fact that a Tyurin degeneration of an elliptic curve can be thought of as a maximal degeneration at a large complex structure limit. A main difference shows up, for example, when we consider a type II degeneration of an abelian surface which is neither a maximal nor a toric degeneration [12]. However the essential mechanism of the DHT conjecture is already apparent in the case of elliptic curves: gluing the base affine manifolds and constructing theta functions from the Landau–Ginzburg superpotentials.

5 Further research direction

A key idea of our proof is to glue the two different affine manifolds B_1 and B_2 along the boundaries to obtain a compact affine manifold $\mathbb{R}/\text{Im}(\tau)\mathbb{Z}$. This idea is not new and a similar construction (doubling) was already suggested by Auroux [3]. However, it is in general a very hard problem to glue together higher dimensional affine manifolds along the boundaries. The difficulty is closely related to a choice of an anti-canonical divisor $Z \in |-K_X|$ of a quasi-Fano manifold X . More precisely, we need a special Lagrangian fibration $\phi: X \rightarrow B$ which is compatible with Z , and then smoothness of Z is likely to be proportional to that of the boundary ∂B . On the other hand, mirror symmetry for (X, Z) tends to be harder as Z gets less singular because we need to trade singularities of ∂B with discriminant loci of the interior of B . This seems the main obstruction to generalizing our discussion in higher dimensions.

A notable feature of the DHT conjecture is that it bridges a gap between mirror symmetry for Calabi–Yau manifolds and that for Fano manifolds. The conjecture naturally suggests a construction of a mirror Calabi–Yau manifold from a more general degeneration of a Calabi–Yau manifold. For instance, the recent Gross–Siebert program [8] is a powerful algebro-geometric program to construct a mirror Calabi–Yau manifold from a given toric degeneration of a Calabi–Yau manifold, which serves as a maximal degeneration of a Calabi–Yau manifold. However, it

is not yet clear how we should make use of the mirror of a more general degeneration limit of a Calabi–Yau manifold. An expectation is that a mirror Calabi–Yau manifold should come equipped with a map W , which could be obtained by *gluing* (or taking the *linear system* of) the mirror Landau–Ginzburg superpotentials of the irreducible components of the degeneration limit. Then the glueability condition (the *weight* of the superpotentials being the same) should be mirror to a Kawamata–Namikawa type log smoothability condition. In general W will not provide a polarization with the mirror manifold (see also [6, Section 5.1]). For example it is the case if the number of the irreducible components is less than the dimension of the Calabi–Yau manifold plus 1.

Lastly, the DHT conjecture and the above speculation can be investigated from many different perspectives and it would be of interest to ask, for example, what they mean in the Lagrangian Floer theory and theoretical physics.

Acknowledgements

The author would like to thank Yu-Wei Fan, Andrew Harder, Hansol Hong and Siu-Cheong Lau for useful conversations on related topics. Special thanks go to the referees for their valuable comments and improvements to this article. This research was supported by the Kyoto University Hakubi Project. Part of this work was carried out during the author’s stay at BIRS in the fall of 2016.

References

- [1] Aspinwall P.S., Bridgeland T., Craw A., Douglas M.R., Gross M., Kapustin A., Moore G.W., Segal G., Szendrői B., Wilson P.M.H., Dirichlet branes and mirror symmetry, *Clay Mathematics Monographs*, Vol. 4, Amer. Math. Soc., Providence, RI, Clay Mathematics Institute, Cambridge, MA, 2009.
- [2] Auroux D., Mirror symmetry and T -duality in the complement of an anticanonical divisor, *J. Gökova Geom. Topol. GGT* **1** (2007), 51–91, [arXiv:0706.3207](#).
- [3] Auroux D., Special Lagrangian fibrations, mirror symmetry and Calabi–Yau double covers, *Astérisque* **321** (2008), 99–128, [arXiv:0803.2734](#).
- [4] Cho C.-H., Oh Y.-G., Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds, *Asian J. Math.* **10** (2006), 773–814, [math.SG/0308225](#).
- [5] Dolgachev I.V., Mirror symmetry for lattice polarized $K3$ surfaces, *J. Math. Sci.* **81** (1996), 2599–2630, [alg-geom/9502005](#).
- [6] Doran C.F., Harder A., Thompson A., Mirror symmetry, Tyurin degenerations, and fibrations on Calabi–Yau manifolds, to appear in Proceedings of String-Math 2015, [arXiv:1601.08110](#).
- [7] Fukaya K., Oh Y.-G., Ohta H., Ono K., Lagrangian intersection Floer theory: anomaly and obstruction, Part I, Part II, *AMS/IP Studies in Advanced Mathematics*, Vol. 46, Amer. Math. Soc., Providence, RI, International Press, Somerville, MA, 2009.
- [8] Gross M., Siebert B., From real affine geometry to complex geometry, *Ann. of Math.* **174** (2011), 1301–1428, [math.AG/0703822](#).
- [9] Harder A., The geometry of Landau–Ginzburg models, Ph.D. Thesis, University of Alberta, 2016.
- [10] Hitchin N.J., The moduli space of special Lagrangian submanifolds, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **25** (1997), 503–515, [dg-ga/9711002](#).
- [11] Hori K., Vafa C., Mirror symmetry, [hep-th/0002222](#).
- [12] Kanazawa A., Doran–Harder–Thompson conjecture via integral affine geometry, in preparation.
- [13] Kanazawa A., Lau S.-C., Calabi–Yau manifolds of \tilde{A} and open Yau–Zaslow formula via SYZ mirror symmetry, [arXiv:1605.00342](#).
- [14] Katzarkov L., Kontsevich M., Pantev T., Bogomolov–Tian–Todorov theorems for Landau–Ginzburg models, *J. Differential Geom.* **105** (2017), 55–117, [arXiv:1409.5996](#).

-
- [15] Kawamata Y., Namikawa Y., Logarithmic deformations of normal crossing varieties and smoothing of degenerate Calabi–Yau varieties, *Invent. Math.* **118** (1994), 395–409.
 - [16] Kontsevich M., Homological algebra of mirror symmetry, in Proceedings of the International Congress of Mathematicians, Vols. 1, 2 (Zürich, 1994), Birkhäuser, Basel, 1995, 120–139, [alg-geom/9411018](#).
 - [17] Leung N.C., Mirror symmetry without corrections, *Comm. Anal. Geom.* **13** (2005), 287–331, [math.DG/0009235](#).
 - [18] McLean R.C., Deformations of calibrated submanifolds, *Comm. Anal. Geom.* **6** (1998), 705–747.
 - [19] Seidel P., More about vanishing cycles and mutation, in Symplectic Geometry and Mirror Symmetry (Seoul, 2000), *World Sci. Publ.*, River Edge, NJ, 2001, 429–465, [math.SG/0010032](#).
 - [20] Strominger A., Yau S.-T., Zaslow E., Mirror symmetry is T -duality, *Nuclear Phys. B* **479** (1996), 243–259, [hep-th/9606040](#).
 - [21] Tyurin A.N., Fano versus Calabi–Yau, in The Fano Conference, University Torino, Turin, 2004, 701–734, [math.AG/0302101](#).