

# A Note on Convergence of Level Sets

F. Camilli

**Abstract.** Given a sequence of functions  $f_n$  converging in some topology to a function  $f$ , in general the 0-level set of  $f_n$  does not give a good approximation of the one of  $f$ . In this paper we show that, if we consider an appropriate perturbation of the 0-level set of  $f_n$ , we get a sequence of sets converging to the 0-level set of  $f$ , where the type of set convergence depends on the type of convergence of  $f_n$  to  $f$ .

**Keywords:** *Perturbed level sets, set convergence, capacity*

**AMS subject classification:** 28 A 12, 46 E 35

## 1. Introduction

In several fields (phase transition, free boundary problems, front propagation, etc.), a set of interest for the solution of the problem is represented by a level or a sublevel set of a function  $f$ . Let us suppose that by means of some approximation technique (f.e. discretization, regularization, rescaling of an order parameter) we get a sequence of functions converging in some topology to  $f$ . In general, no matter how strong is the convergence of  $f_n$  to  $f$ , the level sets of  $f_n$  do not give a good approximation of the ones of  $f$ .

Pursuing an idea used in Baiocchi and Pozzi [1], we show that appropriately perturbing the level sets of  $f_n$  (the same can be done for the sublevels or the superlevels), we get a sequence of sets defined by means of  $f_n$  converging to the level set of  $f$ . The type of set convergence is the convergence to zero of the measure of the symmetric difference between the level set of  $f_n$  and the corresponding one of  $f$ , and the measure depends on the type of convergence of the sequence  $f_n$ .

We analyze the case of convergence in  $L^p$  and in  $W^{1,p}$ , but this technique could be useful in other situations.

The paper is organized as follows. In Section 2, we analyze the case of convergence in  $L^\infty$  and  $W^{1,\infty}$  and the associated convergence of perturbed level sets in set-theoretical sense. In Section 3 we first consider the case of convergence in  $L^p$ , which gives the convergence in the sense of Lebesgue measure. Then we analyze the case of convergence in  $W^{1,p}$  and the corresponding set convergence in the sense of capacity and Hausdorff measure.

---

F. Camilli: Univ. di Torino, Dip. di Matematica, Via Carlo Alberto 10, 10123 Torino (Italy)  
New address: Univ. dell' Aquila, Dip. di Energetica, 67040 Roio Poggio (AQ), Italy  
e-mail: camilli@ing.univaq.it

## 2. The case $p = \infty$

In this section we will study (extending the result given in [1]) the case of the convergence in  $L^\infty$ . We will see that the natural set convergence associated to the  $L^\infty$  convergence is the convergence in set-theoretical sense.

**Definition 2.1.** Given a sequence of sets  $\{A_n\}_{n \in \mathbb{N}}$ , we set

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{n=1}^{\infty} \bigcup_{m=n}^{\infty} A_m \quad \text{and} \quad \liminf_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} \bigcap_{m=n}^{\infty} A_m.$$

We say that  $\{A_n\}_{n \in \mathbb{N}}$  converges to  $A$  in *set-theoretical sense* and write  $A = \lim_{n \rightarrow \infty} A_n$  if

$$A = \limsup_{n \rightarrow \infty} A_n = \liminf_{n \rightarrow \infty} A_n.$$

We have the following result.

**Proposition 2.1.** *Let  $f_n$  and  $f$  be continuous functions on  $\mathbb{R}^N$  such that*

$$\|f - f_n\|_{L^\infty(\mathbb{R}^N)} = \varepsilon_n \tag{2.1}$$

where  $\varepsilon_n \rightarrow 0$  for  $n \rightarrow \infty$ . Let  $\{\delta_n\}_{n \in \mathbb{N}}$  be a sequence such that

$$\left. \begin{array}{l} \delta_n > 0 \quad (n \in \mathbb{N}) \\ \delta_n \rightarrow 0 \quad (n \rightarrow \infty) \\ \frac{\varepsilon_n}{\delta_n} \rightarrow 0 \quad (n \rightarrow \infty). \end{array} \right\} \tag{2.2}$$

Set, for any  $n \in \mathbb{N}$ ,

$$\begin{aligned} \Gamma &= \{x \in \mathbb{R}^N : f(x) = 0\} \\ \Gamma_n &= \{x \in \mathbb{R}^N : |f_n(x)| \leq \delta_n\}. \end{aligned} \tag{2.3}$$

Then  $\Gamma \subset \Gamma_n$ , for  $n$  sufficiently large, and

$$\Gamma = \lim_{n \rightarrow \infty} \Gamma_n. \tag{2.4}$$

**Proof.** Let  $\bar{n} \in \mathbb{N}$  be such that  $\delta_n \geq \varepsilon_n$  for any  $n \geq \bar{n}$  (recall that  $\frac{\varepsilon_n}{\delta_n} \rightarrow 0$ ). If  $x \in \Gamma$ , then, for  $n \geq \bar{n}$ , we have from (2.1)

$$|f_n(x)| \leq |f(x)| + \|f_n - f\|_{L^\infty(\mathbb{R}^N)} = \varepsilon_n \leq \delta_n,$$

hence  $x \in \Gamma_n$ . Hence  $\Gamma \subset \Gamma_n$  for  $n \geq \bar{n}$  and therefore  $\Gamma \subset \liminf_{n \rightarrow \infty} \Gamma_n$ . Let us prove yet that  $\limsup_{n \rightarrow \infty} \Gamma_n \subset \Gamma$ . If  $x \in \limsup_{n \rightarrow \infty} \Gamma_n$ , then by definition there exists a subsequence  $\{\Gamma_{n_k}\}_{k \geq 1}$  such that  $x \in \Gamma_{n_k}$  for any  $k \in \mathbb{N}$ . It follows that  $|f_{n_k}(x)| \leq \delta_{n_k}$  for any  $k \in \mathbb{N}$  and therefore  $f(x) = \lim_{k \rightarrow \infty} f_{n_k}(x) = 0$  which yields  $x \in \Gamma$  ■

**Remark 2.1** Observe that if  $\Gamma_n$  and  $\Gamma$  are contained in a compact set  $K$ , then the previous proposition gives the convergence to zero of the Hausdorff distance between  $\Gamma_n$  and  $\Gamma$ .

In the next proposition we show that improving the convergence of  $f_n$  to  $f$ , we get some additional information on the type of convergence of  $\Gamma_n$  to  $\Gamma$ .

**Proposition 2.2.** *Let  $f, f_n \in C^1(\mathbb{R}^N)$  ( $n \in \mathbb{N}$ ) be such that*

$$\|f - f_n\|_{W^{1,\infty}(\mathbb{R}^N)} = \varepsilon_n$$

where  $\varepsilon_n \rightarrow 0$  for  $n \rightarrow \infty$ . Let  $\delta_n$  and  $\Gamma$  and  $\Gamma_n$  be defined as in (2.2) – (2.3). Set

$$\Gamma^{reg} = \left\{ x \in \mathbb{R}^N : f(x) = 0 \text{ and } \nabla f(x) \neq 0 \right\}$$

$$\Gamma^{sing} = \left\{ x \in \mathbb{R}^N : f(x) = 0 \text{ and } \nabla f(x) = 0 \right\}$$

and

$$\Gamma_n^{reg} = \left\{ x \in \mathbb{R}^N : |f_n(x)| \leq \delta_n \text{ and } |\nabla f_n(x)| > \delta_n \right\}$$

$$\Gamma_n^{sing} = \left\{ x \in \mathbb{R}^N : |f_n(x)| \leq \delta_n \text{ and } |\nabla f_n(x)| \leq \delta_n \right\}.$$

Then

$$\Gamma^{reg} = \lim_{n \rightarrow \infty} \Gamma_n^{reg} \quad \text{and} \quad \Gamma^{sing} = \lim_{n \rightarrow \infty} \Gamma_n^{sing}.$$

**Proof.** Let  $\bar{n} \in \mathbb{N}$  be such that  $\delta_n \geq \varepsilon_n$  for  $n \geq \bar{n}$ . Then, for  $n \geq \bar{n}$ ,  $\Gamma \subset \Gamma_n$  and, if  $x \in \Gamma^{sing}$ , we have

$$|\nabla f_n(x)| \leq |\nabla f(x)| + \|\nabla f_n - \nabla f\|_{L^\infty(\mathbb{R}^N)} = \varepsilon_n \leq \delta_n.$$

Therefore  $\Gamma^{sing} \subset \Gamma_n^{sing}$  for  $n \geq \bar{n}$ . If  $x \in \limsup_{n \rightarrow \infty} \Gamma_n^{sing}$ , then  $x \in \Gamma_{n_k}^{sing}$  for a subsequence  $\Gamma_{n_k}$ . It follows that  $|f_{n_k}(x)| \leq \delta_{n_k}$  and  $|\nabla f_{n_k}(x)| \leq \delta_{n_k}$  for any  $k \in \mathbb{N}$  and therefore

$$f(x) = \lim_{k \rightarrow \infty} f_{n_k}(x) = 0 \quad \text{and} \quad \nabla f(x) = \lim_{k \rightarrow \infty} \nabla f_{n_k}(x) = 0.$$

Therefore  $x \in \Gamma^{sing}$  and  $\Gamma^{sing} = \lim_{n \rightarrow \infty} \Gamma_n^{sing}$ . Since (2.4) holds, we get also  $\Gamma^{reg} = \lim_{n \rightarrow \infty} \Gamma_n^{reg}$  ■

We conclude this section giving an estimate of the Hausdorff distance between  $\Gamma$  and  $\Gamma_n$  in the case that  $\Gamma$  is regular.

**Proposition 2.3.** *Assume the same hypothesis as in Proposition 2.1, with  $\delta_n$  and  $\Gamma, \Gamma_n$  defined as in (2.2) – (2.3). Moreover, assume that  $\Gamma$  is compact and that  $f$  is differentiable with  $\nabla f \neq 0$  on  $\Gamma$ . Then there exists a constant  $C > 0$  such that*

$$d_{\mathcal{H}}(\Gamma, \Gamma_n) \leq C(\varepsilon_n + \delta_n) \tag{2.5}$$

for  $n$  sufficiently large, where  $d_{\mathcal{H}}$  denotes the Hausdorff distance.

**Proof.** By the assumptions on  $f$  and  $\Gamma$ , there exist  $\eta_0 > 0$  and  $C_0 > 0$  such that  $|\nabla f(x)| \geq C_0$  on  $\Gamma_{\eta_0} = \{x : d(x, \Gamma) \leq \eta_0\}$ . For  $\eta \leq \eta_0$ , consider  $y \in \partial(\Gamma_\eta) = \partial\{x : d(x, \Gamma) \leq \eta\}$  and let  $x \in \Gamma$  be such that  $d(y, \Gamma) = |y - x| = \eta$ . Then

$$|(y - x) \cdot \nabla f(x)| = \eta |\nabla f(x)| \geq C_0 \eta.$$

Since  $f(x) = 0$ , if  $\omega$  is a modulus of continuity of  $\nabla f$  on  $\Gamma_{\eta_0}$ , then

$$|f(y)| \geq |(y - x) \cdot \nabla f(x)| - \omega(|y - x|)|y - x| \geq \eta(C_0 - \omega(\eta)). \tag{2.6}$$

For  $n$  sufficiently large in such a way that  $C_0 - \omega(\delta_n + \varepsilon_n) \geq \frac{C_0}{2}$  and  $2\frac{\delta_n + \varepsilon_n}{C_0} \leq \eta_0$ , from (2.6) with  $\eta = 2\frac{\delta_n + \varepsilon_n}{C_0}$  we get  $|f(y)| \geq \delta_n + \varepsilon_n$  and therefore  $|f_n(y)| \geq \delta_n$  on  $\partial\Gamma_\eta$ . It follows that  $\Gamma_n \subset \Gamma_\eta$ . Since  $\Gamma \subset \Gamma_n$  for  $n$  sufficiently large, we finally get  $d_{\mathcal{H}}(\Gamma, \Gamma_n) \leq d_{\mathcal{H}}(\Gamma, \Gamma_\eta) \leq \eta$  and therefore (2.5), with  $C = \frac{2}{C_0}$  ■

All the results of this section have an analogue in the case of sub- and superlevel sets of  $f_n$  and  $f$ .

### 3. The case $1 \leq p < \infty$

We first analyze the case of convergence in  $L^p(\mathbb{R}^N)$ . We prove that in this case an appropriate notion of set convergence is the convergence to 0 of the Lebesgue measure of  $\Gamma \Delta \Gamma_n$ . In the following,  $\mathcal{L}^N$  denotes the Lebesgue measure on  $\mathbb{R}^N$ .

**Proposition 3.1.** *Let  $f_n, f \in L^p(\mathbb{R}^N)$  ( $1 \leq p < \infty$ ;  $n \in \mathbb{N}$ ) such that*

$$\|f - f_n\|_{L^p(\mathbb{R}^N)} = \varepsilon_n \quad (3.1)$$

where  $\varepsilon_n \rightarrow 0$  for  $n \rightarrow \infty$ . Let  $\{\delta_n\}_{n \in \mathbb{N}}$  be a sequence such that

$$0 < \delta_n \quad (n \in \mathbb{N}) \quad \text{and} \quad \delta_n \rightarrow 0 \quad (n \rightarrow \infty). \quad (3.2)$$

Define, for any  $n \in \mathbb{N}$ ,

$$\begin{aligned} \Gamma &= \{x \in \mathbb{R}^N : f(x) = 0\} \\ \Gamma_n &= \{x \in \mathbb{R}^N : |f_n(x)| \leq \delta_n\}. \end{aligned} \quad (3.3)$$

Then:

(i) If  $\lim_{n \rightarrow \infty} \frac{\varepsilon_n}{\delta_n} = 0$ , we have

$$\lim_{n \rightarrow \infty} \mathcal{L}^N(\Gamma \Delta \Gamma_n) = 0 \quad (3.4)$$

$$\mathcal{L}^N\left(\Gamma \Delta \limsup_{n \rightarrow \infty} \Gamma_n\right) = 0. \quad (3.5)$$

(ii) If

$$\sum_n \left(\frac{\varepsilon_n}{\delta_n}\right)^p < \infty, \quad (3.6)$$

we also have

$$\mathcal{L}^N\left(\Gamma \Delta \liminf_{n \rightarrow \infty} \Gamma_n\right) = 0. \quad (3.7)$$

Therefore  $\Gamma = \lim_{n \rightarrow \infty} \Gamma_n$  up to a set of 0-Lebesgue measure.

**Proof.** We first observe that, since we are considering only the measure of  $\Gamma$  and  $\Gamma_n$ , we can assume that these sets are defined by means of any element in the class of equivalence of  $f$  and  $f_n$ . We have

$$\Gamma \Delta \Gamma_n = (\Gamma \setminus \Gamma_n) \cup (\Gamma_n \setminus \Gamma)$$

and

$$\begin{aligned} \Gamma \setminus \Gamma_n &= \left\{x \in \mathbb{R}^N : f(x) = 0 \text{ and } |f_n(x)| > \delta_n\right\} \\ \Gamma_n \setminus \Gamma &= \left\{x \in \mathbb{R}^N : f(x) \neq 0 \text{ and } |f_n(x)| \leq \delta_n\right\} \end{aligned}$$

(the previous and all the others inclusions in this proof are intended up to sets of null Lebesgue measure).

Since  $\Gamma \subset \Gamma_n \subset \{x \in \mathbb{R}^N : |f(x) - f_n(x)| > \delta_n\}$ , from the Cebycev inequality we get

$$\mathcal{L}^N(\Gamma \setminus \Gamma_n) \leq \frac{1}{\delta_n^p} \int_{\mathbb{R}^N} |f(x) - f_n(x)|^p dx = \left(\frac{\varepsilon_n}{\delta_n}\right)^p \quad (3.8)$$

and therefore

$$\lim_{n \rightarrow \infty} \mathcal{L}^N(\Gamma \setminus \Gamma_n) = 0. \quad (3.9)$$

Let us prove that

$$\mathcal{L}^N\left(\limsup_{n \rightarrow \infty}(\Gamma_n \setminus \Gamma)\right) = 0. \quad (3.10)$$

Set  $\tilde{\Gamma} = \limsup_{n \rightarrow \infty}(\Gamma_n \setminus \Gamma)$  and let  $x \in \tilde{\Gamma}$ . Then there exists a subsequence  $\{n_k\}_{k \geq 1}$  such that  $|f_{n_k}(x)| \leq \delta_{n_k}$  for any  $k \in \mathbb{N}$ . It follows that,  $\mathcal{L}^N$ -a.e. on  $\tilde{\Gamma}$ ,

$$\liminf_{n \rightarrow \infty} f_n(x) = 0 \quad \text{and} \quad \liminf_{n \rightarrow \infty} |f_n(x) - f(x)|^p = |f(x)|^p.$$

Applying the Fatou Lemma we get

$$\int_{\tilde{\Gamma}} |f(x)|^p dx \leq \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} |f(x) - f_n(x)|^p dx = 0.$$

Since  $|f(x)| > 0$  on  $\tilde{\Gamma}$ , we get (3.10).

Since for any sequence  $\{A_n\}_{n \geq 1}$  of measurable sets we have

$$\limsup_{n \rightarrow \infty} \mathcal{L}^N(A_n) \leq \mathcal{L}^N\left(\limsup_{n \rightarrow \infty} A_n\right)$$

it follows from (3.10) that  $\lim_{n \rightarrow \infty} \mathcal{L}^N(\Gamma_n \setminus \Gamma) = 0$  and therefore, together with (3.9), also (3.4) holds. From (3.10) and

$$\mathcal{L}^N\left(\Gamma \setminus \limsup_{n \rightarrow \infty} \Gamma_n\right) = \mathcal{L}^N\left(\liminf_{n \rightarrow \infty}(\Gamma \setminus \Gamma_n)\right) \leq \lim_{n \rightarrow \infty} \mathcal{L}^N(\Gamma \setminus \Gamma_n) = 0$$

we get (3.5).

Let us prove now statement (ii). Estimate (3.8) gives

$$\mathcal{L}^N\left(\bigcup_{m=n}^{\infty} (\Gamma \setminus \Gamma_m)\right) \leq \sum_{m=n}^{\infty} \left(\frac{\varepsilon_m}{\delta_m}\right)^p,$$

and therefore, for any  $n \in \mathbb{N}$ ,

$$\mathcal{L}^N\left(\Gamma \setminus \liminf_{n \rightarrow \infty} \Gamma_n\right) = \mathcal{L}^N\left(\limsup_{n \rightarrow \infty}(\Gamma \setminus \Gamma_n)\right) \leq \sum_{m=n}^{\infty} \left(\frac{\varepsilon_m}{\delta_m}\right)^p.$$

For (3.6) we get  $\mathcal{L}^N(\Gamma \setminus \liminf_{n \rightarrow \infty} \Gamma_n) = 0$ . Since (3.10) yields  $\mathcal{L}^N(\liminf_{n \rightarrow \infty} \Gamma_n \setminus \Gamma) = 0$  we get (3.7) ■

**Remark 3.1.** Since we have

$$|\mathcal{L}^N(\Gamma) - \mathcal{L}^N(\Gamma_n)| \leq \mathcal{L}^N(\Gamma \Delta \Gamma_n)$$

then  $\mathcal{L}^N(\Gamma \Delta \Gamma_n) \rightarrow 0$  implies that  $\mathcal{L}^N(\Gamma_n) \rightarrow \mathcal{L}^N(\Gamma)$ . The vice versa in general is not true. The result  $\mathcal{L}^N(\Gamma \Delta \Gamma_n) \rightarrow 0$  gives a more complete information respect to the convergence of the measure of  $\Gamma_n$  to the measure of  $\Gamma$ . In fact, it shows that the measure of the part of  $\Gamma_n$  which does not approximate  $\Gamma$  tends to 0, while the measure of  $\Gamma \setminus \Gamma_n$  can be estimated by means of (3.8).

If we know that  $\|f_n - f\|_{W^{1,p}(\mathbb{R}^N)} = \varepsilon_n$ , then we can prove a result similar to Proposition 2.2 for the convergence of regular and singular parts of  $\Gamma_n$  to  $\Gamma$ . In this case, a more accurate way of studying properties of sets defined through Sobolev functions is given by the notion of *capacity*. We will show that, in the case of convergence in  $W^{1,p}(\mathbb{R}^N)$  ( $1 \leq p < N$ ) we get convergence of  $\Gamma_n$  to  $\Gamma$  up to sets of 0 capacity. Let us recall the definition and some basic properties of the capacity we will need in the following (see [2 - 4] for more details).

**Definition 3.1.** Let  $1 \leq p < N$  and set

$$K^p = \left\{ \varphi : \mathbb{R}^N \rightarrow \mathbb{R} \mid 0 \leq \varphi \in L^{p^*}(\mathbb{R}^N) \text{ with } \nabla \varphi \in L^p(\mathbb{R}^N, \mathbb{R}^N) \right\}$$

where  $p^* = \frac{Np}{N-p}$ . For  $A \subset \mathbb{R}^N$ , we define

$$\text{Cap}_p(A) = \inf \left\{ \int_{\mathbb{R}^N} |\nabla \varphi| dy \mid \varphi \in K^p \text{ with } A \subset \{\varphi \geq 1\}^\circ \right\}.$$

It is possible to prove that  $\text{Cap}_p$  is an exterior measure on subsets of  $\mathbb{R}^N$ . For a function  $\varphi \in L^1_{loc}(\mathbb{R}^N)$ , the *precise representative*  $\varphi^*$  of  $\varphi$  is defined by

$$\varphi^*(x) = \begin{cases} \lim_{r \rightarrow 0^+} \int_{B(x,r)} \varphi(y) dy & \text{if the limit exists} \\ 0 & \text{otherwise} \end{cases}$$

where  $\int_{B(x,r)} \varphi(y) dy = \int_{B(x,r)} \varphi(y) dy / \mathcal{L}^N(B(x,r))$ . We have (see [2: Theorem 4.8.1]) the following

**Theorem 3.1.** Let  $\varphi \in W^{1,p}(\mathbb{R}^N)$  ( $1 \leq p < N$ ). Then:

(i) There is a Borel set  $E \subset \mathbb{R}^N$  such that  $\text{Cap}_p(E) = 0$  and

$$\lim_{r \rightarrow 0^+} \int_{B(x,r)} \varphi(y) dy = \varphi^*(x) \quad (x \in \mathbb{R}^N \setminus E).$$

(ii) In addition,

$$\lim_{r \rightarrow 0^+} \int_{B(x,r)} |\varphi(y) - \varphi^*(x)|^{p^*} dy = 0 \quad (x \in \mathbb{R}^N \setminus E).$$

(iii) The precise representative  $\varphi^*$  is quasi-continuous.

Because of the previous theorem, any function in the space  $W^{1,p}(\mathbb{R}^N)$  admits a quasi-continuous representative. We have the following convergence result for the perturbed level sets.

**Proposition 3.2.** *Let  $f, f_n \in W^{1,p}(\mathbb{R}^N)$  ( $1 \leq p < N$ ) be such that*

$$\|f - f_n\|_{W^{1,p}(\mathbb{R}^N)} = \varepsilon_n$$

where  $\varepsilon_n \rightarrow 0$  for  $n \rightarrow \infty$ . Let  $\delta_n$  and  $\Gamma, \Gamma_n$  be defined as in (3.2) – (3.3) by means of the precise representatives of  $f$  and  $f_n$ . Then:

(i) *If  $\lim_{n \rightarrow \infty} \frac{\varepsilon_n}{\delta_n} = 0$ , then*

$$\text{Cap}_p \left( \limsup_{n \rightarrow \infty} \Gamma_n \Delta \Gamma \right) = 0. \quad (3.11)$$

(ii) *If*

$$\sum_n \left( \frac{\varepsilon_n}{\delta_n} \right)^p < \infty, \quad (3.12)$$

then we have also

$$\text{Cap}_p \left( \Gamma \Delta \liminf_{n \rightarrow \infty} \Gamma_n \right) = 0 \quad (3.13)$$

and therefore  $\Gamma = \lim_{n \rightarrow \infty} \Gamma_n$  up to a set of zero capacity.

**Proof.** Let us prove statement (i). Since the sets  $\Gamma$  and  $\Gamma_n$  are defined by means of the precise representatives of  $f$  and  $f_n$ , then they are well defined, i.e. up to sets of zero capacity. In the following all the relations involving  $\Gamma$  and  $\Gamma_n$  are intended to be satisfied  $\text{Cap}_p$ -a.e. We have

$$\Gamma \setminus \Gamma_n = \left\{ x \in \mathbb{R}^N : f(x) = 0 \text{ and } |f_n(x)| > \delta_n \right\}.$$

Let us prove that, defining

$$B_n = \left\{ x \in \mathbb{R}^N \mid \int_{B(x,r)} |f_n - f| dy > \delta_n \text{ for some } r > 0 \right\}, \quad (3.14)$$

then

$$\text{Cap}_p(\Gamma \setminus \Gamma_n) \leq \text{Cap}_p(B_n). \quad (3.15)$$

In fact, if  $x \in \Gamma \setminus \Gamma_n$ , then, up to a set of zero capacity, we have

$$\lim_{r \rightarrow 0} \int_{B(x,r)} |f_n - f| dy = |f(x) - f_n(x)| > \delta_n.$$

Therefore there exists  $r_0 > 0$  such that  $\int_{B(x,r_0)} |f_n - f| dy > \delta_n$  and so (3.15) holds. Recall that (see [2: Lemma 4.8.1]), if  $\varphi \in K^p$ , then there exists a constant  $C$ , depending only on  $N$  and  $p$ , such that for any  $\eta > 0$

$$\text{Cap}_p \left( \left\{ x \in \mathbb{R}^N \mid \int_{B(x,r)} \varphi(y) dy > \eta \text{ for some } r > 0 \right\} \right) \leq \frac{C}{\eta^p} \int_{\mathbb{R}^N} |D\varphi|^p dy. \quad (3.16)$$

From (3.14) and (3.16) we get

$$\text{Cap}_p(\Gamma \setminus \Gamma_n) \leq \frac{C}{\delta_n^p} \int_{\mathbb{R}^N} |\nabla f - \nabla f_n|^p dy \leq C \left( \frac{\varepsilon_n}{\delta_n} \right)^p \quad (3.17)$$

and therefore  $\lim_{n \rightarrow \infty} \text{Cap}_p(\Gamma \setminus \Gamma_n) = 0$ . From the previous equality and the properties of the capacity, we get

$$\text{Cap}_p\left(\Gamma \setminus \limsup_{n \rightarrow \infty} \Gamma_n\right) = \text{Cap}_p\left(\liminf_{n \rightarrow \infty} (\Gamma \setminus \Gamma_n)\right) \leq \lim_{n \rightarrow \infty} \text{Cap}_p(\Gamma \setminus \Gamma_n) = 0. \quad (3.18)$$

Let  $A$  be the set

$$A = \left\{ x \in \mathbb{R}^N \mid \limsup_{r \rightarrow 0^+} \frac{1}{r^{n-p}} \int_{B(x,r)} |\nabla f(y)|^p dy > 0 \right\}.$$

Then  $\text{Cap}_p(A) = 0$  (see [2: Theorem 2.4.3]) and from the Poincaré inequality we have

$$\lim_{r \rightarrow 0} \int_{B(x,r)} |f(y) - (f)_{x,r}|^p dy = 0 \quad (x \in \mathbb{R}^N \setminus A) \quad (3.19)$$

where  $(f)_{x,r} = \int_{B(x,r)} f(y) dy$ . From Theorem 3.1, for any  $n \in \mathbb{N}$  there exists a Borel set  $E_n$  such that  $\text{Cap}_p(E_n) = 0$  and

$$\lim_{r \rightarrow 0^+} \int_{B(x,r)} |f_n(y) - f_n(x)|^p dy = 0 \quad (x \in \mathbb{R}^N \setminus E_n). \quad (3.20)$$

Set  $\Delta_n = B_n \cup E_n \cup A$ , where  $B_n$  has been defined in (3.14). If  $x \in \Gamma_n \setminus \Delta_n$ , then from Theorem 3.1, (3.14) and (3.19) - (3.20) we get

$$\begin{aligned} \limsup_{r \rightarrow 0} |(f)_{x,r}| &\leq \limsup_{r \rightarrow 0} |(f)_{x,r} - f_n(x)| + \delta_n \\ &\leq \limsup_{r \rightarrow 0} \left\{ \int_{B(x,r)} |f - (f)_{x,r}| dy \right. \\ &\quad \left. + \int_{B(x,r)} |f - f_n| dy + \int_{B(x,r)} |f_n - f_n(x)| dy \right\} + \delta_n \\ &\leq 2\delta_n. \end{aligned} \quad (3.21)$$

Moreover, inequality (3.16) gives

$$\text{Cap}_p(\Delta_n) \leq \text{Cap}_p(B_n) + \text{Cap}_p(E_n) + \text{Cap}_p(A) \leq C \left( \frac{\varepsilon_n}{\delta_n} \right)^p. \quad (3.22)$$

Set  $\Delta = \liminf_{n \rightarrow \infty} \Delta_n$  and  $\tilde{\Gamma} = \limsup_{n \rightarrow \infty} (\Gamma_n \setminus \Gamma)$ . From (3.21) - (3.22) it follows that if  $x \in \tilde{\Gamma} \setminus \Delta$ , then  $\lim_{r \rightarrow 0^+} (f)_{x,r} = 0$ . Therefore from Theorem 3.1 we get  $\tilde{\Gamma} \setminus \Delta \subset \Gamma$  and, since  $\text{Cap}_p(\Delta) \leq \liminf_{n \rightarrow \infty} \text{Cap}_p(\Delta_n) = 0$ , it follows also that

$$\text{Cap}_p\left(\limsup_{n \rightarrow \infty} (\Gamma_n \setminus \Gamma)\right) = 0.$$

The previous equality and (3.18) imply (3.11).

Let us prove statement (ii). If  $x \in \Gamma \setminus B_n$ , then

$$\limsup_{r \rightarrow 0^+} \int_{B(x,r)} |f_n| dy \leq \limsup_{r \rightarrow 0^+} \left( \int_{B(x,r)} |f| dy + \int_{B(x,r)} |f - f_n| dy \right) \leq \delta_n. \quad (3.23)$$

Thus (3.23) yields  $\Gamma \setminus B_n \subset \Gamma_n$  for any  $n$  and therefore

$$\liminf_{n \rightarrow \infty} (\Gamma \setminus B_n) = \Gamma \setminus \limsup_{n \rightarrow \infty} B_n \subset \liminf_{n \rightarrow \infty} \Gamma_n.$$

Set  $B = \limsup_{n \rightarrow \infty} B_n$ . Then, for any  $n \in \mathbb{N}$ ,

$$\text{Cap}_p(B) \leq \sum_{m=n}^{\infty} \text{Cap}_p(B_m) \leq \sum_{m=n}^{\infty} \left( \frac{\varepsilon_m}{\delta_m} \right)^p$$

and, for hypothesis (3.12), we get  $\text{Cap}_p(B) = 0$  and  $(\Gamma \setminus B) \subset \liminf_{n \rightarrow \infty} \Gamma_n$ . From statement (i) we get (3.13) ■

**Remark 3.2.** For the capacity we do not have an analogy of property (3.4). While, as we have proved,  $\lim_{n \rightarrow \infty} \text{Cap}_p(\Gamma \setminus \Gamma_n) = 0$  in general, it is *not* true that  $\lim_{n \rightarrow \infty} \text{Cap}_p(\Gamma_n \setminus \Gamma) = 0$  as it can be easily seen taking  $f_n \equiv f$ .

Taking into account the relation between capacity and Hausdorff measure (see [2, 3]), from the previous proposition we get the following result about convergence in the sense of the Hausdorff measure.

**Corollary 3.1.** *Under the same hypothesis of Proposition 3.2, we have the following:*

(i) *If  $\lim_{n \rightarrow \infty} \frac{\varepsilon_n}{\delta_n} = 0$ , then for any  $\sigma > 0$*

$$\mathcal{H}^{N-p+\sigma} \left( \limsup_{n \rightarrow \infty} \Gamma_n \Delta \Gamma \right) = 0. \quad (3.24)$$

(ii) *If  $\sum_n \left( \frac{\varepsilon_n}{\delta_n} \right)^p < \infty$ , then we also have, for any  $\sigma > 0$ ,*

$$\mathcal{H}^{N-p+\sigma} \left( \Gamma \Delta \liminf_{n \rightarrow \infty} \Gamma_n \right) = 0. \quad (3.25)$$

*If  $p = 1$ , then (3.24) – (3.25) hold also for  $\sigma = 0$ .*

If  $p > N$ , since  $W^{1,p}(\mathbb{R}^N) \subset L^\infty(\mathbb{R}^N)$  with continuous immersion, we can apply the results of Section 2 to the continuous representatives of  $f$  and  $f_n$ . Therefore, from the convergence of  $f_n$  to  $f$  we get the convergence in the set theoretical sense of  $\Gamma_n$  to  $\Gamma$ .

**Acknowledgement.** The author wishes to thank Prof. C. Baiocchi of the University of Roma “La Sapienza” for suggesting the problem.

## References

- [1] Baiocchi, C. and G. A. Pozzi: *Error estimates and free-boundary convergence for a finite difference discretization of a parabolic variational inequality*. RAIRO Anal. Numér. 11 (1977), 315 – 353.
- [2] Evans, L. C. and R. F. Gariepy: *Measure Theory and Fine  $p$  Properties of Functions*. Boca Raton: CRC Press 1992.
- [3] Federer, H. and W. Ziemer: *The Lebesgue set of a function whose distribution derivatives are  $p$ -th power summable*. Indiana Univ. Math. J. 22 (1972), 139 – 158.
- [4] Ziemer, W.: *Weakly Differentiable Functions*. New York: Springer-Verlag 1989.

Received 22.03.1998