

Examples of NOT OK using car package

Kyun-Seop Bae MD PhD

2024-03-25

Contents

1	Tested Version and Books used for the Validation	3
1.1	Packages Used	3
1.2	Books and Articles used for the Test	3
2	Snee EMS ANOVA 1974	4
3	Goodnight	9
3.1	p33	9
4	SAS for Linear Models 4e	11
4.1	p403	11
4.2	p417	14
4.3	p431	15
5	Sahai - Unbalanced	19
5.1	Table 15.3	19
5.2	Table 16.3	23
6	Federer - Variations	27
6.1	Example 2.2	27
6.2	Example 3.1	30
6.3	Example 5.1	39
6.4	Example 7.1	44
6.5	Example 7.3	48
6.6	Example 8.1	54
6.7	Example 9.2	57
6.8	Example 10.1	60
7	Hinkelmann & Kempthorne - Volume 1	74
7.1	p410	74
8	Searle - Linear Models 2e	77
8.1	7.2 (p390, 59%)	77
8.2	7.2 (p393, 60%)	78
9	Web site examples	80

9.1	https://github.com/djnavarro/psyr	80
10	Bioequivalence (BE) data example	82
11	Session Information	84

1 Tested Version and Books used for the Validation

1.1 Packages Used

- ‘sasLM’ version: 0.10.3
- ‘SAS’ version: 9.4 Licensed and University Edition
- ‘car’ version: 3.1.2
- R version: R version 4.3.3 (2024-02-29 ucrt)

The ‘car’ package is not necessary for ‘sasLM.’ It is used for the comparison of the results.

If you see any difference between ‘car’ and ‘sasLM’, ‘SAS’ results coincide with ‘sasLM’, not with ‘car’.

Before ‘sasLM’ is available on CRAN, you can download using the following command in R.

```
install.packages("sasLM", repos="http://r.acr.kr")
```

1.2 Books and Articles used for the Test

1. Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. *J Qual Tech.* 1974;6(3):128-137.
2. Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User’s Group, SAS Institute, Raleigh, N.C. 1976.
3. Littell RC, Stroup WW, Freund RJ. *SAS for Linear Models 4e*. John Wiley & Sons Inc. 2002.
4. Sahai H, Ojeda MM. *Analysis of Variance for Random Models Volume 2 Unbalanced Data*. 2005.
5. Federer WT, King F. *Variations on Split Plot and Split Block Experiment Designs*. John Wiley & Sons Inc. 2007.
6. Hinkelmann K, Kempthorne O. *Design and Analysis of Experiments Volume 1 Introduction to Experimental Design*. 2e. John Wiley & Sons Inc. 2008.
7. Searle SR, Gruber MHJ. *Linear Models 2e*, Kindle Edition. John Wiley & Sons Inc. 2016.

2 Snee EMS ANOVA 1974

Reference

- Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. J Qual Tech. 1974;6(3):128-137.

(1) MODEL

```
Snee = read.csv("http://r.acr.kr/Snee_EMS_ANOVA1974.csv")
Snee = af(Snee, c("Machine", "Analyst", "Test", "Day"))
Snee
```

	Machine	Analyst	Test	Day	Y
1	1	1	1	1	6.1
2	1	1	1	2	8.5
3	1	1	1	3	8.6
4	1	1	1	4	9.3
5	1	1	1	5	8.1
6	1	1	1	6	8.5
7	1	1	1	7	9.8
8	1	1	1	8	9.0
9	1	1	1	9	11.0
10	1	1	1	10	9.7
11	1	1	1	11	10.5
12	1	1	1	12	8.3
13	1	1	1	13	8.4
14	1	1	1	14	10.2
15	1	1	1	15	9.3
16	1	1	1	16	7.1
17	1	1	1	17	5.8
18	1	1	1	18	8.9
19	1	1	1	19	11.5
20	1	1	1	20	10.3
21	1	1	1	21	9.1
22	1	1	1	22	5.7
23	1	1	1	23	8.5
24	1	1	1	24	9.6
25	1	1	1	25	9.4
26	1	1	1	26	10.3
27	1	1	1	27	7.0
28	1	1	1	28	11.5
29	1	1	1	29	6.0
30	1	1	1	30	8.0
31	1	1	1	31	13.4
32	1	1	1	32	12.1
33	1	1	1	33	14.2
34	1	1	1	34	10.0
35	1	1	1	35	6.5
36	1	1	1	36	6.5

37	1	1	1	37	9.2
38	1	1	1	38	11.0
39	1	1	1	39	8.6
40	1	1	1	40	8.9
41	1	1	1	41	6.6
42	1	1	1	42	8.4
43	1	1	2	1	6.6
44	1	1	2	2	9.6
45	1	1	2	3	6.7
46	1	1	2	4	7.2
47	1	1	2	5	7.1
48	1	1	2	6	9.0
49	1	1	2	7	9.8
50	1	1	2	8	8.0
51	1	1	2	9	10.9
52	1	1	2	10	10.6
53	1	1	2	11	8.4
54	1	1	2	12	10.6
55	1	1	2	13	7.2
56	1	1	2	14	8.0
57	1	1	2	15	8.7
58	1	1	2	16	8.7
59	1	1	2	17	6.8
60	1	1	2	18	6.6
61	1	1	2	19	7.1
62	1	1	2	20	10.0
63	1	1	2	21	9.5
64	1	1	2	22	7.7
65	1	1	2	23	8.8
66	1	1	2	24	12.2
67	1	1	2	25	10.4
68	1	1	2	26	10.6
69	1	1	2	27	10.6
70	1	1	2	28	7.3
71	1	1	2	29	7.0
72	1	1	2	30	7.0
73	1	1	2	31	9.2
74	1	1	2	32	11.7
75	1	1	2	33	10.6
76	1	1	2	34	10.4
77	1	1	2	35	8.4
78	1	1	2	36	6.8
79	1	1	2	37	10.1
80	1	1	2	38	11.0
81	1	1	2	39	10.0
82	1	1	2	40	8.0
83	1	1	2	41	7.2
84	1	1	2	42	8.8

85	1	2	1	1	6.6
86	1	2	1	2	8.2
87	1	2	1	3	8.0
88	1	2	1	4	6.5
89	1	2	1	5	2.3
90	1	2	1	6	4.0
91	1	2	1	7	11.7
92	1	2	1	8	6.8
93	1	2	1	9	10.5
94	1	2	1	10	10.3
95	1	2	1	11	10.0
96	1	2	1	12	8.8
97	1	2	1	13	6.7
98	1	2	1	14	8.9
99	1	2	1	15	9.9
100	1	2	1	16	8.2
101	1	2	1	17	7.5
102	1	2	1	18	6.6
103	1	2	1	19	3.1
104	1	2	1	20	7.2
105	1	2	1	21	10.7
106	1	2	1	22	8.4
107	1	2	1	23	7.6
108	1	2	1	24	12.6
109	1	2	1	25	9.6
110	1	2	1	26	12.6
111	1	2	1	27	10.8
112	1	2	1	28	5.1
113	1	2	1	29	6.6
114	1	2	1	30	8.6
115	1	2	1	31	12.5
116	1	2	1	32	10.4
117	1	2	1	33	10.6
118	1	2	1	34	7.2
119	1	2	1	35	7.8
120	1	2	1	36	4.4
121	1	2	1	37	8.7
122	1	2	1	38	11.2
123	1	2	1	39	10.3
124	1	2	1	40	7.0
125	1	2	1	41	7.7
126	1	2	1	42	7.6
127	2	1	1	1	8.8
128	2	1	1	2	8.1
129	2	1	1	3	7.4
130	2	1	1	4	8.0
131	2	1	1	5	9.5
132	2	1	1	6	9.2

133	2	1	1	7	12.8
134	2	1	1	8	9.2
135	2	1	1	9	11.3
136	2	1	1	10	9.3
137	2	1	1	11	4.0
138	2	1	1	12	9.7
139	2	1	1	13	4.6
140	2	1	1	14	2.1
141	2	1	1	15	9.7
142	2	1	1	16	10.0
143	2	1	1	17	10.2
144	2	1	1	18	9.2
145	2	1	1	19	10.8
146	2	1	1	20	9.4
147	2	1	1	21	10.3
148	2	1	1	22	10.3
149	2	1	1	23	8.3
150	2	1	1	24	11.6
151	2	1	1	25	9.4
152	2	1	1	26	11.3
153	2	1	1	27	11.4
154	2	1	1	28	9.6
155	2	1	1	29	2.2
156	2	1	1	30	6.6
157	2	1	1	31	11.5
158	2	1	1	32	9.1
159	2	1	1	33	4.6
160	2	1	1	34	7.9
161	2	1	1	35	9.0
162	2	1	1	36	8.1
163	2	1	1	37	9.4
164	2	1	1	38	10.9
165	2	1	1	39	9.0
166	2	1	1	40	7.8
167	2	1	1	41	9.3
168	2	1	1	42	6.8

```
GLM(Y ~ Day/Machine/Analyst/Test, Snee)
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      167 751.27  4.4986
RESIDUALS   0    0.00
CORRECTED TOTAL 167 751.27
```

```
$Fitness
Root MSE    Y Mean Coef Var R-square
```

```

NA 8.736905      NA      1

$`Type I`
    Df Sum Sq Mean Sq F value Pr(>F)
Day           41 365.58 8.9166
Day:Machine   42 196.59 4.6807
Day:Machine:Analyst 42 118.80 2.8285
Day:Machine:Analyst:Test 42 70.30 1.6739

$`Type II`
    Df Sum Sq Mean Sq F value Pr(>F)
Day           41 365.58 8.9166
Day:Machine   42 196.59 4.6807
Day:Machine:Analyst 42 118.80 2.8285
Day:Machine:Analyst:Test 42 70.30 1.6739

$`Type III`
    Df Sum Sq Mean Sq F value Pr(>F)
Day           41 359.44 8.7669
Day:Machine   42 199.40 4.7477
Day:Machine:Analyst 42 118.80 2.8285
Day:Machine:Analyst:Test 42 70.30 1.6739

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Day/Machine/Analyst/Test, Snee), type=3, singular.ok=TRUE)
# NOT WORKING

```

3 Goodnight

Reference

- Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User's Group, SAS Institute, Raleigh, N.C. 1976.

3.1 p33

(2) MODEL

```
p33 = read.csv("http://r.acr.kr/Goodnight-p33.csv")
p33 = af(p33, c("A", "B"))
p33
```

```
A B      y
1 1 1 2.96
2 1 2 7.90
3 2 1 4.79
4 2 2 9.55
5 3 3 9.53
```

```
GLM(y ~ A + B + A:B, p33) # p35
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      4 34.905 8.7261
RESIDUALS   0  0.000
CORRECTED TOTAL 4 34.905
```

```
$Fitness
Root MSE y Mean Coef Var R-square
      NA  6.946      NA        1
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
A       2 11.3739 5.6870
B       1 23.5225 23.5225
A:B    1  0.0081  0.0081
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
A       1  3.0276  3.0276
B       1 23.5225 23.5225
A:B    1  0.0081  0.0081
```

```
$`Type III`
CAUTION: Singularity Exists !
          Df Sum Sq Mean Sq F value Pr(>F)
```

```
A      1  3.0276  3.0276
B      1 23.5225 23.5225
A:B    1  0.0081  0.0081
```

```
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(y ~ A + B + A:B, p33), type=3, singular.ok=TRUE) # NOT WORKING
```

4 SAS for Linear Models 4e

Reference

- Littell RC, Stroup WW, Freund RJ. SAS for Linear Models 4e. John Wiley & Sons Inc. 2002.

4.1 p403

(3) MODEL

```
p403 = read.table("http://r.acr.kr/sas4lm/p403.txt", header=TRUE)
p403 = af(p403, c("PATIENT", "VISIT"))
p403
```

	PATIENT	SEQUENCE	VISIT	BASEHR	HR	DRUG	RESIDT	RESIDS
1	1	B	2	86	86	placebo	0	0
2	1	B	3	86	106	test	-1	-1
3	1	B	4	62	79	standard	1	0
4	2	F	2	48	66	test	0	0
5	2	F	3	58	56	placebo	1	0
6	2	F	4	74	79	standard	-1	-1
7	3	B	2	78	84	placebo	0	0
8	3	B	3	78	76	test	-1	-1
9	3	B	4	82	91	standard	1	0
10	4	D	2	66	79	standard	0	0
11	4	D	3	72	100	test	0	1
12	4	D	4	90	82	placebo	1	0
13	5	C	2	74	74	test	0	0
14	5	C	3	90	71	standard	1	0
15	5	C	4	66	62	placebo	0	1
16	6	B	2	62	64	placebo	0	0
17	6	B	3	74	90	test	-1	-1
18	6	B	4	58	85	standard	1	0
19	7	A	2	94	75	standard	0	0
20	7	A	3	72	82	placebo	0	1
21	7	A	4	100	102	test	-1	-1
22	8	A	2	54	63	standard	0	0
23	8	A	3	54	58	placebo	0	1
24	8	A	4	66	62	test	-1	-1
25	9	D	2	82	91	standard	0	0
26	9	D	3	96	86	test	0	1
27	9	D	4	78	88	placebo	1	0
28	10	C	2	86	82	test	0	0
29	10	C	3	70	71	standard	1	0
30	10	C	4	58	62	placebo	0	1
31	11	F	2	82	80	test	0	0
32	11	F	3	80	78	placebo	1	0
33	11	F	4	72	75	standard	-1	-1
34	12	E	2	96	90	placebo	0	0

35	12	E	3	92	93	standard	-1	-1
36	12	E	4	82	88	test	0	1
37	13	D	2	78	87	standard	0	0
38	13	D	3	72	80	test	0	1
39	13	D	4	76	78	placebo	1	0
40	14	F	2	98	86	test	0	0
41	14	F	3	86	86	placebo	1	0
42	14	F	4	70	79	standard	-1	-1
43	15	A	2	86	71	standard	0	0
44	15	A	3	66	70	placebo	0	1
45	15	A	4	74	90	test	-1	-1
46	16	E	2	86	86	placebo	0	0
47	16	E	3	90	103	standard	-1	-1
48	16	E	4	82	86	test	0	1
49	17	A	2	66	83	standard	0	0
50	17	A	3	82	86	placebo	0	1
51	17	A	4	86	102	test	-1	-1
52	18	F	2	66	82	test	0	0
53	18	F	3	78	80	placebo	1	0
54	18	F	4	74	95	standard	-1	-1
55	19	E	2	74	80	placebo	0	0
56	19	E	3	78	79	standard	-1	-1
57	19	E	4	70	74	test	0	1
58	20	B	2	66	70	placebo	0	0
59	20	B	3	74	62	test	-1	-1
60	20	B	4	62	67	standard	1	0
61	21	C	2	82	90	test	0	0
62	21	C	3	90	103	standard	1	0
63	21	C	4	76	82	placebo	0	1
64	22	C	2	82	82	test	0	0
65	22	C	3	66	83	standard	1	0
66	22	C	4	90	82	placebo	0	1
67	23	E	2	82	66	placebo	0	0
68	23	E	3	74	87	standard	-1	-1
69	23	E	4	82	82	test	0	1
70	24	D	2	72	75	standard	0	0
71	24	D	3	82	86	test	0	1
72	24	D	4	74	82	placebo	1	0

GLM(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT, p403)

\$ANOVA

Response : HR

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	29	6408.7	220.99	3.912	3.127e-05 ***
RESIDUALS	42	2372.6	56.49		
CORRECTED TOTAL	71	8781.3			

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE  HR Mean Coef Var R-square Adj R-sq
7.515988 80.80556 9.301326 0.7298134 0.543256

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
SEQUENCE      5 508.9 101.79 1.8019 0.133346
SEQUENCE:PATIENT 18 4692.3 260.69 4.6147 2.21e-05 ***
VISIT         2 146.8 73.39 1.2991 0.283499
DRUG          2 668.8 334.39 5.9194 0.005435 **
RESIDS        1 391.0 391.02 6.9219 0.011854 *
RESIDT        1     0.8     0.84 0.0149 0.903511
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
SEQUENCE      5 701.2 140.237 2.4825 0.04665 *
SEQUENCE:PATIENT 18 4692.3 260.685 4.6147 2.21e-05 ***
VISIT         2 146.8 73.389 1.2991 0.28350
DRUG          2 344.0 171.975 3.0443 0.05826 .
RESIDS        1 309.2 309.174 5.4731 0.02414 *
RESIDT        1     0.8     0.840 0.0149 0.90351
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
SEQUENCE      5 701.2 140.237 2.4825 0.04665 *
SEQUENCE:PATIENT 18 4692.3 260.685 4.6147 2.21e-05 ***
VISIT         2 146.8 73.389 1.2991 0.28350
DRUG          2 344.0 171.975 3.0443 0.05826 .
RESIDS        1 309.2 309.174 5.4731 0.02414 *
RESIDT        1     0.8     0.840 0.0149 0.90351
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT,
p403), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: HR

```

          Sum Sq Df F values   Pr(>F)
SEQUENCE      0.0  0
VISIT        146.8  2  1.2991  0.28350
DRUG         343.9  2  3.0443  0.05826 .
RESIDS       309.2  1  5.4731  0.02414 *
RESIDT        0.8  1  0.0149  0.90351
SEQUENCE:PATIENT 4692.3 18  4.6147 2.21e-05 ***
Residuals    2372.6 42
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

4.2 p417

(4) MODEL

```

p417 = read.table("http://r.acr.kr/sas4lm/p417.txt", header=TRUE)
p417 = af(p417, c("TRT", "POT", "PLANT"))
p417

```

	Obs	TRT	POT	PLANT	Y
1	1	1	1	1	15
2	2	1	1	2	13
3	3	1	1	3	16
4	4	1	2	1	17
5	5	1	2	2	19
6	6	1	3	1	12
7	7	2	1	1	20
8	8	2	1	2	21
9	9	2	2	1	20
10	10	2	2	2	23
11	11	2	2	3	19
12	12	2	2	4	19
13	13	3	1	1	12
14	14	3	1	2	13
15	15	3	1	3	14
16	16	3	2	1	11
17	17	3	3	1	12
18	18	3	3	2	13
19	19	3	3	3	15
20	20	3	3	4	11
21	21	3	3	5	9

```

GLM(Y ~ TRT + POT %in% TRT, p417) # p418 Output 11.28

```

```

$ANOVA
Response : Y
          Df  Sum Sq Mean Sq F value   Pr(>F)
MODEL      7 267.226 38.175 12.433 7.522e-05 ***
RESIDUALS 13  39.917  3.071

```

```

CORRECTED TOTAL 20 307.143
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE   Y Mean Coef Var R-square Adj R-sq
1.752288 15.42857 11.35742 0.8700388 0.8000596

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
TRT      2 236.921 118.460 38.580 3.412e-06 ***
TRT:POT  5  30.306   6.061   1.974     0.1499
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
TRT      2 236.921 118.460 38.580 3.412e-06 ***
TRT:POT  5  30.306   6.061   1.974     0.1499
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
TRT      2 200.111 100.055 32.586 8.626e-06 ***
TRT:POT  5  30.306   6.061   1.974     0.1499
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ TRT + POT %in% TRT, p417), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: Y
      Sum Sq Df F values Pr(>F)
TRT      22.310  1    7.266 0.01835 *
TRT:POT  30.306  5    1.974 0.14991
Residuals 39.917 13
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

4.3 p431

(5) MODEL

```

p431 = read.table("http://r.acr.kr/sas4lm/p431.txt", header=TRUE)
p431 = af(p431, c("line", "sire", "agedam", "steerno"))
p431

```

	Obs	line	sire	agedam	steerno	age	intlwt	avdlygn
1	1	1	1	3	1	192	390	2.24
2	2	1	1	3	2	154	403	2.65
3	3	1	1	4	3	185	432	2.41
4	4	1	1	4	4	193	457	2.25
5	5	1	1	5	5	186	483	2.58
6	6	1	1	5	6	177	469	2.67
7	7	1	1	5	7	177	428	2.71
8	8	1	1	5	8	163	439	2.47
9	9	1	2	4	9	188	439	2.29
10	10	1	2	4	10	178	407	2.26
11	11	1	2	5	11	198	498	1.97
12	12	1	2	5	12	193	459	2.14
13	13	1	2	5	13	186	459	2.44
14	14	1	2	5	14	175	375	2.52
15	15	1	2	5	15	171	382	1.72
16	16	1	2	5	16	168	417	2.75
17	17	1	3	3	17	154	389	2.38
18	18	1	3	4	18	184	414	2.46
19	19	1	3	5	19	174	483	2.29
20	20	1	3	5	20	170	430	2.30
21	21	1	3	5	21	169	443	2.94
22	22	2	4	3	22	158	381	2.50
23	23	2	4	3	23	158	365	2.44
24	24	2	4	4	24	169	386	2.44
25	25	2	4	4	25	144	339	2.15
26	26	2	4	5	26	159	419	2.54
27	27	2	4	5	27	152	469	2.74
28	28	2	4	5	28	149	379	2.50
29	29	2	4	5	29	149	375	2.54
30	30	2	5	3	30	189	395	2.65
31	31	2	5	4	31	187	447	2.52
32	32	2	5	4	32	165	430	2.67
33	33	2	5	5	33	181	453	2.79
34	34	2	5	5	34	177	385	2.33
35	35	2	5	5	35	151	414	2.67
36	36	2	5	5	36	147	353	2.69
37	37	3	6	4	37	184	411	3.00
38	38	3	6	4	38	184	420	2.49
39	39	3	6	5	39	187	427	2.25
40	40	3	6	5	40	184	409	2.49
41	41	3	6	5	41	183	337	2.02
42	42	3	6	5	42	177	352	2.31

43	43	3	7	3	43	205	472	2.57
44	44	3	7	3	44	193	340	2.37
45	45	3	7	4	45	162	375	2.64
46	46	3	7	5	46	206	451	2.37
47	47	3	7	5	47	205	472	2.22
48	48	3	7	5	48	187	402	1.90
49	49	3	7	5	49	178	464	2.61
50	50	3	7	5	50	175	414	2.13
51	51	3	8	3	51	200	466	2.16
52	52	3	8	3	52	184	356	2.33
53	53	3	8	3	53	175	449	2.52
54	54	3	8	4	54	178	360	2.45
55	55	3	8	5	55	189	385	1.44
56	56	3	8	5	56	184	431	1.72
57	57	3	8	5	57	183	401	2.17
58	58	3	9	3	58	166	404	2.68
59	59	3	9	4	59	187	482	2.43
60	60	3	9	4	60	186	350	2.36
61	61	3	9	4	61	184	483	2.44
62	62	3	9	5	62	180	425	2.66
63	63	3	9	5	63	177	420	2.46
64	64	3	9	5	64	175	440	2.52
65	65	3	9	5	65	164	405	2.42

```
GLM(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlw, p431)
```

\$ANOVA

Response : avdlygn

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	16	2.5275	0.157966	3.1437	0.001091 **
RESIDUALS	48	2.4119	0.050248		
CORRECTED TOTAL	64	4.9394			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

	Root MSE	avdlygn	Mean Coef	Var R-square	Adj R-sq
	0.2241612	2.411385	9.295956	0.511696	0.348928

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
line	2	0.38009	0.190046	3.7821	0.02983 *
line:sire	6	0.92634	0.154391	3.0726	0.01260 *
agedam	2	0.11894	0.059471	1.1835	0.31497
line:agedam	4	0.64889	0.162222	3.2284	0.02000 *
age	1	0.18349	0.183487	3.6516	0.06200 .
intlw	1	0.26970	0.269704	5.3674	0.02483 *

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

line       2 0.05526 0.02763  0.5498 0.580636  

line:sire   6 0.97389 0.16231  3.2303 0.009543 **  

agedam     2 0.33106 0.16553  3.2943 0.045640 *  

line:agedam 4 0.45343 0.11336  2.2560 0.076821 .  

age        1 0.38128 0.38128  7.5878 0.008277 **  

intlw      1 0.26970 0.26970  5.3674 0.024830 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

line       2 0.13620 0.06810  1.3553 0.267560  

line:sire   6 0.97389 0.16231  3.2303 0.009543 **  

agedam     2 0.13011 0.06505  1.2946 0.283392  

line:agedam 4 0.45343 0.11336  2.2560 0.076821 .  

age        1 0.38128 0.38128  7.5878 0.008277 **  

intlw      1 0.26970 0.26970  5.3674 0.024830 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

# p433 Output 11.40

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlw, p431),
      type=3, singular.ok=TRUE) # NOT OK for line

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: avdlygn

	Sum Sq	Df	F values	Pr(>F)
line	0.00000	0		
agedam	0.13011	2	1.2946	0.283392
age	0.38128	1	7.5878	0.008277 **
intlw	0.26970	1	5.3674	0.024830 *
line:sire	0.97389	6	3.2303	0.009543 **
line:agedam	0.45343	4	2.2560	0.076821 .
Residuals	2.41192	48		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5 Sahai - Unbalanced

Reference

- Sahai H, Ojeda MM. Analysis of Variance for Random Models Volume 2 Unbalanced Data. 2005.

5.1 Table 15.3

(6) MODEL

```
T15.3 = read.table("http://r.acr.kr/sahai/T15.3.txt")
colnames(T15.3) = c("Dam", "Sire", "pH")
T15.3 = af(T15.3, c("Dam", "Sire"))
T15.3
```

	Dam	Sire	pH
1	1	1	7.48
2	1	1	7.48
3	1	1	7.52
4	1	1	7.54
5	6	1	7.54
6	6	1	7.36
7	6	1	7.36
8	6	1	7.40
9	11	1	7.52
10	11	1	7.54
11	11	1	7.52
12	11	1	7.56
13	11	1	7.53
14	1	2	7.48
15	1	2	7.53
16	1	2	7.43
17	1	2	7.39
18	6	2	7.44
19	6	2	7.47
20	6	2	7.48
21	6	2	7.48
22	11	2	7.56
23	11	2	7.39
24	11	2	7.52
25	11	2	7.49
26	11	2	7.48
27	2	1	7.45
28	2	1	7.43
29	2	1	7.49
30	2	1	7.40
31	2	1	7.40
32	6	3	7.43
33	6	3	7.52

34	6	3 7.50
35	6	3 7.46
36	6	3 7.39
37	12	1 7.50
38	12	1 7.45
39	12	1 7.43
40	12	1 7.44
41	12	1 7.49
42	2	2 7.50
43	2	2 7.45
44	2	2 7.43
45	2	2 7.36
46	7	1 7.41
47	7	1 7.42
48	7	1 7.36
49	7	1 7.47
50	12	2 7.52
51	12	2 7.43
52	12	2 7.38
53	12	2 7.33
54	3	1 7.40
55	3	1 7.45
56	3	1 7.42
57	3	1 7.48
58	7	2 7.47
59	7	2 7.36
60	7	2 7.43
61	7	2 7.38
62	7	2 7.41
63	13	1 7.39
64	13	1 7.37
65	13	1 7.33
66	13	1 7.43
67	13	1 7.42
68	3	2 7.45
69	3	2 7.33
70	3	2 7.40
71	3	2 7.46
72	7	3 7.53
73	7	3 7.40
74	7	3 7.44
75	7	3 7.40
76	7	3 7.45
77	13	2 7.43
78	13	2 7.38
79	13	2 7.44
80	3	3 7.40
81	3	3 7.47

82	3	3 7.40
83	3	3 7.47
84	3	3 7.47
85	8	1 7.52
86	8	1 7.53
87	8	1 7.48
88	13	3 7.46
89	13	3 7.44
90	13	3 7.37
91	13	3 7.54
92	4	1 7.38
93	4	1 7.48
94	4	1 7.46
95	8	2 7.40
96	8	2 7.48
97	8	2 7.50
98	8	2 7.40
99	8	2 7.51
100	14	1 7.50
101	14	1 7.53
102	14	1 7.51
103	14	1 7.43
104	4	2 7.37
105	4	2 7.31
106	4	2 7.45
107	4	2 7.41
108	9	1 7.40
109	9	1 7.34
110	9	1 7.37
111	9	1 7.45
112	14	2 7.44
113	14	2 7.45
114	14	2 7.39
115	14	2 7.52
116	5	1 7.44
117	5	1 7.51
118	5	1 7.49
119	5	1 7.51
120	5	1 7.52
121	9	2 7.42
122	9	2 7.37
123	9	2 7.46
124	9	2 7.40
125	14	3 7.42
126	14	3 7.48
127	14	3 7.45
128	14	3 7.51
129	14	3 7.48

```

130   5   2 7.49
131   5   2 7.49
132   5   2 7.49
133   5   2 7.50
134  10   1 7.39
135  10   1 7.31
136  10   1 7.30
137  10   1 7.41
138  10   1 7.48
139  15   1 7.47
140  15   1 7.49
141  15   1 7.45
142  15   1 7.43
143  15   1 7.42
144   5   3 7.48
145   5   3 7.59
146   5   3 7.59
147  10   2 7.50
148  10   2 7.44
149  10   2 7.40
150  10   2 7.45
151  15   2 7.45
152  15   2 7.42
153  15   2 7.52
154  15   2 7.51
155  15   2 7.32
156  15   3 7.51
157  15   3 7.51
158  15   3 7.53
159  15   3 7.45
160  15   3 7.51

```

```
GLM(pH ~ Dam/Sire, T15.3) # p301
```

\$ANOVA

Response : pH

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	36	0.25804	0.0071678	2.8977	7.2e-06 ***
RESIDUALS	123	0.30425	0.0024736		
CORRECTED TOTAL	159	0.56229			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	pH	Mean	Coef	Var	R-square	Adj R-sq
0.04973534	7.449813	0.6676053	0.4589074	0.3005388		

\$`Type I`

```

          Df   Sum Sq   Mean Sq F value    Pr(>F)
Dam        14 0.178017 0.0127155  5.1405 1.563e-07 ***
Dam:Sire  22 0.080024 0.0036374  1.4705  0.09662 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df   Sum Sq   Mean Sq F value    Pr(>F)
Dam        14 0.178017 0.0127155  5.1405 1.563e-07 ***
Dam:Sire  22 0.080024 0.0036374  1.4705  0.09662 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df   Sum Sq   Mean Sq F value    Pr(>F)
Dam        14 0.179405 0.0128146  5.1805 1.347e-07 ***
Dam:Sire  22 0.080024 0.0036374  1.4705  0.09662 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(pH ~ Dam/Sire, T15.3), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: pH
          Sum Sq   Df F values    Pr(>F)
Dam        0.081011   6  5.4584 4.898e-05 ***
Dam:Sire  0.080024  22  1.4705  0.09662 .
Residuals 0.304253 123
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.2 Table 16.3

(7) MODEL

```

T16.3 = read.csv("http://r.acr.kr/sahai/T16.3.csv")
colnames(T16.3) = c("Plot", "Sample", "Subsample", "Residue")
T16.3 = af(T16.3, c("Plot", "Sample", "Subsample"))
T16.3

```

	Plot	Sample	Subsample	Residue
1	1	1	1	0.52
2	1	1	1	0.43
3	1	1	2	0.40
4	1	1	2	0.52

5	1	2	1	0.26
6	1	2	2	0.54
7	1	3	1	0.52
8	2	1	1	0.50
9	2	1	1	0.59
10	2	1	2	0.47
11	2	1	2	0.50
12	2	2	1	0.04
13	2	2	2	0.43
14	2	3	1	1.08
15	3	1	1	0.34
16	3	1	1	0.26
17	3	1	2	0.32
18	3	1	2	0.45
19	3	2	1	0.25
20	3	2	2	0.38
21	3	3	1	0.29
22	4	1	1	0.18
23	4	1	1	0.24
24	4	1	2	0.31
25	4	1	2	0.29
26	4	2	1	0.13
27	4	2	2	0.25
28	4	3	1	0.10
29	5	1	1	1.05
30	5	1	1	0.66
31	5	1	2	0.60
32	5	1	2	0.51
33	5	2	1	0.95
34	5	2	2	0.84
35	5	3	1	0.92
36	6	1	1	0.52
37	6	1	1	0.66
38	6	1	2	0.55
39	6	1	2	0.40
40	6	2	1	0.33
41	6	2	2	0.26
42	6	3	1	0.41
43	7	1	1	0.77
44	7	1	1	0.56
45	7	1	2	0.51
46	7	1	2	0.60
47	7	2	1	0.44
48	7	2	2	0.50
49	7	3	1	0.44
50	8	1	1	0.89
51	8	1	1	0.92
52	8	1	2	0.75

53	8	1	2	0.58
54	8	2	1	0.64
55	8	2	2	0.54
56	8	3	1	0.36
57	9	1	1	0.50
58	9	1	1	0.67
59	9	1	2	0.60
60	9	1	2	0.53
61	9	2	1	0.60
62	9	2	2	0.71
63	9	3	1	0.92
64	10	1	1	0.58
65	10	1	1	0.52
66	10	1	2	0.56
67	10	1	2	0.44
68	10	2	1	0.46
69	10	2	2	0.52
70	10	3	1	0.52
71	11	1	1	0.24
72	11	1	1	0.36
73	11	1	2	0.48
74	11	1	2	0.30
75	11	2	1	0.53
76	11	2	2	0.50
77	11	3	1	0.39

```
GLM(Residue ~ Plot/Sample/Subsample, T16.3) # p344
```

\$ANOVA

Response : Residue

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	54	3.1897	0.059069	5.8842	1.476e-05 ***
RESIDUALS	22	0.2208	0.010039		
CORRECTED TOTAL	76	3.4106			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root	MSE	Residue	Mean	Coef	Var	R-square	Adj R-sq
0.100193	0.5023377	19.94535	0.9352456	0.776303			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Plot	10	1.84041	0.184041	18.3332	1.929e-08 ***
Plot:Sample	22	0.99175	0.045079	4.4906	0.0004209 ***
Plot:Sample:Subsample	22	0.35757	0.016253	1.6191	0.1330632

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

$`Type II` 
          Df  Sum Sq  Mean Sq F value    Pr(>F)
Plot           10 1.84041 0.184041 18.3332 1.929e-08 ***
Plot:Sample     22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
          Df  Sum Sq  Mean Sq F value    Pr(>F)
Plot           10 1.78686 0.178686 17.7998 2.547e-08 ***
Plot:Sample     22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(Residue ~ Plot/Sample/Subsample, T16.3), type=3, singular.ok=TRUE)

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Residue
          Sum Sq Df F values   Pr(>F)
Plot        0.00000  0
Plot:Sample 0.36613 11  3.3156 0.00805 **
Plot:Sample:Subsample 0.35758 22  1.6191 0.13306
Residuals   0.22085 22
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
# NOT OK

```

6 Federer - Variations

Reference

- Federer WT, King F. Variations on Split Plot and Split Block Experiment Designs. John Wiley & Sons Inc. 2007.

6.1 Example 2.2

(8) MODEL

```
ex2.2 = read.table("http://r.acr.kr/split/sbex2_2.txt", header=TRUE)
ex2.2 = af(ex2.2, c("Row", "Column", "R", "S"))
ex2.2
```

	Row	Column	R	S	Y	
1	1		1	1	1027.85	
2	1		1	1	2	982.74
3	1		1	1	3	1007.24
4	1		1	1	4	1008.47
5	1		2	2	1	1004.33
6	1		2	2	2	977.86
7	1		2	2	3	999.15
8	1		2	2	4	990.86
9	1		3	3	1	992.57
10	1		3	3	2	993.71
11	1		3	3	3	1012.57
12	1		3	3	4	968.25
13	1		4	4	1	994.60
14	1		4	4	2	1021.81
15	1		4	4	3	995.03
16	1		4	4	4	1002.17
17	1		5	5	1	1019.89
18	1		5	5	2	1017.48
19	1		5	5	3	987.82
20	1		5	5	4	995.63
21	2		4	1	1	996.18
22	2		4	1	2	981.96
23	2		4	1	3	985.63
24	2		4	1	4	965.80
25	2		5	2	1	996.61
26	2		5	2	2	1011.94
27	2		5	2	3	972.76
28	2		5	2	4	1011.99
29	2		2	3	1	1021.61
30	2		2	3	2	1014.46
31	2		2	3	3	980.03
32	2		2	3	4	1014.80
33	2		3	4	1	1028.78

34	2	3 4 2	1006.01
35	2	3 4 3	1015.04
36	2	3 4 4	1000.72
37	2	1 5 1	994.91
38	2	1 5 2	999.91
39	2	1 5 3	1010.29
40	2	1 5 4	1018.49
41	3	5 1 1	985.72
42	3	5 1 2	1012.60
43	3	5 1 3	984.62
44	3	5 1 4	973.47
45	3	1 2 1	1013.52
46	3	1 2 2	1017.40
47	3	1 2 3	996.63
48	3	1 2 4	989.91
49	3	4 3 1	1003.92
50	3	4 3 2	999.33
51	3	4 3 3	995.70
52	3	4 3 4	988.14
53	3	2 4 1	1010.08
54	3	2 4 2	997.66
55	3	2 4 3	1012.12
56	3	2 4 4	1019.53
57	3	3 5 1	1004.83
58	3	3 5 2	983.86
59	3	3 5 3	1018.60
60	3	3 5 4	1020.95
61	4	2 1 1	991.79
62	4	2 1 2	979.47
63	4	2 1 3	1004.70
64	4	2 1 4	1032.75
65	4	3 2 1	1004.52
66	4	3 2 2	996.53
67	4	3 2 3	1016.95
68	4	3 2 4	983.79
69	4	1 3 1	990.17
70	4	1 3 2	972.21
71	4	1 3 3	1002.17
72	4	1 3 4	1017.56
73	4	5 4 1	1006.13
74	4	5 4 2	1005.57
75	4	5 4 3	1003.18
76	4	5 4 4	992.21
77	4	4 5 1	1011.02
78	4	4 5 2	982.79
79	4	4 5 3	1018.23
80	4	4 5 4	976.68
81	5	3 1 1	993.54

```

82   5      3 1 2 1006.80
83   5      3 1 3 1001.24
84   5      3 1 4 1010.73
85   5      4 2 1  985.04
86   5      4 2 2  987.54
87   5      4 2 3  990.53
88   5      4 2 4  982.68
89   5      5 3 1 1012.14
90   5      5 3 2  999.32
91   5      5 3 3 1005.51
92   5      5 3 4  998.86
93   5      1 4 1  985.12
94   5      1 4 2  984.14
95   5      1 4 3 1010.74
96   5      1 4 4 1004.63
97   5      2 5 1  967.39
98   5      2 5 2 1009.78
99   5      2 5 3 1027.49
100  5      2 5 4 1001.61

```

```
GLM(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2)
```

\$ANOVA

```

Response : Y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL          99 22310 225.36
RESIDUALS       0      0
CORRECTED TOTAL 99 22310

```

\$Fitness

Root MSE	Y Mean	Coef	Var	R-square
NA	1000.098	NA		1

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Row	4	147.4	36.86		
R	4	1159.8	289.94		
S	3	351.9	117.29		
R:S	12	826.0	68.83		
Row:R	16	3979.8	248.74		
S:Column	12	3863.3	321.94		
R:S:Column	48	11982.3	249.63		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Row	0				
R	4	1159.8	289.94		
S	3	351.9	117.29		

```

R:S          12   826.0   68.83
Row:R         0
S:Column     12  3863.3  321.94
R:S:Column  48 11982.3  249.63

$`Type III` 
CAUTION: Singularity Exists !
      Df  Sum Sq Mean Sq F value Pr(>F)
Row          0
R            4   1159.8   289.94
S            3    351.9   117.29
R:S          12   826.0   68.83
Row:R        0
S:Column    12  3863.3  321.94
R:S:Column  48 11982.3  249.63

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2), type=3,
singular.ok=TRUE) # NOT WORKING

```

6.2 Example 3.1

(9) MODEL

```

ex3.1a = read.table("http://r.acr.kr/split/Ex3.1-example.txt", header=TRUE)
ex3.1a = af(ex3.1a, c("row", "P", "column", "R", "S"))
ex3.1a

```

	row	P	column	R	S	height	
1	1	1		1	3	4	103
2	1	1		1	3	2	98
3	1	1		1	3	3	101
4	1	1		1	3	1	101
5	1	1		2	4	2	100
6	1	1		2	4	3	98
7	1	1		2	4	1	100
8	1	1		2	4	4	99
9	1	1		3	5	3	99
10	1	1		3	5	1	99
11	1	1		3	5	2	100
12	1	1		3	5	4	97
13	1	1		4	2	2	99
14	1	1		4	2	1	102
15	1	1		4	2	3	99
16	1	1		4	2	4	100
17	1	1		5	1	1	102
18	1	1		5	1	2	107
19	1	1		5	1	3	98
20	1	1		5	1	4	99

21	1 2	1 3 4	101
22	1 2	1 3 2	101
23	1 2	1 3 3	99
24	1 2	1 3 1	100
25	1 2	2 4 2	97
26	1 2	2 4 3	85
27	1 2	2 4 1	99
28	1 2	2 4 4	97
29	1 2	3 5 3	98
30	1 2	3 5 1	96
31	1 2	3 5 2	88
32	1 2	3 5 4	98
33	1 2	4 2 2	95
34	1 2	4 2 1	90
35	1 2	4 2 3	99
36	1 2	4 2 4	87
37	1 2	5 1 1	98
38	1 2	5 1 2	98
39	1 2	5 1 3	99
40	1 2	5 1 4	89
41	2 1	1 2 4	99
42	2 1	1 2 2	97
43	2 1	1 2 3	98
44	2 1	1 2 1	95
45	2 1	2 3 2	99
46	2 1	2 3 3	98
47	2 1	2 3 1	96
48	2 1	2 3 4	93
49	2 1	3 1 3	97
50	2 1	3 1 1	99
51	2 1	3 1 2	95
52	2 1	3 1 4	98
53	2 1	4 4 2	97
54	2 1	4 4 1	95
55	2 1	4 4 3	99
56	2 1	4 4 4	94
57	2 1	5 5 1	98
58	2 1	5 5 2	93
59	2 1	5 5 3	98
60	2 1	5 5 4	96
61	2 2	1 2 4	99
62	2 2	1 2 2	89
63	2 2	1 2 3	98
64	2 2	1 2 1	94
65	2 2	2 3 2	98
66	2 2	2 3 3	91
67	2 2	2 3 1	97
68	2 2	2 3 4	96

69	2 2	3 1 3	94
70	2 2	3 1 1	97
71	2 2	3 1 2	98
72	2 2	3 1 4	96
73	2 2	4 4 2	99
74	2 2	4 4 1	89
75	2 2	4 4 3	97
76	2 2	4 4 4	98
77	2 2	5 5 1	99
78	2 2	5 5 2	96
79	2 2	5 5 3	93
80	2 2	5 5 4	98
81	3 1	1 4 4	99
82	3 1	1 4 2	88
83	3 1	1 4 3	98
84	3 1	1 4 1	96
85	3 1	2 5 2	98
86	3 1	2 5 3	99
87	3 1	2 5 1	92
88	3 1	2 5 4	88
89	3 1	3 2 3	98
90	3 1	3 2 1	85
91	3 1	3 2 2	88
92	3 1	3 2 4	95
93	3 1	4 1 2	97
94	3 1	4 1 1	87
95	3 1	4 1 3	96
96	3 1	4 1 4	88
97	3 1	5 3 1	88
98	3 1	5 3 2	85
99	3 1	5 3 3	78
100	3 1	5 3 4	78
101	3 2	1 4 4	88
102	3 2	1 4 2	85
103	3 2	1 4 3	78
104	3 2	1 4 1	80
105	3 2	2 5 2	80
106	3 2	2 5 3	79
107	3 2	2 5 1	77
108	3 2	2 5 4	78
109	3 2	3 2 3	90
110	3 2	3 2 1	91
111	3 2	3 2 2	92
112	3 2	3 2 4	93
113	3 2	4 1 2	99
114	3 2	4 1 1	97
115	3 2	4 1 3	98
116	3 2	4 1 4	99

117	3 2	5 3 1	80
118	3 2	5 3 2	81
119	3 2	5 3 3	82
120	3 2	5 3 4	83
121	4 1	1 1 4	80
122	4 1	1 1 2	81
123	4 1	1 1 3	84
124	4 1	1 1 1	80
125	4 1	2 2 2	90
126	4 1	2 2 3	90
127	4 1	2 2 1	90
128	4 1	2 2 4	90
129	4 1	3 3 3	99
130	4 1	3 3 1	98
131	4 1	3 3 2	97
132	4 1	3 3 4	99
133	4 1	4 5 2	95
134	4 1	4 5 1	95
135	4 1	4 5 3	95
136	4 1	4 5 4	96
137	4 1	5 4 1	99
138	4 1	5 4 2	95
139	4 1	5 4 3	98
140	4 1	5 4 4	98
141	4 2	1 1 4	98
142	4 2	1 1 2	99
143	4 2	1 1 3	97
144	4 2	1 1 1	99
145	4 2	2 2 2	88
146	4 2	2 2 3	87
147	4 2	2 2 1	88
148	4 2	2 2 4	86
149	4 2	3 3 3	99
150	4 2	3 3 1	97
151	4 2	3 3 2	96
152	4 2	3 3 4	95
153	4 2	4 5 2	89
154	4 2	4 5 1	88
155	4 2	4 5 3	87
156	4 2	4 5 4	85
157	4 2	5 4 1	90
158	4 2	5 4 2	90
159	4 2	5 4 3	90
160	4 2	5 4 4	97
161	5 1	1 5 4	98
162	5 1	1 5 2	98
163	5 1	1 5 3	99
164	5 1	1 5 1	97

165	5 1	2 1 2	98
166	5 1	2 1 3	97
167	5 1	2 1 1	98
168	5 1	2 1 4	89
169	5 1	3 4 3	88
170	5 1	3 4 1	87
171	5 1	3 4 2	88
172	5 1	3 4 4	88
173	5 1	4 3 2	98
174	5 1	4 3 1	95
175	5 1	4 3 3	97
176	5 1	4 3 4	99
177	5 1	5 2 1	98
178	5 1	5 2 2	98
179	5 1	5 2 3	95
180	5 1	5 2 4	99
181	5 2	1 5 4	88
182	5 2	1 5 2	87
183	5 2	1 5 3	99
184	5 2	1 5 1	98
185	5 2	2 1 2	99
186	5 2	2 1 3	95
187	5 2	2 1 1	99
188	5 2	2 1 4	90
189	5 2	3 4 3	98
190	5 2	3 4 1	99
191	5 2	3 4 2	99
192	5 2	3 4 4	92
193	5 2	4 3 2	88
194	5 2	4 3 1	86
195	5 2	4 3 3	87
196	5 2	4 3 4	83
197	5 2	5 2 1	99
198	5 2	5 2 2	96
199	5 2	5 2 3	98
200	5 2	5 2 4	99

```
GLM(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
    S:R:P + R:S:P:row, ex3.1a)
```

```
$ANOVA
Response : height
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      199 7534.8 37.863
RESIDUALS   0     0.0
CORRECTED TOTAL 199 7534.8
```

\$Fitness

Root	MSE	height	Mean	Coef	Var	R-square
NA	93.965		NA		1	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
row	4	2017.03	504.26		
R	4	90.63	22.66		
P	1	253.12	253.12		
S	3	16.38	5.46		
R:S	12	195.05	16.25		
row:P	4	167.25	41.81		
R:P	4	504.95	126.24		
row:R:P	32	2933.52	91.67		
P:S	3	14.29	4.76		
row:P:S	24	234.68	9.78		
R:P:S	12	100.33	8.36		
row:R:P:S	96	1007.52	10.49		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
row	4	2017.03	504.26		
R	4	90.63	22.66		
P	1	253.12	253.12		
S	3	16.38	5.46		
R:S	12	195.05	16.25		
row:P	4	167.25	41.81		
R:P	4	504.95	126.24		
row:R:P	32	2933.52	91.67		
P:S	3	14.29	4.76		
row:P:S	24	234.68	9.78		
R:P:S	12	100.33	8.36		
row:R:P:S	96	1007.52	10.49		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
row	4	2017.03	504.26		
R	4	90.63	22.66		
P	1	253.13	253.13		
S	3	16.38	5.46		
R:S	12	195.05	16.25		
row:P	4	167.25	41.81		
R:P	4	504.95	126.24		
row:R:P	32	2933.52	91.67		
P:S	3	14.30	4.77		
row:P:S	24	234.68	9.78		
R:P:S	12	100.33	8.36		
row:R:P:S	96	1007.52	10.49		

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +
         S:P:row + S:R:P + R:S:P:row, ex3.1a), type=3, singular.ok=TRUE)
# NOT WORKING

alias(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
      S:R:P + R:S:P:row, ex3.1a) # NO ALIAS

```

Model :

```

height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +
      S:P:row + S:R:P + R:S:P:row

```

(10) MODEL

- p94 Appendix 3.1

```

ex3.1b = read.table("http://r.acr.kr/split/spexvar3.txt", header=TRUE)
ex3.1b = af(ex3.1b, c("rep", "var", "nit", "row", "col"))
ex3.1b

```

	row	col	rep	var	nit	set	reps	yield
1	1	1	1	3	3	1	1	156
2	1	2	1	3	2	1	1	118
3	1	3	4	3	2	2	1	109
4	1	4	4	3	3	2	1	99
5	2	1	1	3	1	1	1	140
6	2	2	1	3	4	1	1	105
7	2	3	4	3	4	2	1	63
8	2	4	4	3	1	2	1	70
9	3	1	1	1	4	1	1	111
10	3	2	1	1	1	1	1	130
11	3	3	4	2	4	2	1	80
12	3	4	4	2	2	2	1	94
13	4	1	1	1	3	1	1	174
14	4	2	1	1	2	1	1	157
15	4	3	4	2	3	2	1	126
16	4	4	4	2	1	2	1	82
17	5	1	1	2	4	1	1	117
18	5	2	1	2	1	1	1	114
19	5	3	4	1	1	2	1	90
20	5	4	4	1	2	2	1	100
21	6	1	1	2	2	1	1	161
22	6	2	1	2	3	1	1	141
23	6	3	4	1	3	2	1	116
24	6	4	4	1	4	2	1	62
25	7	1	2	3	2	1	2	104
26	7	2	2	3	4	1	2	70
27	7	3	5	2	3	2	2	96
28	7	4	5	2	4	2	2	60
29	8	1	2	3	1	1	2	89

30	8	2	2	3	3	1	2	117
31	8	3	5	2	2	2	2	89
32	8	4	5	2	1	2	2	102
33	9	1	2	1	3	1	2	122
34	9	2	2	1	4	1	2	74
35	9	3	5	1	2	2	2	112
36	9	4	5	1	3	2	2	86
37	10	1	2	1	1	1	2	89
38	10	2	2	1	2	1	2	81
39	10	3	5	1	4	2	2	68
40	10	4	5	1	1	2	2	64
41	11	1	2	2	1	1	2	103
42	11	2	2	2	4	1	2	64
43	11	3	5	3	2	2	2	132
44	11	4	5	3	3	2	2	124
45	12	1	2	2	2	1	2	132
46	12	2	2	2	3	1	2	133
47	12	3	5	3	1	2	2	129
48	12	4	5	3	4	2	2	89
49	13	1	3	2	1	1	3	108
50	13	2	3	2	2	1	3	126
51	13	3	6	1	2	2	3	118
52	13	4	6	1	4	2	3	53
53	14	1	3	2	3	1	3	149
54	14	2	3	2	4	1	3	70
55	14	3	6	1	3	2	3	113
56	14	4	6	1	1	2	3	74
57	15	1	3	3	3	1	3	144
58	15	2	3	3	1	1	3	124
59	15	3	6	2	3	2	3	104
60	15	4	6	2	2	2	3	86
61	16	1	3	3	2	1	3	121
62	16	2	3	3	4	1	3	96
63	16	3	6	2	4	2	3	89
64	16	4	6	2	1	2	3	82
65	17	1	3	1	4	1	3	61
66	17	2	3	1	3	1	3	100
67	17	3	6	3	4	2	3	97
68	17	4	6	3	1	2	3	99
69	18	1	3	1	1	1	3	91
70	18	2	3	1	2	1	3	97
71	18	3	6	3	2	2	3	119
72	18	4	6	3	3	2	3	121

```
GLM(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b)
```

\$ANOVA

Response : yield

```

              Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          37 48090 1299.7 11.341 6.734e-11 ***
RESIDUALS      34  3896   114.6
CORRECTED TOTAL 71 51986
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE yield Mean Coef Var R-square Adj R-sq
10.70513 103.9722 10.29615 0.9250491 0.8434848

$`Type I`
              Df Sum Sq Mean Sq F value    Pr(>F)
rep            5 15875.3 3175.1 27.7056 4.391e-11 ***
var            2 1786.4   893.2  7.7939 0.0016359 **
rep:var        10 6013.3   601.3  5.2472 0.0001207 ***
nit            3 20020.5 6673.5 58.2331 1.754e-13 ***
var:nit        6   321.7    53.6  0.4679 0.8271333
row            9   900.9   100.1  0.8734 0.5575581
col            2   3171.5 1585.7 13.8373 4.012e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
              Df Sum Sq Mean Sq F value    Pr(>F)
rep            2 5942.5 2971.3 25.9273 1.449e-07 ***
var            2 2799.8 1399.9 12.2155 0.0001005 ***
rep:var        4   997.8    249.4  2.1767 0.0926008 .
nit            3 12559.3 4186.4 36.5308 9.683e-11 ***
var:nit        6   477.8    79.6  0.6949 0.6553307
row            9   945.0   105.0  0.9162 0.5230151
col            2   3171.5 1585.7 13.8373 4.012e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
              Df Sum Sq Mean Sq F value    Pr(>F)
rep            2 5942.5 2971.3 25.9273 1.449e-07 ***
var            2 2799.8 1399.9 12.2155 0.0001005 ***
rep:var        4   997.8    249.4  2.1767 0.0926008 .
nit            3 11977.9 3992.6 34.8397 1.775e-10 ***
var:nit        6   477.8    79.6  0.6949 0.6553307
row            9   945.0   105.0  0.9162 0.5230151
col            2   3171.5 1585.7 13.8373 4.012e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b),
      type=3, singular.ok=TRUE) # NOT OK for var

```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: yield
            Sum Sq Df F values    Pr(>F)
rep        5942.5  2 25.9273 1.449e-07 ***
var         0.0   0
nit       11977.9  3 34.8397 1.775e-10 ***
row        945.0   9  0.9162    0.5230
col       3171.5  2 13.8373 4.012e-05 ***
rep:var    997.8  4  2.1767    0.0926 .
var:nit    477.8  6  0.6949    0.6553
Residuals 3896.4 34
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.3 Example 5.1

(11) MODEL

```

ex5.1 = read.table("http://r.acr.kr/split/sbsp.txt", header=TRUE)
ex5.1 = af(ex5.1, c("R", "A", "C", "B", "Tx"))
ex5.1

```

	R	A	C	B	Tx	Y
1	1	1	1	2	1	2
2	1	1	1	1	2	5
3	1	1	2	2	4	6
4	1	1	2	1	3	9
5	1	1	3	1	6	8
6	1	1	3	2	5	5
7	1	2	1	2	4	9
8	1	2	1	1	3	7
9	1	2	2	2	6	8
10	1	2	2	1	5	4
11	1	2	3	1	1	3
12	1	2	3	2	2	5
13	2	2	1	2	6	8
14	2	2	1	1	5	5
15	2	2	2	2	1	3
16	2	2	2	1	2	5
17	2	2	3	1	4	9
18	2	2	3	2	3	7

```

19 2 1 1 2 3 3
20 2 1 1 1 6 4
21 2 1 2 2 5 3
22 2 1 2 1 1 0
23 2 1 3 1 2 1
24 2 1 3 2 4 2
25 3 1 1 2 5 5
26 3 1 1 1 1 5
27 3 1 2 2 2 5
28 3 1 2 1 4 9
29 3 1 3 1 3 7
30 3 1 3 2 6 8
31 3 2 1 2 2 6
32 3 2 1 1 4 8
33 3 2 2 2 3 7
34 3 2 2 1 6 8
35 3 2 3 1 5 6
36 3 2 3 2 1 3

```

```
GLM(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	24	196.238	8.1766	7.0476	0.0008758 ***
RESIDUALS	11	12.762	1.1602		
CORRECTED TOTAL	35	209.000			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

	Root MSE	Y Mean	Coef Var	R-square	Adj R-sq
	1.077122	5.5	19.58405	0.9389372	0.8057093

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	2	33.500	16.7500	14.4373	0.0008391 ***
A	1	16.000	16.0000	13.7908	0.0034197 **
R:A	2	32.167	16.0833	13.8626	0.0009856 ***
C	2	0.500	0.2500	0.2155	0.8094766
B	1	1.778	1.7778	1.5323	0.2415358
C:B	2	0.389	0.1944	0.1676	0.8478141
Tx	5	103.333	20.6667	17.8131	6.055e-05 ***
A:Tx	5	6.521	1.3042	1.1241	0.4027183
B:Tx	4	2.050	0.5126	0.4418	0.7761730

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R      2   23.116 11.5581  9.9622  0.003396 **  

A      1   12.375 12.3751 10.6664  0.007519 **  

R:A    2   27.426 13.7132 11.8197  0.001820 **  

C      2     0.970  0.4850  0.4180  0.668392  

B      1     1.757  1.7574  1.5148  0.244080  

C:B    2     0.085  0.0424  0.0366  0.964202  

Tx     5 103.333 20.6667 17.8131 6.055e-05 ***  

A:Tx   4     2.655  0.6636  0.5720  0.688652  

B:Tx   4     2.050  0.5126  0.4418  0.776173  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

CAUTION: Singularity Exists !  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R      2   22.186 11.0928  9.5611  0.003924 **  

A      1   15.185 15.1853 13.0886  0.004042 **  

R:A    2   27.426 13.7132 11.8197  0.001820 **  

C      2     1.010  0.5049  0.4352  0.657839  

B      1     1.792  1.7922  1.5448  0.239751  

C:B    2     0.085  0.0424  0.0366  0.964202  

Tx     5 103.333 20.6667 17.8131 6.055e-05 ***  

A:Tx   4     2.655  0.6636  0.5720  0.688652  

B:Tx   4     2.050  0.5126  0.4418  0.776173  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
alias(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)
```

```
Model :  

Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx
```

```
Complete :  

  (Intercept) R1     R2     A1     C1     C2     B1     Tx1    Tx2    Tx3    Tx4    Tx5    R1:A1  

B1:Tx5      0       0     0 -1/5     0     0 -1/5     0     0     0     0     0     0  

          R2:A1  C1:B1  C2:B1 A1:Tx1 A1:Tx2 A1:Tx3 A1:Tx4 A1:Tx5 B1:Tx1 B1:Tx2 B1:Tx3  

B1:Tx5      0       0     0  1/5    1/5    1/5    1/5     -1    1/5    1/5    1/5  

          B1:Tx4  

B1:Tx5    1/5
```

```
options(contrasts=c("contr.sum", "contr.poly"))  

Anova(lm(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1),  

      type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: Y
          Sum Sq Df F values    Pr(>F)
R           22.186  2   9.5611  0.003924 ***
A           0.000  0
C           1.010  2   0.4352  0.657839
B           0.000  0
Tx          103.333  5  17.8131 6.055e-05 ***
R:A         27.426  2  11.8197  0.001820 **
C:B         0.085  2   0.0366  0.964202
A:Tx        2.655  4   0.5720  0.688652
B:Tx        2.050  4   0.4418  0.776173
Residuals  12.762 11
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(12) MODEL

```
GLM(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)
```

```

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      28 204.2  7.2929 10.635  0.001719 ***
RESIDUALS  7   4.8  0.6857
CORRECTED TOTAL 35 209.0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE Y Mean Coef Var  R-square  Adj R-sq
0.8280787  5.5 15.05598 0.9770335 0.8851675

```

```

$`Type I` 
          Df Sum Sq Mean Sq F value    Pr(>F)
R           2 33.500 16.7500 24.4271 0.0006969 ***
A           1 16.000 16.0000 23.3333 0.0018985 **
R:A         2 32.167 16.0833 23.4549 0.0007889 ***
C           2   0.500  0.2500  0.3646 0.7069339
B           1   1.778  1.7778  2.5926 0.1513998
C:B         2   0.389  0.1944  0.2836 0.7613494
Tx          5 103.333 20.6667 30.1389 0.0001357 ***
A:Tx        5   6.521  1.3042  1.9019 0.2123307
B:Tx        4   2.050  0.5126  0.7475 0.5896365
A:B:Tx     4   7.962  1.9905  2.9029 0.1038803
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type II`
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
R	2	31.838	15.9191	23.2153	0.0008139 ***						
A	1	12.375	12.3751	18.0470	0.0038017 **						
R:A	1	2.017	2.0174	2.9420	0.1300172						
C	2	0.500	0.2500	0.3645	0.7069558						
B	1	1.757	1.7574	2.5629	0.1534298						
C:B	1	0.644	0.6445	0.9399	0.3646045						
Tx	5	103.333	20.6667	30.1389	0.0001357 ***						
A:Tx	4	2.655	0.6636	0.9678	0.4812226						
B:Tx	4	2.050	0.5126	0.7475	0.5896365						
A:B:Tx	4	7.962	1.9905	2.9029	0.1038803						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'. '	0.1	' '	1

\$`Type III`

CAUTION: Singularity Exists !

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	2	28.112	14.0562	20.4986	0.0011846 **
A	1	14.655	14.6551	21.3720	0.0024176 **
R:A	1	2.017	2.0174	2.9420	0.1300172
C	2	0.471	0.2356	0.3436	0.7205632
B	1	1.769	1.7694	2.5804	0.1522328
C:B	1	0.644	0.6445	0.9399	0.3646045
Tx	5	103.815	20.7630	30.2793	0.0001336 ***
A:Tx	4	2.951	0.7378	1.0760	0.4358837
B:Tx	4	3.553	0.8882	1.2954	0.3579988
A:B:Tx	4	7.962	1.9905	2.9029	0.1038803

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '. ' 0.1 ' ' 1

```
alias(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)
```

Model :

Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx

Complete :

	(Intercept)	R1	R2	A1	C1	C2	B1	Tx1	Tx2	Tx3	Tx4	Tx5
B1:Tx5	0		0	0	-1/5	0	0	-1/5	0	0	0	0
A1:B1:Tx5	-1/6		0	0	0	0	0	1/6	1/6	1/6	1/6	-5/6
A1:B1:Tx6	0		2/3	0	4/45	2/3	-2/3	4/45	-1/3	1/3	-1/3	0
	R1:A1	R2:A1	C1:B1	C2:B1	A1:Tx1	A1:Tx2	A1:Tx3	A1:Tx4	A1:Tx5	B1:Tx1		
B1:Tx5	0	0	0	0	1/5	1/5	1/5	1/5	-1	1/5		
A1:B1:Tx5	0	0	0	0	0	0	0	0	0	0		
A1:B1:Tx6	-2/9	4/9	-2/9	-2/9	-1/5	-1/5	-1/5	4/5	0	-1/5		
	B1:Tx2	B1:Tx3	B1:Tx4	A1:B1:Tx1	A1:B1:Tx2	A1:B1:Tx3	A1:B1:Tx4					
B1:Tx5	1/5	1/5	1/5	0	0	0	0	0	0	0		
A1:B1:Tx5	0	0	0	0	0	0	0	0	0	0		
A1:B1:Tx6	-1/5	-1/5	4/5	1	-1		1		0	0		

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F values	Pr(>F)
R	11.643	1	16.9793	0.004456 **
A	0.000	0		
C	0.002	1	0.0025	0.961483
B	0.000	0		
Tx	89.178	3	43.3503	6.87e-05 ***
R:A	2.017	1	2.9420	0.130017
C:B	0.644	1	0.9399	0.364604
A:Tx	0.543	3	0.2640	0.849381
B:Tx	3.384	3	1.6451	0.264128
A:B:Tx	7.962	4	2.9029	0.103880
Residuals	4.800	7		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.4 Example 7.1

(13) MODEL

```

ex7.1 = read.table("http://r.acr.kr/split/asped.txt", header=TRUE)
ex7.1 = af(ex7.1, c("R", "G", "F"))
ex7.1

```

	Y	R	G	F
1	2	1	25	1
2	4	1	25	2
3	6	1	25	3
4	1	1	26	1
5	3	1	26	2
6	5	1	26	3
7	9	1	27	1
8	9	1	27	2
9	8	1	27	3
10	9	1	28	1
11	9	1	28	2
12	7	1	28	3
13	2	1	1	1
14	5	1	1	2
15	7	1	1	3

16	3	1	2	1
17	6	1	2	2
18	5	1	2	3
19	4	1	3	1
20	7	1	3	2
21	6	1	3	3
22	5	1	4	1
23	8	1	4	2
24	4	1	4	3
25	6	1	5	1
26	8	1	5	2
27	8	1	5	3
28	7	1	6	1
29	8	1	6	2
30	7	1	6	3
31	3	2	25	1
32	3	2	25	2
33	7	2	25	3
34	2	2	26	1
35	2	2	26	2
36	4	2	26	3
37	8	2	27	1
38	8	2	27	2
39	8	2	27	3
40	7	2	28	1
41	8	2	28	2
42	9	2	28	3
43	1	2	7	1
44	2	2	7	2
45	3	2	7	3
46	2	2	8	1
47	3	2	8	2
48	5	2	8	3
49	3	2	9	1
50	4	2	9	2
51	4	2	9	3
52	4	2	10	1
53	4	2	10	2
54	5	2	10	3
55	8	2	11	1
56	8	2	11	2
57	8	2	11	3
58	3	2	12	1
59	5	2	12	2
60	7	2	12	3
61	4	3	25	1
62	6	3	25	2
63	8	3	25	3

64 2 3 26 1
65 5 3 26 2
66 7 3 26 3
67 8 3 27 1
68 7 3 27 2
69 9 3 27 3
70 7 3 28 1
71 7 3 28 2
72 9 3 28 3
73 7 3 13 1
74 7 3 13 2
75 9 3 13 3
76 5 3 14 1
77 6 3 14 2
78 8 3 14 3
79 3 3 15 1
80 5 3 15 2
81 6 3 15 3
82 7 3 16 1
83 7 3 16 2
84 9 3 16 3
85 6 3 17 1
86 8 3 17 2
87 8 3 17 3
88 5 3 18 1
89 7 3 18 2
90 8 3 18 3
91 4 4 25 1
92 5 4 25 2
93 6 4 25 3
94 5 4 26 1
95 2 4 26 2
96 5 4 26 3
97 9 4 27 1
98 9 4 27 2
99 9 4 27 3
100 9 4 28 1
101 8 4 28 2
102 7 4 28 3
103 5 4 19 1
104 8 4 19 2
105 9 4 19 3
106 6 4 20 1
107 6 4 20 2
108 8 4 20 3
109 7 4 21 1
110 4 4 21 2
111 8 4 21 3

```

112 8 4 22 1
113 7 4 22 2
114 9 4 22 3
115 9 4 23 1
116 8 4 23 2
117 9 4 23 3
118 9 4 24 1
119 8 4 24 2
120 9 4 24 3

```

```
GLM(Y ~ R + G + R:G + F + F:G, ex7.1)
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	95	577.82	6.0824	5.3082	1.068e-05 ***
RESIDUALS	24	27.50	1.1458		
CORRECTED TOTAL	119	605.32			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	MSE	Y Mean	Coef	Var	R-square	Adj R-sq
1.070436	6.175	17.335	0.9545699	0.7747422		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	3	84.76	28.2528	24.6570	1.655e-07 ***
G	27	343.48	12.7216	11.1025	4.286e-08 ***
R:G	9	11.75	1.3056	1.1394	0.3749
F	2	59.85	29.9250	26.1164	9.481e-07 ***
G:F	54	77.98	1.4441	1.2603	0.2718

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	3	5.75	1.9167	1.6727	0.1994
G	27	343.48	12.7216	11.1025	4.286e-08 ***
R:G	9	11.75	1.3056	1.1394	0.3749
F	2	59.85	29.9250	26.1164	9.481e-07 ***
G:F	54	77.98	1.4441	1.2603	0.2718

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	3	5.75	1.9167	1.6727	0.1994

```

G    27 343.48 12.7216 11.1025 4.286e-08 ***
R:G   9   11.75  1.3056  1.1394     0.3749
F     2   50.50 25.2525 22.0385 3.686e-06 ***
G:F  54   77.98  1.4441  1.2603     0.2718
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + G + R:G + F + F:G, ex7.1), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F values	Pr(>F)
R	0.000	0		
G	202.417	3	58.8848	3.258e-11 ***
F	50.505	2	22.0385	3.686e-06 ***
R:G	11.750	9	1.1394	0.3749
G:F	77.983	54	1.2603	0.2718
Residuals	27.500	24		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.5 Example 7.3

(14) MODEL

```

ex7.3 = read.table("http://r.acr.kr/split/assped.txt", header=TRUE)
ex7.3 = af(ex7.3, c("R", "T", "G", "F"))
ex7.3

```

	Y	R	T	G	F
1	2	1	1	1	1
2	4	1	1	1	2
3	6	1	1	1	3
4	3	1	1	2	1
5	5	1	1	2	2
6	7	1	1	2	3
7	7	1	1	3	1
8	7	1	1	3	2
9	9	1	1	3	3
10	8	1	1	4	1
11	8	1	1	4	2
12	9	1	1	4	3
13	8	1	1	5	1
14	8	1	1	5	2
15	9	1	1	5	3

16	2	1	1	21	1
17	5	1	1	21	2
18	7	1	1	21	3
19	4	1	1	22	1
20	6	1	1	22	2
21	7	1	1	22	3
22	6	1	1	23	1
23	7	1	1	23	2
24	8	1	1	23	3
25	3	1	2	1	1
26	4	1	2	1	2
27	5	1	2	1	3
28	4	1	2	2	1
29	6	1	2	2	2
30	8	1	2	2	3
31	7	1	2	3	1
32	8	1	2	3	2
33	9	1	2	3	3
34	9	1	2	4	1
35	8	1	2	4	2
36	9	1	2	4	3
37	7	1	2	5	1
38	9	1	2	5	2
39	9	1	2	5	3
40	3	1	2	21	1
41	6	1	2	21	2
42	7	1	2	21	3
43	5	1	2	22	1
44	7	1	2	22	2
45	8	1	2	22	3
46	6	1	2	23	1
47	7	1	2	23	2
48	8	1	2	23	3
49	4	2	1	6	1
50	5	2	1	6	2
51	6	2	1	6	3
52	6	2	1	7	1
53	7	2	1	7	2
54	8	2	1	7	3
55	7	2	1	8	1
56	8	2	1	8	2
57	9	2	1	8	3
58	7	2	1	9	1
59	8	2	1	9	2
60	9	2	1	9	3
61	3	2	1	10	1
62	5	2	1	10	2
63	6	2	1	10	3

64 3 2 1 21 1
65 5 2 1 21 2
66 7 2 1 21 3
67 5 2 1 22 1
68 5 2 1 22 2
69 7 2 1 22 3
70 6 2 1 23 1
71 7 2 1 23 2
72 9 2 1 23 3
73 5 2 2 6 1
74 6 2 2 6 2
75 7 2 2 6 3
76 6 2 2 7 1
77 7 2 2 7 2
78 7 2 2 7 3
79 7 2 2 8 1
80 9 2 2 8 2
81 8 2 2 8 3
82 7 2 2 9 1
83 7 2 2 9 2
84 9 2 2 9 3
85 4 2 2 10 1
86 5 2 2 10 2
87 7 2 2 10 3
88 2 2 2 21 1
89 4 2 2 21 2
90 5 2 2 21 3
91 6 2 2 22 1
92 7 2 2 22 2
93 8 2 2 22 3
94 6 2 2 23 1
95 7 2 2 23 2
96 8 2 2 23 3
97 4 3 1 11 1
98 5 3 1 11 2
99 6 3 1 11 3
100 7 3 1 12 1
101 8 3 1 12 2
102 8 3 1 12 3
103 6 3 1 13 1
104 7 3 1 13 2
105 7 3 1 13 3
106 7 3 1 14 1
107 7 3 1 14 2
108 9 3 1 14 3
109 2 3 1 15 1
110 3 3 1 15 2
111 4 3 1 15 3

112 4 3 1 21 1
113 5 3 1 21 2
114 5 3 1 21 3
115 6 3 1 22 1
116 7 3 1 22 2
117 8 3 1 22 3
118 7 3 1 23 1
119 8 3 1 23 2
120 8 3 1 23 3
121 5 3 2 11 1
122 5 3 2 11 2
123 6 3 2 11 3
124 8 3 2 12 1
125 8 3 2 12 2
126 9 3 2 12 3
127 7 3 2 13 1
128 7 3 2 13 2
129 9 3 2 13 3
130 7 3 2 14 1
131 8 3 2 14 2
132 8 3 2 14 3
133 4 3 2 15 1
134 5 3 2 15 2
135 7 3 2 15 3
136 3 3 2 21 1
137 6 3 2 21 2
138 6 3 2 21 3
139 7 3 2 22 1
140 7 3 2 22 2
141 9 3 2 22 3
142 7 3 2 23 1
143 8 3 2 23 2
144 9 3 2 23 3
145 1 4 1 16 1
146 3 4 1 16 2
147 5 4 1 16 3
148 2 4 1 17 1
149 4 4 1 17 2
150 5 4 1 17 3
151 3 4 1 18 1
152 4 4 1 18 2
153 6 4 1 18 3
154 4 4 1 19 1
155 5 4 1 19 2
156 7 4 1 19 3
157 5 4 1 20 1
158 5 4 1 20 2
159 7 4 1 20 3

```

160 5 4 1 21 1
161 6 4 1 21 2
162 8 4 1 21 3
163 5 4 1 22 1
164 7 4 1 22 2
165 7 4 1 22 3
166 6 4 1 23 1
167 8 4 1 23 2
168 9 4 1 23 3
169 2 4 2 16 1
170 2 4 2 16 2
171 4 4 2 16 3
172 3 4 2 17 1
173 5 4 2 17 2
174 6 4 2 17 3
175 4 4 2 18 1
176 6 4 2 18 2
177 7 4 2 18 3
178 5 4 2 19 1
179 7 4 2 19 2
180 7 4 2 19 3
181 6 4 2 20 1
182 7 4 2 20 2
183 8 4 2 20 3
184 4 4 2 21 1
185 6 4 2 21 2
186 7 4 2 21 3
187 7 4 2 22 1
188 8 4 2 22 2
189 8 4 2 22 3
190 7 4 2 23 1
191 8 4 2 23 2
192 9 4 2 23 3

```

```
GLM(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3)
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	155	656.12	4.2330	13.446	3.997e-14 ***
RESIDUALS	36	11.33	0.3148		
CORRECTED TOTAL	191	667.45			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	Y Mean	Coef	Var	R-square	Adj R-sq
0.5610836	6.265625	8.95495	0.98302	0.9099118	

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R       3  27.06   9.019  28.6489 1.203e-09 ***  

T       1  10.55  10.547  33.5018 1.334e-06 ***  

R:T     3   2.97   0.991   3.1489  0.036705 *  

G      22 389.01  17.682  56.1668 < 2.2e-16 ***  

T:G    22  18.42   0.837   2.6601  0.004445 **  

R:T:G  12   8.78   0.731   2.3235  0.025315 *  

F       2 164.28  82.141  260.9173 < 2.2e-16 ***  

T:F    2   0.84   0.422   1.3401  0.274574  

G:F    44  23.47   0.533   1.6943  0.053191 .  

T:G:F  44  10.74   0.244   0.7753  0.790640  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R       3  12.49   4.162  13.2206 5.655e-06 ***  

T       1  10.55  10.547  33.5018 1.334e-06 ***  

R:T     3   1.15   0.384   1.2206  0.316281  

G      22 389.01  17.682  56.1668 < 2.2e-16 ***  

T:G    22  18.42   0.837   2.6601  0.004445 **  

R:T:G  12   8.78   0.731   2.3235  0.025315 *  

F       2 164.28  82.141  260.9173 < 2.2e-16 ***  

T:F    2   0.84   0.422   1.3401  0.274574  

G:F    44  23.47   0.533   1.6943  0.053191 .  

T:G:F  44  10.74   0.244   0.7753  0.790640  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R       3  12.49   4.162  13.2206 5.655e-06 ***  

T       1  11.16  11.158  35.4430 8.021e-07 ***  

R:T     3   1.15   0.384   1.2206  0.316281  

G      22 389.01  17.682  56.1668 < 2.2e-16 ***  

T:G    22  18.42   0.837   2.6601  0.004445 **  

R:T:G  12   8.78   0.731   2.3235  0.025315 *  

F       2 120.56  60.282 191.4828 < 2.2e-16 ***  

T:F    2   0.82   0.411   1.3060  0.283432  

G:F    44  23.47   0.533   1.6943  0.053191 .  

T:G:F  44  10.74   0.244   0.7753  0.790640  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3),

```

```
type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F values	Pr(>F)							
R	0.000	0									
T	0.000	0									
G	73.444	2	116.6471 < 2.2e-16 ***								
F	120.563	2	191.4828 < 2.2e-16 ***								
R:T	0.000	0									
T:G	5.778	2	9.1765 0.0006018 ***								
T:F	0.822	2	1.3060 0.2834316								
G:F	23.469	44	1.6943 0.0531910 .								
R:T:G	8.778	12	2.3235 0.0253153 *								
T:G:F	10.740	44	0.7753 0.7906401								
Residuals	11.333	36									

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'. '	0.1	' '	1

6.6 Example 8.1

(15) MODEL

```
ex8.1 = read.table("http://r.acr.kr/split/asbed.txt", header=TRUE)
ex8.1 = af(ex8.1, c("R", "A", "B"))
ex8.1
```

	Y	R	A	B
1	9	1	1	1
2	2	1	1	2
3	8	1	1	7
4	7	1	1	8
5	5	1	1	9
6	9	1	2	1
7	7	1	2	2
8	3	1	2	7
9	5	1	2	8
10	4	1	2	9
11	9	1	3	1
12	2	1	3	2
13	8	1	3	7
14	7	1	3	8
15	5	1	3	9
16	9	1	10	1
17	1	1	10	2

18	9	1	10	7
19	7	1	10	8
20	5	1	10	9
21	9	1	11	1
22	7	1	11	2
23	3	1	11	7
24	5	1	11	8
25	4	1	11	9
26	9	1	12	1
27	2	1	12	2
28	8	1	12	7
29	7	1	12	8
30	5	1	12	9
31	9	1	13	1
32	7	1	13	2
33	3	1	13	7
34	5	1	13	8
35	4	1	13	9
36	9	2	4	3
37	7	2	4	4
38	13	2	4	7
39	8	2	4	8
40	8	2	4	9
41	9	2	5	3
42	12	2	5	4
43	8	2	5	7
44	7	2	5	8
45	8	2	5	9
46	9	2	6	3
47	7	2	6	4
48	13	2	6	7
49	9	2	6	8
50	12	2	6	9
51	9	2	10	3
52	11	2	10	4
53	9	2	10	7
54	7	2	10	8
55	5	2	10	9
56	9	2	11	3
57	7	2	11	4
58	13	2	11	7
59	5	2	11	8
60	4	2	11	9
61	9	2	12	3
62	12	2	12	4
63	8	2	12	7
64	7	2	12	8
65	5	2	12	9

```

66  9 2 13 3
67  7 2 13 4
68  13 2 13 7
69  5 2 13 8
70  4 2 13 9
71  19 3  7 5
72  17 3  7 6
73  13 3  7 7
74  15 3  7 8
75  14 3  7 9
76  19 3  8 5
77  12 3  8 6
78  18 3  8 7
79  17 3  8 8
80  45 3  8 9
81  19 3  9 5
82  17 3  9 6
83  13 3  9 7
84  25 3  9 8
85  34 3  9 9
86  15 3 10 5
87  9 3 10 6
88  11 3 10 7
89  10 3 10 8
90  10 3 10 9
91  9 3 11 5
92  17 3 11 6
93  13 3 11 7
94  15 3 11 8
95  14 3 11 9
96  9 3 12 5
97  12 3 12 6
98  8 3 12 7
99  17 3 12 8
100 15 3 12 9
101 9 3 13 5
102 17 3 13 6
103 13 3 13 7
104 15 3 13 8
105 14 3 13 9

```

```
GLM(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1)
```

```
$ANOVA
Response : Y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL          104 3951.8 37.999
RESIDUALS       0    0.0
```

```
CORRECTED TOTAL 104 3951.8
```

```
$Fitness
  Root MSE  Y Mean Coef Var R-square
    NA 10.0381      NA       1

$`Type I`
  Df  Sum Sq Mean Sq F value Pr(>F)
R     2 1787.68 893.84
A    12 601.24 50.10
R:A    6  24.93  4.16
B     8 156.87 19.61
R:B    4 319.87 79.97
A:B   60 1012.26 16.87
R:A:B 12  49.00  4.08

$`Type II`
  Df  Sum Sq Mean Sq F value Pr(>F)
R     2 372.22 186.111
A    12 601.24 50.103
R:A    6  50.00  8.333
B     8 156.87 19.609
R:B    4  87.44 21.861
A:B   60 1012.26 16.871
R:A:B 12  49.00  4.083

$`Type III`
  Df  Sum Sq Mean Sq F value Pr(>F)
R     2 372.22 186.111
A    12 572.31 47.692
R:A    6  50.00  8.333
B     8 185.85 23.231
R:B    4  87.44 21.861
A:B   60 1012.26 16.871
R:A:B 12  49.00  4.083

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1), type="III",
singular.ok=TRUE) # NOT WORKING
```

6.7 Example 9.2

(16) MODEL

```
ex9.2 = read.table("http://r.acr.kr/split/Ex9.2-sbex.txt", header=TRUE)
ex9.2 = af(ex9.2, c("rep", "hyb", "gen"))
ex9.2
```

yield rep hyb gen

1	48	1	3	1
2	46	1	3	3
3	43	1	3	2
4	46	1	8	1
5	45	1	8	3
6	42	1	8	2
7	46	1	2	1
8	44	1	2	3
9	42	1	2	2
10	42	1	1	1
11	46	1	1	3
12	44	1	1	2
13	43	1	6	1
14	45	1	6	3
15	44	1	6	2
16	47	1	7	1
17	49	1	7	3
18	47	1	7	2
19	48	1	0	1
20	45	1	0	3
21	45	1	0	2
22	46	1	9	1
23	48	1	9	3
24	47	1	9	2
25	46	1	4	1
26	48	1	4	3
27	47	1	4	2
28	49	1	5	1
29	49	1	5	3
30	48	1	5	2
31	46	2	4	2
32	48	2	4	3
33	42	2	4	1
34	45	2	3	2
35	44	2	3	3
36	42	2	3	1
37	46	2	9	2
38	46	2	9	3
39	44	2	9	1
40	45	2	5	2
41	45	2	5	3
42	43	2	5	1
43	43	2	1	2
44	50	2	1	3
45	44	2	1	1
46	48	2	7	2
47	51	2	7	3
48	48	2	7	1

```

49   44   2   2   2
50   48   2   2   3
51   47   2   2   1
52   44   2   8   2
53   46   2   8   3
54   46   2   8   1
55   47   2   6   2
56   48   2   6   3
57   44   2   6   1

GLM(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2)

$ANOVA
Response : yield
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      40 247.813  6.1953  4.4606 0.001119 **
RESIDUALS  16  22.222  1.3889
CORRECTED TOTAL 56 270.035
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE yield Mean Coef Var  R-square  Adj R-sq
1.178511  45.77193 2.574747 0.9177062 0.7119716

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
rep      1  0.239  0.2388  0.1719 0.6839085
hyb      9 66.796  7.4218  5.3437 0.0018370 **
rep:hyb  8 67.000  8.3750  6.0300 0.0011569 **
gen      2 36.351 18.1754 13.0863 0.0004293 ***
rep:gen   2 16.923  8.4616  6.0924 0.0107858 *
hyb:gen  18 60.504  3.3613  2.4201 0.0408545 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
rep      1  0.167  0.1667  0.1200 0.7335481
hyb      9 66.796  7.4218  5.3437 0.0018370 **
rep:hyb  8 67.000  8.3750  6.0300 0.0011569 **
gen      2 36.351 18.1754 13.0863 0.0004293 ***
rep:gen   2 12.111  6.0556  4.3600 0.0308015 *
hyb:gen  18 60.504  3.3613  2.4201 0.0408545 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
```

```

          Df Sum Sq Mean Sq F value    Pr(>F)
rep        1  0.167  0.1667  0.1200 0.7335481
hyb        9 66.796  7.4218  5.3437 0.0018370 **
rep:hyb   8 67.000  8.3750  6.0300 0.0011569 **
gen        2 30.671 15.3356 11.0416 0.0009707 ***
rep:gen   2 12.111  6.0556  4.3600 0.0308015 *
hyb:gen  18 60.504  3.3613  2.4201 0.0408545 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2), type=3,
      singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: yield
          Sum Sq Df F values    Pr(>F)
rep        0.000  0
hyb       66.704  8 6.0033 0.0011847 **
gen       30.671  2 11.0416 0.0009707 ***
rep:hyb   67.000  8 6.0300 0.0011569 **
rep:gen   12.111  2 4.3600 0.0308015 *
hyb:gen  60.504 18 2.4201 0.0408545 *
Residuals 22.222 16
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.8 Example 10.1

(17) MODEL

```

ex10.1 = read.table("http://r.acr.kr/split/Ex10.1-New.txt", header=TRUE)
ex10.1 = af(ex10.1, c("Site", "Block", "A", "B", "C"))
ex10.1

```

	Obs	Site	Block	A	B	C	Yield
1	1	1	R1	A1	B1	C1	6979
2	2	1	R1	A1	B1	C2	7272
3	3	1	R1	A1	B1	C3	7565
4	4	1	R1	A1	B1	C4	7827
5	5	1	R1	A1	B2	C1	8113
6	6	1	R1	A1	B2	C2	7025
7	7	1	R1	A1	B2	C3	7340
8	8	1	R1	A1	B2	C4	7637
9	9	1	R1	A2	B1	C1	7910
10	10	1	R1	A2	B1	C2	8250

11	11	1	R1	A2	B1	C3	8611
12	12	1	R1	A2	B1	C4	8865
13	13	1	R1	A2	B2	C1	9090
14	14	1	R1	A2	B2	C2	9453
15	15	1	R1	A2	B2	C3	9762
16	16	1	R1	A2	B2	C4	8440
17	17	1	R1	A3	B1	C1	8785
18	18	1	R1	A3	B1	C2	8963
19	19	1	R1	A3	B1	C3	9278
20	20	1	R1	A3	B1	C4	11100
21	21	1	R1	A3	B2	C1	10800
22	22	1	R1	A3	B2	C2	10600
23	23	1	R1	A3	B2	C3	10200
24	24	1	R1	A3	B2	C4	10100
25	25	1	R1	A4	B1	C1	9834
26	26	1	R1	A4	B1	C2	10200
27	27	1	R1	A4	B1	C3	10400
28	28	1	R1	A4	B1	C4	10900
29	29	1	R1	A4	B2	C1	11000
30	30	1	R1	A4	B2	C2	12600
31	31	1	R1	A4	B2	C3	12400
32	32	1	R1	A4	B2	C4	12100
33	33	1	R1	A5	B1	C1	11900
34	34	1	R1	A5	B1	C2	11500
35	35	1	R1	A5	B1	C3	11800
36	36	1	R1	A5	B1	C4	12100
37	37	1	R1	A5	B2	C1	12400
38	38	1	R1	A5	B2	C2	12700
39	39	1	R1	A5	B2	C3	12800
40	40	1	R1	A5	B2	C4	13300
41	41	1	R2	A1	B1	C1	7132
42	42	1	R2	A1	B1	C2	7412
43	43	1	R2	A1	B1	C3	7659
44	44	1	R2	A1	B1	C4	7947
45	45	1	R2	A1	B2	C1	8241
46	46	1	R2	A1	B2	C2	7273
47	47	1	R2	A1	B2	C3	7493
48	48	1	R2	A1	B2	C4	7837
49	49	1	R2	A2	B1	C1	8050
50	50	1	R2	A2	B1	C2	8398
51	51	1	R2	A2	B1	C3	8700
52	52	1	R2	A2	B1	C4	8954
53	53	1	R2	A2	B2	C1	9380
54	54	1	R2	A2	B2	C2	9478
55	55	1	R2	A2	B2	C3	10000
56	56	1	R2	A2	B2	C4	8498
57	57	1	R2	A3	B1	C1	8944
58	58	1	R2	A3	B1	C2	9070

59	59	1	R2	A3	B1	C3	9388
60	60	1	R2	A3	B1	C4	11300
61	61	1	R2	A3	B2	C1	10900
62	62	1	R2	A3	B2	C2	10600
63	63	1	R2	A3	B2	C3	10400
64	64	1	R2	A3	B2	C4	10100
65	65	1	R2	A4	B1	C1	10100
66	66	1	R2	A4	B1	C2	10300
67	67	1	R2	A4	B1	C3	10500
68	68	1	R2	A4	B1	C4	10900
69	69	1	R2	A4	B2	C1	11200
70	70	1	R2	A4	B2	C2	12800
71	71	1	R2	A4	B2	C3	12600
72	72	1	R2	A4	B2	C4	12300
73	73	1	R2	A5	B1	C1	11900
74	74	1	R2	A5	B1	C2	11700
75	75	1	R2	A5	B1	C3	11800
76	76	1	R2	A5	B1	C4	12200
77	77	1	R2	A5	B2	C1	12500
78	78	1	R2	A5	B2	C2	12800
79	79	1	R2	A5	B2	C3	12900
80	80	1	R2	A5	B2	C4	13500
81	81	1	R3	A1	B1	C1	6794
82	82	1	R3	A1	B1	C2	7055
83	83	1	R3	A1	B1	C3	7368
84	84	1	R3	A1	B1	C4	7664
85	85	1	R3	A1	B2	C1	7918
86	86	1	R3	A1	B2	C2	6842
87	87	1	R3	A1	B2	C3	7215
88	88	1	R3	A1	B2	C4	7454
89	89	1	R3	A2	B1	C1	7768
90	90	1	R3	A2	B1	C2	7976
91	91	1	R3	A2	B1	C3	8356
92	92	1	R3	A2	B1	C4	8555
93	93	1	R3	A2	B2	C1	8885
94	94	1	R3	A2	B2	C2	9164
95	95	1	R3	A2	B2	C3	9592
96	96	1	R3	A2	B2	C4	8204
97	97	1	R3	A3	B1	C1	8464
98	98	1	R3	A3	B1	C2	8901
99	99	1	R3	A3	B1	C3	9021
100	100	1	R3	A3	B1	C4	11000
101	101	1	R3	A3	B2	C1	10700
102	102	1	R3	A3	B2	C2	10400
103	103	1	R3	A3	B2	C3	10200
104	104	1	R3	A3	B2	C4	9949
105	105	1	R3	A4	B1	C1	9642
106	106	1	R3	A4	B1	C2	9990

107	107	1	R3	A4	B1	C3	10300
108	108	1	R3	A4	B1	C4	10500
109	109	1	R3	A4	B2	C1	10900
110	110	1	R3	A4	B2	C2	12400
111	111	1	R3	A4	B2	C3	12200
112	112	1	R3	A4	B2	C4	11900
113	113	1	R3	A5	B1	C1	11600
114	114	1	R3	A5	B1	C2	11400
115	115	1	R3	A5	B1	C3	11600
116	116	1	R3	A5	B1	C4	11800
117	117	1	R3	A5	B2	C1	12200
118	118	1	R3	A5	B2	C2	12400
119	119	1	R3	A5	B2	C3	12700
120	120	1	R3	A5	B2	C4	13200
121	121	2	R1	A1	B1	C1	6940
122	122	2	R1	A1	B1	C2	7267
123	123	2	R1	A1	B1	C3	7475
124	124	2	R1	A1	B1	C4	7868
125	125	2	R1	A1	B2	C1	8077
126	126	2	R1	A1	B2	C2	7078
127	127	2	R1	A1	B2	C3	7299
128	128	2	R1	A1	B2	C4	7643
129	129	2	R1	A2	B1	C1	7916
130	130	2	R1	A2	B1	C2	8193
131	131	2	R1	A2	B1	C3	8653
132	132	2	R1	A2	B1	C4	8873
133	133	2	R1	A2	B2	C1	9036
134	134	2	R1	A2	B2	C2	9449
135	135	2	R1	A2	B2	C3	9770
136	136	2	R1	A2	B2	C4	8316
137	137	2	R1	A3	B1	C1	8793
138	138	2	R1	A3	B1	C2	8943
139	139	2	R1	A3	B1	C3	9291
140	140	2	R1	A3	B1	C4	11100
141	141	2	R1	A3	B2	C1	10900
142	142	2	R1	A3	B2	C2	10600
143	143	2	R1	A3	B2	C3	10200
144	144	2	R1	A3	B2	C4	9879
145	145	2	R1	A4	B1	C1	9861
146	146	2	R1	A4	B1	C2	10200
147	147	2	R1	A4	B1	C3	10300
148	148	2	R1	A4	B1	C4	10800
149	149	2	R1	A4	B2	C1	10900
150	150	2	R1	A4	B2	C2	12600
151	151	2	R1	A4	B2	C3	12400
152	152	2	R1	A4	B2	C4	12100
153	153	2	R1	A5	B1	C1	11800
154	154	2	R1	A5	B1	C2	11500

155	155	2	R1	A5	B1	C3	11600
156	156	2	R1	A5	B1	C4	12100
157	157	2	R1	A5	B2	C1	12400
158	158	2	R1	A5	B2	C2	12600
159	159	2	R1	A5	B2	C3	12800
160	160	2	R1	A5	B2	C4	13300
161	161	2	R2	A1	B1	C1	6819
162	162	2	R2	A1	B1	C2	7137
163	163	2	R2	A1	B1	C3	7398
164	164	2	R2	A1	B1	C4	7680
165	165	2	R2	A1	B2	C1	7903
166	166	2	R2	A1	B2	C2	6968
167	167	2	R2	A1	B2	C3	7172
168	168	2	R2	A1	B2	C4	7494
169	169	2	R2	A2	B1	C1	7811
170	170	2	R2	A2	B1	C2	8000
171	171	2	R2	A2	B1	C3	8350
172	172	2	R2	A2	B1	C4	8730
173	173	2	R2	A2	B2	C1	8956
174	174	2	R2	A2	B2	C2	9195
175	175	2	R2	A2	B2	C3	9547
176	176	2	R2	A2	B2	C4	8183
177	177	2	R2	A3	B1	C1	8484
178	178	2	R2	A3	B1	C2	8865
179	179	2	R2	A3	B1	C3	9115
180	180	2	R2	A3	B1	C4	11100
181	181	2	R2	A3	B2	C1	10700
182	182	2	R2	A3	B2	C2	10400
183	183	2	R2	A3	B2	C3	10000
184	184	2	R2	A3	B2	C4	9830
185	185	2	R2	A4	B1	C1	9789
186	186	2	R2	A4	B1	C2	9977
187	187	2	R2	A4	B1	C3	10200
188	188	2	R2	A4	B1	C4	10500
189	189	2	R2	A4	B2	C1	10900
190	190	2	R2	A4	B2	C2	12500
191	191	2	R2	A4	B2	C3	12300
192	192	2	R2	A4	B2	C4	11800
193	193	2	R2	A5	B1	C1	11600
194	194	2	R2	A5	B1	C2	11300
195	195	2	R2	A5	B1	C3	11500
196	196	2	R2	A5	B1	C4	12000
197	197	2	R2	A5	B2	C1	12100
198	198	2	R2	A5	B2	C2	12600
199	199	2	R2	A5	B2	C3	12700
200	200	2	R2	A5	B2	C4	13100
201	201	2	R3	A1	B1	C1	7189
202	202	2	R3	A1	B1	C2	7371

203	203	2	R3	A1	B1	C3	7700
204	204	2	R3	A1	B1	C4	8047
205	205	2	R3	A1	B2	C1	8337
206	206	2	R3	A1	B2	C2	7327
207	207	2	R3	A1	B2	C3	7595
208	208	2	R3	A1	B2	C4	7867
209	209	2	R3	A2	B1	C1	8105
210	210	2	R3	A2	B1	C2	8396
211	211	2	R3	A2	B1	C3	8807
212	212	2	R3	A2	B1	C4	8953
213	213	2	R3	A2	B2	C1	9390
214	214	2	R3	A2	B2	C2	9733
215	215	2	R3	A2	B2	C3	9858
216	216	2	R3	A2	B2	C4	8640
217	217	2	R3	A3	B1	C1	9035
218	218	2	R3	A3	B1	C2	9194
219	219	2	R3	A3	B1	C3	9442
220	220	2	R3	A3	B1	C4	11400
221	221	2	R3	A3	B2	C1	11000
222	222	2	R3	A3	B2	C2	10800
223	223	2	R3	A3	B2	C3	10600
224	224	2	R3	A3	B2	C4	10200
225	225	2	R3	A4	B1	C1	9976
226	226	2	R3	A4	B1	C2	10300
227	227	2	R3	A4	B1	C3	10600
228	228	2	R3	A4	B1	C4	11000
229	229	2	R3	A4	B2	C1	11200
230	230	2	R3	A4	B2	C2	12800
231	231	2	R3	A4	B2	C3	12600
232	232	2	R3	A4	B2	C4	12200
233	233	2	R3	A5	B1	C1	11900
234	234	2	R3	A5	B1	C2	11700
235	235	2	R3	A5	B1	C3	11800
236	236	2	R3	A5	B1	C4	12300
237	237	2	R3	A5	B2	C1	12600
238	238	2	R3	A5	B2	C2	12900
239	239	2	R3	A5	B2	C3	13000
240	240	2	R3	A5	B2	C4	13500
241	241	3	R1	A1	B1	C1	7035
242	242	3	R1	A1	B1	C2	7161
243	243	3	R1	A1	B1	C3	7590
244	244	3	R1	A1	B1	C4	7909
245	245	3	R1	A1	B2	C1	8123
246	246	3	R1	A1	B2	C2	7088
247	247	3	R1	A1	B2	C3	7270
248	248	3	R1	A1	B2	C4	7705
249	249	3	R1	A2	B1	C1	7992
250	250	3	R1	A2	B1	C2	8293

251	251	3	R1	A2	B1	C3	8574
252	252	3	R1	A2	B1	C4	8872
253	253	3	R1	A2	B2	C1	9159
254	254	3	R1	A2	B2	C2	9451
255	255	3	R1	A2	B2	C3	9779
256	256	3	R1	A2	B2	C4	8399
257	257	3	R1	A3	B1	C1	8683
258	258	3	R1	A3	B1	C2	8991
259	259	3	R1	A3	B1	C3	9314
260	260	3	R1	A3	B1	C4	11300
261	261	3	R1	A3	B2	C1	10800
262	262	3	R1	A3	B2	C2	10600
263	263	3	R1	A3	B2	C3	10400
264	264	3	R1	A3	B2	C4	10100
265	265	3	R1	A4	B1	C1	9803
266	266	3	R1	A4	B1	C2	10100
267	267	3	R1	A4	B1	C3	10500
268	268	3	R1	A4	B1	C4	10700
269	269	3	R1	A4	B2	C1	11100
270	270	3	R1	A4	B2	C2	12600
271	271	3	R1	A4	B2	C3	12500
272	272	3	R1	A4	B2	C4	12100
273	273	3	R1	A5	B1	C1	11900
274	274	3	R1	A5	B1	C2	11600
275	275	3	R1	A5	B1	C3	11700
276	276	3	R1	A5	B1	C4	12000
277	277	3	R1	A5	B2	C1	12400
278	278	3	R1	A5	B2	C2	12600
279	279	3	R1	A5	B2	C3	12900
280	280	3	R1	A5	B2	C4	13400
281	281	3	R2	A1	B1	C1	7007
282	282	3	R2	A1	B1	C2	7311
283	283	3	R2	A1	B1	C3	7557
284	284	3	R2	A1	B1	C4	7935
285	285	3	R2	A1	B2	C1	8209
286	286	3	R2	A1	B2	C2	7048
287	287	3	R2	A1	B2	C3	7322
288	288	3	R2	A1	B2	C4	7783
289	289	3	R2	A2	B1	C1	8055
290	290	3	R2	A2	B1	C2	8247
291	291	3	R2	A2	B1	C3	8590
292	292	3	R2	A2	B1	C4	8901
293	293	3	R2	A2	B2	C1	9210
294	294	3	R2	A2	B2	C2	9521
295	295	3	R2	A2	B2	C3	9746
296	296	3	R2	A2	B2	C4	8480
297	297	3	R2	A3	B1	C1	8766
298	298	3	R2	A3	B1	C2	9014

299	299	3	R2	A3	B1	C3	9370
300	300	3	R2	A3	B1	C4	11200
301	301	3	R2	A3	B2	C1	11000
302	302	3	R2	A3	B2	C2	10700
303	303	3	R2	A3	B2	C3	10300
304	304	3	R2	A3	B2	C4	10100
305	305	3	R2	A4	B1	C1	9872
306	306	3	R2	A4	B1	C2	10100
307	307	3	R2	A4	B1	C3	10400
308	308	3	R2	A4	B1	C4	10800
309	309	3	R2	A4	B2	C1	11100
310	310	3	R2	A4	B2	C2	12600
311	311	3	R2	A4	B2	C3	12500
312	312	3	R2	A4	B2	C4	12200
313	313	3	R2	A5	B1	C1	11900
314	314	3	R2	A5	B1	C2	11600
315	315	3	R2	A5	B1	C3	11700
316	316	3	R2	A5	B1	C4	12100
317	317	3	R2	A5	B2	C1	12400
318	318	3	R2	A5	B2	C2	12700
319	319	3	R2	A5	B2	C3	12900
320	320	3	R2	A5	B2	C4	13400
321	321	3	R3	A1	B1	C1	7108
322	322	3	R3	A1	B1	C2	7295
323	323	3	R3	A1	B1	C3	7675
324	324	3	R3	A1	B1	C4	7948
325	325	3	R3	A1	B2	C1	8220
326	326	3	R3	A1	B2	C2	7142
327	327	3	R3	A1	B2	C3	7413
328	328	3	R3	A1	B2	C4	7826
329	329	3	R3	A2	B1	C1	8038
330	330	3	R3	A2	B1	C2	8358
331	331	3	R3	A2	B1	C3	8718
332	332	3	R3	A2	B1	C4	9000
333	333	3	R3	A2	B2	C1	9410
334	334	3	R3	A2	B2	C2	9520
335	335	3	R3	A2	B2	C3	9812
336	336	3	R3	A2	B2	C4	8452
337	337	3	R3	A3	B1	C1	8894
338	338	3	R3	A3	B1	C2	9137
339	339	3	R3	A3	B1	C3	9409
340	340	3	R3	A3	B1	C4	11300
341	341	3	R3	A3	B2	C1	10900
342	342	3	R3	A3	B2	C2	10700
343	343	3	R3	A3	B2	C3	10400
344	344	3	R3	A3	B2	C4	10100
345	345	3	R3	A4	B1	C1	9975
346	346	3	R3	A4	B1	C2	10200

347	347	3	R3	A4	B1	C3	10500
348	348	3	R3	A4	B1	C4	10900
349	349	3	R3	A4	B2	C1	11200
350	350	3	R3	A4	B2	C2	12700
351	351	3	R3	A4	B2	C3	12500
352	352	3	R3	A4	B2	C4	12200
353	353	3	R3	A5	B1	C1	11900
354	354	3	R3	A5	B1	C2	11600
355	355	3	R3	A5	B1	C3	11800
356	356	3	R3	A5	B1	C4	12300
357	357	3	R3	A5	B2	C1	12500
358	358	3	R3	A5	B2	C2	12800
359	359	3	R3	A5	B2	C3	12900
360	360	3	R3	A5	B2	C4	13500
361	361	4	R1	A1	B1	C1	6995
362	362	4	R1	A1	B1	C2	7287
363	363	4	R1	A1	B1	C3	7580
364	364	4	R1	A1	B1	C4	7774
365	365	4	R1	A1	B2	C1	8150
366	366	4	R1	A1	B2	C2	7026
367	367	4	R1	A1	B2	C3	7322
368	368	4	R1	A1	B2	C4	7698
369	369	4	R1	A2	B1	C1	7970
370	370	4	R1	A2	B1	C2	8243
371	371	4	R1	A2	B1	C3	8520
372	372	4	R1	A2	B1	C4	8812
373	373	4	R1	A2	B2	C1	9088
374	374	4	R1	A2	B2	C2	9508
375	375	4	R1	A2	B2	C3	9718
376	376	4	R1	A2	B2	C4	8326
377	377	4	R1	A3	B1	C1	8744
378	378	4	R1	A3	B1	C2	9061
379	379	4	R1	A3	B1	C3	9310
380	380	4	R1	A3	B1	C4	11300
381	381	4	R1	A3	B2	C1	10900
382	382	4	R1	A3	B2	C2	10600
383	383	4	R1	A3	B2	C3	10200
384	384	4	R1	A3	B2	C4	9971
385	385	4	R1	A4	B1	C1	9832
386	386	4	R1	A4	B1	C2	10200
387	387	4	R1	A4	B1	C3	10500
388	388	4	R1	A4	B1	C4	10700
389	389	4	R1	A4	B2	C1	11000
390	390	4	R1	A4	B2	C2	12600
391	391	4	R1	A4	B2	C3	12500
392	392	4	R1	A4	B2	C4	12100
393	393	4	R1	A5	B1	C1	11800
394	394	4	R1	A5	B1	C2	11600

395	395	4	R1	A5	B1	C3	11800
396	396	4	R1	A5	B1	C4	12100
397	397	4	R1	A5	B2	C1	12300
398	398	4	R1	A5	B2	C2	12600
399	399	4	R1	A5	B2	C3	12900
400	400	4	R1	A5	B2	C4	13300
401	401	4	R2	A1	B1	C1	6796
402	402	4	R2	A1	B1	C2	7122
403	403	4	R2	A1	B1	C3	7489
404	404	4	R2	A1	B1	C4	7695
405	405	4	R2	A1	B2	C1	8050
406	406	4	R2	A1	B2	C2	7010
407	407	4	R2	A1	B2	C3	7324
408	408	4	R2	A1	B2	C4	7540
409	409	4	R2	A2	B1	C1	7933
410	410	4	R2	A2	B1	C2	8130
411	411	4	R2	A2	B1	C3	8423
412	412	4	R2	A2	B1	C4	8674
413	413	4	R2	A2	B2	C1	9138
414	414	4	R2	A2	B2	C2	9380
415	415	4	R2	A2	B2	C3	9704
416	416	4	R2	A2	B2	C4	8313
417	417	4	R2	A3	B1	C1	8584
418	418	4	R2	A3	B1	C2	8890
419	419	4	R2	A3	B1	C3	9246
420	420	4	R2	A3	B1	C4	11100
421	421	4	R2	A3	B2	C1	10700
422	422	4	R2	A3	B2	C2	10500
423	423	4	R2	A3	B2	C3	10200
424	424	4	R2	A3	B2	C4	9882
425	425	4	R2	A4	B1	C1	9785
426	426	4	R2	A4	B1	C2	10100
427	427	4	R2	A4	B1	C3	10300
428	428	4	R2	A4	B1	C4	10800
429	429	4	R2	A4	B2	C1	11000
430	430	4	R2	A4	B2	C2	12500
431	431	4	R2	A4	B2	C3	12400
432	432	4	R2	A4	B2	C4	12100
433	433	4	R2	A5	B1	C1	11700
434	434	4	R2	A5	B1	C2	11500
435	435	4	R2	A5	B1	C3	11700
436	436	4	R2	A5	B1	C4	12100
437	437	4	R2	A5	B2	C1	12300
438	438	4	R2	A5	B2	C2	12600
439	439	4	R2	A5	B2	C3	12800
440	440	4	R2	A5	B2	C4	13300
441	441	4	R3	A1	B1	C1	7125
442	442	4	R3	A1	B1	C2	7505

```

443 443    4   R3 A1 B1 C3  7752
444 444    4   R3 A1 B1 C4  8099
445 445    4   R3 A1 B2 C1  8409
446 446    4   R3 A1 B2 C2  7332
447 447    4   R3 A1 B2 C3  7512
448 448    4   R3 A1 B2 C4  7917
449 449    4   R3 A2 B1 C1  8176
450 450    4   R3 A2 B1 C2  8382
451 451    4   R3 A2 B1 C3  8861
452 452    4   R3 A2 B1 C4  9056
453 453    4   R3 A2 B2 C1  9419
454 454    4   R3 A2 B2 C2  9700
455 455    4   R3 A2 B2 C3  10000
456 456    4   R3 A2 B2 C4  8573
457 457    4   R3 A3 B1 C1  8953
458 458    4   R3 A3 B1 C2  9278
459 459    4   R3 A3 B1 C3  9538
460 460    4   R3 A3 B1 C4  11400
461 461    4   R3 A3 B2 C1  11100
462 462    4   R3 A3 B2 C2  10800
463 463    4   R3 A3 B2 C3  10600
464 464    4   R3 A3 B2 C4  10300
465 465    4   R3 A4 B1 C1  10000
466 466    4   R3 A4 B1 C2  10400
467 467    4   R3 A4 B1 C3  10700
468 468    4   R3 A4 B1 C4  11000
469 469    4   R3 A4 B2 C1  11200
470 470    4   R3 A4 B2 C2  12900
471 471    4   R3 A4 B2 C3  12600
472 472    4   R3 A4 B2 C4  12400
473 473    4   R3 A5 B1 C1  12000
474 474    4   R3 A5 B1 C2  11700
475 475    4   R3 A5 B1 C3  12000
476 476    4   R3 A5 B1 C4  12300
477 477    4   R3 A5 B2 C1  12500
478 478    4   R3 A5 B2 C2  12900
479 479    4   R3 A5 B2 C3  13000
480 480    4   R3 A5 B2 C4  13700

f10.1 = Yield ~ Site/Block + A/Site + B/Site + A:B + A:B:Site + A:B:Site:Block +
          C + A:C + B:C + A:B:C + C:Site + A:C:Site + B:C:Site + A:B:C:Site
GLM(f10.1, ex10.1)

```

```

$ANOVA
Response : Yield
              Df     Sum Sq Mean Sq F value    Pr(>F)
MODEL           239 1639561484 6860090      2162 < 2.2e-16 ***
RESIDUALS       240    761522    3173

```

CORRECTED TOTAL 479 1640323006

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root	MSE	Yield	Mean	Coef	Var	R-square	Adj R-sq
56.32947	9967.354	0.5651396	0.9995357	0.9990734			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	552717	184239	5.8064e+01	< 2e-16 ***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16 ***
A	4	1387680917	346920229	1.0933e+05	< 2e-16 ***
Site:A	12	34068	2839	8.9470e-01	0.55301
B	1	100939695	100939695	3.1812e+04	< 2e-16 ***
Site:B	3	1618	539	1.6990e-01	0.91662
A:B	4	31444008	7861002	2.4775e+03	< 2e-16 ***
Site:A:B	12	33737	2811	8.8600e-01	0.56185
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155
C	3	19356264	6452088	2.0334e+03	< 2e-16 ***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16 ***
B:C	3	23901388	7967129	2.5109e+03	< 2e-16 ***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16 ***
Site:C	9	47625	5292	1.6677e+00	0.09747 .
Site:A:C	36	104110	2892	9.1140e-01	0.61768
Site:B:C	9	61111	6790	2.1400e+00	0.02701 *
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	552717	184239	5.8064e+01	< 2e-16 ***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16 ***
A	4	1387680917	346920229	1.0933e+05	< 2e-16 ***
Site:A	12	34068	2839	8.9470e-01	0.55301
B	1	100939695	100939695	3.1812e+04	< 2e-16 ***
Site:B	3	1618	539	1.6990e-01	0.91662
A:B	4	31444008	7861002	2.4775e+03	< 2e-16 ***
Site:A:B	12	33737	2811	8.8600e-01	0.56185
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155
C	3	19356264	6452088	2.0334e+03	< 2e-16 ***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16 ***
B:C	3	23901388	7967129	2.5109e+03	< 2e-16 ***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16 ***
Site:C	9	47625	5292	1.6677e+00	0.09747 .
Site:A:C	36	104110	2892	9.1140e-01	0.61768
Site:B:C	9	61111	6790	2.1400e+00	0.02701 *

```

Site:A:B:C      36      82475      2291 7.2200e-01 0.87941
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
          Df   Sum Sq  Mean Sq    F value   Pr(>F)
Site           3     552717   184239 5.8064e+01 < 2e-16 ***
Site:Block     8     7062320   882790 2.7822e+02 < 2e-16 ***
A              4    1387680917 346920229 1.0933e+05 < 2e-16 ***
Site:A         12    34068     2839 8.9470e-01 0.55301
B              1    100939695 100939695 3.1812e+04 < 2e-16 ***
Site:B         3      1618      539 1.6990e-01 0.91662
A:B            4    31444008   7861002 2.4775e+03 < 2e-16 ***
Site:A:B       12    33737     2811 8.8600e-01 0.56185
Site:Block:A:B 72    186911     2596 8.1810e-01 0.84155
C              3    19356264   6452088 2.0334e+03 < 2e-16 ***
A:C            12    26075792   2172983 6.8483e+02 < 2e-16 ***
B:C            3    23901387   7967129 2.5109e+03 < 2e-16 ***
A:B:C          12    41996729   3499727 1.1030e+03 < 2e-16 ***
Site:C          9     47625     5292 1.6677e+00 0.09747 .
Site:A:C        36    104110     2892 9.1140e-01 0.61768
Site:B:C        9     61111     6790 2.1400e+00 0.02701 *
Site:A:B:C      36    82475     2291 7.2200e-01 0.87941
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(f10.1, ex10.1), type=3, singular.ok=TRUE) # NOT OK for Site:Block

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: Yield
          Sum Sq  Df   F values   Pr(>F)
Site           552717  3 5.8064e+01 < 2e-16 ***
A              1387680917  4 1.0933e+05 < 2e-16 ***
B              100939695  1 3.1812e+04 < 2e-16 ***
C              19356264  3 2.0334e+03 < 2e-16 ***
Site:Block      0  0
Site:A          34068  12 8.9470e-01 0.55301
Site:B          1618  3 1.6990e-01 0.91662
A:B            31444008  4 2.4775e+03 < 2e-16 ***
A:C            26075792  12 6.8483e+02 < 2e-16 ***
B:C            23901388  3 2.5109e+03 < 2e-16 ***
Site:C          47625  9 1.6677e+00 0.09747 .
Site:A:B        33737  12 8.8600e-01 0.56185
A:B:C          41996729  12 1.1030e+03 < 2e-16 ***

```

```
Site:A:C      104110  36 9.1140e-01 0.61768
Site:B:C      61111   9 2.1400e+00 0.02701 *
Site:Block:A:B 186911  72 8.1810e-01 0.84155
Site:A:B:C     82475   36 7.2200e-01 0.87941
Residuals      761522 240
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7 Hinkelmann & Kempthorne - Volume 1

Reference

- Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 1 Introduction to Experimental Design. 2e. John Wiley & Sons Inc. 2008.

7.1 p410

(18) MODEL

```
v1p410 = read.table("http://r.acr.kr/kemp/v1p410.txt", head=TRUE)
v1p410$carry = ifelse(v1p410$carry == 0, 3, v1p410$carry)
v1p410 = af(v1p410,c("period", "sequence", "steer", "trt", "carry"))
v1p410
```

	period	sequence	steer	trt	carry	y
1	1	1	1	1	3	50
2	2	1	1	2	1	61
3	3	1	1	3	2	53
4	1	1	2	1	3	55
5	2	1	2	2	1	63
6	3	1	2	3	2	57
7	1	2	3	2	3	44
8	2	2	3	3	2	42
9	3	2	3	1	3	57
10	1	2	4	2	3	51
11	2	2	4	3	2	46
12	3	2	4	1	3	59
13	1	3	5	3	3	35
14	2	3	5	1	3	55
15	3	3	5	2	1	47
16	1	3	6	3	3	41
17	2	3	6	1	3	56
18	3	3	6	2	1	50
19	1	4	7	1	3	54
20	2	4	7	3	1	48
21	3	4	7	2	3	51
22	1	4	8	1	3	58
23	2	4	8	3	1	51
24	3	4	8	2	3	54
25	1	5	9	2	3	50
26	2	5	9	1	2	57
27	3	5	9	3	1	51
28	1	5	10	2	3	55
29	2	5	10	1	2	59
30	3	5	10	3	1	55
31	1	6	11	3	3	41
32	2	6	11	2	3	56

```

33      3       6     11   1     2 58
34      1       6     12   3     3 46
35      2       6     12   2     3 58
36      3       6     12   1     2 61

GLM(y ~ period + sequence + steer:sequence + trt + carry, v1p410) # OK

$ANOVA
Response : y
          Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      17 1302.51  76.618  8.7402 1.572e-05 ***
RESIDUALS   18 157.79   8.766
CORRECTED TOTAL 35 1460.31
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    y Mean Coef Var R-square Adj R-sq
2.960778 52.36111 5.654535 0.8919461 0.7898953

$`Type I`
          Df  Sum Sq Mean Sq F value    Pr(>F)
period      2 292.06 146.028 16.6580 8.038e-05 ***
sequence    5 326.47  65.294  7.4484 0.0006072 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt         2 549.06 274.528 31.3166 1.377e-06 ***
carry       2 16.43   8.215  0.9372 0.4100385
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df  Sum Sq Mean Sq F value    Pr(>F)
period      2 172.31  86.154  9.8279 0.0013030 **
sequence    5 318.69  63.738  7.2709 0.0006954 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt         2 440.61 220.304 25.1311 6.164e-06 ***
carry       2 16.43   8.215  0.9372 0.4100385
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df  Sum Sq Mean Sq F value    Pr(>F)
period      2 172.31  86.154  9.8279 0.0013030 **
sequence    5 318.69  63.738  7.2709 0.0006954 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt         2 440.61 220.304 25.1311 6.164e-06 ***
carry       2 16.43   8.215  0.9372 0.4100385
---

```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(y ~ period + sequence + steer:sequence + trt + carry, v1p410), type=3,
      singular.ok=TRUE) # NOT OK for sequence
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: y

	Sum Sq	Df	F values	Pr(>F)
period	172.31	2	9.8279	0.001303 **
sequence	0.00	0		
trt	440.61	2	25.1311	6.164e-06 ***
carry	16.43	2	0.9372	0.410038
sequence:steer	118.50	6	2.2530	0.084912 .
Residuals	157.79	18		

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8 Searle - Linear Models 2e

Reference

- Searle SR, Gruber MHJ. Linear Models 2e, Kindle Edition. John Wiley & Sons Inc. 2016.

8.1 7.2 (p390, 59%)

(19) MODEL

```
weight = c(8,13,9,12,7,11,6,12,12,14,9,7,14,16,10,14,11,13)
treatment = c("ta","ta","ta","ta","ta","tb","tb","tb","tb","tc","tc","tc",
             "tc","tc","tc")
variety = c("va","va","va","vc","vd","vd","va","vb","vb","vb","vb","vc",
           "vc","vd","vd","vd")
d1 = data.frame(weight, treatment, variety)
GLM(weight ~ treatment*variety, d1)

$ANOVA
Response : weight
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7   82    11.714  2.0918  0.14
RESIDUALS      10   56     5.600
CORRECTED TOTAL 17  138

$Fitness
Root MSE weight Mean Coef Var R-square Adj R-sq
2.366432           11 21.51302 0.5942029 0.3101449

$`Type I`
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2 10.500   5.250  0.9375 0.42348
variety        3 36.786  12.262  2.1896 0.15232
treatment:variety  2 34.714  17.357  3.0995 0.08965 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2   9.486  4.7429  0.8469 0.45731
variety        3  36.786 12.2619  2.1896 0.15232
treatment:variety  2 34.714 17.3571  3.0995 0.08965 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2 12.471  6.2353  1.1134 0.36595
variety        3 34.872 11.6240  2.0757 0.16719
```

```

treatment:variety 2 34.714 17.3571 3.0995 0.08965 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(weight ~ treatment*variety, d1), type=3, singular.ok=TRUE) # NOT OK

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: weight
          Sum Sq Df F values Pr(>F)
treatment      0.000  0
variety        0.000  0
treatment:variety 34.714  2   3.0995 0.08965 .
Residuals     56.000 10
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.2 7.2 (p393, 60%)

(20) MODEL

```

percent = c(31,33,44,36,38,26,37,59,42,42,34,42,28,39,36,32,38,42,36,22,42,46,
          26,37,43)
refinery = c(rep("g",9),rep("n",8),rep("s",8))
process = as.factor(c(1,1,1,1,1,1,2,2,2,1,1,1,2,2,2,2,1,1,1,2,2,2,2,2))
source0 = c("t","t","t","t","o","m","t","t","o","m","i","i","i","t","o","m","m",
           "t","o","i","o","o","m","i","i")
d2 = data.frame(percent, refinery, process, source=source0)
GLM(percent ~ refinery*source, d2)

$ANOVA
Response : percent
          Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       10  442.56  44.256  0.6361 0.7616
RESIDUALS    14  974.00  69.571
CORRECTED TOTAL 24 1416.56

$Fitness
Root MSE percent Mean Coef Var R-square Adj R-sq
8.340949      37.24 22.39782 0.3124188 -0.1787106

$`Type I`
          Df  Sum Sq Mean Sq F value Pr(>F)
refinery      2  20.963  10.481  0.1507 0.8615
source        3 266.124  88.708  1.2751 0.3212
refinery:source 5 155.474  31.095  0.4469 0.8086

```

```
$`Type II`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

refinery       2  25.535 12.767  0.1835 0.8343  

source         3 266.124 88.708  1.2751 0.3212  

refinery:source 5 155.474 31.095  0.4469 0.8086  

$`Type III`  

      Df  Sum Sq Mean Sq F value Pr(>F)  

refinery       2  10.766  5.383  0.0774 0.9259  

source         3 282.633  94.211  1.3542 0.2972  

refinery:source 5 155.474  31.095  0.4469 0.8086  

options(contrasts=c("contr.sum", "contr.poly"))  

Anova(lm(percent ~ refinery*source, d2), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

Response: percent

	Sum Sq	Df	F values	Pr(>F)
refinery	2.52	1	0.0362	0.8518
source	268.19	2	1.9275	0.1822
refinery:source	155.47	5	0.4469	0.8086
Residuals	974.00	14		

9 Web site examples

9.1 <https://github.com/djnavarro/psyr>

```
(21) MODEL  
d21 = read.csv("http://r.acr.kr/psyr/coffee.csv")  
GLM(babble ~ sugar*milk - 1, d21)  
  
$ANOVA  
Response : babble  
          Df Sum Sq Mean Sq F value    Pr(>F)  
MODEL       6 472.54 78.756 298.84 2.39e-12 ***  
RESIDUALS   12  3.16   0.264  
UNCORRECTED TOTAL 18 475.70  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
  
$Fitness  
Root MSE babble Mean Coef Var R-square Adj R-sq  
0.5133631 5.066667 10.13217 0.9933519 0.9900279  
  
$`Type I`  
          Df Sum Sq Mean Sq F value    Pr(>F)  
sugar      3 465.64 155.213 588.9486 2.756e-13 ***  
milk       1  0.96   0.956   3.6279  0.081061 .  
sugar:milk 2  5.94   2.972  11.2769  0.001754 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
  
$`Type II`  
          Df Sum Sq Mean Sq F value    Pr(>F)  
sugar      2 3.0696 1.53482  5.8238 0.017075 *  
milk       1 0.9561 0.95611  3.6279 0.081061 .  
sugar:milk 2 5.9439 2.97193 11.2769 0.001754 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
  
$`Type III`  
CAUTION: Singularity Exists !  
          Df Sum Sq Mean Sq F value    Pr(>F)  
sugar      2 2.1318  1.0659  4.0446 0.045426 *  
milk       1 1.0041  1.0041  3.8102 0.074672 .  
sugar:milk 2 5.9439  2.9719 11.2769 0.001754 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
options(contrasts=c("contr.sum", "contr.poly"))  
r21 = lm(babble ~ sugar*milk - 1, d21)
```

```
anova(r21) # Type I SS OK
```

Analysis of Variance Table

Response: babble

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
sugar	3	465.64	155.213	588.9486	2.756e-13 ***
milk	1	0.96	0.956	3.6279	0.081061 .
sugar:milk	2	5.94	2.972	11.2769	0.001754 **
Residuals	12	3.16	0.264		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
Anova(r21, type=2) # NOT OK
```

Anova Table (Type II tests)

Response: babble

	Sum Sq	Df	F value	Pr(>F)
sugar	453.76	3	573.9233	3.214e-13 ***
milk	0.96	1	3.6279	0.081061 .
sugar:milk	5.94	2	11.2769	0.001754 **
Residuals	3.16	12		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
Anova(r21, type=3) # NOT OK
```

Anova Table (Type III tests)

Response: babble

	Sum Sq	Df	F value	Pr(>F)
sugar	454.77	3	575.1970	3.172e-13 ***
milk	1.00	1	3.8102	0.074672 .
sugar:milk	5.94	2	11.2769	0.001754 **
Residuals	3.16	12		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10 Bioequivalence (BE) data example

(22) MODEL

```
GLM(log(CMAX) ~ SEQ/SUBJ + PRD + TRT, BEdata) # a BE dataset in sasLM package
```

\$ANOVA

```
Response : log(CMAX)
            Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL          48 23.1924 0.48317  5.6278 4.395e-08 ***
RESIDUALS      42  3.6059 0.08585
CORRECTED TOTAL 90 26.7983
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
Root MSE log(CMAX) Mean Coef Var R-square Adj R-sq
0.2930098      6.071036 4.826355 0.8654428 0.7116631
```

\$`Type I`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.6454 0.64544  7.5178  0.008938 **
SEQ:SUBJ 45 22.4395 0.49866  5.8081 3.359e-08 ***
PRD     1  0.0969 0.09686  1.1281  0.294242
TRT     1  0.0106 0.01057  0.1231  0.727410
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.6440 0.64395  7.5005  0.009011 **
SEQ:SUBJ 45 22.5232 0.50052  5.8298 3.173e-08 ***
PRD     1  0.0996 0.09958  1.1599  0.287632
TRT     1  0.0106 0.01057  0.1231  0.727410
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.3368 0.33679  3.9228  0.05421 .
SEQ:SUBJ 45 22.5232 0.50052  5.8298 3.173e-08 ***
PRD     1  0.0996 0.09958  1.1599  0.28763
TRT     1  0.0106 0.01057  0.1231  0.72741
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(log(CMAX) ~ SEQ/SUBJ + PRD + TRT, BEdata), type=3, singular.ok=TRUE)
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: log(CMAX)

	Sum Sq	Df	F values	Pr(>F)
SEQ	0.0000	0		
PRD	0.0996	1	1.1599	0.2876
TRT	0.0106	1	0.1231	0.7274
SEQ:SUBJ	22.5232	45	5.8298	3.173e-08 ***
Residuals	3.6059	42		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1				

11 Session Information

```
R version 4.3.3 (2024-02-29 ucrt)
Platform: x86_64-w64-mingw32/x64 (64-bit)
Running under: Windows 10 x64 (build 19044)
```

```
Matrix products: default
```

```
locale:
```

```
[1] LC_COLLATE=Korean_Korea.utf8  LC_CTYPE=Korean_Korea.utf8
[3] LC_MONETARY=Korean_Korea.utf8 LC_NUMERIC=C
[5] LC_TIME=Korean_Korea.utf8
```

```
time zone: Asia/Seoul
```

```
tzcode source: internal
```

```
attached base packages:
```

```
[1] stats      graphics   grDevices utils      datasets  methods   base
```

```
other attached packages:
```

```
[1] car_3.1-2      carData_3.0-5  sasLM_0.10.3   mvtnorm_1.2-4  rmarkdown_2.25
```

```
loaded via a namespace (and not attached):
```

```
[1] digest_0.6.34    fastmap_1.1.1    xfun_0.41       abind_1.4-5
[5] knitr_1.45       htmltools_0.5.7  tinytex_0.49    cli_3.6.2
[9] compiler_4.3.3   tools_4.3.3     evaluate_0.23  yaml_2.3.8
[13] rlang_1.1.3     MASS_7.3-60.0.1
```