

Validation of ‘sasLM’ Package

Kyun-Seop Bae MD PhD

2024-03-25

Contents

1	Tested Version and Books used for the Validation	4
1.1	Packages Used	4
1.2	Books and Articles used for the Test	4
2	ARS20-8	5
2.1	p8	5
2.2	p42	5
2.3	p101	7
3	Snee EMS ANOVA 1974	10
4	Goodnight	11
4.1	Type I SS	11
4.2	Type II SS	15
4.3	Type III SS	17
5	SAS for Linear Models 4e	20
5.1	Chapter 2	20
5.2	Chapter 3	24
5.3	Chapter 4	28
5.4	Chapter 5	32
5.5	Chapter 6	34
5.6	Chapter 7	38
5.7	Chapter 8	49
5.8	Chapter 11	50
6	Sahai - Unbalanced	65
6.1	Table 11.2	65
6.2	Table 12.6	65
6.3	Table 13.6	66
6.4	Table 14.2	67
6.5	Table 15.3	68
6.6	Table 16.3	69
7	Federer - Variations	72
7.1	Example 1.1	72

7.2	Example 1.2	73
7.3	Example 2.1	74
7.4	Example 2.2	75
7.5	Example 3.1	78
7.6	Example 4.1	85
7.7	Example 5.1	88
7.8	Example 7.1	94
7.9	Example 7.2	96
7.10	Example 7.3	97
7.11	Example 8.1	99
7.12	Example 9.1	100
7.13	Example 9.2	101
7.14	Example 10.1	102
7.15	Example 10.2	105
7.16	Example 11.1	106
7.17	Example 11.2	109
7.18	Example 11.3	114
8	Hinkelmann & Kempthorne - Volume 1	118
8.1	Chapter 6	118
8.2	Chapter 7	119
8.3	Chapter 8	121
8.4	Chapter 9	124
8.5	Chapter 10	129
8.6	Chapter 11	132
8.7	Chapter 12	138
8.8	Chapter 13	141
8.9	Chapter 14	143
9	Hinkelmann & Kempthorne - Volume 2	145
9.1	Chapter 1	145
9.2	Chapter 2	146
9.3	Chapter 6	148
9.4	Chapter 7	150
9.5	Chapter 8	153
9.6	Chapter 9	155
9.7	Chapter 10	160
9.8	Chapter 14	161
9.9	Chapter 16	165
9.10	Chapter 17	170
9.11	Chapter 19	172
10	Lawson - DAE with SAS	175
10.1	Chapter 2	175
10.2	Chapter 3	177
10.3	Chapter 4	184
10.4	Chapter 5	187
10.5	Chapter 7	189

10.6 Chapter 8	192
10.7 Chapter 9	197
10.8 Chapter 11	202
10.9 Chapter 12	205
11 Searle - Linear Models 2e	215
11.1 7.2 (p390, 59%)	215
11.2 7.2 (p393, 60%)	216
12 Web site examples	218
12.1 https://github.com/djnavarro/psyr	218
13 Bioequivalence (BE) data example	220
14 Test Summary	222
15 Session Information	223

1 Tested Version and Books used for the Validation

1.1 Packages Used

- ‘sasLM’ version: 0.10.3
- ‘SAS’ version: 9.4 Licensed and University Edition
- ‘car’ version: 3.1.2
- R version: R version 4.3.3 (2024-02-29 ucrt)

The ‘car’ package is not necessary for ‘sasLM.’ It is used for the comparison of the results.

If you see any difference between ‘car’ and ‘sasLM’, ‘SAS’ results coincide with ‘sasLM’, not with ‘car’.

Before ‘sasLM’ is available on CRAN, you can download using the following command in R.

```
install.packages("sasLM", repos="http://r.acr.kr")
```

1.2 Books and Articles used for the Test

1. Harvey WR. Least-Squares Analysis of Data with Unequal Subclass Frequencies. USDA, Agriculture Research Service, ARS 20-8. 1960. reprinted with corrections as ARS H-4, 1975, also reprinted 1979.
2. Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. J Qual Tech. 1974;6(3):128-137.
3. Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User’s Group, SAS Institute, Raleigh, N.C. 1976.
4. Littell RC, Stroup WW, Freund RJ. SAS for Linear Models 4e. John Wiley & Sons Inc. 2002.
5. Sahai H, Ojeda MM. Analysis of Variance for Random Models Volume 2 Unbalanced Data. 2005.
6. Federer WT, King F. Variations on Split Plot and Split Block Experiment Designs. John Wiley & Sons Inc. 2007.
7. Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 1 Introduction to Experimental Design. 2e. John Wiley & Sons Inc. 2008.
8. Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 2 Advanced Experimental Design. John Wiley & Sons Inc. 2005.
9. Lawson J. Design and Analysis of Experiments with SAS. Taylor and Francis Group. 2010.
10. Searle SR, Gruber MHJ. Linear Models 2e, Kindle Edition. John Wiley & Sons Inc. 2016.

2 ARS20-8

Reference

- Harvey WR. Least-Squares Analysis of Data with Unequal Subclass Frequencies. USDA, Agriculture Research Service, ARS 20-8. 1960. reprinted with corrections as ARS H-4, 1975, also reprinted 1979.

2.1 p8

(1) MODEL

```
p8 = read.csv("C:/G/Rt/ANOVA/ARS20-8p8.csv")
p8 = af(p8, c("PigNo", "Ration"))
GLM(Barrow ~ Ration, p8)
```

```
$ANOVA
Response : Barrow
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       2 11.111  5.5556  1.2626 0.3113
RESIDUALS   15 66.000  4.4000
CORRECTED TOTAL 17 77.111
```

```
$Fitness
Root MSE Barrow Mean Coef Var  R-square    Adj R-sq
2.097618     5.222222 40.16715 0.1440922 0.02997118
```

```
$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
Ration   2 11.111  5.5556  1.2626 0.3113
```

```
$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
Ration   2 11.111  5.5556  1.2626 0.3113
```

```
$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
Ration   2 11.111  5.5556  1.2626 0.3113
```

2.2 p42

(2) MODEL

```
p42 = read.csv("C:/G/Rt/ANOVA/ARS20-8p42.csv")
p42 = af(p42, c("Ration", "Pig", "Sire"))
GLM(Y ~ Sire + Ration, p42)
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       3 20.819  6.9397  1.7259 0.2075
```

```
RESIDUALS      14 56.292  4.0209
CORRECTED TOTAL 17 77.111
```

```
$Fitness
Root MSE   Y Mean Coef Var  R-square  Adj R-sq
  2.00521 5.222222 38.39764 0.2699867 0.1135553
```

```
$`Type I` 
  Df  Sum Sq Mean Sq F value Pr(>F)
Sire    2 11.1111  5.5556  1.3817 0.2834
Ration  1  9.7079  9.7079  2.4144 0.1425
```

```
$`Type II` 
  Df  Sum Sq Mean Sq F value Pr(>F)
Sire    2 15.6829  7.8414  1.9502 0.1790
Ration  1  9.7079  9.7079  2.4144 0.1425
```

```
$`Type III` 
  Df  Sum Sq Mean Sq F value Pr(>F)
Sire    2 15.6829  7.8414  1.9502 0.1790
Ration  1  9.7079  9.7079  2.4144 0.1425
```

(3) MODEL

```
GLM(Y ~ Sire + Ration + Sire:Ration, p42)
```

```
$ANOVA
Response : Y
  Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      5 51.044 10.2089  4.6997 0.01311 *
RESIDUALS 12 26.067  2.1722
CORRECTED TOTAL 17 77.111
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE   Y Mean Coef Var  R-square  Adj R-sq
  1.473846 5.222222 28.22258 0.6619597 0.5211095
```

```
$`Type I` 
  Df  Sum Sq Mean Sq F value   Pr(>F)
Sire       2 11.1111  5.5556  2.5575 0.118799
Ration     1  9.7079  9.7079  4.4691 0.056129 .
Sire:Ration 2 30.2255 15.1127  6.9573 0.009859 **
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II` 
  Df  Sum Sq Mean Sq F value   Pr(>F)
```

```

Sire          2 15.6829  7.8414  3.6099 0.059238 .
Ration        1  9.7079  9.7079  4.4691 0.056129 .
Sire:Ration  2 30.2255 15.1127  6.9573 0.009859 **

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
Sire          2 21.0007 10.5004  4.8339 0.028853 *
Ration        1  3.5919  3.5919  1.6535 0.222736
Sire:Ration  2 30.2255 15.1127  6.9573 0.009859 **

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

2.3 p101

(4) MODEL

```

p101 = read.csv("C:/G/Rt/ANOVA/ARS20-8p101.csv")
p101 = af(p101, c("Line", "Sire", "Dam", "Steer"))
GLM(Gain ~ Line + Sire + Dam + Line:Dam + Age + Weight, p101)

```

```

$ANOVA
Response : Gain
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       16 2.4972 0.156073  3.0675 0.001364 **
RESIDUALS   48 2.4422 0.050879
CORRECTED TOTAL 64 4.9394

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE Gain Mean Coef Var  R-square  Adj R-sq
  0.2255642 2.411385 9.354136 0.5055646 0.3407528

```

```

$`Type I` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
Line       2 0.38009 0.190046  3.7352 0.03107 *
Sire       6 0.92634 0.154391  3.0345 0.01347 *
Dam        2 0.11894 0.059471  1.1689 0.31940
Line:Dam   4 0.64889 0.162222  3.1884 0.02113 *
Age        1 0.16462 0.164622  3.2356 0.07835 .
Weight     1 0.25828 0.258283  5.0764 0.02886 *

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
Line      0

```

```

Sire      6 0.95299 0.15883 3.1217 0.01155 *
Dam       2 0.32039 0.16019 3.1485 0.05190 .
Line:Dam  4 0.46516 0.11629 2.2856 0.07373 .
Age       1 0.34830 0.34830 6.8456 0.01185 *
Weight    1 0.25828 0.25828 5.0764 0.02886 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

CAUTION: Singularity Exists !
      Df  Sum Sq Mean Sq F value Pr(>F)
Line     0
Sire      6 0.95299 0.15883 3.1217 0.01155 *
Dam       2 0.12469 0.06234 1.2253 0.30268
Line:Dam  4 0.46516 0.11629 2.2856 0.07373 .
Age       1 0.34830 0.34830 6.8456 0.01185 *
Weight    1 0.25828 0.25828 5.0764 0.02886 *
---

```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(5) MODEL

```
GLM(Gain ~ Sire + Dam + Line:Dam, p101)
```

```

$ANOVA
Response : Gain
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL   14 2.0743 0.148162 2.5856 0.006996 **
RESIDUALS 50 2.8651 0.057302
CORRECTED TOTAL 64 4.9394
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE Gain Mean Coef Var R-square Adj R-sq
0.2393787 2.411385 9.927022 0.4199453 0.25753

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value Pr(>F)
Sire     8 1.30644 0.163305 2.8499 0.01089 *
Dam      2 0.11894 0.059471 1.0379 0.36172
Dam:Line 4 0.64889 0.162222 2.8310 0.03412 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value Pr(>F)
Sire     6 1.06000 0.176667 3.0831 0.01202 *
Dam      2 0.11894 0.059471 1.0379 0.36172

```

```
Dam:Line 4 0.64889 0.162222 2.8310 0.03412 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  
CAUTION: Singularity Exists !
      Df  Sum Sq  Mean Sq F value Pr(>F)
Sire       6 1.06000 0.176667  3.0831 0.01202 *
Dam        2 0.02569 0.012844  0.2242 0.79999
Dam:Line  4 0.64889 0.162222  2.8310 0.03412 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

3 Snee EMS ANOVA 1974

Reference

- Snee RD. Computation and Use of Expected Mean Squares in Analysis of Variance. J Qual Tech. 1974;6(3):128-137.

(6) MODEL

```
Snee = read.csv("C:/G/Rt/ANOVA/Snee_EMSS_ANOVA1974.csv")
Snee = af(Snee, c("Machine", "Analyst", "Test", "Day"))
GLM(Y ~ Day/Machine/Analyst/Test, Snee)
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      167 751.27 4.4986
RESIDUALS   0    0.00
CORRECTED TOTAL 167 751.27
```

```
$Fitness
Root MSE   Y Mean Coef Var R-square
NA 8.736905     NA       1
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
Day           41 365.58 8.9166
Day:Machine   42 196.59 4.6807
Day:Machine:Analyst 42 118.80 2.8285
Day:Machine:Analyst:Test 42 70.30 1.6739
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
Day           41 365.58 8.9166
Day:Machine   42 196.59 4.6807
Day:Machine:Analyst 42 118.80 2.8285
Day:Machine:Analyst:Test 42 70.30 1.6739
```

```
$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
Day           41 359.44 8.7669
Day:Machine   42 199.40 4.7477
Day:Machine:Analyst 42 118.80 2.8285
Day:Machine:Analyst:Test 42 70.30 1.6739
```

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Day/Machine/Analyst/Test, Snee), type=3, singular.ok=TRUE)
# NOT WORKING
```

4 Goodnight

Reference

- Goodnight JH. The General Linear Models Procedure, Proceedings of the First International SAS User's Group, SAS Institute, Raleigh, N.C. 1976.

4.1 Type I SS

4.1.1 p7

(7) MODEL

```
p7 = read.csv("C:/G/Rt/ANOVA/Goodnight-p7.csv")
p7 = af(p7, c("A", "B"))
GLM(y ~ A + B + A:B, p7)
```

```
$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       3 13.6027  4.5342   2.807 0.1721
RESIDUALS    4  6.4613  1.6153
CORRECTED TOTAL 7 20.0639
```

```
$Fitness
Root MSE y Mean Coef Var R-square Adj R-sq
1.270954 5.4725 23.22438 0.6779647 0.4364382
```

```
$`Type I`
      Df  Sum Sq Mean Sq F value Pr(>F)
A       1 10.8113 10.8113  6.6929 0.06087 .
B       1  1.3122  1.3122  0.8123 0.41839
A:B     1  1.4792  1.4792  0.9157 0.39279
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
      Df  Sum Sq Mean Sq F value Pr(>F)
A       1 10.8113 10.8113  6.6929 0.06087 .
B       1  1.3122  1.3122  0.8123 0.41839
A:B     1  1.4792  1.4792  0.9157 0.39279
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
      Df  Sum Sq Mean Sq F value Pr(>F)
A       1 10.8113 10.8113  6.6929 0.06087 .
B       1  1.3122  1.3122  0.8123 0.41839
A:B     1  1.4792  1.4792  0.9157 0.39279
---
```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(8) MODEL

GLM(y ~ A + A:B + B, p7)

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	13.6027	4.5342	2.807	0.1721
RESIDUALS	4	6.4613	1.6153		
CORRECTED TOTAL	7	20.0639			

\$Fitness

Root	MSE	y	Mean	Coef	Var	R-square	Adj R-sq
1.270954	5.4725	23.22438	0.6779647	0.4364382			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	10.8113	10.8113	6.6929	0.06087 .
A:B	2	2.7914	1.3957	0.8640	0.48764
B	0				

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	10.8113	10.8113	6.6929	0.06087 .
A:B	1	1.4792	1.4792	0.9157	0.39279
B	1	1.3122	1.3122	0.8123	0.41839

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	10.8113	10.8113	6.6929	0.06087 .
A:B	1	1.4792	1.4792	0.9157	0.39279
B	1	1.3122	1.3122	0.8123	0.41839

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(9) MODEL

GLM(y ~ B + A + A:B, p7)

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	13.6027	4.5342	2.807	0.1721
RESIDUALS	4	6.4613	1.6153		

CORRECTED TOTAL 7 20.0639

\$Fitness

Root	MSE	y	Mean	Coef	Var	R-square	Adj	R-sq
1.270954	5.4725	23.22438	0.6779647	0.4364382				

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
A	1	10.8113	10.8113	6.6929	0.06087 .
B:A	1	1.4792	1.4792	0.9157	0.39279

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
A	1	10.8113	10.8113	6.6929	0.06087 .
B:A	1	1.4792	1.4792	0.9157	0.39279

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.41839
A	1	10.8113	10.8113	6.6929	0.06087 .
B:A	1	1.4792	1.4792	0.9157	0.39279

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(10) MODEL

GLM(y ~ B + A:B + A, p7)

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	13.6027	4.5342	2.807	0.1721
RESIDUALS	4	6.4613	1.6153		
CORRECTED TOTAL	7	20.0639			

\$Fitness

Root	MSE	y	Mean	Coef	Var	R-square	Adj	R-sq
1.270954	5.4725	23.22438	0.6779647	0.4364382				

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
B	1	1.3122	1.3122	0.8123	0.4184
B:A	2	12.2905	6.1452	3.8043	0.1187

```

A      0

$`Type II`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

B     1   1.3122  1.3122  0.8123 0.41839  

B:A   1   1.4792  1.4792  0.9157 0.39279  

A     1 10.8113 10.8113  6.6929 0.06087 .  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

B     1   1.3122  1.3122  0.8123 0.41839  

B:A   1   1.4792  1.4792  0.9157 0.39279  

A     1 10.8113 10.8113  6.6929 0.06087 .  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(11) MODEL  

GLM(y ~ A:B + A + B, p7)

$ANOVA  

Response : y  

  Df  Sum Sq Mean Sq F value Pr(>F)  

MODEL          3 13.6027  4.5342   2.807 0.1721  

RESIDUALS       4  6.4613  1.6153  

CORRECTED TOTAL 7 20.0639

$Fitness  

Root MSE y Mean Coef Var R-square Adj R-sq  

1.270954 5.4725 23.22438 0.6779647 0.4364382

$`Type I`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A:B   3 13.603  4.5342   2.807 0.1721  

A      0  

B      0

$`Type II`  

  Df  Sum Sq Mean Sq F value Pr(>F)  

A:B   1   1.4792  1.4792  0.9157 0.39279  

A     1 10.8113 10.8113  6.6929 0.06087 .  

B     1   1.3122  1.3122  0.8123 0.41839  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df  Sum Sq Mean Sq F value Pr(>F)

```

```

A:B 1 1.4792 1.4792 0.9157 0.39279
A 1 10.8113 10.8113 6.6929 0.06087 .
B 1 1.3122 1.3122 0.8123 0.41839
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(12) MODEL

```
GLM(y ~ A:B + A + B, p7)
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      3 13.6027 4.5342 2.807 0.1721
RESIDUALS  4 6.4613 1.6153
CORRECTED TOTAL 7 20.0639
```

```
$Fitness
Root MSE y Mean Coef Var R-square Adj R-sq
1.270954 5.4725 23.22438 0.6779647 0.4364382
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
A:B 3 13.603 4.5342 2.807 0.1721
A 0
B 0
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
A:B 1 1.4792 1.4792 0.9157 0.39279
A 1 10.8113 10.8113 6.6929 0.06087 .
B 1 1.3122 1.3122 0.8123 0.41839
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
A:B 1 1.4792 1.4792 0.9157 0.39279
A 1 10.8113 10.8113 6.6929 0.06087 .
B 1 1.3122 1.3122 0.8123 0.41839
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

4.2 Type II SS

4.2.1 p14

(13) MODEL

```

GLM(y ~ A + B + A:B, p7[-8,]) # p16

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       3 12.7672 4.2557 2.0088 0.2906
RESIDUALS    3  6.3555 2.1185
CORRECTED TOTAL 6 19.1227

$Fitness
Root MSE   y Mean Coef Var R-square Adj R-sq
1.455507 5.342857 27.24211 0.6676471 0.3352941

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
A     1 9.9567 9.9567 4.6999 0.1187
B     1 1.9225 1.9225 0.9075 0.4111
A:B   1 0.8880 0.8880 0.4192 0.5635

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
A     1 11.1715 11.1715 5.2733 0.1053
B     1 1.9225 1.9225 0.9075 0.4111
A:B   1 0.8880 0.8880 0.4192 0.5635

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
A     1 9.5258 9.5258 4.4965 0.1241
B     1 1.3690 1.3690 0.6462 0.4803
A:B   1 0.8880 0.8880 0.4192 0.5635

```

4.2.2 p24

(14) MODEL

```

p24 = read.csv("C:/G/Rt/ANOVA/Goodnight-p24.csv")
p24 = af(p24, c("A", "B", "C"))
GLM(Y ~ A + B + C, p24) # p27

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       6 45.924 7.6540 9.1615 0.00499 ***
RESIDUALS    7  5.848 0.8354
CORRECTED TOTAL 13 51.772
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE   Y Mean Coef Var  R-square  Adj R-sq
  0.9140295 6.159286 14.83986 0.8870405 0.7902181

$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 4.724 4.7235 5.6538 0.04904 *
B 3 37.998 12.6660 15.1606 0.00191 **
C 2 3.203 1.6013 1.9167 0.21686
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
A 0
B 2 0.4424 0.2212 0.2648 0.7747
C 2 3.2025 1.6013 1.9167 0.2169

$`Type III`
CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value Pr(>F)
A 0
B 2 0.4424 0.2212 0.2648 0.7747
C 2 3.2025 1.6013 1.9167 0.2169

```

4.3 Type III SS

4.3.1 p27

(15) MODEL

```

p27 = read.csv("C:/G/Rt/ANOVA/Goodnight-p27.csv")
p27 = af(p27, c("A", "B"))
GLM(y ~ A + B + A:B, p27) # p29

```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      5 128.193 25.6386 53.469 6.77e-05 ***
RESIDUALS  6    2.877  0.4795
CORRECTED TOTAL 11 131.070
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE y Mean Coef Var  R-square  Adj R-sq
  0.6924594 9.34 7.413912 0.9780499 0.9597582

```

\$`Type I`

```

      Df Sum Sq Mean Sq F value    Pr(>F)
A      2 89.580 44.790 93.4102 3.013e-05 ***
B      2 38.542 19.271 40.1901 0.0003351 ***
A:B    1  0.071   0.071  0.1471 0.7145464
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
A      2 126.778 63.389 132.1977 1.093e-05 ***
B      2 38.542 19.271 40.1901 0.0003351 ***
A:B    1  0.071   0.071  0.1471 0.7145464
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value    Pr(>F)
A      2 126.778 63.389 132.1977 1.093e-05 ***
B      2 38.542 19.271 40.1901 0.0003351 ***
A:B    1  0.071   0.071  0.1471 0.7145464
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

4.3.2 p33

(16) MODEL

```

p33 = read.csv("C:/G/Rt/ANOVA/Goodnight-p33.csv")
p33 = af(p33, c("A", "B"))
GLM(y ~ A + B + A:B, p33) # p35

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL        4 34.905  8.7261
RESIDUALS     0  0.000
CORRECTED TOTAL 4 34.905

```

```

$Fitness
Root MSE y Mean Coef Var R-square
      NA  6.946       NA         1

```

```

$`Type I` 
      Df Sum Sq Mean Sq F value Pr(>F)
A      2 11.3739  5.6870
B      1 23.5225 23.5225
A:B    1  0.0081  0.0081

```

```
$`Type II`
```

```

Df   Sum Sq Mean Sq F value Pr(>F)
A     1  3.0276  3.0276
B     1 23.5225 23.5225
A:B   1  0.0081  0.0081

$`Type III`  

CAUTION: Singularity Exists !
Df   Sum Sq Mean Sq F value Pr(>F)
A     1  3.0276  3.0276
B     1 23.5225 23.5225
A:B   1  0.0081  0.0081

options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(y ~ A + B + A:B, p33), type=3, singular.ok=TRUE) # NOT WORKING

```

5 SAS for Linear Models 4e

Reference

- Littell RC, Stroup WW, Freund RJ. SAS for Linear Models 4e. John Wiley & Sons Inc. 2002.

5.1 Chapter 2

5.1.1 p5

(17) MODEL

```
p5 = read.table("C:/G/Rt/SAS4lm/p5.txt", head=TRUE)
GLM(COST ~ CATTLE, p5) # p6 Output 2.2
```

\$ANOVA

Response : COST

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	1	6582.1	6582.1	59.34	6.083e-07 ***
RESIDUALS	17	1885.7	110.9		
CORRECTED TOTAL	18	8467.8			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	MSE	COST	Mean Coef	Var	R-square	Adj R-sq
10.53198	35.29342	29.84119	0.7773107	0.7642113		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
CATTLE	1	6582.1	6582.1	59.34	6.083e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
CATTLE	1	6582.1	6582.1	59.34	6.083e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
CATTLE	1	6582.1	6582.1	59.34	6.083e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5.1.2 p12

(18) MODEL

```

p12 = read.table("C:/G/Rt/SAS4lm/p12.txt", head=TRUE)
GLM(COST ~ CATTLE + CALVES + HOGS + SHEEP, p12)

$ANOVA
Response : COST
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       4 7936.7 1984.18   52.31 2.885e-08 ***
RESIDUALS   14  531.0   37.93
CORRECTED TOTAL 18 8467.8
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE COST Mean Coef Var R-square Adj R-sq
6.158842 35.29342 17.4504 0.9372871 0.9193691

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 6582.1 6582.1 173.5265 2.801e-09 ***
CALVES     1 186.7   186.7   4.9213 0.0435698 *
HOGS       1 489.9   489.9  12.9145 0.0029351 **
SHEEP      1 678.1   678.1  17.8773 0.0008431 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 2200.71 2200.71 58.0183 2.413e-06 ***
CALVES     1 136.08 136.08  3.5876 0.0790616 .
HOGS       1 113.66 113.66  2.9964 0.1054198
SHEEP      1 678.11 678.11 17.8773 0.0008431 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE     1 2200.71 2200.71 58.0183 2.413e-06 ***
CALVES     1 136.08 136.08  3.5876 0.0790616 .
HOGS       1 113.66 113.66  2.9964 0.1054198
SHEEP      1 678.11 678.11 17.8773 0.0008431 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(19) MODEL
GLM(COST ~ CATTLE + CALVES + SHEEP, p12)

```

\$ANOVA
Response : COST

```

          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL        3 7823.1 2607.69  60.673 1.281e-08 ***
RESIDUALS     15 644.7   42.98
CORRECTED TOTAL 18 8467.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE COST Mean Coef Var  R-square Adj R-sq
6.555887 35.29342 18.57538 0.9238649 0.9086379

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE      1 6582.1 6582.1 153.1443 2.835e-09 ***
CALVES      1 186.7   186.7   4.3432 0.0546701 .
SHEEP       1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE      1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES      1 260.6   260.6   6.0634 0.0263909 *
SHEEP       1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE      1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES      1 260.6   260.6   6.0634 0.0263909 *
SHEEP       1 1054.3 1054.3  24.5306 0.0001735 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(20) MODEL
GLM(COST ~ CATTLE + CALVES + offset(1*HOGS) + SHEEP, p12)

$ANOVA
Response : COST
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL        3 7823.1 2607.69  60.673 1.281e-08 ***
RESIDUALS     15 644.7   42.98
CORRECTED TOTAL 18 8467.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness

```

```

Root MSE COST Mean Coef Var R-square Adj R-sq
6.555887 35.29342 18.57538 0.9238649 0.9086379

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE   1 6582.1 6582.1 153.1443 2.835e-09 ***
CALVES   1 186.7 186.7  4.3432 0.0546701 .
SHEEP    1 1054.3 1054.3 24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE   1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES   1 260.6 260.6  6.0634 0.0263909 *
SHEEP    1 1054.3 1054.3 24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III` 
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE   1 2519.8 2519.8 58.6265 1.471e-06 ***
CALVES   1 260.6 260.6  6.0634 0.0263909 *
SHEEP    1 1054.3 1054.3 24.5306 0.0001735 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(21) MODEL

```
GLM(COST ~ CATTLE + CALVES + I(HOGS + SHEEP), p12)
```

```

$ANOVA
Response : COST
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          3 7936.7 2645.6 74.726 3.011e-09 ***
RESIDUALS      15 531.1   35.4
CORRECTED TOTAL 18 8467.8
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE COST Mean Coef Var R-square Adj R-sq
5.950105 35.29342 16.85896 0.937285 0.924742

```

```

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
CATTLE          1 6582.1 6582.1 185.9151 7.406e-10 ***
CALVES          1 186.7 186.7  5.2726  0.03649 *
I(HOGS + SHEEP) 1 1168.0 1168.0 32.9896 3.883e-05 ***

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

CATTLE        1 2215.48 2215.48 62.5775 9.887e-07 ***  

CALVES        1 155.03 155.03  4.3788   0.0538 .  

I(HOGS + SHEEP) 1 1167.96 1167.96 32.9896 3.883e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

CATTLE        1 2215.48 2215.48 62.5775 9.887e-07 ***  

CALVES        1 155.03 155.03  4.3788   0.0538 .  

I(HOGS + SHEEP) 1 1167.96 1167.96 32.9896 3.883e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(22) MODEL  

REG(COST ~ CATTLE + CALVES + I(HOGS + SHEEP) - 1, p12)

$ANOVA  

Response : COST  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL          3 31586 10528.5 306.83 2.398e-14 ***  

RESIDUALS       16    549     34.3  

UNCORRECTED TOTAL 19  32135  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness  

Root MSE COST Mean Coef Var R-square Adj R-sq    PRESS    R2pred  

5.857788 35.29342 16.59739 0.9829151 0.9797116 1365.162 0.9575175

$Coefficients  

      Estimate Std. Error Df Lower CL Upper CL t value    Pr(>|t|)  

CATTLE        3.3000    0.38314 16  2.48782   4.1123 8.6131 2.100e-07 ***  

CALVES        1.9672    0.59108 16  0.71414   3.2202 3.3281 0.004259 **  

I(HOGS + SHEEP) 0.8068    0.13800 16  0.51428   1.0994 5.8466 2.479e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.2 Chapter 3

5.2.1 p63

(23) MODEL

```

p63w = read.table("C:/G/Rt/SAS4lm/p63.txt", header=TRUE)
p63l = reshape(p63w,
  direction = "long",
  varying = list(names(p63w)[2:9]),
  v.names = "fruitwt",
  idvar = c("irrig"),
  timevar = "bloc",
  times = 1:8)
p63l = af(p63l, c("bloc"))
GLM(fruitwt ~ bloc + irrig, p63l) # p64

$ANOVA
Response : fruitwt
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL     11 445334   40485   12.04 6.643e-08 ***
RESIDUALS 28  94147    3362
CORRECTED TOTAL 39 539481
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE fruitwt Mean Coef Var R-square Adj R-sq
57.98607      267.075 21.71153 0.8254864 0.7569274

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
bloc    7 401308   57330 17.0503 1.452e-08 ***
irrig   4  44026   11006  3.2734   0.02539 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
bloc    7 401308   57330 17.0503 1.452e-08 ***
irrig   4  44026   11006  3.2734   0.02539 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
bloc    7 401308   57330 17.0503 1.452e-08 ***
irrig   4  44026   11006  3.2734   0.02539 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.2.2 p72

(24) MODEL

```

p72 = read.table("C:/G/Rt/SAS4lm/p72.txt", header=TRUE)
p72 = af(p72, c("run", "pos", "mat"))
GLM(wtloss ~ run + pos + mat, p72) # p73

$ANOVA
Response : wtloss
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       9 7076.5  786.28  12.837 0.002828 ***
RESIDUALS     6  367.5   61.25
CORRECTED TOTAL 15 7444.0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE wtloss Mean Coef Var  R-square  Adj R-sq
7.826238      239.5  3.26774 0.9506314 0.8765785

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3 986.5  328.83  5.3687 0.0390130 *
pos   3 1468.5  489.50  7.9918 0.0161685 *
mat   3 4621.5 1540.50 25.1510 0.0008498 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3 986.5  328.83  5.3687 0.0390130 *
pos   3 1468.5  489.50  7.9918 0.0161685 *
mat   3 4621.5 1540.50 25.1510 0.0008498 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
run   3 986.5  328.83  5.3687 0.0390130 *
pos   3 1468.5  489.50  7.9918 0.0161685 *
mat   3 4621.5 1540.50 25.1510 0.0008498 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
GLM(shrink ~ run + pos + mat, p72) # p73

```

```

$ANOVA
Response : shrink
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       9 265.75  29.528  9.8426 0.005775 ***
RESIDUALS     6  18.00   3.000

```

```

CORRECTED TOTAL 15 283.75
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE shrink Mean Coef Var R-square Adj R-sq
1.732051      47.125 3.675439 0.9365639 0.8414097

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
run  3 33.25 11.083 3.6944 0.081254 .
pos  3 60.25 20.083 6.6944 0.024212 *
mat  3 172.25 57.417 19.1389 0.001786 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
run  3 33.25 11.083 3.6944 0.081254 .
pos  3 60.25 20.083 6.6944 0.024212 *
mat  3 172.25 57.417 19.1389 0.001786 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
run  3 33.25 11.083 3.6944 0.081254 .
pos  3 60.25 20.083 6.6944 0.024212 *
mat  3 172.25 57.417 19.1389 0.001786 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.2.3 p75

(25) MODEL

```

p75w = read.table("C:/G/Rt/SAS4lm/p75.txt", header=TRUE)
p75l = reshape(p75w,
               direction = "long",
               varying = list(names(p75w)[4:9]),
               v.names = "Y",
               idvar = c("method", "variety", "trt"),
               timevar = "yield",
               times = 1:6)
p75l = af(p75l, c("variety", "yield"))
GLM(Y ~ method*variety, p75l) # p78

```

```

$ANOVA
Response : Y

```

```

              Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          14 1339.0  95.645  4.8674 2.723e-06 ***
RESIDUALS      75 1473.8  19.650
CORRECTED TOTAL 89 2812.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    Y Mean Coef Var R-square Adj R-sq
4.432857 18.43778 24.04225 0.4760484 0.3782441

$`Type I`
              Df Sum Sq Mean Sq F value    Pr(>F)
method         2 953.16  476.58 24.2531 7.525e-09 ***
variety        4   11.38    2.85  0.1448   0.96476
method:variety 8 374.49   46.81   2.3822   0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
              Df Sum Sq Mean Sq F value    Pr(>F)
method         2 953.16  476.58 24.2531 7.525e-09 ***
variety        4   11.38    2.85  0.1448   0.96476
method:variety 8 374.49   46.81   2.3822   0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
              Df Sum Sq Mean Sq F value    Pr(>F)
method         2 953.16  476.58 24.2531 7.525e-09 ***
variety        4   11.38    2.85  0.1448   0.96476
method:variety 8 374.49   46.81   2.3822   0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3 Chapter 4

5.3.1 p94

(26) MODEL

```
p94w = read.table("C:/G/Rt/SAS4lm/p94.txt", head=TRUE)
p94l = reshape(p94w,
               direction = "long",
               varying = list(names(p94w)[3:8]),
               v.names = "ct",
               idvar = c("package"),
               timevar = "sample",
               times = 1:6)
```

```

p941$sampleA = floor((p941$sample + 1)/2)
p941$sampleB = 2 - (p941$sample) %% 2
p941$logct = log10(p941$ct)
p941 = af(p941, c("sample", "sampleA", "sampleB", "package"))
GLM(logct ~ package + sampleA %in% package, p941) # p97

$ANOVA
Response : logct
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      59 50.463 0.85531  22.229 < 2.2e-16 ***
RESIDUALS   60  2.309 0.03848
CORRECTED TOTAL 119 52.772
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE logct Mean Coef Var  R-square  Adj R-sq
0.196156  3.049459 6.432487 0.9562528 0.9132347

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
package     19 30.529 1.60680  41.760 < 2.2e-16 ***
package:sampleA 40 19.934 0.49836  12.952 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
package     19 30.529 1.60680  41.760 < 2.2e-16 ***
package:sampleA 40 19.934 0.49836  12.952 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
package     19 30.529 1.60680  41.760 < 2.2e-16 ***
package:sampleA 40 19.934 0.49836  12.952 < 2.2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3.2 p116

(27) MODEL

```
GLM(Y ~ method + variety + method:variety, p751) # p116
```

```
$ANOVA
Response : Y
```

```

                  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL             14 1339.0  95.645  4.8674 2.723e-06 ***
RESIDUALS        75 1473.8  19.650
CORRECTED TOTAL  89 2812.8
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE      Y Mean Coef Var R-square   Adj R-sq
4.432857 18.43778 24.04225 0.4760484 0.3782441

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
method          2 953.16  476.58 24.2531 7.525e-09 ***
variety         4   11.38    2.85  0.1448   0.96476
method:variety  8 374.49   46.81   2.3822  0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
method          2 953.16  476.58 24.2531 7.525e-09 ***
variety         4   11.38    2.85  0.1448   0.96476
method:variety  8 374.49   46.81   2.3822  0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
method          2 953.16  476.58 24.2531 7.525e-09 ***
variety         4   11.38    2.85  0.1448   0.96476
method:variety  8 374.49   46.81   2.3822  0.02409 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3.3 p122

(28) MODEL

```

p122 = read.table("C:/G/Rt/SAS4lm/p122.txt", header=TRUE)
p122 = af(p122, c("et", "wafer", "pos"))
GLM(resista ~ et + wafer %in% et + pos + et:pos, p122)

```

```

$ANOVA
Response : resista
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       23  9.3250  0.40544  3.6477 0.001263 **
RESIDUALS    24  2.6676  0.11115
CORRECTED TOTAL 47 11.9926

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE resista Mean Coef Var R-square Adj R-sq
  0.3333906    6.002917 5.553811 0.7775641 0.5643963

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
et        3 3.1122 1.03739 9.3333 0.0002851 ***
et:wafer  8 4.2745 0.53431 4.8071 0.0012742 **
pos       3 1.1289 0.37630 3.3855 0.0345139 *
et:pos    9 0.8095 0.08994 0.8092 0.6125279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
et        3 3.1122 1.03739 9.3333 0.0002851 ***
et:wafer  8 4.2745 0.53431 4.8071 0.0012742 **
pos       3 1.1289 0.37630 3.3855 0.0345139 *
et:pos    9 0.8095 0.08994 0.8092 0.6125279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
et        3 3.1122 1.03739 9.3333 0.0002851 ***
et:wafer  8 4.2745 0.53431 4.8071 0.0012742 **
pos       3 1.1289 0.37630 3.3855 0.0345139 *
et:pos    9 0.8095 0.08994 0.8092 0.6125279
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.3.4 p136

(29) MODEL

```

p136 = read.table("C:/G/Rt/SAS4lm/p136.txt", header=TRUE)
p136 = af(p136, "rep")
GLM(drywt ~ rep + cult + rep:cult + inoc + cult:inoc, p136)

```

```

$ANOVA
Response : drywt
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      11 157.208 14.2917   20.26 4.594e-06 ***
RESIDUALS   12   8.465  0.7054
CORRECTED TOTAL 23 165.673
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE drywt Mean Coef Var R-square Adj R-sq
  0.8398909 30.41667 2.761285 0.9489055 0.9020688

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      3 25.320  8.440 11.9646 0.0006428 ***
cult     1  2.407  2.407  3.4117 0.0895283 .
rep:cult 3  9.480  3.160  4.4796 0.0249095 *
inoc     2 118.176 59.088 83.7631 8.919e-08 ***
cult:inoc 2  1.826  0.913  1.2942 0.3097837
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      3 25.320  8.440 11.9646 0.0006428 ***
cult     1  2.407  2.407  3.4117 0.0895283 .
rep:cult 3  9.480  3.160  4.4796 0.0249095 *
inoc     2 118.176 59.088 83.7631 8.919e-08 ***
cult:inoc 2  1.826  0.913  1.2942 0.3097837
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      3 25.320  8.440 11.9646 0.0006428 ***
cult     1  2.407  2.407  3.4117 0.0895283 .
rep:cult 3  9.480  3.160  4.4796 0.0249095 *
inoc     2 118.176 59.088 83.7631 8.919e-08 ***
cult:inoc 2  1.826  0.913  1.2942 0.3097837
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.4 Chapter 5

5.4.1 p142

(30) MODEL

```

p142 = read.table("C:/G/Rt/SAS4lm/p142.txt", header=TRUE, na.strings=".")
p142 = af(p142, c("STUDY", "PATIENT"))
GLM(FLUSH ~ STUDY + TRT, p142) # Incomplete data, 56 lines are truncated.

```

```

$ANOVA
Response : FLUSH
      Df Sum Sq Mean Sq F value    Pr(>F)

```

```

MODEL           5  3619.9   723.98    2.392 0.04607 *
RESIDUALS      71 21489.2   302.67
CORRECTED TOTAL 76 25109.1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE FLUSH Mean Coef Var  R-square   Adj R-sq
17.39728   23.12697  75.2251 0.1441665 0.08389657

$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
STUDY  4 3553.9  888.46  2.9355 0.02638 *
TRT     1    66.0    66.04  0.2182 0.64185
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
STUDY  4 3599.4  899.85  2.9731 0.02496 *
TRT     1    66.0    66.04  0.2182 0.64185
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
STUDY  4 3599.4  899.85  2.9731 0.02496 *
TRT     1    66.0    66.04  0.2182 0.64185
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(31) MODEL
GLM(GLUSH ~ TRT + STUDY + TRT:STUDY, p142) # Different data

```

```

$ANOVA
Response : GLUSH
Df  Sum Sq Mean Sq F value Pr(>F)
MODEL          9  4093.7  454.86  1.4501 0.1851
RESIDUALS      67 21015.4  313.66
CORRECTED TOTAL 76 25109.1

```

```

$Fitness
Root MSE FLUSH Mean Coef Var  R-square   Adj R-sq
17.71054   23.12697  76.57962 0.1630364 0.05060842

```

```

$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
TRT     1    20.5    20.49  0.0653 0.79906

```

```

STUDY      4 3599.4  899.85  2.8688 0.02956 *
TRT:STUDY 4  473.8   118.45  0.3776 0.82383
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

TRT      1   66.0   66.04  0.2105 0.64783  

STUDY     4 3599.4  899.85  2.8688 0.02956 *  

TRT:STUDY 4  473.8   118.45  0.3776 0.82383
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

TRT      1    1.9    1.93  0.0062 0.9377  

STUDY     4 3339.4  834.85  2.6616 0.0400 *  

TRT:STUDY 4  473.8   118.45  0.3776 0.8238
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.5 Chapter 6

5.5.1 p171

(32) MODEL

```
p171 = read.table("C:/G/Rt/SAS4lm/p171.txt", header=TRUE)
GLM(score2 ~ teach, p171) # p173 Output 6.2, p174 Output 6.5
```

```
$ANOVA
Response : score2
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      2   49.74   24.868  0.5598 0.5776
RESIDUALS  28 1243.94   44.426
CORRECTED TOTAL 30 1293.68
```

```
$Fitness
Root MSE score2 Mean Coef Var   R-square   Adj R-sq
  6.66532    73.54839 9.062496 0.03844533 -0.03023714
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)
teach   2 49.736   24.868  0.5598 0.5776
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)
teach   2 49.736   24.868  0.5598 0.5776
```

```
$`Type III`  
Df Sum Sq Mean Sq F value Pr(>F)  
teach 2 49.736 24.868 0.5598 0.5776
```

5.5.2 p188

(33) MODEL

```
p188 = read.table("C:/G/Rt/SAS4lm/p188.txt", header=TRUE)  
p188 = af(p188, c("a", "b"))  
GLM(y ~ a + b + a:b, p188) # p189
```

```
$ANOVA  
Response : y  
Df Sum Sq Mean Sq F value Pr(>F)  
MODEL 5 63.711 12.7422 5.866 0.005724 **  
RESIDUALS 12 26.067 2.1722  
CORRECTED TOTAL 17 89.778  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  
Root MSE y Mean Coef Var R-square Adj R-sq  
1.473846 5.111111 28.83612 0.7096535 0.5886757
```

```
$`Type I`  
Df Sum Sq Mean Sq F value Pr(>F)  
a 1 7.803 7.8028 3.5921 0.082395 .  
b 2 20.492 10.2459 4.7168 0.030798 *  
a:b 2 35.416 17.7082 8.1521 0.005807 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  
Df Sum Sq Mean Sq F value Pr(>F)  
a 1 15.850 15.850 7.2968 0.019265 *  
b 2 20.492 10.246 4.7168 0.030798 *  
a:b 2 35.416 17.708 8.1521 0.005807 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  
Df Sum Sq Mean Sq F value Pr(>F)  
a 1 9.641 9.6407 4.4382 0.056865 .  
b 2 30.866 15.4330 7.1047 0.009212 **  
a:b 2 35.416 17.7082 8.1521 0.005807 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.5.3 p203

(34) MODEL

```
GLM(y ~ a + b + a:b, p188[-8,])
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      4 45.816 11.4539  5.2729 0.01097 *
RESIDUALS  12 26.067  2.1722
CORRECTED TOTAL 16 71.882
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE   y Mean Coef Var R-square Adj R-sq
1.473846 5.352941 27.53339 0.6373704 0.5164939

$`Type I` 
          Df Sum Sq Mean Sq F value Pr(>F)
a      1 2.9252 2.9252  1.3466 0.268432
b      2 13.3224 6.6612  3.0665 0.083997 .
a:b    1 29.5681 29.5681 13.6119 0.003095 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
          Df Sum Sq Mean Sq F value Pr(>F)
a      1 5.5652 5.5652  2.5620 0.135442
b      2 13.3224 6.6612  3.0665 0.083997 .
a:b    1 29.5681 29.5681 13.6119 0.003095 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
          Df Sum Sq Mean Sq F value Pr(>F)
a      1 0.3507 0.3507  0.1615 0.694881
b      2 16.0733 8.0367  3.6997 0.056021 .
a:b    1 29.5681 29.5681 13.6119 0.003095 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.5.4 p215

(35) MODEL

```
p215 = read.table("C:/G/Rt/SAS4lm/p215.txt", header=TRUE)
p215 = af(p215, c("irrig", "reps"))
```

```
GLM(yield ~ irrig/reps + cult + irrig:cult, p215) # p216 Book is wrong.
```

```
$ANOVA
Response : yield
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      11 67.662 6.1511 0.6253 0.7636
RESIDUALS   6 59.023 9.8372
CORRECTED TOTAL 17 126.685
```

```
$Fitness
Root MSE yield Mean Coef Var R-square Adj R-sq
3.136435 30.91667 10.1448 0.5340937 -0.3200677
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
irrig       2 7.320 3.6600 0.3721 0.7042
irrig:reps  6 59.870 9.9783 1.0143 0.4933
cult        1 0.467 0.4672 0.0475 0.8347
irrig:cult   2 0.004 0.0022 0.0002 0.9998
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
irrig       2 7.320 3.6600 0.3721 0.7042
irrig:reps  6 59.870 9.9783 1.0143 0.4933
cult        1 0.467 0.4672 0.0475 0.8347
irrig:cult   2 0.004 0.0022 0.0002 0.9998
```

```
$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
irrig       2 7.320 3.6600 0.3721 0.7042
irrig:reps  6 59.870 9.9783 1.0143 0.4933
cult        1 0.467 0.4672 0.0475 0.8347
irrig:cult   2 0.004 0.0022 0.0002 0.9998
```

```
# Compare with SAS output
```

(36) MODEL

```
GLM(yield ~ reps + irrig + reps:irrig + cult + cult:irrig, p215)
```

```
$ANOVA
Response : yield
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      11 67.662 6.1511 0.6253 0.7636
RESIDUALS   6 59.023 9.8372
CORRECTED TOTAL 17 126.685
```

```
$Fitness
Root MSE yield Mean Coef Var R-square Adj R-sq
```

```

3.136435 30.91667 10.1448 0.5340937 -0.3200677

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

reps       2 49.703 24.8517  2.5263 0.1600  

irrig      2  7.320  3.6600  0.3721 0.7042  

reps:irrig 4 10.167  2.5417  0.2584 0.8944  

cult       1  0.467  0.4672  0.0475 0.8347  

irrig:cult 2  0.004  0.0022  0.0002 0.9998

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

reps       2 49.703 24.8517  2.5263 0.1600  

irrig      2  7.320  3.6600  0.3721 0.7042  

reps:irrig 4 10.167  2.5417  0.2584 0.8944  

cult       1  0.467  0.4672  0.0475 0.8347  

irrig:cult 2  0.004  0.0022  0.0002 0.9998

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

reps       2 49.703 24.8517  2.5263 0.1600  

irrig      2  7.320  3.6600  0.3721 0.7042  

reps:irrig 4 10.167  2.5417  0.2584 0.8944  

cult       1  0.467  0.4672  0.0475 0.8347  

irrig:cult 2  0.004  0.0022  0.0002 0.9998

```

5.6 Chapter 7

5.6.1 p232

(37) MODEL

```

p232 = read.table("C:/G/Rt/SAS4lm/p232.txt", header=TRUE)
p232 = af(p232, c("trt", "rep"))
GLM(final ~ trt + initial, p232) # p233

```

```

$ANOVA
Response : final
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 354.45  70.889  235.05 5.493e-13 ***
RESIDUALS 14   4.22   0.302
CORRECTED TOTAL 19 358.67
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE final Mean Coef Var  R-square  Adj R-sq
0.5491762     30.845 1.780438 0.9882278 0.9840235

```

```
$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

trt      4 198.41  49.602 164.47 1.340e-11 ***  

initial  1 156.04 156.040 517.38 1.867e-12 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

trt      4 12.089   3.022 10.021 0.0004819 ***  

initial  1 156.040 156.040 517.384 1.867e-12 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

trt      4 12.089   3.022 10.021 0.0004819 ***  

initial  1 156.040 156.040 517.384 1.867e-12 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.6.2 p240

(38) MODEL

```
GLM(final ~ initial + trt + trt:initial, p232) # p240
```

```
$ANOVA  

Response : final  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL          9 355.84  39.537 139.51 2.572e-09 ***  

RESIDUALS     10   2.83   0.283  

CORRECTED TOTAL 19 358.67  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

Root MSE	final	Mean Coef	Var	R-square	Adj R-sq
0.5323541	30.845	1.725901	0.9920985	0.9849872	

```
$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

initial      1 342.36  342.36 1208.0336 9.211e-12 ***  

trt         4 12.09    3.02   10.6645  0.001247 **  

initial:trt  4   1.39    0.35    1.2247  0.360175  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```

          Df  Sum Sq Mean Sq F value    Pr(>F)
initial      1 156.040 156.040 550.5987 4.478e-10 ***
trt         4 12.089   3.022 10.6645  0.001247 **
initial:trt  4  1.388   0.347  1.2247  0.360175
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df  Sum Sq Mean Sq F value    Pr(>F)
initial      1 68.529 68.529 241.8091 2.472e-08 ***
trt         4  1.696   0.424   1.4963   0.2752
initial:trt  4  1.388   0.347   1.2247   0.3602
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.3 p241

(39) MODEL

```

p241 = read.table("C:/G/Rt/SAS4lm/p241.txt", header=TRUE)
p241 = af(p241, c("STORE", "DAY"))
GLM(Q1 ~ P1 + DAY + P1:DAY, p241) # p242

```

```

$ANOVA
Response : Q1
          Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      11 1111.52 101.048  4.6445 0.0008119 ***
RESIDUALS   24  522.15  21.756
CORRECTED TOTAL 35 1633.68
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  Q1 Mean Coef Var  R-square  Adj R-sq
4.664374 10.21711 45.65257 0.6803814 0.5338895

```

```

$`Type I`
          Df  Sum Sq Mean Sq F value    Pr(>F)
P1         1 516.59  516.59 23.7444 5.739e-05 ***
DAY        5 430.54   86.11  3.9578  0.009275 **
P1:DAY     5 164.39   32.88  1.5112  0.223566
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
          Df  Sum Sq Mean Sq F value    Pr(>F)
P1         1 696.73  696.73 32.0243 7.925e-06 ***
DAY        5 430.54   86.11  3.9578  0.009275 **
P1:DAY     5 164.39   32.88  1.5112  0.223566

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
P1      1 554.79  554.79 25.4999 3.665e-05 ***
DAY     5 201.17   40.23  1.8493   0.1412
P1:DAY  5 164.39   32.88  1.5112   0.2236
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.6.4 p243

(40) MODEL

```
GLM(Q1 ~ DAY + DAY:P1, p241)
```

```

$ANOVA
Response : Q1
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       11 1111.52 101.048 4.6445 0.0008119 ***
RESIDUALS    24 522.15  21.756
CORRECTED TOTAL 35 1633.68
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```

$Fitness
Root MSE  Q1 Mean Coef Var R-square Adj R-sq
4.664374 10.21711 45.65257 0.6803814 0.5338895
```

```

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
DAY      5 250.40  50.079  2.3018 0.0764717 .
DAY:P1   6 861.13 143.521  6.5967 0.0003239 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
DAY      5 250.40  50.079  2.3018 0.0764717 .
DAY:P1   6 861.13 143.521  6.5967 0.0003239 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
DAY      5 201.17   40.234  1.8493 0.1411648
DAY:P1   6 861.13 143.521  6.5967 0.0003239 ***
---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
REG(Q1 ~ DAY + DAY:P1 - 1, p241) # Output 7.10

```

```

$ANOVA
Response : Q1
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      12 4869.5 405.79 18.652 2.638e-09 ***
RESIDUALS   24  522.2   21.76
UNCORRECTED TOTAL 36 5391.7
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  Q1 Mean Coef Var R-square Adj R-sq PRESS R2pred
4.664374 10.21711 45.65257 0.903156 0.854734 1431.83 0.7344377

```

```

$Coefficients
   Estimate Std. Error Df Lower CL Upper CL t value Pr(>|t|)
DAY1     18.675    14.4110 24  -11.067   48.418  1.2959 0.2073286
DAY2     38.487    15.1094 24     7.303   69.671  2.5472 0.0176863 *
DAY3     45.330    26.1576 24    -8.657   99.316  1.7329 0.0959384 .
DAY4     49.149    16.6092 24    14.870   83.429  2.9592 0.0068366 **
DAY5     77.899    27.5007 24    21.140  134.658  2.8326 0.0092034 **
DAY6     73.273    13.4837 24    45.444  101.102  5.4341 1.39e-05 ***
DAY1:P1  -0.220    0.2915 24    -0.822   0.381 -0.7562 0.4568599
DAY2:P1  -0.624    0.2978 24    -1.238  -0.009 -2.0940 0.0470031 *
DAY3:P1  -0.611    0.5049 24    -1.653   0.431 -1.2102 0.2379998
DAY4:P1  -0.796    0.3193 24    -1.455  -0.137 -2.4914 0.0200350 *
DAY5:P1  -1.196    0.5049 24    -2.238  -0.154 -2.3683 0.0262648 *
DAY6:P1  -1.225    0.2652 24    -1.773  -0.678 -4.6199 0.0001092 ***

```

```

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(41) MODEL

```

GLM(Q1 ~ P1 + DAY + P1:DAY, p241)

```

```

$ANOVA
Response : Q1
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      11 1111.52 101.048 4.6445 0.0008119 ***
RESIDUALS   24  522.15   21.756
CORRECTED TOTAL 35 1633.68
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  Q1 Mean Coef Var R-square Adj R-sq
4.664374 10.21711 45.65257 0.6803814 0.5338895

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

P1       1 516.59  516.59 23.7444 5.739e-05 ***  

DAY      5 430.54   86.11  3.9578  0.009275 **  

P1:DAY   5 164.39   32.88  1.5112  0.223566  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

P1       1 696.73  696.73 32.0243 7.925e-06 ***  

DAY      5 430.54   86.11  3.9578  0.009275 **  

P1:DAY   5 164.39   32.88  1.5112  0.223566  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

P1       1 554.79  554.79 25.4999 3.665e-05 ***  

DAY      5 201.17   40.23  1.8493   0.1412  

P1:DAY   5 164.39   32.88  1.5112   0.2236  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(42) MODEL  

GLM(Q1 ~ STORE + DAY + P1 + P2, p241)

$ANOVA  

Response : Q1  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      12 1225.37 102.114  5.7521 0.0001688 ***  

RESIDUALS   23  408.31  17.753  

CORRECTED TOTAL 35 1633.68  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

Root MSE  Q1 Mean Coef Var  R-square  Adj R-sq  

4.213375 10.21711 41.23842 0.7500678 0.6196683

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

STORE   5 313.42   62.68  3.5310   0.01629 *  

DAY     5 250.40   50.08  2.8210   0.03957 *  

P1      1 622.01  622.01 35.0377 4.924e-06 ***  

P2      1  39.54   39.54  2.2274   0.14917  

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

STORE  5 223.83   44.77  2.5217  0.058346 .  

DAY    5 433.10   86.62  4.8793  0.003456 **  

P1     1 538.17  538.17 30.3150 1.342e-05 ***  

P2     1 39.54   39.54  2.2274  0.149171  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

STORE  5 223.83   44.77  2.5217  0.058346 .  

DAY    5 433.10   86.62  4.8793  0.003456 **  

P1     1 538.17  538.17 30.3150 1.342e-05 ***  

P2     1 39.54   39.54  2.2274  0.149171  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.5 p250

(43) MODEL

```

p250 = read.table("C:/G/Rt/SAS4lm/p250.txt", header=TRUE)
p250 = af(p250, c("variety", "spacing", "plant"))
GLM(lint ~ bollwt + variety + spacing + variety:spacing + variety:spacing:plant,
p250) # p252 Output 7.18, Parameter is different due to different order

```

```

$ANOVA
Response : lint
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          8 31.160  3.8950 80.704 < 2.2e-16 ***
RESIDUALS       40  1.931   0.0483
CORRECTED TOTAL 48 33.091
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE lint Mean Coef Var  R-square  Adj R-sq
  0.2196884  1.77551 12.37325 0.9416596 0.9299915

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt          1 29.0693 29.0693 602.3107 < 2.2e-16 ***  

variety         1  1.2635  1.2635  26.1802 8.158e-06 ***  

spacing         1  0.4666  0.4666   9.6689  0.003447 **  

variety:spacing 1  0.0933  0.0933   1.9325  0.172169  

variety:spacing:plant 4  0.2673  0.0668   1.3847  0.256548

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt       1 11.1186 11.1186 230.3745 < 2.2e-16 ***  

variety      1  1.1973  1.1973  24.8084 1.259e-05 ***  

spacing       1  0.4666  0.4666   9.6689  0.003447 **  

variety:spacing  1  0.0933  0.0933   1.9325  0.172169  

variety:spacing:plant 4  0.2673  0.0668   1.3847  0.256548  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt       1 11.1186 11.1186 230.3745 < 2.2e-16 ***  

variety      1  0.9424  0.9424  19.5269 7.379e-05 ***  

spacing       1  0.3748  0.3748   7.7666  0.008101 **  

variety:spacing  1  0.0479  0.0479   0.9915  0.325350  

variety:spacing:plant 4  0.2673  0.0668   1.3847  0.256548  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.6 p254 Output 7.20

(44) MODEL

```
GLM(lint ~ bollwt + variety + spacing, p250)
```

```
$ANOVA  

Response : lint  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL        3 30.799 10.2665 201.65 < 2.2e-16 ***  

RESIDUALS     45  2.291  0.0509  

CORRECTED TOTAL 48 33.091  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

Root MSE  lint  Mean Coef  Var  R-square  Adj R-sq  

0.2256406 1.77551 12.70849 0.9307624 0.9261466
```

```
$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

bollwt      1 29.0693 29.0693 570.9531 < 2.2e-16 ***  

variety     1  1.2635  1.2635  24.8172 9.777e-06 ***  

spacing     1  0.4666  0.4666   9.1655  0.004072 **  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bollwt   1 11.5717 11.5717 227.2815 < 2.2e-16 ***  

variety  1  1.1973  1.1973  23.5168 1.516e-05 ***  

spacing   1  0.4666  0.4666   9.1655  0.004072 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bollwt   1 11.5717 11.5717 227.2815 < 2.2e-16 ***  

variety  1  1.1973  1.1973  23.5168 1.516e-05 ***  

spacing   1  0.4666  0.4666   9.1655  0.004072 **  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.6.7 p256

(45) MODEL

```
p256 = read.table("C:/G/Rt/SAS4lm/p256.txt", header=TRUE)  

p256b = af(p256, c("bloc", "type", "logdose"))  

GLM(y ~ bloc + type + logdose + type:logdose, p256b) # p258 Output 7.22
```

```
$ANOVA  

Response : y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      8 816.50 102.063  6.0641 0.0014 **  

RESIDUALS  15 252.46  16.831  

CORRECTED TOTAL 23 1068.96  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

Root MSE  y Mean Coef Var R-square Adj R-sq  

4.102506 54.95833 7.464757 0.7638277 0.6378692
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

bloc       3 538.79 179.597 10.6709 0.0005223 ***  

type       1 12.04  12.042  0.7155 0.4109264  

logdose    2 121.58  60.792  3.6120 0.0524231 .  

type:logdose 2 144.08  72.042  4.2804 0.0338265 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)
```

```

bloc      3 538.79 179.597 10.6709 0.0005223 ***
type      1 12.04 12.042 0.7155 0.4109264
logdose   2 121.58 60.792 3.6120 0.0524231 .
type:logdose 2 144.08 72.042 4.2804 0.0338265 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

Df Sum Sq Mean Sq F value    Pr(>F)  

bloc      3 538.79 179.597 10.6709 0.0005223 ***  

type      1 12.04 12.042 0.7155 0.4109264  

logdose   2 121.58 60.792 3.6120 0.0524231 .  

type:logdose 2 144.08 72.042 4.2804 0.0338265 *  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.8 p261 Output 7.27

(46) MODEL

```

p256 = af(p256, c("bloc", "type"))
p256$logd2 = (p256$logdose)^2
GLM(y ~ bloc + type + logdose + logd2 + type:logdose + type:logd2, p256)

```

```

$ANOVA
Response : y
Df  Sum Sq Mean Sq F value Pr(>F)
MODEL     8  816.50 102.062  6.0641 0.0014 **
RESIDUALS 15  252.46  16.831
CORRECTED TOTAL 23 1068.96
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$Fitness

Root MSE	y	Mean Coef	Var	R-square	Adj R-sq
4.102506	54.95833	7.464757	0.7638277	0.6378692	

```

$`Type I`  

Df Sum Sq Mean Sq F value    Pr(>F)  

bloc      3 538.79 179.597 10.6709 0.0005223 ***  

type      1 12.04 12.042 0.7155 0.4109264  

logdose   1 115.56 115.562 6.8662 0.0193005 *  

logd2     1  6.02  6.021 0.3577 0.5586917  

type:logdose 1 138.06 138.062 8.2031 0.0118242 *  

type:logd2  1  6.02  6.021 0.3577 0.5586917  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$`Type II`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
bloc        3 538.79 179.597 10.6709 0.0005223 ***
type        1 12.04 12.042  0.7155 0.4109264
logdose     1  0.39  0.389  0.0231 0.8811262
logd2       1  6.02  6.021  0.3577 0.5586917
type:logdose 1  0.81  0.812  0.0483 0.8290541
type:logd2   1  6.02  6.021  0.3577 0.5586917
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
bloc        3 538.79 179.597 10.6709 0.0005223 ***
type        1 28.12 28.125  1.6711 0.2156736
logdose     1  0.39  0.389  0.0231 0.8811262
logd2       1  6.02  6.021  0.3577 0.5586917
type:logdose 1  0.81  0.812  0.0483 0.8290541
type:logd2   1  6.02  6.021  0.3577 0.5586917
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.6.9 p262 Output 7.28

(47) MODEL

```
GLM(y ~ bloc + type + type:logdose, p256b)
```

```

$ANOVA
Response : y
          Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      8  816.50 102.063  6.0641 0.0014 **
RESIDUALS 15  252.46 16.831
CORRECTED TOTAL 23 1068.96
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$Fitness

Root MSE	y	Mean Coef	Var	R-square	Adj R-sq
4.102506	54.95833	7.464757	0.7638277	0.6378692	

```

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
bloc        3 538.79 179.597 10.6709 0.0005223 ***
type        1 12.04 12.042  0.7155 0.4109264
type:logdose 4 265.67 66.417  3.9462 0.0220552 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$`Type II`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
bloc       3 538.79 179.597 10.6709 0.0005223 ***
type      1 12.04 12.042  0.7155 0.4109264
type:logdose 4 265.67 66.417  3.9462 0.0220552 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
bloc       3 538.79 179.597 10.6709 0.0005223 ***
type      1 12.04 12.042  0.7155 0.4109264
type:logdose 4 265.67 66.417  3.9462 0.0220552 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.7 Chapter 8

5.7.1 p269

(48) MODEL

```

p269 = read.csv("C:/G/Rt/SAS4lm/fev1uni.csv")
p269 = af(p269, c("drug", "hour", "patient"))
GLM(fev1 ~ drug + patient %in% drug + hour + drug:hour, p269) # p271 Output 8.3

```

```

$ANOVA
Response : fev1
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      92 296.65 3.2244 51.078 < 2.2e-16 ***
RESIDUALS  483 30.49  0.0631
CORRECTED TOTAL 575 327.14
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE fev1 Mean Coef Var R-square Adj R-sq
  0.2512505 3.087049 8.138859 0.9067963 0.8890432

```

```

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
drug        2 25.783 12.8913 204.212 < 2.2e-16 ***
drug:patient 69 247.412 3.5857 56.801 < 2.2e-16 ***
hour        7 17.170 2.4529 38.857 < 2.2e-16 ***
drug:hour     14 6.280 0.4486   7.106 1.923e-13 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)

```

```

drug           2  25.783 12.8913 204.212 < 2.2e-16 ***
drug:patient 69 247.412  3.5857  56.801 < 2.2e-16 ***
hour          7  17.170  2.4529  38.857 < 2.2e-16 ***
drug:hour    14   6.280  0.4486   7.106 1.923e-13 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

drug           2  25.783 12.8913 204.212 < 2.2e-16 ***  

drug:patient 69 247.412  3.5857  56.801 < 2.2e-16 ***  

hour          7  17.170  2.4529  38.857 < 2.2e-16 ***  

drug:hour    14   6.280  0.4486   7.106 1.923e-13 ***  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8 Chapter 11

5.8.1 p390

(49) MODEL

```

p390 = read.table("C:/G/Rt/SAS4lm/p390.txt", header=TRUE)
p390$ca = ifelse(p390$a == 0, -1, 1)
p390$cb = ifelse(p390$b == 0, -1, 1)
p390$cc = ifelse(p390$c == 0, -1, 1)
p390 = af(p390, c("rep", "blk", "a", "b", "c"))
GLM(y ~ rep/blk + ca*cb*cc, p390)

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL       12  81.75  6.8125 33.601 6.618e-07 ***  

RESIDUALS    11    2.23  0.2027  

CORRECTED TOTAL 23  83.98
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE  y Mean Coef Var R-square Adj R-sq
  0.4502714 2.37375 18.96878 0.9734438 0.9444733

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep        2  0.051   0.025   0.1256 0.8832237  

rep:blk   3  7.432   2.477  12.2194 0.0007966 ***  

ca         1 21.075  21.075 103.9487 6.090e-07 ***  

cb         1  0.005   0.005   0.0224 0.8837872  

ca:cb     1  1.723   1.723   8.4969 0.0140640 *

```

```

cc      1 37.776 37.776 186.3209 3.063e-08 ***
ca:cc   1  2.318   2.318  11.4332 0.0061285 **
cb:cc   1 11.340  11.340  55.9328 1.232e-05 ***
ca:cb:cc 1  0.031   0.031   0.1511 0.7049490
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep      2  0.051   0.025   0.1256  0.883224  

rep:blk  3  1.668   0.556   2.7416  0.093789 .  

ca       1 21.075  21.075 103.9487 6.090e-07 ***  

cb       1  0.005   0.005   0.0224  0.883787  

ca:cb    1  1.723   1.723   8.4969  0.014064 *  

cc      1 37.776 37.776 186.3209 3.063e-08 ***  

ca:cc   1  2.318   2.318  11.4332 0.006129 **  

cb:cc   1 11.340  11.340  55.9328 1.232e-05 ***  

ca:cb:cc 1  0.031   0.031   0.1511 0.704949  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep      2  0.051   0.025   0.1256  0.883224  

rep:blk  3  1.668   0.556   2.7416  0.093789 .  

ca       1 21.075  21.075 103.9487 6.090e-07 ***  

cb       1  0.005   0.005   0.0224  0.883787  

ca:cb    1  1.723   1.723   8.4969  0.014064 *  

cc      1 37.776 37.776 186.3209 3.063e-08 ***  

ca:cc   1  2.318   2.318  11.4332 0.006129 **  

cb:cc   1 11.340  11.340  55.9328 1.232e-05 ***  

ca:cb:cc 1  0.031   0.031   0.1511 0.704949  

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.2 p394

(50) MODEL

```

p394 = read.table("C:/G/Rt/SAS4lm/p394.txt", header=TRUE)
p394 = af(p394, c("a", "b", "c", "d"))
GLM(y ~ ca*cb*cc*cd, p394)

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      7 6.3559 0.90798
RESIDUALS   0 0.0000
CORRECTED TOTAL 7 6.3559

```

```

$Fitness
Root MSE  y Mean Coef Var R-square
      NA 2.68875       NA       1

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
ca        1 2.07061 2.07061
cb        1 0.59951 0.59951
ca:cb     1 0.00031 0.00031
cc        1 0.00551 0.00551
ca:cc     1 0.80011 0.80011
cb:cc     1 2.82031 2.82031
ca:cb:cc  1 0.05951 0.05951
cd        0
ca:cd     0
cb:cd     0
ca:cb:cd  0
cc:cd     0
ca:cc:cd  0
cb:cc:cd  0
ca:cb:cc:cd 0

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
ca        0
cb        0
ca:cb     0
cc        0
ca:cc     0
cb:cc     0
ca:cb:cc  0
cd        0
ca:cd     0
cb:cd     0
ca:cb:cd  0
cc:cd     0
ca:cc:cd  0
cb:cc:cd  0
ca:cb:cc:cd 0

$`Type III`
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value Pr(>F)
ca        0
cb        0
ca:cb     0
cc        0

```

```

ca:cc      0
cb:cc      0
ca:cb:cc   0
cd         0
ca:cd      0
cb:cd      0
ca:cb:cd   0
cc:cd      0
ca:cc:cd   0
cb:cc:cd   0
ca:cb:cc:cd 0

```

(51) MODEL

```
GLM(y ~ a*b*c*d, p394)
```

```
$ANOVA
Response : y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7 6.3559 0.90798
RESIDUALS      0 0.0000
CORRECTED TOTAL 7 6.3559
```

```
$Fitness
Root MSE  y Mean Coef Var R-square
        NA 2.68875      NA       1
```

```
$`Type I`
              Df Sum Sq Mean Sq F value Pr(>F)
a            1 2.07061 2.07061
b            1 0.59951 0.59951
a:b          1 0.00031 0.00031
c            1 0.00551 0.00551
a:c          1 0.80011 0.80011
b:c          1 2.82031 2.82031
a:b:c        1 0.05951 0.05951
d            0
a:d          0
b:d          0
a:b:d        0
c:d          0
a:c:d        0
b:c:d        0
a:b:c:d     0
```

```
$`Type II`
              Df Sum Sq Mean Sq F value Pr(>F)
a            0
b            0
```

```

a:b      0
c      0
a:c      0
b:c      0
a:b:c    0
d      0
a:d      0
b:d      0
a:b:d    0
c:d      0
a:c:d    0
b:c:d    0
a:b:c:d  0

$`Type III`  

CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value Pr(>F)
a      0
b      0
a:b    0
c      0
a:c    0
b:c    0
a:b:c  0
d      0
a:d    0
b:d    0
a:b:d  0
c:d    0
a:c:d  0
b:c:d  0
a:b:c:d 0

```

5.8.3 p399

(52) MODEL

```

p399 = read.table("C:/G/Rt/SAS4lm/p399.txt", header=TRUE)
p399 = af(p399, c("blk", "trt"))
GLM(y ~ trt + blk, p399)

```

```

$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        8 281.128 35.141 40.822 0.005606 ***
RESIDUALS     3   2.583   0.861
CORRECTED TOTAL 11 283.710
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE y Mean Coef Var R-square Adj R-sq
0.927811 9.75 9.516011 0.9908974 0.9666238

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt  3 102.26 34.086 39.596 0.006515 ***
blk  5 178.87 35.774 41.558 0.005691 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt  3 59.018 19.673 22.853 0.014388 *
blk  5 178.871 35.774 41.558 0.005691 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt  3 59.018 19.673 22.853 0.014388 *
blk  5 178.871 35.774 41.558 0.005691 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.4 p403

```

(53) MODEL

p403 = read.table("C:/G/Rt/SAS4lm/p403.txt", header=TRUE)
p403 = af(p403, c("PATIENT", "VISIT"))
GLM(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT, p403)

$ANOVA
Response : HR
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      29 6408.7 220.99   3.912 3.127e-05 ***
RESIDUALS  42 2372.6   56.49
CORRECTED TOTAL 71 8781.3
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE HR Mean Coef Var R-square Adj R-sq
7.515988 80.80556 9.301326 0.7298134 0.543256

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)

```

```

SEQUENCE      5  508.9  101.79  1.8019 0.133346
SEQUENCE:PATIENT 18 4692.3  260.69  4.6147 2.21e-05 ***
VISIT         2   146.8   73.39  1.2991 0.283499
DRUG          2   668.8  334.39  5.9194 0.005435 **
RESIDS        1   391.0  391.02  6.9219 0.011854 *
RESIDT        1     0.8    0.84  0.0149 0.903511
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
SEQUENCE      5  701.2 140.237  2.4825  0.04665 *
SEQUENCE:PATIENT 18 4692.3 260.685  4.6147 2.21e-05 ***
VISIT         2   146.8  73.389  1.2991 0.28350
DRUG          2   344.0 171.975  3.0443  0.05826 .
RESIDS        1   309.2 309.174  5.4731  0.02414 *
RESIDT        1     0.8    0.840  0.0149 0.90351
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df Sum Sq Mean Sq F value    Pr(>F)
SEQUENCE      5  701.2 140.237  2.4825  0.04665 *
SEQUENCE:PATIENT 18 4692.3 260.685  4.6147 2.21e-05 ***
VISIT         2   146.8  73.389  1.2991 0.28350
DRUG          2   344.0 171.975  3.0443  0.05826 .
RESIDS        1   309.2 309.174  5.4731  0.02414 *
RESIDT        1     0.8    0.840  0.0149 0.90351
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(HR ~ SEQUENCE + PATIENT %in% SEQUENCE + VISIT + DRUG + RESIDS + RESIDT,
          p403), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: HR

	Sum Sq	Df	F values	Pr(>F)
SEQUENCE	0.0	0		
VISIT	146.8	2	1.2991	0.28350
DRUG	343.9	2	3.0443	0.05826 .
RESIDS	309.2	1	5.4731	0.02414 *
RESIDT	0.8	1	0.0149	0.90351
SEQUENCE:PATIENT	4692.3	18	4.6147	2.21e-05 ***
Residuals	2372.6	42		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5.8.5 p409 11.5

(54) MODEL

```
p409 = read.table("C:/G/Rt/SAS4lm/p409.txt", header=TRUE)
GLM(TS ~ SOURCE*AMT, p409) # p410 Output 11.21
```

\$ANOVA
Response : TS

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	5	258.727	51.745	263.71	1.785e-09 ***
RESIDUALS	9	1.766	0.196		
CORRECTED TOTAL	14	260.493			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	TS	Mean Coef	Var	R-square	Adj R-sq
0.4429698	16.03333	2.762805	0.9932206	0.9894542	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
SOURCE	2	98.001	49.001	249.720	1.306e-08 ***
AMT	1	138.245	138.245	704.534	7.392e-10 ***
SOURCE:AMT	2	22.481	11.240	57.284	7.595e-06 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
SOURCE	2	98.001	49.001	249.720	1.306e-08 ***
AMT	1	138.245	138.245	704.534	7.392e-10 ***
SOURCE:AMT	2	22.481	11.240	57.284	7.595e-06 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
SOURCE	2	0.070	0.035	0.179	0.839
AMT	1	138.245	138.245	704.534	7.392e-10 ***
SOURCE:AMT	2	22.481	11.240	57.284	7.595e-06 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5.8.6 p412

(55) MODEL

```
p412 = read.table("C:/G/Rt/SAS4lm/p412.txt", header=TRUE)
GLM(ts ~ source:amt, p412) # p413 Output 11.24
```

\$ANOVA

Response : ts

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	393.01	131.002	903.34	< 2.2e-16 ***
RESIDUALS	16	2.32	0.145		
CORRECTED TOTAL	19	395.33			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	ts	Mean Coef	Var	R-square	Adj R-sq
0.380815	14.535	2.619986	0.9941306	0.9930301	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
source:amt	3	393.01	131	903.34	< 2.2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
source:amt	3	393.01	131	903.34	< 2.2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
source:amt	3	393.01	131	903.34	< 2.2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

5.8.7 p414

(56) MODEL

```
p414 = read.table("C:/G/Rt/SAS4lm/p414.txt", header=TRUE)
p414 = af(p414, c("lackofit"))
GLM(loglivcu ~ level + lackofit, p414) # p415 Output 11.26
```

\$ANOVA

Response : loglivcu

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	3	5.2310	1.74365	155.47	5.018e-14 ***

```

RESIDUALS      20 0.2243 0.01122
CORRECTED TOTAL 23 5.4553
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE loglivcu Mean Coef Var  R-square  Adj R-sq
  0.1059034      1.750172 6.051026 0.9588819 0.9527142

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
level      1 4.9859  4.9859 444.555 3.997e-15 ***
lackofit   2 0.2450  0.1225 10.924 0.0006216 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
level      0
lackofit   2 0.24504 0.12252 10.924 0.0006216 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value    Pr(>F)
level      0
lackofit   2 0.24504 0.12252 10.924 0.0006216 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.8 p417

(57) MODEL

```

p417 = read.table("C:/G/Rt/SAS4lm/p417.txt", header=TRUE)
p417 = af(p417, c("TRT", "POT", "PLANT"))
GLM(Y ~ TRT + POT %in% TRT, p417) # p418 Output 11.28

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          7 267.226  38.175 12.433 7.522e-05 ***
RESIDUALS     13  39.917   3.071
CORRECTED TOTAL 20 307.143
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness

```

```

Root MSE      Y Mean Coef Var  R-square  Adj R-sq
1.752288  15.42857 11.35742 0.8700388 0.8000596

$`Type I` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
TRT       2 236.921 118.460 38.580 3.412e-06 ***
TRT:POT  5  30.306   6.061   1.974     0.1499
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
TRT       2 236.921 118.460 38.580 3.412e-06 ***
TRT:POT  5  30.306   6.061   1.974     0.1499
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
TRT       2 200.111 100.055 32.586 8.626e-06 ***
TRT:POT  5  30.306   6.061   1.974     0.1499
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ TRT + POT %in% TRT, p417), type=3, singular.ok=TRUE) # NOT OK

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y
      Sum Sq Df F values  Pr(>F)
TRT      22.310  1    7.266 0.01835 *
TRT:POT  30.306  5    1.974 0.14991
Residuals 39.917 13
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

5.8.9 p431

(58) MODEL

```

p431 = read.table("C:/G/Rt/SAS4lm/p431.txt", header=TRUE)
p431 = af(p431, c("line", "sire", "agedam", "steerno"))
GLM(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlw, p431)

```

```

$ANOVA
Response : avdlygn

```

```

              Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          16 2.5275 0.157966  3.1437 0.001091 ***
RESIDUALS      48 2.4119 0.050248
CORRECTED TOTAL 64 4.9394
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE avdlygn Mean Coef Var R-square Adj R-sq
0.2241612     2.411385 9.295956 0.511696 0.348928

$`Type I`
              Df Sum Sq Mean Sq F value    Pr(>F)
line           2 0.38009 0.190046  3.7821 0.02983 *
line:sire      6 0.92634 0.154391  3.0726 0.01260 *
agedam         2 0.11894 0.059471  1.1835 0.31497
line:agedam    4 0.64889 0.162222  3.2284 0.02000 *
age            1 0.18349 0.183487  3.6516 0.06200 .
intlwt         1 0.26970 0.269704  5.3674 0.02483 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
              Df Sum Sq Mean Sq F value    Pr(>F)
line           2 0.05526 0.02763  0.5498 0.580636
line:sire      6 0.97389 0.16231  3.2303 0.009543 **
agedam         2 0.33106 0.16553  3.2943 0.045640 *
line:agedam    4 0.45343 0.11336  2.2560 0.076821 .
age            1 0.38128 0.38128  7.5878 0.008277 **
intlwt         1 0.26970 0.26970  5.3674 0.024830 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
              Df Sum Sq Mean Sq F value    Pr(>F)
line           2 0.13620 0.06810  1.3553 0.267560
line:sire      6 0.97389 0.16231  3.2303 0.009543 **
agedam         2 0.13011 0.06505  1.2946 0.283392
line:agedam    4 0.45343 0.11336  2.2560 0.076821 .
age            1 0.38128 0.38128  7.5878 0.008277 **
intlwt         1 0.26970 0.26970  5.3674 0.024830 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
# p433 Output 11.40

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(avdlygn ~ line + line:sire + agedam + line:agedam + age + intlwt, p431),

```

```
type=3, singular.ok=TRUE) # NOT OK for line
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: avdlygn

	Sum Sq	Df	F values	Pr(>F)
line	0.00000	0		
agedam	0.13011	2	1.2946	0.283392
age	0.38128	1	7.5878	0.008277 **
intlwt	0.26970	1	5.3674	0.024830 *
line:sire	0.97389	6	3.2303	0.009543 **
line:agedam	0.45343	4	2.2560	0.076821 .
Residuals	2.41192	48		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(59) MODEL

```
GLM(avdlygn ~ sire + agedam, p431) # # p434 Output 11.41
```

\$ANOVA

Response : avdlygn

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	10	1.4254	0.142538	2.1904	0.03237 *
RESIDUALS	54	3.5140	0.065074		
CORRECTED TOTAL	64	4.9394			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	avdlygn	Mean Coef	Var	R-square	Adj R-sq
0.2550961	2.411385	10.57882	0.2885747	0.1568292	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
sire	8	1.30644	0.163305	2.5095	0.02138 *
agedam	2	0.11894	0.059471	0.9139	0.40707

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
sire	8	1.33017	0.166271	2.5551	0.01937 *
agedam	2	0.11894	0.059471	0.9139	0.40707

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$`Type III`  

      Df  Sum Sq  Mean Sq F value  Pr(>F)  

sire     8 1.33017 0.166271  2.5551  0.01937 *  

agedam  2 0.11894 0.059471  0.9139  0.40707  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

5.8.10 p437 ABSORB option in SAS

(60) MODEL

```
GLM(avdlygn ~ line + sire + agedam + line:agedam + age + intlw, p431)
```

```
$ANOVA  

Response : avdlygn  

      Df  Sum Sq  Mean Sq F value  Pr(>F)  

MODEL      16 2.5275 0.157966  3.1437 0.001091 **  

RESIDUALS   48 2.4119 0.050248  

CORRECTED TOTAL 64 4.9394  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  Root MSE avdlygn Mean Coef Var R-square Adj R-sq  

  0.2241612    2.411385 9.295956 0.511696 0.348928
```

```
$`Type I`  

      Df  Sum Sq  Mean Sq F value  Pr(>F)  

line       2 0.38009 0.190046  3.7821 0.02983 *  

sire       6 0.92634 0.154391  3.0726 0.01260 *  

agedam    2 0.11894 0.059471  1.1835 0.31497  

line:agedam 4 0.64889 0.162222  3.2284 0.02000 *  

age        1 0.18349 0.183487  3.6516 0.06200 .  

intlw      1 0.26970 0.269704  5.3674 0.02483 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df  Sum Sq  Mean Sq F value  Pr(>F)  

line       0  

sire       6 0.97389 0.16231  3.2303 0.009543 **  

agedam    2 0.33106 0.16553  3.2943 0.045640 *  

line:agedam 4 0.45343 0.11336  2.2560 0.076821 .  

age        1 0.38128 0.38128  7.5878 0.008277 **  

intlw      1 0.26970 0.26970  5.3674 0.024830 *  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  
CAUTION: Singularity Exists !  
          Df  Sum Sq Mean Sq F value    Pr(>F)  
line        0  
sire        6 0.97389 0.16231  3.2303 0.009543 **  
agedam      2 0.13011 0.06505  1.2946 0.283392  
line:agedam 4 0.45343 0.11336  2.2560 0.076821 .  
age         1 0.38128 0.38128  7.5878 0.008277 **  
intlwt       1 0.26970 0.26970  5.3674 0.024830 *  
---  
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
# p437 Output 11.43
```

6 Sahai - Unbalanced

Reference

- Sahai H, Ojeda MM. Analysis of Variance for Random Models Volume 2 Unbalanced Data. 2005.

6.1 Table 11.2

(61) MODEL

```
T11.2 = read.table("C:/G/Rt/ANOVA/T11.2.txt")
colnames(T11.2) = c("Group", "Y")
T11.2 = af(T11.2, "Group")
GLM(Y ~ Group, T11.2) # p115

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       4   80.401  20.1003  5.9884 0.0004103 ***
RESIDUALS   59  198.036   3.3565
CORRECTED TOTAL 63  278.438
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE   Y Mean Coef Var R-square Adj R-sq
1.832089 64.15625 2.855667 0.2887583 0.2405385

$`Type I`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Group       4   80.401   20.1   5.9884 0.0004103 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Group       4   80.401   20.1   5.9884 0.0004103 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value    Pr(>F)
Group       4   80.401   20.1   5.9884 0.0004103 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

6.2 Table 12.6

(62) MODEL

```

T12.6 = read.table("C:/G/Rt/ANOVA/T12.6.txt")
colnames(T12.6) = c("Location", "Family", "Y")
T12.6 = af(T12.6, c("Location", "Family"))
GLM(Y ~ Location + Family, T12.6) # p184

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL           7 1.6144 0.230636 8.9562 7.223e-07 ***
RESIDUALS      45 1.1588 0.025752
CORRECTED TOTAL 52 2.7733
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE      Y Mean Coef Var R-square   Adj R-sq
0.160473 0.6279434 25.55532 0.5821469 0.5171475

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
Location  3 0.74036 0.24679 9.5833 5.219e-05 ***
Family    4 0.87410 0.21852 8.4859 3.436e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
Location  3 0.83765 0.27921 10.8426 1.753e-05 ***
Family    4 0.87410 0.21852 8.4859 3.436e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
Location  3 0.83765 0.27921 10.8426 1.753e-05 ***
Family    4 0.87410 0.21852 8.4859 3.436e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.3 Table 13.6

(63) MODEL

```

T13.6 = read.table("C:/G/Rt/ANOVA/T13.6.txt")
colnames(T13.6) = c("Site", "Worker", "Y")
T13.6 = af(T13.6, c("Site", "Worker"))
GLM(Y ~ Site + Worker + Site:Worker, T13.6)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       11 2643.11 240.283  60.323 < 2.2e-16 ***
RESIDUALS    35 139.42   3.983
CORRECTED TOTAL 46 2782.52
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    Y Mean Coef Var R-square Adj R-sq
1.995817 84.18936 2.370629 0.9498962 0.9341493

$`Type I` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
Site        2 1281.55 640.77 160.866 < 2.2e-16 ***
Worker      3 399.27 133.09 33.412 2.234e-10 ***
Site:Worker 6 962.29 160.38 40.264 2.720e-14 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
Site        2 1322.24 661.12 165.973 < 2.2e-16 ***
Worker      3 399.27 133.09 33.412 2.234e-10 ***
Site:Worker 6 962.29 160.38 40.264 2.720e-14 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
Site        2 804.83 402.42 101.026 2.887e-15 ***
Worker      3 430.88 143.63 36.058 8.310e-11 ***
Site:Worker 6 962.29 160.38 40.264 2.720e-14 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.4 Table 14.2

(64) MODEL

```

T14.2 = read.csv("C:/G/Rt/ANOVA/T14.2.csv")
T14.2 = T14.2[!is.na(T14.2$Y),]
T14.2 = af(T14.2, c("Day", "Machine", "Operator"))
GLM(Y ~ Day + Machine + Operator, T14.2)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)

```

```

MODEL           7  6345.4  906.48  8.1297 5.931e-08 ***
RESIDUALS      110 12265.3  111.50
CORRECTED TOTAL 117 18610.6
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    Y Mean Coef Var R-square   Adj R-sq
10.55946 192.1373 5.495791 0.340954 0.2990147

$`Type I` 
  Df Sum Sq Mean Sq F value    Pr(>F)
Day      2 3737.8 1868.90 16.7611 4.426e-07 ***
Machine  2 2440.7 1220.33 10.9445 4.625e-05 ***
Operator 3 166.9   55.63  0.4989     0.6838
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
  Df Sum Sq Mean Sq F value    Pr(>F)
Day      2 3795.1 1897.56 17.0181 3.636e-07 ***
Machine  2 2464.8 1232.39 11.0526 4.227e-05 ***
Operator 3 166.9   55.63  0.4989     0.6838
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df Sum Sq Mean Sq F value    Pr(>F)
Day      2 3795.1 1897.56 17.0181 3.636e-07 ***
Machine  2 2464.8 1232.39 11.0526 4.227e-05 ***
Operator 3 166.9   55.63  0.4989     0.6838
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

6.5 Table 15.3

(65) MODEL

```

T15.3 = read.table("C:/G/Rt/ANOVA/T15.3.txt")
colnames(T15.3) = c("Dam", "Sire", "pH")
T15.3 = af(T15.3, c("Dam", "Sire"))
GLM(pH ~ Dam/Sire, T15.3) # p301

```

```

$ANOVA
Response : pH
  Df  Sum Sq  Mean Sq F value    Pr(>F)
MODEL       36 0.25804 0.0071678 2.8977 7.2e-06 ***
RESIDUALS   123 0.30425 0.0024736
CORRECTED TOTAL 159 0.56229

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE  pH Mean Coef Var R-square Adj R-sq
  0.04973534 7.449813 0.6676053 0.4589074 0.3005388

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
Dam        14 0.178017 0.0127155 5.1405 1.563e-07 ***
Dam:Sire  22 0.080024 0.0036374 1.4705  0.09662 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
Dam        14 0.178017 0.0127155 5.1405 1.563e-07 ***
Dam:Sire  22 0.080024 0.0036374 1.4705  0.09662 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
Dam        14 0.179405 0.0128146 5.1805 1.347e-07 ***
Dam:Sire  22 0.080024 0.0036374 1.4705  0.09662 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(pH ~ Dam/Sire, T15.3), type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

Response: pH

	Sum Sq	Df	F values	Pr(>F)
Dam	0.081011	6	5.4584	4.898e-05 ***
Dam:Sire	0.080024	22	1.4705	0.09662 .
Residuals	0.304253	123		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

6.6 Table 16.3

(66) MODEL

```

T16.3 = read.csv("C:/G/Rt/ANOVA/T16.3.csv")
colnames(T16.3) = c("Plot", "Sample", "Subsample", "Residue")
T16.3 = af(T16.3, c("Plot", "Sample", "Subsample"))
GLM(Residue ~ Plot/Sample/Subsample, T16.3) # p344

$ANOVA
Response : Residue
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      54 3.1897 0.059069  5.8842 1.476e-05 ***
RESIDUALS   22 0.2208 0.010039
CORRECTED TOTAL 76 3.4106
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE Residue Mean Coef Var R-square Adj R-sq
0.100193 0.5023377 19.94535 0.9352456 0.776303

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
Plot      10 1.84041 0.184041 18.3332 1.929e-08 ***
Plot:Sample 22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
Plot      10 1.84041 0.184041 18.3332 1.929e-08 ***
Plot:Sample 22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
Plot      10 1.78686 0.178686 17.7998 2.547e-08 ***
Plot:Sample 22 0.99175 0.045079  4.4906 0.0004209 ***
Plot:Sample:Subsample 22 0.35757 0.016253  1.6191 0.1330632
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(Residue ~ Plot/Sample/Subsample, T16.3), type=3, singular.ok=TRUE)

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```
Response: Residue
          Sum Sq Df F values Pr(>F)
Plot        0.00000  0
Plot:Sample 0.36613 11  3.3156 0.00805 **
Plot:Sample:Subsample 0.35758 22  1.6191 0.13306
Residuals   0.22085 22
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
# NOT OK
```

7 Federer - Variations

Reference

- Federer WT, King F. Variations on Split Plot and Split Block Experiment Designs. John Wiley & Sons Inc. 2007.

7.1 Example 1.1

(67) MODEL

```
ex1.1 = read.table("C:/G/Rt/Split/Ex1.1-spex1.txt", header=TRUE)
ex1.1 = af(ex1.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + A:B, ex1.1)
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	27	4905.7	181.694	10.75	1.994e-10 ***
RESIDUALS	36	608.5	16.902		
CORRECTED TOTAL	63	5514.2			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	Y Mean	Coef Var	R-square	Adj R-sq
4.111227	66.14375	6.215594	0.8896527	0.8068923

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	3	223.8	74.60	4.4138	0.00963 **
A	3	194.6	64.85	3.8370	0.01756 *
R:A	9	158.2	17.58	1.0402	0.42842
B	3	4107.4	1369.13	81.0030	4.441e-16 ***
A:B	9	221.7	24.64	1.4577	0.20117

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	3	223.8	74.60	4.4138	0.00963 **
A	3	194.6	64.85	3.8370	0.01756 *
R:A	9	158.2	17.58	1.0402	0.42842
B	3	4107.4	1369.13	81.0030	4.441e-16 ***
A:B	9	221.7	24.64	1.4577	0.20117

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
R       3  223.8   74.60  4.4138  0.00963 ***
A       3  194.6   64.85  3.8370  0.01756 *
R:A     9  158.2   17.58  1.0402  0.42842
B      34 107.4 1369.13 81.0030 4.441e-16 ***
A:B     9  221.7   24.64  1.4577  0.20117
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.2 Example 1.2

(68) MODEL

```

ex1.2 = read.table("C:/G/Rt/Split/Ex1.2-spex2.txt", header=TRUE)
ex1.2 = af(ex1.2, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + A:B, ex1.2)

```

```

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL        47 35573  756.88  31.243 < 2.2e-16 ***
RESIDUALS     48   1163   24.23
CORRECTED TOTAL 95 36736
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE   Y Mean Coef Var R-square Adj R-sq
 4.92196 25.30208 19.45279 0.9683464 0.9373523

```

```

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
R       2   38.6    19.3   0.7963 0.4568480
A       7  763.2   109.0   4.5003 0.0006418 ***
R:A    14 1377.2    98.4   4.0608 0.0001343 ***
B       3 30774.3 10258.1 423.4386 < 2.2e-16 ***
A:B    21 2620.1   124.8   5.1502 1.327e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
R       2   38.6    19.3   0.7963 0.4568480
A       7  763.2   109.0   4.5003 0.0006418 ***
R:A    14 1377.2    98.4   4.0608 0.0001343 ***
B       3 30774.3 10258.1 423.4386 < 2.2e-16 ***
A:B    21 2620.1   124.8   5.1502 1.327e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     2   38.6   19.3   0.7963 0.4568480  

A     7  763.2   109.0   4.5003 0.0006418 ***  

R:A  14 1377.2    98.4   4.0608 0.0001343 ***  

B     3 30774.3 10258.1 423.4386 < 2.2e-16 ***  

A:B  21 2620.1   124.8   5.1502 1.327e-06 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7.3 Example 2.1

(69) MODEL

```
ex2.1 = read.table("C:/G/Rt/Split/sbex.txt", header=TRUE)  

colnames(ex2.1) = c("Y", "R", "A", "B")  

ex2.1 = af(ex2.1, c("R", "A", "B"))  

GLM(Y ~ R + A + R:A + B + R:B + A:B, ex2.1)
```

```
$ANOVA  

Response : Y  

  Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL          41 274.750  6.7012  5.1475 0.0002305 ***  

RESIDUALS      18  23.433  1.3019  

CORRECTED TOTAL 59 298.183  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

Root MSE    Y Mean Coef Var R-square  Adj R-sq  

  1.140987 45.61667 2.501251 0.921413 0.7424093
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     1   2.817   2.8167  2.1636 0.1585807  

A     9  77.683   8.6315  6.6302 0.0003456 ***  

R:A  9 81.017   9.0019  6.9147 0.0002658 ***  

B     2 35.433  17.7167 13.6088 0.0002510 ***  

R:B  2 16.233   8.1167  6.2347 0.0087635 **  

A:B 18 61.567   3.4204  2.6273 0.0236253 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     1   2.817   2.8167  2.1636 0.1585807  

A     9  77.683   8.6315  6.6302 0.0003456 ***  

R:A  9 81.017   9.0019  6.9147 0.0002658 ***
```

```

B      2 35.433 17.7167 13.6088 0.0002510 ***
R:B    2 16.233  8.1167  6.2347 0.0087635 **
A:B   18 61.567  3.4204  2.6273 0.0236253 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df Sum Sq Mean Sq F value Pr(>F)
R     1  2.817  2.8167  2.1636 0.1585807
A     9 77.683  8.6315  6.6302 0.0003456 ***
R:A   9 81.017  9.0019  6.9147 0.0002658 ***
B     2 35.433 17.7167 13.6088 0.0002510 ***
R:B    2 16.233  8.1167  6.2347 0.0087635 **
A:B   18 61.567  3.4204  2.6273 0.0236253 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.4 Example 2.2

(70) MODEL

```

ex2.2 = read.table("C:/G/Rt/Split/sbex2_2.txt", header=TRUE)
ex2.2 = af(ex2.2, c("Row", "Column", "R", "S"))
GLM(Y ~ Column + R + R:Column + S + S:Column + R:S, ex2.2)

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      51 10328  202.51  0.8112 0.7688
RESIDUALS   48 11982  249.63
CORRECTED TOTAL 99 22310

```

```

$Fitness
Root MSE   Y Mean Coef Var  R-square   Adj R-sq
15.79971 1000.098 1.579816 0.4629279 -0.1077112

```

```

$`Type I` 
  Df Sum Sq Mean Sq F value Pr(>F)
Column     4 1318.6  329.66  1.3206 0.2758
R          4 1159.8  289.94  1.1615 0.3396
Column:R 16 2808.6  175.54  0.7032 0.7766
S          3  351.9  117.29  0.4699 0.7047
Column:S 12 3863.3  321.94  1.2897 0.2555
R:S       12  826.0   68.83  0.2757 0.9906

```

```

$`Type II` 
  Df Sum Sq Mean Sq F value Pr(>F)
Column     4 1318.6  329.66  1.3206 0.2758
R          4 1159.8  289.94  1.1615 0.3396

```

```

Column:R 16 2808.6 175.54 0.7032 0.7766
S         3  351.9  117.29 0.4699 0.7047
Column:S 12 3863.3 321.94 1.2897 0.2555
R:S      12  826.0   68.83 0.2757 0.9906

```

```

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Column     4 1318.6 329.66 1.3206 0.2758  

R          4 1159.8 289.94 1.1615 0.3396  

Column:R 16 2808.6 175.54 0.7032 0.7766  

S         3  351.9  117.29 0.4699 0.7047  

Column:S 12 3863.3 321.94 1.2897 0.2555  

R:S      12  826.0   68.83 0.2757 0.9906

```

(71) MODEL

```
GLM(Y ~ Row + R + Row:R + S + Column:S + R:S + Column:R:S, ex2.2)
```

```

$ANOVA  

Response : Y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      99 22310 225.36  

RESIDUALS    0      0  

CORRECTED TOTAL 99 22310

```

```

$Fitness  

Root MSE   Y Mean Coef Var R-square  

      NA 1000.098      NA        1

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Row       4   147.4   36.86  

R          4 1159.8 289.94  

Row:R     16 3979.8 248.74  

S          3   351.9  117.29  

S:Column  12 3863.3 321.94  

R:S       12   826.0   68.83  

R:S:Column 48 11982.3 249.63

```

```

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

Row       0  

R          4 1159.8 289.94  

Row:R     0  

S          3   351.9  117.29  

S:Column  12 3863.3 321.94  

R:S       12   826.0   68.83  

R:S:Column 48 11982.3 249.63

```

```
$`Type III`  
CAUTION: Singularity Exists !  
          Df  Sum Sq Mean Sq F value Pr(>F)  
Row          0  
R            4  1159.8  289.94  
Row:R        0  
S            3   351.9  117.29  
S:Column     12  3863.3  321.94  
R:S          12   826.0   68.83  
R:S:Column  48 11982.3  249.63
```

(72) MODEL

```
GLM(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2)
```

```
$ANOVA  
Response : Y  
          Df  Sum Sq Mean Sq F value Pr(>F)  
MODEL       99  22310  225.36  
RESIDUALS    0      0  
CORRECTED TOTAL 99  22310
```

```
$Fitness  
Root MSE    Y Mean Coef Var R-square  
           NA 1000.098      NA         1
```

```
$`Type I`  
          Df  Sum Sq Mean Sq F value Pr(>F)  
Row          4   147.4   36.86  
R            4  1159.8  289.94  
S            3   351.9  117.29  
R:S          12   826.0   68.83  
Row:R        16  3979.8  248.74  
S:Column     12  3863.3  321.94  
R:S:Column  48 11982.3  249.63
```

```
$`Type II`  
          Df  Sum Sq Mean Sq F value Pr(>F)  
Row          0  
R            4  1159.8  289.94  
S            3   351.9  117.29  
R:S          12   826.0   68.83  
Row:R        0  
S:Column     12  3863.3  321.94  
R:S:Column  48 11982.3  249.63
```

```
$`Type III`  
CAUTION: Singularity Exists !  
          Df  Sum Sq Mean Sq F value Pr(>F)
```

```

Row          0
R           4  1159.8  289.94
S           3   351.9  117.29
R:S         12   826.0   68.83
Row:R       0
S:Column    12  3863.3  321.94
R:S:Column  48 11982.3  249.63

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ Row + R + S + R:S + Row:R + Column:S + Column:R:S, ex2.2), type=3,
      singular.ok=TRUE) # NOT WORKING

```

7.5 Example 3.1

(73) MODEL

```

ex3.1 = read.table("C:/G/Rt/Split/spedsite.txt", header=TRUE)
ex3.1 = af(ex3.1, c("Site", "A", "B", "C", "Block"))
GLM(Yield ~ Site + Site:Block + A + B + A:B + A:Site + B:Site + A:B:Site +
     A:B:Site:Block + C + A:C + B:C + A:B:C + C:Site + A:C:Site + B:C:Site +
     A:B:C:Site, ex3.1)

```

```

$ANOVA
Response : Yield
            Df   Sum Sq Mean Sq F value    Pr(>F)
MODEL        239 2724374186 11399055  23.682 < 2.2e-16 ***
RESIDUALS    240   115521933   481341
CORRECTED TOTAL 479 2839896119
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE Yield Mean Coef Var  R-square Adj R-sq
693.7877 8290.769 8.368195 0.9593218 0.918813

```

```

$`Type I`
            Df   Sum Sq Mean Sq F value    Pr(>F)
Site         3   621230991 207076997 430.2082 < 2e-16 ***
Site:Block   8  1305369943 163171243 338.9928 < 2e-16 ***
A            1   1333205   1333205   2.7698  0.09737 .
B            4   47928577  11982144  24.8932 < 2e-16 ***
A:B          4    14849     3712   0.0077  0.99988
Site:A       3    33010     11003   0.0229  0.99531
Site:B       12    37932      3161   0.0066  1.00000
Site:A:B     12    11494      958   0.0020  1.00000
Site:Block:A:B 72   8239680   114440   0.2378  1.00000
C            3   739890389 246630130 512.3809 < 2e-16 ***
A:C          3     3233      1078   0.0022  0.99985
B:C          12    34961     2913   0.0061  1.00000

```

A:B:C	12	11077	923	0.0019	1.00000
Site:C	9	25983	2887	0.0060	1.00000
Site:A:C	9	22227	2470	0.0051	1.00000
Site:B:C	36	88610	2461	0.0051	1.00000
Site:A:B:C	36	98025	2723	0.0057	1.00000

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	621230991	207076997	430.2082	< 2e-16 ***
Site:Block	8	1305369943	163171243	338.9928	< 2e-16 ***
A	1	1333205	1333205	2.7698	0.09737 .
B	4	47928577	11982144	24.8932	< 2e-16 ***
A:B	4	14849	3712	0.0077	0.99988
Site:A	3	33010	11003	0.0229	0.99531
Site:B	12	37932	3161	0.0066	1.00000
Site:A:B	12	11494	958	0.0020	1.00000
Site:Block:A:B	72	8239680	114440	0.2378	1.00000
C	3	739890389	246630130	512.3809	< 2e-16 ***
A:C	3	3233	1078	0.0022	0.99985
B:C	12	34961	2913	0.0061	1.00000
A:B:C	12	11077	923	0.0019	1.00000
Site:C	9	25983	2887	0.0060	1.00000
Site:A:C	9	22227	2470	0.0051	1.00000
Site:B:C	36	88610	2461	0.0051	1.00000
Site:A:B:C	36	98025	2723	0.0057	1.00000

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	621230991	207076997	430.2082	< 2e-16 ***
Site:Block	8	1305369943	163171243	338.9928	< 2e-16 ***
A	1	1333205	1333205	2.7698	0.09737 .
B	4	47928577	11982144	24.8932	< 2e-16 ***
A:B	4	14849	3712	0.0077	0.99988
Site:A	3	33010	11003	0.0229	0.99531
Site:B	12	37932	3161	0.0066	1.00000
Site:A:B	12	11494	958	0.0020	1.00000
Site:Block:A:B	72	8239680	114440	0.2378	1.00000
C	3	739890389	246630130	512.3809	< 2e-16 ***
A:C	3	3233	1078	0.0022	0.99985
B:C	12	34961	2913	0.0061	1.00000
A:B:C	12	11077	923	0.0019	1.00000
Site:C	9	25983	2887	0.0060	1.00000
Site:A:C	9	22227	2470	0.0051	1.00000
Site:B:C	36	88610	2461	0.0051	1.00000

```

Site:A:B:C      36      98025      2723    0.0057 1.00000
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(74) MODEL

ex3.1a = read.table("C:/G/Rt/Split/Ex3.1-example.txt", header=TRUE)
ex3.1a = af(ex3.1a, c("row", "P", "column", "R", "S"))
GLM(height ~ P + column + column:P + R + P:R + column:R + column:R:P + S +
     P:S + column:S + column:S:P + R:S + R:S:column + R:S:P + R:S:P:column, ex3.1a)

$ANOVA
Response : height
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL       199 7534.8 37.863
RESIDUALS      0    0.0
CORRECTED TOTAL 199 7534.8

$Fitness
Root MSE height Mean Coef Var R-square
      NA      93.965      NA      1

$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
P           1 253.1 253.125
column      4 109.4 27.357
P:column    4 207.9 51.987
R           4   90.6 22.657
P:R         4 505.0 126.238
column:R    16 3357.8 209.864
P:column:R  16 1442.6 90.163
S           3   16.4  5.458
P:S         3   14.3  4.765
column:S    12 265.5 22.121
P:column:S  12   96.5  8.044
R:S         12 195.1 16.254
column:R:S  48 365.5  7.615
P:R:S       12 100.3  8.361
P:column:R:S 48 514.7 10.723

$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
P           1 253.1 253.125
column      4 109.4 27.358
P:column    4 207.9 51.987
R           4   90.6 22.657
P:R         4 505.0 126.238
column:R    16 3357.8 209.864
P:column:R  16 1442.6 90.162

```

```

S           3   16.4   5.458
P:S         3   14.3   4.765
column:S    12  265.4  22.121
P:column:S  12   96.5   8.044
R:S         12  195.0  16.254
column:R:S  48  365.5   7.615
P:R:S       12  100.3   8.361
P:column:R:S 48  514.7  10.723

$`Type III`  

          Df Sum Sq Mean Sq F value Pr(>F)  

P           1 253.1 253.125  

column      4 109.4 27.358  

P:column    4 207.9 51.987  

R           4  90.6 22.657  

P:R         4 505.0 126.238  

column:R    16 3357.8 209.864  

P:column:R  16 1442.6 90.163  

S           3   16.4   5.458  

P:S         3   14.3   4.765  

column:S    12  265.4  22.121  

P:column:S  12   96.5   8.044  

R:S         12  195.0  16.254  

column:R:S  48  365.5   7.615  

P:R:S       12  100.3   8.361  

P:column:R:S 48  514.7  10.723

```

(75) MODEL

```
GLM(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
     S:R:P + R:S:P:row, ex3.1a)
```

```
$ANOVA  

Response : height  

          Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      199 7534.8 37.863  

RESIDUALS   0     0.0  

CORRECTED TOTAL 199 7534.8
```

```
$Fitness  

Root MSE height Mean Coef Var R-square  

      NA     93.965      NA      1
```

```
$`Type I`  

          Df Sum Sq Mean Sq F value Pr(>F)  

row        4 2017.03 504.26  

R          4   90.63  22.66  

P          1  253.12 253.12  

S          3   16.38   5.46
```

```

R:S      12 195.05 16.25
row:P     4 167.25 41.81
R:P      4 504.95 126.24
row:R:P   32 2933.52 91.67
P:S      3 14.29 4.76
row:P:S  24 234.68 9.78
R:P:S    12 100.33 8.36
row:R:P:S 96 1007.52 10.49

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

row      4 2017.03 504.26  

R        4  90.63 22.66  

P        1 253.12 253.12  

S        3 16.38 5.46  

R:S      12 195.05 16.25  

row:P    4 167.25 41.81  

R:P      4 504.95 126.24  

row:R:P   32 2933.52 91.67  

P:S      3 14.29 4.76  

row:P:S  24 234.68 9.78  

R:P:S    12 100.33 8.36  

row:R:P:S 96 1007.52 10.49

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

row      4 2017.03 504.26  

R        4  90.63 22.66  

P        1 253.13 253.13  

S        3 16.38 5.46  

R:S      12 195.05 16.25  

row:P    4 167.25 41.81  

R:P      4 504.95 126.24  

row:R:P   32 2933.52 91.67  

P:S      3 14.30 4.77  

row:P:S  24 234.68 9.78  

R:P:S    12 100.33 8.36  

row:R:P:S 96 1007.52 10.49

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +
          S:P:row + S:R:P + R:S:P:row, ex3.1a), type=3, singular.ok=TRUE)
# NOT WORKING

alias(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
      S:R:P + R:S:P:row, ex3.1a) # NO ALIAS

Model :
height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P +

```

```

S:P:row + S:R:P + R:S:P:row

(76) MODEL
  • p94 Appendix 3.1

ex3.1b = read.table("C:/G/Rt/Split/spexvar3.txt", header=TRUE)
ex3.1b = af(ex3.1b, c("rep", "var", "nit", "row", "col"))
GLM(yield ~ rep + var + rep:var + nit + var:nit, ex3.1b)

$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       26  44017 1692.97  9.5603 4.779e-11 ***
RESIDUALS   45   7969  177.08
CORRECTED TOTAL 71   51986
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE yield Mean Coef Var  R-square  Adj R-sq
13.30727 103.9722 12.79887 0.8467134 0.7581478

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      5 15875.3 3175.1 17.9297 9.525e-10 ***
var      2 1786.4   893.2  5.0438  0.010557 *
rep:var 10 6013.3   601.3  3.3957  0.002251 **
nit      3 20020.5 6673.5 37.6856 2.458e-12 ***
var:nit  6   321.7    53.6  0.3028  0.932199
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      5 15875.3 3175.1 17.9297 9.525e-10 ***
var      2 1786.4   893.2  5.0438  0.010557 *
rep:var 10 6013.3   601.3  3.3957  0.002251 **
nit      3 20020.5 6673.5 37.6856 2.458e-12 ***
var:nit  6   321.7    53.6  0.3028  0.932199
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      5 15875.3 3175.1 17.9297 9.525e-10 ***
var      2 1786.4   893.2  5.0438  0.010557 *
rep:var 10 6013.3   601.3  3.3957  0.002251 **
nit      3 20020.5 6673.5 37.6856 2.458e-12 ***

```

```

var:nit  6    321.7    53.6  0.3028  0.932199
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
(77) MODEL
GLM(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b)

$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       37 48090 1299.7 11.341 6.734e-11 ***
RESIDUALS   34   3896   114.6
CORRECTED TOTAL 71 51986
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE yield Mean Coef Var R-square Adj R-sq
10.70513 103.9722 10.29615 0.9250491 0.8434848

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      5 15875.3 3175.1 27.7056 4.391e-11 ***
var      2 1786.4   893.2  7.7939 0.0016359 **
rep:var 10 6013.3   601.3  5.2472 0.0001207 ***
nit      3 20020.5 6673.5 58.2331 1.754e-13 ***
var:nit  6   321.7    53.6  0.4679 0.8271333
row      9   900.9    100.1  0.8734 0.5575581
col      2 3171.5 1585.7 13.8373 4.012e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 5942.5 2971.3 25.9273 1.449e-07 ***
var      2 2799.8 1399.9 12.2155 0.0001005 ***
rep:var  4   997.8    249.4  2.1767 0.0926008 .
nit      3 12559.3 4186.4 36.5308 9.683e-11 ***
var:nit  6   477.8     79.6  0.6949 0.6553307
row      9   945.0    105.0  0.9162 0.5230151
col      2 3171.5 1585.7 13.8373 4.012e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 5942.5 2971.3 25.9273 1.449e-07 ***

```

```

var      2  2799.8  1399.9 12.2155 0.0001005 ***
rep:var  4   997.8   249.4  2.1767 0.0926008 .
nit      3 11977.9  3992.6 34.8397 1.775e-10 ***
var:nit  6   477.8    79.6  0.6949 0.6553307
row      9   945.0   105.0  0.9162 0.5230151
col      2  3171.5  1585.7 13.8373 4.012e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + var + rep:var + nit + var:nit + row + col, ex3.1b),
      type=3, singular.ok=TRUE) # NOT OK for var

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: yield
      Sum Sq Df F values    Pr(>F)
rep      5942.5  2 25.9273 1.449e-07 ***
var       0.0  0
nit     11977.9  3 34.8397 1.775e-10 ***
row      945.0  9  0.9162    0.5230
col     3171.5  2 13.8373 4.012e-05 ***
rep:var   997.8  4  2.1767    0.0926 .
var:nit   477.8  6  0.6949    0.6553
Residuals 3896.4 34
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.6 Example 4.1

```

(78) MODEL

ex4.1 = read.table("C:/G/Rt/Split/Ex4.1-example.txt", header=TRUE)
ex4.1 = af(ex4.1, c("row", "P", "column", "R", "S"))
GLM(height ~ P + column + column:P + R + P:R + column:R + column:R:P + S +
     P:S + column:S + column:S:P + R:S + R:S:column + R:S:P + R:S:P:column, ex4.1)

$ANOVA
Response : height
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      199 1710.2  8.5937
RESIDUALS      0    0.0
CORRECTED TOTAL 199 1710.2

$Fitness
Root MSE height Mean Coef Var R-square
      NA       6.815      NA        1

```

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	28.12	28.1250		
column	4	34.33	8.5825		
P:column	4	91.45	22.8625		
R	4	31.03	7.7575		
P:R	4	48.95	12.2375		
column:R	16	467.92	29.2450		
P:column:R	16	350.10	21.8813		
S	3	3.77	1.2583		
P:S	3	3.29	1.0983		
column:S	12	74.55	6.2125		
P:column:S	12	47.03	3.9192		
R:S	12	36.65	3.0542		
column:R:S	48	197.40	4.1125		
P:R:S	12	26.33	2.1942		
P:column:R:S	48	269.22	5.6087		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	28.12	28.1250		
column	4	34.33	8.5825		
P:column	4	91.45	22.8625		
R	4	31.03	7.7575		
P:R	4	48.95	12.2375		
column:R	16	467.92	29.2450		
P:column:R	16	350.10	21.8812		
S	3	3.77	1.2583		
P:S	3	3.30	1.0983		
column:S	12	74.55	6.2125		
P:column:S	12	47.03	3.9192		
R:S	12	36.65	3.0542		
column:R:S	48	197.40	4.1125		
P:R:S	12	26.33	2.1942		
P:column:R:S	48	269.22	5.6087		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	1	28.12	28.1250		
column	4	34.33	8.5825		
P:column	4	91.45	22.8625		
R	4	31.03	7.7575		
P:R	4	48.95	12.2375		
column:R	16	467.92	29.2450		
P:column:R	16	350.10	21.8813		
S	3	3.77	1.2583		
P:S	3	3.29	1.0983		

```

column:S      12  74.55  6.2125
P:column:S   12  47.03  3.9192
R:S          12  36.65  3.0542
column:R:S    48 197.40  4.1125
P:R:S        12  26.33  2.1942
P:column:R:S 48 269.22  5.6087

```

(79) MODEL

```

GLM(height ~ row + R + P + S + S:R + row:P + R:P + row:R:P + S:P + S:P:row +
     S:R:P + R:S:P:row, ex4.1)

```

\$ANOVA

```

Response : height
           Df Sum Sq Mean Sq F value Pr(>F)
MODEL       199 1710.2 8.5937
RESIDUALS   0     0.0
CORRECTED TOTAL 199 1710.2

```

\$Fitness

	Root	MSE	height	Mean	Coef	Var	R-square
	NA		6.815		NA		1

\$`Type I`

	Df	Sum Sq	Mean Sq	F	value	Pr(>F)
row	4	309.43	77.357			
R	4	31.03	7.758			
P	1	28.12	28.125			
S	3	3.77	1.258			
R:S	12	36.65	3.054			
row:P	4	130.25	32.563			
R:P	4	48.95	12.237			
row:R:P	32	504.12	15.754			
P:S	3	3.29	1.098			
row:P:S	24	171.28	7.137			
R:P:S	12	26.33	2.194			
row:R:P:S	96	416.92	4.343			

\$`Type II`

	Df	Sum Sq	Mean Sq	F	value	Pr(>F)
row	4	309.43	77.357			
R	4	31.03	7.757			
P	1	28.12	28.125			
S	3	3.78	1.258			
R:S	12	36.65	3.054			
row:P	4	130.25	32.563			
R:P	4	48.95	12.237			
row:R:P	32	504.12	15.754			
P:S	3	3.30	1.098			

```

row:P:S    24 171.28   7.137
R:P:S      12 26.33   2.194
row:R:P:S  96 416.92   4.343

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

row        4 309.43 77.357  

R          4  31.03  7.758  

P          1  28.12 28.125  

S          3   3.77  1.258  

R:S        12 36.65  3.054  

row:P      4 130.25 32.562  

R:P       4  48.95 12.238  

row:R:P    32 504.12 15.754  

P:S       3   3.29  1.098  

row:P:S   24 171.28   7.137  

R:P:S    12 26.33   2.194
row:R:P:S 96 416.92   4.343

```

7.7 Example 5.1

(80) MODEL

```

ex5.1 = read.table("C:/G/Rt/Split/sbsp.txt", header=TRUE)
ex5.1 = af(ex5.1, c("R", "A", "C", "B", "Tx"))
GLM(Y ~ R + A + R:A + C + B + C:B + Tx + B:Tx, ex5.1)

```

```

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value     Pr(>F)
MODEL      20 193.583 9.6792 9.4176 2.969e-05 ***
RESIDUALS  15  15.417 1.0278
CORRECTED TOTAL 35 209.000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE Y Mean Coef Var R-square   Adj R-sq
1.013794   5.5 18.43261 0.926236 0.8278841

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value     Pr(>F)
R      2 33.500 16.7500 16.2973 0.0001734 ***
A      1 16.000 16.0000 15.5676 0.0012951 **
R:A    2 32.167 16.0833 15.6486 0.0002133 ***
C      2   0.500  0.2500  0.2432 0.7871141
B      1   1.778  1.7778  1.7297 0.2081966
C:B   2   0.389  0.1944  0.1892 0.8295745
Tx    5 103.333 20.6667 20.1081  3.63e-06 ***

```

```

B:Tx 5 5.917 1.1833 1.1514 0.3770453
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R     2 23.047 11.5236 11.2122 0.0010520 **  

A     1 12.375 12.3751 12.0406 0.0034285 **  

R:A   2 27.164 13.5819 13.2148 0.0004907 ***  

C     2  0.500  0.2500  0.2432 0.7871141  

B     1  1.778  1.7778  1.7297 0.2081966  

C:B   2  0.389  0.1944  0.1892 0.8295745  

Tx    5 103.333 20.6667 20.1081 3.63e-06 ***  

B:Tx  5 5.917  1.1833  1.1514 0.3770453
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

R     2 22.451 11.2254 10.9220 0.0011828 **  

A     1 15.001 15.0013 14.5958 0.0016719 **  

R:A   2 27.164 13.5819 13.2148 0.0004907 ***  

C     2  0.500  0.2500  0.2432 0.7871141  

B     1  1.778  1.7778  1.7297 0.2081966  

C:B   2  0.389  0.1944  0.1892 0.8295745  

Tx    5 103.333 20.6667 20.1081 3.63e-06 ***  

B:Tx  5 5.917  1.1833  1.1514 0.3770453
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(81) MODEL  

GLM(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx, ex5.1)  

$ANOVA  

Response : Y  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL        20 194.188  9.7094  9.8323 2.254e-05 ***  

RESIDUALS    15  14.813  0.9875  

CORRECTED TOTAL 35 209.000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness  

  Root MSE Y Mean Coef Var R-square Adj R-sq  

0.9937303   5.5 18.06782 0.9291268 0.8346292

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)

```

```

R      2   33.500 16.7500 16.9620 0.0001410 ***
A      1   16.000 16.0000 16.2025 0.0011013 **
R:A    2   32.167 16.0833 16.2869 0.0001739 ***
C      2     0.500  0.2500  0.2532  0.7795913
B      1     1.778  1.7778  1.8003  0.1996385
C:B    2     0.389  0.1944  0.1969  0.8233570
Tx     5 103.333 20.6667 20.9283 2.813e-06 ***
A:Tx   5     6.521  1.3042  1.3207  0.3078554
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R      2   33.500 16.7500 16.9620 0.0001410 ***  

A      1   16.000 16.0000 16.2025 0.0011013 **  

R:A    2   32.167 16.0833 16.2869 0.0001739 ***  

C      2     0.807  0.4037  0.4088  0.6716130  

B      1     1.757  1.7574  1.7797  0.2020905  

C:B    2     0.030  0.0150  0.0152  0.9849064  

Tx     5 103.333 20.6667 20.9283 2.813e-06 ***  

A:Tx   5     6.521  1.3042  1.3207  0.3078554
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

R      2   33.500 16.7500 16.9620 0.0001410 ***  

A      1   16.000 16.0000 16.2025 0.0011013 **  

R:A    2   32.167 16.0833 16.2869 0.0001739 ***  

C      2     0.780  0.3902  0.3952  0.6803789  

B      1     1.776  1.7756  1.7980  0.1999029  

C:B    2     0.030  0.0150  0.0152  0.9849064  

Tx     5 103.333 20.6667 20.9283 2.813e-06 ***  

A:Tx   5     6.521  1.3042  1.3207  0.3078554
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(82) MODEL

```
GLM(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)
```

```

$ANOVA  

Response : Y  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL        24 196.238  8.1766  7.0476 0.0008758 ***  

RESIDUALS    11 12.762   1.1602  

CORRECTED TOTAL 35 209.000
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE Y Mean Coef Var R-square Adj R-sq
  1.077122   5.5 19.58405 0.9389372 0.8057093

$`Type I` 
  Df Sum Sq Mean Sq F value    Pr(>F)
R     2 33.500 16.7500 14.4373 0.0008391 ***
A     1 16.000 16.0000 13.7908 0.0034197 **
R:A    2 32.167 16.0833 13.8626 0.0009856 ***
C     2  0.500  0.2500  0.2155  0.8094766
B     1  1.778  1.7778  1.5323  0.2415358
C:B    2  0.389  0.1944  0.1676  0.8478141
Tx    5 103.333 20.6667 17.8131 6.055e-05 ***
A:Tx   5  6.521  1.3042  1.1241  0.4027183
B:Tx   4  2.050  0.5126  0.4418  0.7761730
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
  Df Sum Sq Mean Sq F value    Pr(>F)
R     2 23.116 11.5581 9.9622 0.003396 **
A     1 12.375 12.3751 10.6664 0.007519 **
R:A    2 27.426 13.7132 11.8197 0.001820 **
C     2  0.970  0.4850  0.4180  0.668392
B     1  1.757  1.7574  1.5148  0.244080
C:B    2  0.085  0.0424  0.0366  0.964202
Tx    5 103.333 20.6667 17.8131 6.055e-05 ***
A:Tx   4  2.655  0.6636  0.5720  0.688652
B:Tx   4  2.050  0.5126  0.4418  0.776173
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value    Pr(>F)
R     2 22.186 11.0928 9.5611 0.003924 **
A     1 15.185 15.1853 13.0886 0.004042 **
R:A    2 27.426 13.7132 11.8197 0.001820 **
C     2  1.010  0.5049  0.4352  0.657839
B     1  1.792  1.7922  1.5448  0.239751
C:B    2  0.085  0.0424  0.0366  0.964202
Tx    5 103.333 20.6667 17.8131 6.055e-05 ***
A:Tx   4  2.655  0.6636  0.5720  0.688652
B:Tx   4  2.050  0.5126  0.4418  0.776173
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

alias(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1)

Model :
Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx

Complete :
  (Intercept) R1     R2     A1     C1     C2     B1     Tx1    Tx2    Tx3    Tx4    Tx5    R1:A1
B1:Tx5      0       0     0 -1/5     0       0 -1/5     0       0     0     0     0     0
  R2:A1  C1:B1  C2:B1 A1:Tx1 A1:Tx2 A1:Tx3 A1:Tx4 A1:Tx5 B1:Tx1 B1:Tx2 B1:Tx3
B1:Tx5      0       0     0  1/5    1/5    1/5    1/5     -1    1/5    1/5    1/5
  B1:Tx4
B1:Tx5  1/5

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + B:C + Tx + A:Tx + B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y

	Sum Sq	Df	F values	Pr(>F)
R	22.186	2	9.5611	0.003924 **
A	0.000	0		
C	1.010	2	0.4352	0.657839
B	0.000	0		
Tx	103.333	5	17.8131	6.055e-05 ***
R:A	27.426	2	11.8197	0.001820 **
C:B	0.085	2	0.0366	0.964202
A:Tx	2.655	4	0.5720	0.688652
B:Tx	2.050	4	0.4418	0.776173
Residuals	12.762	11		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(83) MODEL

```
GLM(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	28	204.2	7.2929	10.635	0.001719 **
RESIDUALS	7	4.8	0.6857		
CORRECTED TOTAL	35	209.0			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

$Fitness
  Root MSE Y Mean Coef Var R-square Adj R-sq
  0.8280787   5.5 15.05598 0.9770335 0.8851675

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
R       2 33.500 16.7500 24.4271 0.0006969 ***
A       1 16.000 16.0000 23.3333 0.0018985 **
R:A     2 32.167 16.0833 23.4549 0.0007889 ***
C       2  0.500  0.2500  0.3646 0.7069339
B       1  1.778  1.7778  2.5926 0.1513998
C:B     2  0.389  0.1944  0.2836 0.7613494
Tx      5 103.333 20.6667 30.1389 0.0001357 ***
A:Tx    5  6.521  1.3042  1.9019 0.2123307
B:Tx    4  2.050  0.5126  0.7475 0.5896365
A:B:Tx  4  7.962  1.9905  2.9029 0.1038803
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
R       2 31.838 15.9191 23.2153 0.0008139 ***
A       1 12.375 12.3751 18.0470 0.0038017 **
R:A     1  2.017  2.0174  2.9420 0.1300172
C       2  0.500  0.2500  0.3645 0.7069558
B       1  1.757  1.7574  2.5629 0.1534298
C:B     1  0.644  0.6445  0.9399 0.3646045
Tx      5 103.333 20.6667 30.1389 0.0001357 ***
A:Tx    4  2.655  0.6636  0.9678 0.4812226
B:Tx    4  2.050  0.5126  0.7475 0.5896365
A:B:Tx  4  7.962  1.9905  2.9029 0.1038803
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value    Pr(>F)
R       2 28.112 14.0562 20.4986 0.0011846 **
A       1 14.655 14.6551 21.3720 0.0024176 **
R:A     1  2.017  2.0174  2.9420 0.1300172
C       2  0.471  0.2356  0.3436 0.7205632
B       1  1.769  1.7694  2.5804 0.1522328
C:B     1  0.644  0.6445  0.9399 0.3646045
Tx      5 103.815 20.7630 30.2793 0.0001336 ***
A:Tx    4  2.951  0.7378  1.0760 0.4358837
B:Tx    4  3.553  0.8882  1.2954 0.3579988
A:B:Tx  4  7.962  1.9905  2.9029 0.1038803
---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
alias(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1)

Model :
Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx

Complete :
  (Intercept) R1   R2   A1   C1   C2   B1   Tx1  Tx2  Tx3  Tx4  Tx5
B1:Tx5      0     0 -1/5  0     0 -1/5  0     0     0     0     0     0
A1:B1:Tx5 -1/6    0     0     0     0     0  1/6  1/6  1/6  1/6 -5/6
A1:B1:Tx6   0     2/3  0 4/45  2/3 -2/3  4/45 -1/3  1/3 -1/3   0     0
          R1:A1 R2:A1 C1:B1 C2:B1 A1:Tx1 A1:Tx2 A1:Tx3 A1:Tx4 A1:Tx5 B1:Tx1
B1:Tx5      0     0     0  1/5  1/5  1/5  1/5   -1   1/5
A1:B1:Tx5   0     0     0     0     0     0     0     0     0     0
A1:B1:Tx6 -2/9  4/9 -2/9 -2/9 -1/5 -1/5 -1/5  4/5   0 -1/5
          B1:Tx2 B1:Tx3 B1:Tx4 A1:B1:Tx1 A1:B1:Tx2 A1:B1:Tx3 A1:B1:Tx4
B1:Tx5      1/5  1/5  1/5   0     0     0     0
A1:B1:Tx5   0     0     0     0     0     0     0
A1:B1:Tx6 -1/5 -1/5  4/5   1     -1     1     0

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + A:R + C + B + C:B + Tx + A:Tx + B:Tx + A:B:Tx, ex5.1),
      type=3, singular.ok=TRUE) # NOT OK

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y
  Sum Sq Df F values    Pr(>F)
R     11.643  1 16.9793 0.004456 ***
A     0.000  0
C     0.002  1  0.0025 0.961483
B     0.000  0
Tx    89.178  3 43.3503 6.87e-05 ***
R:A    2.017  1  2.9420 0.130017
C:B    0.644  1  0.9399 0.364604
A:Tx   0.543  3  0.2640 0.849381
B:Tx   3.384  3  1.6451 0.264128
A:B:Tx  7.962  4  2.9029 0.103880
Residuals 4.800  7
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.8 Example 7.1

(84) MODEL

```

ex7.1 = read.table("C:/G/Rt/Split/asped.txt", header=TRUE)
ex7.1 = af(ex7.1, c("R", "G", "F"))
GLM(Y ~ R + G + R:G + F + F:G, ex7.1)

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      95 577.82 6.0824 5.3082 1.068e-05 ***
RESIDUALS   24  27.50 1.1458
CORRECTED TOTAL 119 605.32
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE Y Mean Coef Var  R-square  Adj R-sq
1.070436 6.175 17.335 0.9545699 0.7747422

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3 84.76 28.2528 24.6570 1.655e-07 ***
G     27 343.48 12.7216 11.1025 4.286e-08 ***
R:G    9 11.75 1.3056 1.1394 0.3749
F      2 59.85 29.9250 26.1164 9.481e-07 ***
G:F   54 77.98 1.4441 1.2603 0.2718
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3 5.75  1.9167 1.6727 0.1994
G     27 343.48 12.7216 11.1025 4.286e-08 ***
R:G    9 11.75 1.3056 1.1394 0.3749
F      2 59.85 29.9250 26.1164 9.481e-07 ***
G:F   54 77.98 1.4441 1.2603 0.2718
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
R      3 5.75  1.9167 1.6727 0.1994
G     27 343.48 12.7216 11.1025 4.286e-08 ***
R:G    9 11.75 1.3056 1.1394 0.3749
F      2 50.50 25.2525 22.0385 3.686e-06 ***
G:F   54 77.98 1.4441 1.2603 0.2718
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + G + R:G + F + F:G, ex7.1), type=3, singular.ok=TRUE) # NOT OK

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Y
          Sum Sq Df F values    Pr(>F)
R           0.000  0
G        202.417  3 58.8848 3.258e-11 ***
F         50.505  2 22.0385 3.686e-06 ***
R:G       11.750  9  1.1394    0.3749
G:F      77.983 54  1.2603    0.2718
Residuals 27.500 24
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.9 Example 7.2

(85) MODEL

```

ex7.2 = read.table("C:/G/Rt/Split/aspedt.txt", header=TRUE)
ex7.2 = af(ex7.2, c("R", "T", "G"))
GLM(Y ~ R + T + R:T + G + G:T, ex7.2)

```

```

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      99 538.70  5.4415 5.1892 1.286e-05 ***
RESIDUALS   24  25.17  1.0486
CORRECTED TOTAL 123 563.87
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE    Y Mean Coef Var R-square  Adj R-sq
1.024017 6.032258 16.97569 0.955368 0.7712612

```

```

$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
R      3 73.255 24.4183 23.2863 2.752e-07 ***
T      3 32.000 10.6667 10.1722 0.0001645 ***
R:T     9 28.402  3.1558  3.0095 0.0149568 *
G     21 309.908 14.7575 14.0734 7.158e-09 ***
T:G    63  95.140  1.5102  1.4401 0.1617931
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     3   4.229  1.4097  1.3444  0.2834998  

T     3  32.000 10.6667 10.1722  0.0001645 ***  

R:T   9  10.854  1.2060  1.1501  0.3684706  

G    21 309.908 14.7575 14.0734 7.158e-09 ***  

T:G  63  95.140  1.5102  1.4401  0.1617931  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     3   4.229  1.4097  1.3444  0.283500  

T     3  22.668  7.5559  7.2056  0.001299 **  

R:T   9  10.854  1.2060  1.1501  0.368471  

G    21 309.908 14.7575 14.0734 7.158e-09 ***  

T:G  63  95.140  1.5102  1.4401  0.161793  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7.10 Example 7.3

(86) MODEL

```
ex7.3 = read.table("C:/G/Rt/Split/assped.txt", header=TRUE)
ex7.3 = af(ex7.3, c("R", "T", "G", "F"))
GLM(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3)
```

```
$ANOVA  

Response : Y  

              Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL          155 656.12  4.2330  13.446 3.997e-14 ***  

RESIDUALS      36  11.33  0.3148  

CORRECTED TOTAL 191 667.45  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  Root MSE    Y Mean Coef Var R-square  Adj R-sq  

  0.5610836 6.265625 8.95495 0.98302 0.9099118
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R      3  27.06   9.019  28.6489 1.203e-09 ***  

T      1  10.55  10.547  33.5018 1.334e-06 ***  

R:T    3   2.97   0.991   3.1489  0.036705 *  

G     22 389.01  17.682  56.1668 < 2.2e-16 ***  

T:G   22  18.42   0.837   2.6601  0.004445 **
```

```

R:T:G 12   8.78   0.731   2.3235  0.025315 *
F      2 164.28  82.141  260.9173 < 2.2e-16 ***
T:F    2   0.84   0.422   1.3401  0.274574
G:F    44  23.47   0.533   1.6943  0.053191 .
T:G:F 44  10.74   0.244   0.7753  0.790640
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df Sum Sq Mean Sq  F value    Pr(>F)  

R      3 12.49   4.162  13.2206 5.655e-06 ***  

T      1 10.55   10.547 33.5018 1.334e-06 ***  

R:T    3   1.15   0.384   1.2206  0.316281  

G     22 389.01  17.682  56.1668 < 2.2e-16 ***  

T:G    22 18.42   0.837   2.6601  0.004445 **  

R:T:G 12   8.78   0.731   2.3235  0.025315 *  

F      2 164.28  82.141  260.9173 < 2.2e-16 ***  

T:F    2   0.84   0.422   1.3401  0.274574  

G:F    44  23.47   0.533   1.6943  0.053191 .  

T:G:F 44  10.74   0.244   0.7753  0.790640
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df Sum Sq Mean Sq  F value    Pr(>F)  

R      3 12.49   4.162  13.2206 5.655e-06 ***  

T      1 11.16   11.158 35.4430 8.021e-07 ***  

R:T    3   1.15   0.384   1.2206  0.316281  

G     22 389.01  17.682  56.1668 < 2.2e-16 ***  

T:G    22 18.42   0.837   2.6601  0.004445 **  

R:T:G 12   8.78   0.731   2.3235  0.025315 *  

F      2 120.56  60.282 191.4828 < 2.2e-16 ***  

T:F    2   0.82   0.411   1.3060  0.283432  

G:F    44  23.47   0.533   1.6943  0.053191 .  

T:G:F 44  10.74   0.244   0.7753  0.790640
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + T + R:T + G + G:T + R:T:G + F + F:T + F:G + F:G:T, ex7.3),
      type=3, singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: Y  

      Sum Sq Df F values    Pr(>F)

```

```

R          0.000  0
T          0.000  0
G         73.444  2 116.6471 < 2.2e-16 ***
F        120.563  2 191.4828 < 2.2e-16 ***
R:T       0.000  0
T:G       5.778  2   9.1765 0.0006018 ***
T:F       0.822  2   1.3060 0.2834316
G:F      23.469 44   1.6943 0.0531910 .
R:T:G     8.778 12   2.3235 0.0253153 *
T:G:F    10.740 44   0.7753 0.7906401
Residuals 11.333 36
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.11 Example 8.1

(87) MODEL

```

ex8.1 = read.table("C:/G/Rt/Split/asbed.txt", header=TRUE)
ex8.1 = af(ex8.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1)

```

```

$ANOVA
Response : Y
              Df Sum Sq Mean Sq F value Pr(>F)
MODEL           104 3951.8 37.999
RESIDUALS        0    0.0
CORRECTED TOTAL 104 3951.8

```

```

$Fitness
Root MSE  Y Mean Coef Var R-square
      NA 10.0381      NA        1

```

```

$`Type I`
              Df Sum Sq Mean Sq F value Pr(>F)
R            2 1787.68 893.84
A            12 601.24 50.10
R:A           6  24.93  4.16
B            8 156.87 19.61
R:B           4 319.87 79.97
A:B          60 1012.26 16.87
R:A:B        12  49.00  4.08

```

```

$`Type II`
              Df Sum Sq Mean Sq F value Pr(>F)
R            2 372.22 186.111
A            12 601.24 50.103
R:A           6  50.00  8.333
B            8 156.87 19.609

```

```

R:B      4    87.44  21.861
A:B     60   1012.26  16.871
R:A:B  12    49.00   4.083

$`Type III` 
      Df  Sum Sq Mean Sq F value Pr(>F)
R       2    372.22 186.111
A      12    572.31  47.692
R:A      6     50.00   8.333
B       8    185.85  23.231
R:B      4    87.44  21.861
A:B     60   1012.26  16.871
R:A:B  12    49.00   4.083

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(Y ~ R + A + R:A + B + B:R + A:B + A:B:R, ex8.1), type="III",
singular.ok=TRUE) # NOT WORKING

```

7.12 Example 9.1

(88) MODEL

```

ex9.1 = read.table("C:/G/Rt/Split/Ex9.1-spex1.txt", header=TRUE)
ex9.1 = af(ex9.1, c("R", "A", "B"))
GLM(Y ~ R + A + R:A + B + A:B, ex9.1)

```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	27	4920.8	182.251	10.594	5.927e-10 ***
RESIDUALS	34	584.9	17.203		
CORRECTED TOTAL	61	5505.6			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	Y	Mean	Coef	Var	R-square	Adj R-sq
4.147591	66.19839	6.265396	0.8937663	0.8094043		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
R	3	218.7	72.89	4.2369	0.01199 *
A	3	194.9	64.96	3.7760	0.01930 *
R:A	9	186.9	20.76	1.2070	0.32287
B	3	4087.4	1362.47	79.2018	1.998e-15 ***
A:B	9	233.0	25.88	1.5047	0.18602

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     3 157.8   52.61  3.0583  0.04134 *  

A     3 227.2   75.73  4.4020  0.01014 *  

R:A   9  94.5   10.50  0.6106  0.77932  

B     3 4087.4 1362.47 79.2018 1.998e-15 ***  

A:B   9 233.0   25.88  1.5047  0.18602  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

R     3 171.0   57.01  3.3138  0.03143 *  

A     3 209.7   69.92  4.0643  0.01431 *  

R:A   9  94.5   10.50  0.6106  0.77932  

B     3 4089.9 1363.29 79.2493 1.998e-15 ***  

A:B   9 233.0   25.88  1.5047  0.18602  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

7.13 Example 9.2

(89) MODEL

```
ex9.2 = read.table("C:/G/Rt/Split/Ex9.2-sbex.txt", header=TRUE)  

ex9.2 = af(ex9.2, c("rep", "hyb", "gen"))  

GLM(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2)
```

```
$ANOVA  

Response : yield  

  Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL        40 247.813  6.1953  4.4606 0.001119 **  

RESIDUALS     16  22.222  1.3889  

CORRECTED TOTAL 56 270.035  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

Root MSE yield Mean Coef Var  R-square  Adj R-sq  

  1.178511  45.77193 2.574747 0.9177062 0.7119716
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

rep      1  0.239  0.2388  0.1719  0.6839085  

hyb      9 66.796  7.4218  5.3437  0.0018370 **  

rep:hyb  8 67.000  8.3750  6.0300  0.0011569 **  

gen      2 36.351 18.1754 13.0863  0.0004293 ***  

rep:gen  2 16.923  8.4616  6.0924  0.0107858 *  

hyb:gen 18 60.504  3.3613  2.4201  0.0408545 *
```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep      1  0.167  0.1667  0.1200 0.7335481  

hyb      9 66.796  7.4218  5.3437 0.0018370 **  

rep:hyb  8 67.000  8.3750  6.0300 0.0011569 **  

gen      2 36.351 18.1754 13.0863 0.0004293 ***  

rep:gen   2 12.111  6.0556  4.3600 0.0308015 *  

hyb:gen  18 60.504  3.3613  2.4201 0.0408545 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

rep      1  0.167  0.1667  0.1200 0.7335481  

hyb      9 66.796  7.4218  5.3437 0.0018370 **  

rep:hyb  8 67.000  8.3750  6.0300 0.0011569 **  

gen      2 30.671 15.3356 11.0416 0.0009707 ***  

rep:gen   2 12.111  6.0556  4.3600 0.0308015 *  

hyb:gen  18 60.504  3.3613  2.4201 0.0408545 *  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(yield ~ rep + hyb + rep:hyb + gen + gen:rep + gen:hyb, ex9.2), type=3,
      singular.ok=TRUE) # NOT OK

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: yield  

      Sum Sq Df F values    Pr(>F)  

rep      0.000  0  

hyb     66.704  8  6.0033 0.0011847 **  

gen     30.671  2 11.0416 0.0009707 ***  

rep:hyb 67.000  8  6.0300 0.0011569 **  

rep:gen 12.111  2  4.3600 0.0308015 *  

hyb:gen 60.504 18  2.4201 0.0408545 *  

Residuals 22.222 16  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.14 Example 10.1

(90) MODEL

```

ex10.1 = read.table("C:/G/Rt/Split/Ex10.1-new.txt", header=TRUE)
ex10.1 = af(ex10.1, c("Site", "Block", "A", "B", "C"))
f10.1 = Yield ~ Site/Block + A/Site + B/Site + A:B + A:B:Site + A:B:Site:Block +
          C + A:C + B:C + A:B:C + C:Site + A:C:Site + B:C:Site + A:B:C:Site
GLM(f10.1, ex10.1)

```

\$ANOVA

Response : Yield

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	239	1639561484	6860090	2162	< 2.2e-16 ***
RESIDUALS	240	761522	3173		
CORRECTED TOTAL	479	1640323006			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root	MSE	Yield	Mean	Coef	Var	R-square	Adj R-sq
56.32947	9967.354	0.5651396	0.9995357	0.9990734			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	552717	184239	5.8064e+01	< 2e-16 ***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16 ***
A	4	1387680917	346920229	1.0933e+05	< 2e-16 ***
Site:A	12	34068	2839	8.9470e-01	0.55301
B	1	100939695	100939695	3.1812e+04	< 2e-16 ***
Site:B	3	1618	539	1.6990e-01	0.91662
A:B	4	31444008	7861002	2.4775e+03	< 2e-16 ***
Site:A:B	12	33737	2811	8.8600e-01	0.56185
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155
C	3	19356264	6452088	2.0334e+03	< 2e-16 ***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16 ***
B:C	3	23901388	7967129	2.5109e+03	< 2e-16 ***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16 ***
Site:C	9	47625	5292	1.6677e+00	0.09747 .
Site:A:C	36	104110	2892	9.1140e-01	0.61768
Site:B:C	9	61111	6790	2.1400e+00	0.02701 *
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	552717	184239	5.8064e+01	< 2e-16 ***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16 ***
A	4	1387680917	346920229	1.0933e+05	< 2e-16 ***
Site:A	12	34068	2839	8.9470e-01	0.55301

B	1	100939695	100939695	3.1812e+04	< 2e-16	***
Site:B	3	1618	539	1.6990e-01	0.91662	
A:B	4	31444008	7861002	2.4775e+03	< 2e-16	***
Site:A:B	12	33737	2811	8.8600e-01	0.56185	
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155	
C	3	19356264	6452088	2.0334e+03	< 2e-16	***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16	***
B:C	3	23901388	7967129	2.5109e+03	< 2e-16	***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16	***
Site:C	9	47625	5292	1.6677e+00	0.09747	.
Site:A:C	36	104110	2892	9.1140e-01	0.61768	
Site:B:C	9	61111	6790	2.1400e+00	0.02701	*
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941	

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Site	3	552717	184239	5.8064e+01	< 2e-16 ***
Site:Block	8	7062320	882790	2.7822e+02	< 2e-16 ***
A	4	1387680917	346920229	1.0933e+05	< 2e-16 ***
Site:A	12	34068	2839	8.9470e-01	0.55301
B	1	100939695	100939695	3.1812e+04	< 2e-16 ***
Site:B	3	1618	539	1.6990e-01	0.91662
A:B	4	31444008	7861002	2.4775e+03	< 2e-16 ***
Site:A:B	12	33737	2811	8.8600e-01	0.56185
Site:Block:A:B	72	186911	2596	8.1810e-01	0.84155
C	3	19356264	6452088	2.0334e+03	< 2e-16 ***
A:C	12	26075792	2172983	6.8483e+02	< 2e-16 ***
B:C	3	23901387	7967129	2.5109e+03	< 2e-16 ***
A:B:C	12	41996729	3499727	1.1030e+03	< 2e-16 ***
Site:C	9	47625	5292	1.6677e+00	0.09747
Site:A:C	36	104110	2892	9.1140e-01	0.61768
Site:B:C	9	61111	6790	2.1400e+00	0.02701 *
Site:A:B:C	36	82475	2291	7.2200e-01	0.87941

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(f10.1, ex10.1), type=3, singular.ok=TRUE) # NOT OK for Site:Block
```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

Response: Yield

	Sum Sq	Df	F values	Pr(>F)
Site	552717	3	5.8064e+01	< 2e-16 ***

```

A          1387680917  4 1.0933e+05 < 2e-16 ***
B          100939695   1 3.1812e+04 < 2e-16 ***
C          19356264    3 2.0334e+03 < 2e-16 ***
Site:Block      0 0
Site:A          34068  12 8.9470e-01 0.55301
Site:B          1618   3 1.6990e-01 0.91662
A:B          31444008  4 2.4775e+03 < 2e-16 ***
A:C          26075792  12 6.8483e+02 < 2e-16 ***
B:C          23901388  3 2.5109e+03 < 2e-16 ***
Site:C          47625  9 1.6677e+00 0.09747 .
Site:A:B        33737  12 8.8600e-01 0.56185
A:B:C        41996729  12 1.1030e+03 < 2e-16 ***
Site:A:C        104110 36 9.1140e-01 0.61768
Site:B:C        61111   9 2.1400e+00 0.02701 *
Site:Block:A:B  186911  72 8.1810e-01 0.84155
Site:A:B:C      82475  36 7.2200e-01 0.87941
Residuals       761522 240
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.15 Example 10.2

(91) MODEL

```

ex10.2 = read.table("C:/G/Rt/Split/Ex10.2-spbsite.txt", header=TRUE)
ex10.2 = af(ex10.2, c("Site", "Block", "A", "B"))
GLM(Yield ~ Site + Site:Block + A + A:Site + A:Site:Block + B + B:Site +
     B:Site:Block + A:B + A:B:Site, ex10.2)

```

```

$ANOVA
Response : Yield
            Df Sum Sq Mean Sq F value    Pr(>F)
MODEL           227 6370995084 28066058   10814 < 2.2e-16 ***
RESIDUALS       252    654049      2595
CORRECTED TOTAL 479 6371649132
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE Yield Mean Coef Var R-square Adj R-sq
50.94537 11083.06 0.4596687 0.9998974 0.9998049

```

```

$`Type I`
            Df Sum Sq Mean Sq F value    Pr(>F)
Site           2 523573968 261786984 1.0086e+05 < 2.2e-16 ***
Site:Block     9 3756646710 417405190 1.6082e+05 < 2.2e-16 ***
A              4 29288163   7322041 2.8211e+03 < 2.2e-16 ***
Site:A         8 247899     30987 1.1939e+01 1.998e-14 ***
Site:Block:A  36 1783391    49539 1.9087e+01 < 2.2e-16 ***

```

```

B          7 1937592291 276798899 1.0665e+05 < 2.2e-16 ***
Site:B     14 15903698 1135978 4.3768e+02 < 2.2e-16 ***
Site:Block:B 63 105727288 1678211 6.4660e+02 < 2.2e-16 ***
A:B        28 91141      3255 1.2541e+00   0.1838
Site:A:B    56 140534      2510 9.6690e-01   0.5461
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
Site       2 523573968 261786984 1.0086e+05 < 2.2e-16 ***
Site:Block  9 3756646710 417405190 1.6082e+05 < 2.2e-16 ***
A          4 29288163 7322041 2.8211e+03 < 2.2e-16 ***
Site:A      8 247899    30987 1.1939e+01 1.998e-14 ***
Site:Block:A 36 1783391 49539 1.9087e+01 < 2.2e-16 ***
B          7 1937592291 276798899 1.0665e+05 < 2.2e-16 ***
Site:B     14 15903698 1135978 4.3768e+02 < 2.2e-16 ***
Site:Block:B 63 105727288 1678211 6.4660e+02 < 2.2e-16 ***
A:B        28 91141      3255 1.2541e+00   0.1838
Site:A:B    56 140534      2510 9.6690e-01   0.5461
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
Site       2 523573968 261786984 1.0086e+05 < 2.2e-16 ***
Site:Block  9 3756646710 417405190 1.6082e+05 < 2.2e-16 ***
A          4 29288163 7322041 2.8211e+03 < 2.2e-16 ***
Site:A      8 247899    30987 1.1939e+01 1.998e-14 ***
Site:Block:A 36 1783391 49539 1.9087e+01 < 2.2e-16 ***
B          7 1937592291 276798899 1.0665e+05 < 2.2e-16 ***
Site:B     14 15903698 1135978 4.3768e+02 < 2.2e-16 ***
Site:Block:B 63 105727288 1678211 6.4660e+02 < 2.2e-16 ***
A:B        28 91141      3255 1.2541e+00   0.1838
Site:A:B    56 140534      2510 9.6690e-01   0.5461
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.16 Example 11.1

(92) MODEL

```

ex11.1 = read.table("C:/G/Rt/Split/Ex11.1-cov.txt", header=TRUE)
ex11.1 = af(ex11.1, c("R", "T", "S"))
GLM(Y ~ R + T + R:T + S + S:T, ex11.1)

```

```

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)

```

```

MODEL           11    328 29.8182  3.1948 0.02875 *
RESIDUALS      12    112  9.3333
CORRECTED TOTAL 23    440

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE Y Mean Coef Var  R-square Adj R-sq
 3.05505     7 43.64358 0.7454545 0.5121212

$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
R    2    48    24   2.5714 0.11765
T    1    24    24   2.5714 0.13479
R:T   2    16     8   0.8571 0.44880
S    3   156    52   5.5714 0.01251 *
T:S   3    84    28   3.0000 0.07277 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
R    2    48    24   2.5714 0.11765
T    1    24    24   2.5714 0.13479
R:T   2    16     8   0.8571 0.44880
S    3   156    52   5.5714 0.01251 *
T:S   3    84    28   3.0000 0.07277 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
R    2    48    24   2.5714 0.11765
T    1    24    24   2.5714 0.13479
R:T   2    16     8   0.8571 0.44880
S    3   156    52   5.5714 0.01251 *
T:S   3    84    28   3.0000 0.07277 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(93) MODEL

```
GLM(Z ~ R + T + R:T + S + S:T, ex11.1)
```

```
$ANOVA
Response : Z
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL       11    46  4.1818  2.5091 0.06452 .
RESIDUALS   12    20  1.6667
```

```

CORRECTED TOTAL 23      66
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE Z Mean Coef Var R-square Adj R-sq
1.290994   2.5 51.63978 0.6969697 0.4191919

$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
R    2     9     4.5     2.7 0.1076
T    1     6     6.0     3.6 0.0821 .
R:T   2     1     0.5     0.3 0.7462
S    3     9     3.0     1.8 0.2008
T:S   3    21     7.0     4.2 0.0301 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
R    2     9     4.5     2.7 0.1076
T    1     6     6.0     3.6 0.0821 .
R:T   2     1     0.5     0.3 0.7462
S    3     9     3.0     1.8 0.2008
T:S   3    21     7.0     4.2 0.0301 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
R    2     9     4.5     2.7 0.1076
T    1     6     6.0     3.6 0.0821 .
R:T   2     1     0.5     0.3 0.7462
S    3     9     3.0     1.8 0.2008
T:S   3    21     7.0     4.2 0.0301 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(94) MODEL
GLM(Y ~ R + T + R:T + S + S:T + Z, ex11.1)

$ANOVA
Response : Y
Df Sum Sq Mean Sq F value Pr(>F)
MODEL      12 342.45 28.5375   3.218 0.03116 *
RESIDUALS   11  97.55  8.8682
CORRECTED TOTAL 23 440.00
---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE Y Mean Coef Var R-square Adj R-sq
2.977949      7 42.54213 0.7782955 0.536436

$`Type I`
Df Sum Sq Mean Sq F value Pr(>F)
R   2 48.00 24.00 2.7063 0.11071
T   1 24.00 24.00 2.7063 0.12820
R:T  2 16.00  8.00 0.9021 0.43373
S   3 156.00 52.00 5.8637 0.01211 *
T:S  3 84.00 28.00 3.1574 0.06828 .
Z   1 14.45 14.45 1.6294 0.22807
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
R   2 18.300 9.1500 1.0318 0.38844
T   1 2.679  2.6786 0.3020 0.59359
R:T  2 9.450  4.7250 0.5328 0.60137
S   3 79.196 26.3985 2.9768 0.07822 .
T:S  3 37.474 12.4915 1.4086 0.29234
Z   1 14.450 14.4500 1.6294 0.22807
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
R   2 20.209 10.1043 1.1394 0.35505
T   1 6.104  6.1038 0.6883 0.42439
R:T  2 9.450  4.7250 0.5328 0.60137
S   3 84.243 28.0810 3.1665 0.06782 .
T:S  3 37.474 12.4915 1.4086 0.29234
Z   1 14.450 14.4500 1.6294 0.22807
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.17 Example 11.2

(95) MODEL

```

ex11.2a = read.table("C:/G/Rt/Split/Ex11.2-sp3.txt", header=TRUE)
ex11.2a = af(ex11.2a, "A")
ex11.2a$MY = (ex11.2a$Y1 + ex11.2a$Y2)/sqrt(2)
ex11.2a$Z = 2*ex11.2a$Z/sqrt(2)
GLM(MY ~ Z + A, ex11.2a)

```

```

$ANOVA
Response : MY
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       2 234.639 117.32 9.5696 0.01953 *
RESIDUALS    5  61.298   12.26
CORRECTED TOTAL 7 295.938
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE  MY Mean Coef Var R-square Adj R-sq
3.501377 20.06415 17.45091 0.7928678 0.7100149

$`Type I`
      Df  Sum Sq Mean Sq F value Pr(>F)
Z  1 190.148 190.148 15.5101 0.01098 *
A  1  44.492  44.492  3.6291 0.11512
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df  Sum Sq Mean Sq F value Pr(>F)
Z  1 166.577 166.577 13.5874 0.0142 *
A  1  44.492  44.492  3.6291 0.1151
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df  Sum Sq Mean Sq F value Pr(>F)
Z  1 166.577 166.577 13.5874 0.0142 *
A  1  44.492  44.492  3.6291 0.1151
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(96) MODEL
ex11.2b = read.table("C:/G/Rt/Split/Ex11.2-two.txt", header=TRUE)
ex11.2b = af(ex11.2b, c("sub", "A", "B"))
GLM(Y ~ A + A:sub + B + A:B, ex11.2b)

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value     Pr(>F)
MODEL       9 382.06  42.451 39.954 0.0001135 ***
RESIDUALS    6   6.38   1.062
CORRECTED TOTAL 15 388.44
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  Y Mean Coef Var  R-square  Adj R-sq
1.030776 14.1875 7.265384 0.9835881 0.9589702

$`Type I` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
A       1   68.062  68.062 64.0588 0.0002029 ***
A:sub  6  227.875  37.979 35.7451 0.0001934 ***
B       1   85.562  85.562 80.5294 0.0001070 ***
A:B     1    0.562   0.562  0.5294 0.4942562
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
A       1   68.062  68.062 64.0588 0.0002029 ***
A:sub  6  227.875  37.979 35.7451 0.0001934 ***
B       1   85.562  85.562 80.5294 0.0001070 ***
A:B     1    0.562   0.562  0.5294 0.4942562
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
A       1   68.062  68.062 64.0588 0.0002029 ***
A:sub  6  227.875  37.979 35.7451 0.0001934 ***
B       1   85.562  85.562 80.5294 0.0001070 ***
A:B     1    0.562   0.562  0.5294 0.4942562
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(97) MODEL

ex11.2c = read.table("C:/G/Rt/Split/Ex11.2-spcov2.txt", header=TRUE)
ex11.2c = af(ex11.2c, c("block", "whole", "split"))
GLM(Y ~ block + whole + block:whole + split + split:whole, ex11.2c)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        11    328 29.8182  3.1948 0.02875 *
RESIDUALS    12    112  9.3333
CORRECTED TOTAL 23    440
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  Y Mean Coef Var  R-square  Adj R-sq
3.05505    7 43.64358 0.7454545 0.5121212

```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block       2    48     24  2.5714 0.11765  

whole       1    24     24  2.5714 0.13479  

block:white 2    16      8  0.8571 0.44880  

split       3   156     52 5.5714 0.01251 *  

whole:split  3    84     28 3.0000 0.07277 .  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block       2    48     24  2.5714 0.11765  

whole       1    24     24  2.5714 0.13479  

block:white 2    16      8  0.8571 0.44880  

split       3   156     52 5.5714 0.01251 *  

whole:split  3    84     28 3.0000 0.07277 .  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block       2    48     24  2.5714 0.11765  

whole       1    24     24  2.5714 0.13479  

block:white 2    16      8  0.8571 0.44880  

split       3   156     52 5.5714 0.01251 *  

whole:split  3    84     28 3.0000 0.07277 .  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

(98) MODEL

```
GLM(Z ~ block + whole + block:white + split + split:white, ex11.2c)
```

```
$ANOVA  

Response : Z  

      Df Sum Sq Mean Sq     F value     Pr(>F)  

MODEL          11    38  3.4545 3.5903e+15 < 2.2e-16 ***  

RESIDUALS      12    0  0.0000  

CORRECTED TOTAL 23    38  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

      Root MSE Z Mean     Coef Var R-square Adj R-sq  

3.101924e-08    3.5 8.86264e-07           1         1
```

```
$`Type I`
```

```

          Df Sum Sq Mean Sq   F value Pr(>F)
block       2 36.000 18.0000 1.8707e+16 <2e-16 ***
whole      1  0.667  0.6667 6.9286e+14 <2e-16 ***
block:white 2  1.333  0.6667 6.9286e+14 <2e-16 ***
split      3  0.000  0.0000 0.0000e+00      1
whole:split 3  0.000  0.0000 0.0000e+00      1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq   F value Pr(>F)
block       2 36.000 18.0000 1.8707e+16 <2e-16 ***
whole      1  0.667  0.6667 6.9286e+14 <2e-16 ***
block:white 2  1.333  0.6667 6.9286e+14 <2e-16 ***
split      3  0.000  0.0000 0.0000e+00      1
whole:split 3  0.000  0.0000 0.0000e+00      1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df Sum Sq Mean Sq   F value Pr(>F)
block       2 36.000 18.0000 1.8707e+16 <2e-16 ***
whole      1  0.667  0.6667 6.9286e+14 <2e-16 ***
block:white 2  1.333  0.6667 6.9286e+14 <2e-16 ***
split      3  0.000  0.0000 0.0000e+00      1
whole:split 3  0.000  0.0000 0.0000e+00      1
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(99) MODEL
GLM(Y ~ block + whole + block:white + split + split:white + Z, ex11.2c)

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      11     328 29.8182  3.1948 0.02875 *
RESIDUALS  12     112  9.3333
CORRECTED TOTAL 23     440
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE Y Mean Coef Var R-square Adj R-sq
  3.05505    7 43.64358 0.7454545 0.5121212

$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
block      2     48      24  2.5714 0.11765

```

```

whole      1     24      24  2.5714 0.13479
block:whole 2     16       8  0.8571 0.44880
split      3    156      52  5.5714 0.01251 *
whole:split 3     84      28  3.0000 0.07277 .
Z          0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df  Sum Sq Mean Sq F value Pr(>F)
block      2  13.286   6.643  0.7117 0.51039
whole      1  16.000  16.000  1.7143 0.21495
block:whole 1  16.000  16.000  1.7143 0.21495
split      3 156.000  52.000  5.5714 0.01251 *
whole:split 3  84.000  28.000  3.0000 0.07277 .
Z          0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
      Df  Sum Sq Mean Sq F value Pr(>F)
block      2  13.286   6.643  0.7117 0.51039
whole      1  16.000  16.000  1.7143 0.21495
block:whole 1  16.000  16.000  1.7143 0.21495
split      3 156.000  52.000  5.5714 0.01251 *
whole:split 3  84.000  28.000  3.0000 0.07277 .
Z          0
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

7.18 Example 11.3

(100) MODEL

```

ex11.3 = read.table("C:/G/Rt/Split/Ex11.3-sbcov.txt", header=TRUE)
ex11.3 = af(ex11.3, c("block", "A", "B"))
GLM(Y ~ block + A + block:A + B + block:B + A:B, ex11.3)

```

```

$ANOVA
Response : Y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      17 16.833  0.9902  1.9804 0.2038
RESIDUALS      6  3.000  0.5000
CORRECTED TOTAL 23 19.833

```

```

$Fitness
  Root MSE   Y Mean Coef Var  R-square  Adj R-sq
0.7071068 2.916667 24.24366 0.8487395 0.4201681

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block     3 4.5000 1.5000 3.0000 0.11696  

A         1 1.5000 1.5000 3.0000 0.13397  

block:A   3 0.5000 0.1667 0.3333 0.80220  

B         2 8.3333 4.1667 8.3333 0.01855 *  

block:B   6 1.0000 0.1667 0.3333 0.89648  

A:B       2 1.0000 0.5000 1.0000 0.42188  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block     3 4.5000 1.5000 3.0000 0.11696  

A         1 1.5000 1.5000 3.0000 0.13397  

block:A   3 0.5000 0.1667 0.3333 0.80220  

B         2 8.3333 4.1667 8.3333 0.01855 *  

block:B   6 1.0000 0.1667 0.3333 0.89648  

A:B       2 1.0000 0.5000 1.0000 0.42188  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block     3 4.5000 1.5000 3.0000 0.11696  

A         1 1.5000 1.5000 3.0000 0.13397  

block:A   3 0.5000 0.1667 0.3333 0.80220  

B         2 8.3333 4.1667 8.3333 0.01855 *  

block:B   6 1.0000 0.1667 0.3333 0.89648  

A:B       2 1.0000 0.5000 1.0000 0.42188  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(101) MODEL

```
GLM(Z ~ block + A + block:A + B + block:B + A:B, ex11.3)
```

```
$ANOVA  

Response : Z  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      17 31.167 1.83333    3.3 0.07324 .  

RESIDUALS   6  3.333 0.55556  

CORRECTED TOTAL 23 34.500  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

Root MSE Z Mean Coef Var R-square Adj R-sq
```

```

0.745356 1.75 42.59177 0.9033816 0.6296296

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block   3 6.8333 2.2778     4.1 0.06689 .  

A       1 6.0000 6.0000    10.8 0.01669 *  

block:A 3 1.6667 0.5556     1.0 0.45472  

B       2 13.0000 6.5000    11.7 0.00850 **  

block:B 6 3.6667 0.6111     1.1 0.45542  

A:B     2 0.0000 0.0000     0.0 1.00000  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block   3 6.8333 2.2778     4.1 0.06689 .  

A       1 6.0000 6.0000    10.8 0.01669 *  

block:A 3 1.6667 0.5556     1.0 0.45472  

B       2 13.0000 6.5000    11.7 0.00850 **  

block:B 6 3.6667 0.6111     1.1 0.45542  

A:B     2 0.0000 0.0000     0.0 1.00000  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block   3 6.8333 2.2778     4.1 0.06689 .  

A       1 6.0000 6.0000    10.8 0.01669 *  

block:A 3 1.6667 0.5556     1.0 0.45472  

B       2 13.0000 6.5000    11.7 0.00850 **  

block:B 6 3.6667 0.6111     1.1 0.45542  

A:B     2 0.0000 0.0000     0.0 1.00000  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

(102) MODEL

```
GLM(Y ~ block + A + block:A + B + block:B + A:B + Z, ex11.3)
```

```
$ANOVA  

Response : Y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      18 17.8417 0.99120 2.4884 0.1589  

RESIDUALS    5 1.9917 0.39833  

CORRECTED TOTAL 23 19.8333
```

```
$Fitness  

Root MSE    Y Mean Coef Var R-square Adj R-sq  

0.6311365 2.916667 21.63897 0.8995798 0.5380672
```

```

$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block     3 4.5000 1.5000 3.7657 0.09378 .  

A         1 1.5000 1.5000 3.7657 0.10999  

block:A   3 0.5000 0.1667 0.4184 0.74788  

B         2 8.3333 4.1667 10.4603 0.01634 *  

block:B   6 1.0000 0.1667 0.4184 0.84059  

A:B       2 1.0000 0.5000 1.2552 0.36163  

Z         1 1.0083 1.0083 2.5314 0.17248  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block     3 3.6203 1.20678 3.0296 0.1319  

A         1 0.0000 0.00000 0.0000 1.0000  

block:A   3 0.2583 0.08611 0.2162 0.8813  

B         2 1.0317 0.51587 1.2951 0.3522  

block:B   6 0.4210 0.07017 0.1762 0.9717  

A:B       2 1.0000 0.50000 1.2552 0.3616  

Z         1 1.0083 1.00833 2.5314 0.1725

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

block     3 3.6613 1.22045 3.0639 0.1297  

A         1 0.0054 0.00536 0.0134 0.9122  

block:A   3 0.2583 0.08611 0.2162 0.8813  

B         2 0.7685 0.38427 0.9647 0.4423  

block:B   6 0.4210 0.07017 0.1762 0.9717  

A:B       2 1.0000 0.50000 1.2552 0.3616  

Z         1 1.0083 1.00833 2.5314 0.1725

```

8 Hinkelmann & Kempthorne - Volume 1

Reference

- Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 1 Introduction to Experimental Design. 2e. John Wiley & Sons Inc. 2008.

8.1 Chapter 6

8.1.1 p202

(103) MODEL

```
v1p202 = read.table("C:/G/Rt/Kemp/v1p202.txt", head=TRUE)
v1p202 = af(v1p202,c("brand"))
GLM(miles ~ brand, v1p202) # OK
```

\$ANOVA

Response : miles

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	4	47.234	11.809	15.661	0.004924 **
RESIDUALS	5	3.770	0.754		
CORRECTED TOTAL	9	51.004			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	miles	Mean Coef	Var	R-square	Adj R-sq
0.8683317	26.24	3.309191	0.9260842	0.8669516	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
brand	4	47.234	11.809	15.661	0.004924 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
brand	4	47.234	11.809	15.661	0.004924 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
brand	4	47.234	11.809	15.661	0.004924 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

8.1.2 p205

(104) MODEL

```
v1p205 = read.table("C:/G/Rt/Kemp/v1p205.txt", head=TRUE)
v1p205 = af(v1p205,c("brand", "car"))
GLM(miles ~ brand + car %in% brand, v1p205) # OK

$ANOVA
Response : miles
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      9 140.05 15.561   80.21 1.017e-13 ***
RESIDUALS  20   3.88   0.194
CORRECTED TOTAL 29 143.93
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE miles Mean Coef Var R-square Adj R-sq
0.4404543 26.16667 1.683265 0.9730418 0.9609106

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand      4 133.243 33.311 171.7053 3.553e-15 ***
brand:car  5   6.803   1.361   7.0137 0.0006214 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand      4 133.243 33.311 171.7053 3.553e-15 ***
brand:car  5   6.803   1.361   7.0137 0.0006214 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
brand      4 133.243 33.311 171.7053 3.553e-15 ***
brand:car  5   6.803   1.361   7.0137 0.0006214 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.2 Chapter 7

8.2.1 p232

(105) MODEL

```
v1p232 = read.table("C:/G/Rt/Kemp/v1p232.txt", head=TRUE)
v1p232 = af(v1p232,c("trt"))
```

```

GLM(yield ~ trt, v1p232) # OK

$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      4 59.174 14.793 28.781 0.0012 **
RESIDUALS   5  2.570  0.514
CORRECTED TOTAL 9 61.744
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE yield Mean Coef Var R-square Adj R-sq
0.7169379     15.94 4.497729 0.9583765 0.9250777

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
trt    4 59.174 14.793 28.781 0.0012 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
trt    4 59.174 14.793 28.781 0.0012 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
trt    4 59.174 14.793 28.781 0.0012 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.2.2 p235

(106) MODEL

```

v1p235 = read.table("C:/G/Rt/Kemp/v1p235.txt", head=TRUE)
v1p235 = af(v1p235,c("density"))
GLM(yield ~ density, v1p235) # OK

```

```

$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value     Pr(>F)
MODEL      4 88.007 22.0017 32.198 1.095e-05 ***
RESIDUALS   10  6.833  0.6833
CORRECTED TOTAL 14 94.840
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE yield Mean Coef Var R-square Adj R-sq
0.8266398      16.4 5.040486 0.9279488 0.8991284

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
density  4 88.007 22.002 32.198 1.095e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
density  4 88.007 22.002 32.198 1.095e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
density  4 88.007 22.002 32.198 1.095e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.3 Chapter 8

8.3.1 p265

(107) MODEL

```
v1p265 = read.table("C:/G/Rt/Kemp/v1p265.txt", head=TRUE)
v1p265 = af(v1p265,c("trt"))
GLM(y ~ trt + x, v1p265) # OK
```

```
$ANOVA
Response : y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      3 84.678 28.2260 36.866 4.941e-06 ***
RESIDUALS 11 8.422  0.7656
CORRECTED TOTAL 14 93.100
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
  Root MSE y Mean Coef Var R-square Adj R-sq
0.8750081     9 9.722312 0.9095378 0.8848663
```

```
$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
```

```

trt 2 66.868 33.434 43.668 5.858e-06 ***
x 1 17.810 17.810 23.262 0.0005333 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt 2 83.147 41.573 54.299 1.996e-06 ***
x 1 17.810 17.810 23.262 0.0005333 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt 2 83.147 41.573 54.299 1.996e-06 ***
x 1 17.810 17.810 23.262 0.0005333 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.3.2 p272

(108) MODEL

```
GLM(y ~ trt + x %in% trt, v1p265) # OK
```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 85.711 17.142 20.881 0.0001046 ***
RESIDUALS  9 7.389  0.821
CORRECTED TOTAL 14 93.100
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE y Mean Coef Var R-square Adj R-sq
0.9060697    9 10.06744 0.9206374 0.876547

```

```

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt      2 66.868 33.434 40.7254 3.092e-05 ***
trt:x   3 18.843  6.281  7.6509  0.007578 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt      2 66.868 33.434 40.7254 3.092e-05 ***
trt:x   3 18.843  6.281  7.6509  0.007578 **

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt     2 6.1392  3.0696  3.7390 0.065769 .
trt:x   3 18.8433  6.2811  7.6509 0.007578 **

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.3.3 p273

(109) MODEL

```
GLM(y ~ trt + x + x %in% trt, v1p265) # OK
```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 85.711 17.142  20.881 0.0001046 ***
RESIDUALS   9  7.389   0.821
CORRECTED TOTAL 14 93.100

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```
$Fitness
  Root MSE y Mean Coef Var  R-square Adj R-sq
  0.9060697    9 10.06744 0.9206374 0.876547
```

```

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt     2 66.868 33.434 40.7254 3.092e-05 ***
x       1 17.810 17.810 21.6940  0.001189 **
trt:x   2  1.033   0.517  0.6294  0.554843

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt     2 83.147 41.573 50.6397 1.267e-05 ***
x       1 17.810 17.810 21.6940  0.001189 **
trt:x   2  1.033   0.517  0.6294  0.554843

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
trt     2 6.1392  3.0696  3.7390 0.065769 .
x       1 17.2071 17.2071 20.9597 0.001331 **
```

```

trt:x 2 1.0334 0.5167 0.6294 0.554843
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.4 Chapter 9

8.4.1 p344

(110) MODEL

```

v1p344 = read.table("C:/G/Rt/Kemp/v1p344.txt", head=TRUE)
v1p344 = af(v1p344,c("diet", "litter"))
GLM(gain ~ litter + diet, v1p344)

```

\$ANOVA

Response : gain

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	9	4915.6	546.18	15.544	3.363e-07 ***
RESIDUALS	20	702.8	35.14		
CORRECTED TOTAL	29	5618.4			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	gain	Mean Coef	Var R-square	Adj R-sq
5.927698	68.31333	8.677219	0.874919	0.8186325

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
litter	5	4438.0	887.6	25.2608	5.298e-08 ***
diet	4	477.6	119.4	3.3981	0.02824 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
litter	5	4438.0	887.6	25.2608	5.298e-08 ***
diet	4	477.6	119.4	3.3981	0.02824 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
litter	5	4438.0	887.6	25.2608	5.298e-08 ***
diet	4	477.6	119.4	3.3981	0.02824 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

8.4.2 p349

(111) MODEL

```
v1p349 = read.table("C:/G/Rt/Kemp/v1p349.txt", head=TRUE)
v1p349 = af(v1p349,c("subject", "exercise"))
GLM(diast ~ subject + exercise + subject:exercise, v1p349) # OK

$ANOVA
Response : diast
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      14 1541.5 110.105  28.475 2.953e-08 ***
RESIDUALS   15   58.0   3.867
CORRECTED TOTAL 29 1599.5
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE diast Mean Coef Var R-square Adj R-sq
1.966384 134.5333 1.461633 0.9637379 0.9298933

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
subject      4 905.13 226.283 58.5216 5.672e-09 ***
exercise     2 591.27 295.633 76.4569 1.357e-08 ***
subject:exercise 8 45.07   5.633  1.4569     0.2522
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
subject      4 905.13 226.283 58.5216 5.672e-09 ***
exercise     2 591.27 295.633 76.4569 1.357e-08 ***
subject:exercise 8 45.07   5.633  1.4569     0.2522
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
subject      4 905.13 226.283 58.5216 5.672e-09 ***
exercise     2 591.27 295.633 76.4569 1.357e-08 ***
subject:exercise 8 45.07   5.633  1.4569     0.2522
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.4.3 p354

(112) MODEL

```

v1p354 = read.table("C:/G/Rt/Kemp/v1p354.txt", head=TRUE)
v1p354 = af(v1p354,c("loc", "block", "HSF"))
GLM(height ~ loc + block %in% loc + HSF + loc:HSF + block:loc:HSF, v1p354) # OK

$ANOVA
Response : height
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      23 40782 1773.12 80.444 < 2.2e-16 ***
RESIDUALS   24     529   22.04
CORRECTED TOTAL 47 41311
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE height Mean Coef Var R-square Adj R-sq
4.694855   210.6667 2.228571 0.9871946 0.9749227

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
loc        1 20336.3 20336.3 922.6314 < 2.2e-16 ***
loc:block  6  1462.3   243.7 11.0573 6.408e-06 ***
HSF        2 12170.7  6085.3 276.0832 < 2.2e-16 ***
loc:HSF    2  6511.2  3255.6 147.7013 3.242e-14 ***
loc:block:HSF 12   301.2    25.1   1.1386    0.3769
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
loc        1 20336.3 20336.3 922.6314 < 2.2e-16 ***
loc:block  6  1462.3   243.7 11.0573 6.408e-06 ***
HSF        2 12170.7  6085.3 276.0832 < 2.2e-16 ***
loc:HSF    2  6511.2  3255.6 147.7013 3.242e-14 ***
loc:block:HSF 12   301.2    25.1   1.1386    0.3769
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
loc        1 20336.3 20336.3 922.6314 < 2.2e-16 ***
loc:block  6  1462.3   243.7 11.0573 6.408e-06 ***
HSF        2 12170.7  6085.3 276.0832 < 2.2e-16 ***
loc:HSF    2  6511.2  3255.6 147.7013 3.242e-14 ***
loc:block:HSF 12   301.2    25.1   1.1386    0.3769
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.4.4 p357

(113) MODEL

```
v1p357 = read.table("C:/G/Rt/Kemp/v1p357.txt", head=TRUE)
v1p357 = af(v1p357,c("var", "N"))
GLM(y ~ var + N + var:N, v1p357) # OK
```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	9	4465.4	496.16	14.116	0.000142 ***
RESIDUALS	10	351.5	35.15		
CORRECTED TOTAL	19	4816.9			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root	MSE	y	Mean	Coef	Var	R-square	Adj R-sq
5.928744	137.55	4.310246	0.9270285	0.8613542			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
var	1	140.5	140.45	3.9957	0.073519 .
N	4	3393.7	848.42	24.1373	4.027e-05 ***
var:N	4	931.3	232.82	6.6238	0.007152 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
var	1	140.5	140.45	3.9957	0.073519 .
N	4	3393.7	848.43	24.1373	4.027e-05 ***
var:N	4	931.3	232.82	6.6238	0.007152 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
var	1	140.5	140.45	3.9957	0.073519 .
N	4	3393.7	848.43	24.1373	4.027e-05 ***
var:N	4	931.3	232.82	6.6238	0.007152 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

8.4.5 p361

(114) MODEL

```

v1p361 = read.table("C:/G/Rt/Kemp/v1p361.txt", head=TRUE)
v1p361 = af(v1p361,c("block", "trt"))
GLM(y ~ block + trt, v1p361) # OK

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL       4 241.33  60.333  40.222 0.1176
RESIDUALS    1   1.50   1.500
CORRECTED TOTAL 5 242.83

$Fitness
Root MSE   y Mean Coef Var R-square Adj R-sq
1.224745 19.83333 6.175184 0.9938229 0.9691146

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
block  2 24.333 12.167  8.1111 0.24097
trt    2 217.000 108.500 72.3333 0.08286 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
block  2    108     54.0  36.000 0.11704
trt    2    217     108.5  72.333 0.08286 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
block  2    108     54.0  36.000 0.11704
trt    2    217     108.5  72.333 0.08286 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

y = model.frame(y ~ block + trt, v1p361)[,1]
x = ModelMatrix(y ~ block + trt, v1p361)
rx = lfit(x, y)
K = cbind(rep(1, 3), matrix(1/3, nrow=3, ncol=3), diag(3)) ; K

[,1]      [,2]      [,3]      [,4]      [,5]      [,6]      [,7]
[1,] 1 0.3333333 0.3333333 0.3333333 1 0 0
[2,] 1 0.3333333 0.3333333 0.3333333 0 1 0
[3,] 1 0.3333333 0.3333333 0.3333333 0 0 1

est(K, x$X, rx)

Estimate Lower CL Upper CL Std. Error t value Df Pr(>|t|)

```

```
[1,] 29.5 17.334735 41.66526 0.9574271 30.81175 1 0.02065434
[2,] 16.5 4.334735 28.66526 0.9574271 17.23369 1 0.03689905
[3,] 13.5 1.334735 25.66526 0.9574271 14.10029 1 0.04507394
attr(,"Estimability")
[1] TRUE TRUE TRUE
```

8.5 Chapter 10

8.5.1 p405

(115) MODEL

```
v1p405 = read.table("C:/G/Rt/Kemp/v1p405.txt", head=TRUE)
v1p405 = af(v1p405,c("trt", "Row", "Col"))
GLM(y ~ Row + Col + trt, v1p405) # OK
```

\$ANOVA

```
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      12 4094.7 341.23 2.3416 0.07739 .
RESIDUALS  12 1748.7 145.73
CORRECTED TOTAL 24 5843.4
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
Root MSE y Mean Coef Var R-square Adj R-sq
12.07173 93.32 12.93584 0.7007379 0.4014758
```

\$`Type I`

```
Df Sum Sq Mean Sq F value Pr(>F)
Row 4 514.24 128.56 0.8822 0.50328
Col 4 1711.44 427.86 2.9360 0.06611 .
trt 4 1869.04 467.26 3.2064 0.05229 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
Df Sum Sq Mean Sq F value Pr(>F)
Row 4 514.24 128.56 0.8822 0.50328
Col 4 1711.44 427.86 2.9360 0.06611 .
trt 4 1869.04 467.26 3.2064 0.05229 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```
Df Sum Sq Mean Sq F value Pr(>F)
Row 4 514.24 128.56 0.8822 0.50328
Col 4 1711.44 427.86 2.9360 0.06611 .
```

```

trt 4 1869.04 467.26 3.2064 0.05229 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.5.2 p408

(116) MODEL

```

v1p408 = read.table("C:/G/Rt/Kemp/v1p408.txt", head=TRUE)
v1p408 = af(v1p408,c("breed", "farm", "wclass", "dosage"))
GLM(response ~ breed + breed:farm + wclass + dosage + breed:dosage, v1p408) # OK

```

\$ANOVA

```

Response : response
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       16 4470.3 279.391 140.87 2.039e-13 ***
RESIDUALS   15   29.7   1.983
CORRECTED TOTAL 31 4500.0
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$Fitness

Root MSE	response	Mean Coef	Var	R-square	Adj R-sq
1.408309		155.75	0.904211	0.9933889	0.986337

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
breed	1	3280.5	3280.5	1654.0336	< 2.2e-16 ***
breed:farm	6	9.0	1.5	0.7563	0.6146
wclass	3	466.8	155.6	78.4454	2.142e-09 ***
dosage	3	580.2	193.4	97.5210	4.596e-10 ***
breed:dosage	3	133.8	44.6	22.4790	8.366e-06 ***

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
breed	1	3280.5	3280.5	1654.0336	< 2.2e-16 ***
breed:farm	6	9.0	1.5	0.7563	0.6146
wclass	3	466.8	155.6	78.4454	2.142e-09 ***
dosage	3	580.2	193.4	97.5210	4.596e-10 ***
breed:dosage	3	133.8	44.6	22.4790	8.366e-06 ***

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
breed	1	3280.5	3280.5	1654.0336	< 2.2e-16 ***
breed:farm	6	9.0	1.5	0.7563	0.6146

```
wclass      3  466.8   155.6   78.4454 2.142e-09 ***
dosage      3  580.2   193.4   97.5210 4.596e-10 ***
breed:dosage 3  133.8    44.6   22.4790 8.366e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.5.3 p410

(117) MODEL

```
v1p410 = read.table("C:/G/Rt/Kemp/v1p410.txt", head=TRUE)
v1p410$carry = ifelse(v1p410$carry == 0, 3, v1p410$carry)
v1p410 = af(v1p410,c("period", "sequence", "steer", "trt", "carry"))
GLM(y ~ period + sequence + steer:sequence + trt + carry, v1p410) # OK
```

\$ANOVA

```
Response : y
          Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL       17 1302.51  76.618  8.7402 1.572e-05 ***
RESIDUALS   18  157.79   8.766
CORRECTED TOTAL 35 1460.31
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
Root MSE     y Mean Coef Var R-square Adj R-sq
2.960778 52.36111 5.654535 0.8919461 0.7898953
```

\$`Type I`

```
          Df  Sum Sq Mean Sq F value    Pr(>F)
period       2 292.06 146.028 16.6580 8.038e-05 ***
sequence     5 326.47  65.294  7.4484 0.0006072 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt          2 549.06 274.528 31.3166 1.377e-06 ***
carry         2  16.43   8.215  0.9372 0.4100385
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
          Df  Sum Sq Mean Sq F value    Pr(>F)
period       2 172.31  86.154  9.8279 0.0013030 **
sequence     5 318.69  63.738  7.2709 0.0006954 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt          2 440.61 220.304 25.1311 6.164e-06 ***
carry         2  16.43   8.215  0.9372 0.4100385
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
period        2 172.31  86.154  9.8279 0.0013030 ***
sequence      5 318.69  63.738  7.2709 0.0006954 ***
sequence:steer 6 118.50  19.750  2.2530 0.0849122 .
trt           2 440.61 220.304 25.1311 6.164e-06 ***
carry          2   16.43   8.215  0.9372 0.4100385
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(y ~ period + sequence + steer:sequence + trt + carry, v1p410), type=3,
singular.ok=TRUE) # NOT OK for sequence

```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

```

Response: y
          Sum Sq Df F values    Pr(>F)
period        172.31  2  9.8279 0.001303 ***
sequence      0.00   0
trt           440.61  2 25.1311 6.164e-06 ***
carry          16.43  2  0.9372 0.410038
sequence:steer 118.50  6  2.2530 0.084912 .
Residuals     157.79 18
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.6 Chapter 11

8.6.1 p432

(118) MODEL

```

v1p432 = read.table("C:/G/Rt/Kemp/v1p432.txt", head=TRUE)
v1p432 = af(v1p432,c("V", "Block", "A", "B", "C"))
GLM(Y ~ V + Block:V + A + B + A:B + V:A + V:B + V:A:B + Block:A:V + Block:B:V,
v1p432) # OK

```

```

$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       94 261663 2783.65 30.584 2.065e-14 ***
RESIDUALS    25   2275   91.02
CORRECTED TOTAL 119 263939
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

\$Fitness

	Root	MSE	Y	Mean	Coef	Var	R-square	Adj R-sq
	9.540266	612.9	1.556578	0.991379	0.958964			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	282.2094	< 2.2e-16 ***
V:Block	25	50019	2001	21.9825	1.588e-11 ***
A	1	18451	18451	202.7233	1.692e-13 ***
B	1	78541	78541	862.9280	< 2.2e-16 ***
A:B	1	108	108	1.1899	0.28575
V:A	4	3751	938	10.3023	4.532e-05 ***
V:B	4	307	77	0.8421	0.51168
V:A:B	4	1495	374	4.1058	0.01081 *
V:Block:A	25	3416	137	1.5011	0.15818
V:Block:B	25	2833	113	1.2451	0.29390

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	282.2094	< 2.2e-16 ***
V:Block	25	50019	2001	21.9825	1.588e-11 ***
A	1	18451	18451	202.7233	1.692e-13 ***
B	1	78541	78541	862.9280	< 2.2e-16 ***
A:B	1	108	108	1.1899	0.28575
V:A	4	3751	938	10.3023	4.532e-05 ***
V:B	4	307	77	0.8421	0.51168
V:A:B	4	1495	374	4.1058	0.01081 *
V:Block:A	25	3416	137	1.5011	0.15818
V:Block:B	25	2833	113	1.2451	0.29390

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	282.2094	< 2.2e-16 ***
V:Block	25	50019	2001	21.9825	1.588e-11 ***
A	1	18451	18451	202.7233	1.692e-13 ***
B	1	78541	78541	862.9280	< 2.2e-16 ***
A:B	1	108	108	1.1899	0.28575
V:A	4	3751	938	10.3023	4.532e-05 ***
V:B	4	307	77	0.8421	0.51168
V:A:B	4	1495	374	4.1058	0.01081 *
V:Block:A	25	3416	137	1.5011	0.15818
V:Block:B	25	2833	113	1.2451	0.29390

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

8.6.2 p434

(119) MODEL

```
GLM(Y ~ V + Block:V + A + B + A:B + V:A + V:B + V:A:B, v1p432) # OK
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	44	255415	5804.9	51.075	< 2.2e-16 ***
RESIDUALS	75	8524	113.7		
CORRECTED TOTAL	119	263939			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	MSE	Y	Mean	Coef	Var	R-square	Adj R-sq
10.66088	612.9	1.739417	0.9677043	0.9487575			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
A	1	18451	18451	162.3447	< 2.2e-16 ***
B	1	78541	78541	691.0494	< 2.2e-16 ***
A:B	1	108	108	0.9529	0.33212
V:A	4	3751	938	8.2503	1.435e-05 ***
V:B	4	307	77	0.6744	0.61182
V:A:B	4	1495	374	3.2880	0.01541 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***
A	1	18451	18451	162.3447	< 2.2e-16 ***
B	1	78541	78541	691.0494	< 2.2e-16 ***
A:B	1	108	108	0.9529	0.33212
V:A	4	3751	938	8.2503	1.435e-05 ***
V:B	4	307	77	0.6744	0.61182
V:A:B	4	1495	374	3.2880	0.01541 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
V	4	102743	25686	225.9988	< 2.2e-16 ***
V:Block	25	50019	2001	17.6040	< 2.2e-16 ***

```

A      1 18451  18451 162.3447 < 2.2e-16 ***
B      1 78541  78541 691.0494 < 2.2e-16 ***
A:B    1   108   108  0.9529  0.33212
V:A    4   3751   938  8.2503 1.435e-05 ***
V:B    4    307    77  0.6744  0.61182
V:A:B  4   1495   374  3.2880  0.01541 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.6.3 p438

(120) MODEL

```
GLM(Y ~ V + Block:V + C + V:C, v1p432) # OK
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      44 255415  5804.9 51.075 < 2.2e-16 ***
RESIDUALS  75  8524   113.7
CORRECTED TOTAL 119 263939
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE Y Mean Coef Var R-square Adj R-sq
10.66088 612.9 1.739417 0.9677043 0.9487575
```

```
$`Type I`
          Df Sum Sq Mean Sq F value    Pr(>F)
V         4 102743  25686 225.9988 < 2.2e-16 ***
V:Block  25 50019   2001  17.6040 < 2.2e-16 ***
C         3  97100  32367 284.7823 < 2.2e-16 ***
V:C       12  5552    463   4.0709  7.23e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
          Df Sum Sq Mean Sq F value    Pr(>F)
V         4 102743  25686 225.9988 < 2.2e-16 ***
V:Block  25 50019   2001  17.6040 < 2.2e-16 ***
C         3  97100  32367 284.7823 < 2.2e-16 ***
V:C       12  5552    463   4.0709  7.23e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
          Df Sum Sq Mean Sq F value    Pr(>F)
V         4 102743  25686 225.9988 < 2.2e-16 ***
```

```

V:Block 25 50019      2001 17.6040 < 2.2e-16 ***
C          3 97100     32367 284.7823 < 2.2e-16 ***
V:C       12 5552      463   4.0709  7.23e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.6.4 p444

(121) MODEL

```

v1p444 = v1p432[v1p432$Block==5,]
GLM(Y ~ V + A + B + A:B + V:A, v1p444) # OK

```

```

$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      11 39278  3570.8 59.787 1.897e-06 ***
RESIDUALS    8    478    59.7
CORRECTED TOTAL 19 39756
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE Y Mean Coef Var R-square Adj R-sq
7.728195 630.7 1.225336 0.9879817 0.9714567

```

```

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
V      4 19287.7 4821.9 80.7355 1.674e-06 ***
A      1 3380.0 3380.0 56.5927 6.780e-05 ***
B      1 14045.0 14045.0 235.1612 3.247e-07 ***
A:B    1   115.2   115.2   1.9288  0.202326
V:A    4 2450.5   612.6  10.2574  0.003081 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
V      4 19287.7 4821.9 80.7355 1.674e-06 ***
A      1 3380.0 3380.0 56.5927 6.780e-05 ***
B      1 14045.0 14045.0 235.1612 3.247e-07 ***
A:B    1   115.2   115.2   1.9288  0.202326
V:A    4 2450.5   612.6  10.2574  0.003081 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
V      4 19287.7 4821.9 80.7355 1.674e-06 ***

```

```

A     1 3380.0 3380.0 56.5927 6.780e-05 ***
B     1 14045.0 14045.0 235.1612 3.247e-07 ***
A:B   1    115.2    115.2   1.9288   0.202326
V:A   4   2450.5    612.6   10.2574   0.003081 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.6.5 p482

(122) MODEL

```

v1p482 = read.table("C:/G/Rt/Kemp/v1p482.txt", head=TRUE)
v1p482 = af(v1p482,c("block", "A", "B"))
GLM(y ~ block + A + B + A:B, v1p482) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       8 156.88 19.6094 9.8871 9.377e-05 ***
RESIDUALS   15 29.75  1.9833
CORRECTED TOTAL 23 186.62
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE y Mean Coef Var R-square Adj R-sq
1.408309 7.875 17.88328 0.8405894 0.7555704

```

```

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
block   5 108.38 21.675 10.9286 0.0001415 ***
A       1    4.00    4.000  2.0168 0.1760166
B       1   42.25   42.250 21.3025 0.0003365 ***
A:B    1    2.25    2.250  1.1345 0.3036727
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
block   5 31.417   6.283   3.1681 0.0377804 *
A       1  4.000   4.000   2.0168 0.1760166
B       1 42.250   42.250 21.3025 0.0003365 ***
A:B    1   2.250   2.250  1.1345 0.3036727
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
block   5 31.417   6.283   3.1681 0.0377804 *

```

```

A      1  4.000  4.000  2.0168 0.1760166
B      1 42.250 42.250 21.3025 0.0003365 ***
A:B    1  2.250  2.250  1.1345 0.3036727
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.7 Chapter 12

8.7.1 p525

(123) MODEL

```
v1p525 = read.table("C:/G/Rt/Kemp/v1p525.txt", head=TRUE)
REG(y ~ x1 + x2 + x3, v1p525)
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      3 84.948 28.3158 164.15 5.26e-10 ***
RESIDUALS 12  2.070  0.1725
CORRECTED TOTAL 15 87.017
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE  y Mean Coef Var R-square Adj R-sq PRESS     R2pred
0.4153312 14.2125 2.922295 0.9762117 0.9702646 3.68 0.9577097
```

```
$Coefficients
            Estimate Std. Error Df Lower CL Upper CL t value Pr(>|t|)
(Intercept) 14.2125    0.10383 12  13.9863 14.4387 136.8787 < 2.2e-16 ***
x1          0.7875    0.10383 12   0.5613  1.0137  7.5843 6.465e-06 ***
x2          1.3875    0.10383 12   1.1613  1.6137 13.3628 1.446e-08 ***
x3          1.6625    0.10383 12   1.4363  1.8887 16.0113 1.839e-09 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
GLM(y ~ x1 + x2 + x3, v1p525) # OK
```

```
$ANOVA
Response : y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      3 84.948 28.3158 164.15 5.26e-10 ***
RESIDUALS 12  2.070  0.1725
CORRECTED TOTAL 15 87.017
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE  y Mean Coef Var R-square Adj R-sq
```

```

0.4153312 14.2125 2.922295 0.9762117 0.9702646

$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

x1  1  9.923   9.923  57.522 6.465e-06 ***  

x2  1 30.803  30.803 178.565 1.446e-08 ***  

x3  1 44.223  44.223 256.362 1.839e-09 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

x1  1  9.923   9.923  57.522 6.465e-06 ***  

x2  1 30.803  30.803 178.565 1.446e-08 ***  

x3  1 44.223  44.223 256.362 1.839e-09 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

x1  1  9.923   9.923  57.522 6.465e-06 ***  

x2  1 30.803  30.803 178.565 1.446e-08 ***  

x3  1 44.223  44.223 256.362 1.839e-09 ***  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.7.2 p527

(124) MODEL

```
v1p527 = read.table("C:/G/Rt/Kemp/v1p527.txt", head=TRUE)
GLM(y ~ A + B, v1p527) # OK
```

```
$ANOVA  

Response : y  

  Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      2  22.99 11.4952  4.8917 0.04686 *  

RESIDUALS   7  16.45  2.3499  

CORRECTED TOTAL 9  39.44  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

Root MSE y Mean Coef Var R-square Adj R-sq  

1.532954   5.2 29.47989 0.5829197 0.4637539
```

```
$`Type I`  

  Df Sum Sq Mean Sq F value    Pr(>F)  

A  1 10.364  10.364  4.4103 0.07386 .
```

```

B 1 12.626 12.626 5.3730 0.05355 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 10.364 10.364 4.4103 0.07386 .
B 1 12.626 12.626 5.3730 0.05355 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 10.364 10.364 4.4103 0.07386 .
B 1 12.626 12.626 5.3730 0.05355 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.7.3 p529

(125) MODEL

```
v1p529 = read.table("C:/G/Rt/Kemp/v1p529.txt", head=TRUE)
GLM(y ~ A + B + I(A*A) + I(B*B) + I(A*B), v1p529) # OK
```

```
$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      5 35.713 7.1427 6.7928 0.01857 *
RESIDUALS   6  6.309  1.0515
CORRECTED TOTAL 11 42.023
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE y Mean Coef Var R-square Adj R-sq
1.025434 5.275 19.4395 0.8498641 0.7247508
```

```
$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
A      1 11.6012 11.6012 11.0329 0.01597 *
B      1 12.6263 12.6263 12.0077 0.01338 *
I(A * A) 1  1.7167  1.7167  1.6326 0.24855
I(B * B) 1  5.3593  5.3593  5.0967 0.06476 .
I(A * B) 1  4.4100  4.4100  4.1940 0.08649 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
```

```

          Df  Sum Sq Mean Sq F value Pr(>F)
A           1 11.6012 11.6012 11.0329 0.01597 *
B           1 12.6263 12.6263 12.0077 0.01338 *
I(A * A)   1  5.5468  5.5468  5.2750 0.06137 .
I(B * B)   1  5.3593  5.3593  5.0967 0.06476 .
I(A * B)   1  4.4100  4.4100  4.1940 0.08649 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
          Df  Sum Sq Mean Sq F value Pr(>F)
A           1 11.6012 11.6012 11.0329 0.01597 *
B           1 12.6263 12.6263 12.0077 0.01338 *
I(A * A)   1  5.5468  5.5468  5.2750 0.06137 .
I(B * B)   1  5.3593  5.3593  5.0967 0.06476 .
I(A * B)   1  4.4100  4.4100  4.1940 0.08649 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.8 Chapter 13

8.8.1 p563

(126) MODEL

```
v1p563 = read.table("C:/G/Rt/Kemp/v1p563.txt", head=TRUE)
v1p563 = af(v1p563, c("rep", "A", "B"))
GLM(y ~ rep + A + rep:A + B + A:B, v1p563) # OK
```

```
$ANOVA
Response : y
          Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      14 2097.08 149.792 17.228 8.385e-05 ***
RESIDUALS   9   78.25   8.694
CORRECTED TOTAL 23 2175.33
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE   y Mean Coef Var R-square Adj R-sq
2.948634 31.16667 9.460859 0.9640285 0.9080728
```

```
$`Type I` 
          Df  Sum Sq Mean Sq F value    Pr(>F)
rep      3 1241.00 413.67 47.5783 7.606e-06 ***
A       2  353.08 176.54 20.3051 0.0004613 ***
rep:A   6  192.25  32.04  3.6853 0.0393557 *
B       1  216.00 216.00 24.8435 0.0007550 ***
A:B     2   94.75  47.38  5.4489 0.0281496 *
```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep     3 1241.00  413.67 47.5783 7.606e-06 ***  

A       2   353.08   176.54 20.3051 0.0004613 ***  

rep:A   6   192.25    32.04  3.6853 0.0393557 *  

B       1   216.00   216.00 24.8435 0.0007550 ***  

A:B     2    94.75    47.38  5.4489 0.0281496 *
---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

rep     3 1241.00  413.67 47.5783 7.606e-06 ***  

A       2   353.08   176.54 20.3051 0.0004613 ***  

rep:A   6   192.25    32.04  3.6853 0.0393557 *  

B       1   216.00   216.00 24.8435 0.0007550 ***  

A:B     2    94.75    47.38  5.4489 0.0281496 *
---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

8.8.2 p566

(127) MODEL

```

v1p566 = read.table("C:/G/Rt/Kemp/v1p566.txt", head=TRUE)
v1p566 = af(v1p566, c("subject", "A", "B"))
GLM(y ~ A + B + A:B, v1p566) # OK

```

```

$ANOVA  

Response : y  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL      5 1469.58  293.92     86.2 5.592e-09 ***  

RESIDUALS  12   40.92    3.41  

CORRECTED TOTAL 17 1510.50  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

Root MSE    y Mean Coef Var R-square Adj R-sq  

1.846543 35.83333 5.153144 0.9729118 0.9616251

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

A       2 1390.04  695.02 203.8350 5.466e-10 ***  

B       1   76.06   76.06 22.3055 0.0004945 ***  

A:B     2    3.49    1.74  0.5112 0.6122667

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df  Sum Sq Mean Sq F value    Pr(>F)
A     2 1390.04  695.02 203.8350 5.466e-10 ***
B     1   76.06   76.06  22.3055 0.0004945 ***
A:B   2    3.49    1.74   0.5112 0.6122667
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df  Sum Sq Mean Sq F value    Pr(>F)
A     2 1390.04  695.02 203.8350 5.466e-10 ***
B     1   79.00   79.00  23.1700 0.0004237 ***
A:B   2    3.49    1.74   0.5112 0.6122667
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

8.9 Chapter 14

8.9.1 p581

(128) MODEL

```
v1p581 = read.table("C:/G/Rt/Kemp/v1p581.txt", head=TRUE)
v1p581 = af(v1p581, c("drug", "person", "time"))
GLM(rate ~ drug + person:drug + time + drug:time, v1p581) # OK
```

```
$ANOVA
Response : rate
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      23 2449.5 106.500 12.733 3.469e-11 ***
RESIDUALS  36  301.1   8.364
CORRECTED TOTAL 59 2750.6
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE rate Mean Coef Var R-square Adj R-sq
2.892039      77.7 3.722058 0.890533 0.8205957
```

```
$`Type I`
  Df  Sum Sq Mean Sq F value    Pr(>F)
drug        2  337.60 168.800 20.1820 1.323e-06 ***
drug:person 12 1498.50 124.875 14.9303 1.501e-10 ***
time        3  256.33  85.444 10.2159 5.230e-05 ***
drug:time    6  357.07  59.511  7.1152 4.707e-05 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
drug	2	337.60	168.800	20.1820	1.323e-06 ***
drug:person	12	1498.50	124.875	14.9303	1.501e-10 ***
time	3	256.33	85.444	10.2159	5.230e-05 ***
drug:time	6	357.07	59.511	7.1152	4.707e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
drug	2	337.60	168.800	20.1820	1.323e-06 ***
drug:person	12	1498.50	124.875	14.9303	1.501e-10 ***
time	3	256.33	85.444	10.2159	5.230e-05 ***
drug:time	6	357.07	59.511	7.1152	4.707e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

9 Hinkelmann & Kempthorne - Volume 2

Reference - Hinkelmann K, Kempthorne O. Design and Analysis of Experiments Volume 2 Advanced Experimental Design. 2e. John Wiley & Sons Inc. 2008.

9.1 Chapter 1

9.1.1 p53

(129) MODEL

```
v2p53 = read.table("C:/G/Rt/Kemp/v2p53.txt", head=TRUE)
v2p53 = af(v2p53, c("TRT", "BLOCK"))
GLM(Y ~ BLOCK + TRT, v2p53) # OK
```

\$ANOVA

```
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL    7 518.21  74.030  8.1408 0.1137
RESIDUALS 2   18.19   9.094
CORRECTED TOTAL 9 536.40
```

\$Fitness

```
Root MSE Y Mean Coef Var R-square Adj R-sq
3.015585 19.4 15.54425 0.9660934 0.8474203
```

\$`Type I`

```
      Df Sum Sq Mean Sq F value Pr(>F)
BLOCK  4 261.40  65.350  7.1863 0.12587
TRT    3 256.81  85.604  9.4135 0.09755 .
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
      Df Sum Sq Mean Sq F value Pr(>F)
BLOCK  4 79.146  19.786  2.1758 0.33880
TRT    3 256.812  85.604  9.4135 0.09755 .
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```
      Df Sum Sq Mean Sq F value Pr(>F)
BLOCK  4 79.146  19.786  2.1758 0.33880
TRT    3 256.812  85.604  9.4135 0.09755 .
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.1.2 p62

(130) MODEL

```
GLM(Y ~ TRT + BLOCK, v2p53) # OK
```

```
$ANOVA
Response : Y
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL      7 518.21 74.030 8.1408 0.1137
RESIDUALS   2 18.19  9.094
CORRECTED TOTAL 9 536.40
```

```
$Fitness
Root MSE Y Mean Coef Var R-square Adj R-sq
3.015585 19.4 15.54425 0.9660934 0.8474203
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
TRT       3 439.07 146.356 16.0941 0.05907 .
BLOCK     4  79.15 19.786  2.1758 0.33880
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
TRT       3 256.812 85.604 9.4135 0.09755 .
BLOCK     4  79.146 19.786  2.1758 0.33880
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
TRT       3 256.812 85.604 9.4135 0.09755 .
BLOCK     4  79.146 19.786  2.1758 0.33880
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.2 Chapter 2

9.2.1 p82

(131) MODEL

```
v2p82 = read.table("C:/G/Rt/Kemp/v2p82.txt", head=TRUE)
v2p82 = af(v2p82, c("B", "Tx"))
GLM(Y ~ B + Tx, v2p82) # OK
```

```
$ANOVA
Response : Y
```

```

          Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        14  889.11  63.508  6.3183 0.000518 ***
RESIDUALS     15  150.77  10.052
CORRECTED TOTAL 29 1039.89
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    Y Mean Coef Var R-square Adj R-sq
3.170413 38.46667 8.241975 0.8550104 0.7196867

$`Type I`
          Df  Sum Sq Mean Sq F value    Pr(>F)
B      9 730.39  81.154  8.0738 0.0002454 ***
Tx     5 158.73  31.745  3.1583 0.0381655 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df  Sum Sq Mean Sq F value    Pr(>F)
B      9 595.74  66.193  6.5854 0.0007602 ***
Tx     5 158.73  31.745  3.1583 0.0381655 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
          Df  Sum Sq Mean Sq F value    Pr(>F)
B      9 595.74  66.193  6.5854 0.0007602 ***
Tx     5 158.73  31.745  3.1583 0.0381655 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.2.2 p87

(132) MODEL

```
v2p87 = read.table("C:/G/Rt/Kemp/v2p87.txt", head=TRUE)
GLM(y ~ x1 + x2 + x3 + x4 + x5 + x6, v2p87) # OK
```

Warning in min(abs(rx\$coefficients[rx\$coefficients != 0])): no non-missing arguments to min; returning Inf

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	5	1613.25	322.65	2.2332	0.2282
RESIDUALS	4	577.91	144.48		
CORRECTED TOTAL	9	2191.16			

```

$Fitness
Root MSE y Mean Coef Var R-square Adj R-sq
12.01991 115.4 10.41587 0.7362523 0.4065678

$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
x1  1 1044.48 1044.48 7.2293 0.05473 .
x2  1   89.79   89.79  0.6215 0.47459
x3  1   10.45   10.45  0.0724 0.80124
x4  1  407.08  407.08  2.8176 0.16854
x5  1   61.44   61.44  0.4253 0.54990
x6  0
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
x1  0
x2  0
x3  0
x4  0
x5  0
x6  0

$`Type III`
CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value Pr(>F)
x1  0
x2  0
x3  0
x4  0
x5  0
x6  0

```

9.3 Chapter 6

9.3.1 p217

(133) MODEL

```

v2p217 = read.table("C:/G/Rt/Kemp/v2p217.txt", head=TRUE)
v2p217 = af(v2p217, c("R", "C", "Tx"))
GLM(Y ~ R + C + Tx, v2p217) # OK

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      22 4305.1 195.687  7.5094 0.0002682 ***
RESIDUALS  13  338.8  26.059

```

```

CORRECTED TOTAL 35 4643.9
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE   Y Mean Coef Var  R-square Adj R-sq
5.104813 27.05556 18.86789 0.9270507 0.803598

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
R    3 3951.4 1317.15 50.5446 1.998e-07 ***
C    8 168.9   21.11  0.8101     0.6062
Tx 11 184.8   16.80  0.6446     0.7638
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
R    3 3403.5 1134.51 43.5360 4.83e-07 ***
C    8 112.4   14.05  0.5390     0.8077
Tx 11 184.8   16.80  0.6446     0.7638
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df Sum Sq Mean Sq F value    Pr(>F)
R    3 3403.5 1134.51 43.5360 4.83e-07 ***
C    8 112.4   14.05  0.5390     0.8077
Tx 11 184.8   16.80  0.6446     0.7638
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.3.2 p234

(134) MODEL

```

v2p234 = read.table("C:/G/Rt/Kemp/v2p234.txt", head=TRUE)
v2p234 = af(v2p234, c("R", "C", "Tx"))
GLM(Y ~ C + R + Tx, v2p234) # OK

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      13 426.50 32.808 7.0936 0.1302
RESIDUALS   2   9.25   4.625
CORRECTED TOTAL 15 435.75

```

```

$Fitness
Root MSE   Y Mean Coef Var  R-square Adj R-sq

```

```

2.150581 29.625 7.259346 0.9787722 0.8407917

$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)  

C   3  16.25  5.417  1.1712 0.49129  

R   3 357.25 119.083 25.7477 0.03762 *  

Tx  7  53.00  7.571  1.6371 0.43052  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df Sum Sq Mean Sq F value Pr(>F)  

C   3  10.25  3.417  0.7387 0.6189  

R   3 285.50  95.167 20.5766 0.0467 *  

Tx  7  53.00  7.571  1.6371 0.4305  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

C   3  10.25  3.417  0.7387 0.6189  

R   3 285.50  95.167 20.5766 0.0467 *  

Tx  7  53.00  7.571  1.6371 0.4305  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.4 Chapter 7

9.4.1 p268

(135) MODEL

```

v2p268 = read.table("C:/G/Rt/Kemp/v2p268.txt", head=TRUE)
v2p268 = af(v2p268, c("A", "B", "C"))
GLM(y ~ block + A*B*C, v2p268) # OK

```

```

$ANOVA
Response : y
  Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL        8 1026.00 128.250  24.981 0.0001765 ***
RESIDUALS     7   35.94   5.134
CORRECTED TOTAL 15 1061.94
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  y Mean Coef Var  R-square  Adj R-sq
2.265817 25.5625 8.863833 0.9661586 0.9274826

```

```
$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

block  1 715.56  715.56 139.3791 7.093e-06 ***  

A      1  68.06   68.06 13.2574 0.0082753 **  

B      1  0.06   0.06  0.0122 0.9152401  

A:B    1  0.56   0.56  0.1096 0.7503276  

C      1 232.56  232.56 45.2991 0.0002698 ***  

A:C    1  0.06   0.06  0.0122 0.9152401  

B:C    1  7.56   7.56  1.4730 0.2642229  

A:B:C  1  1.56   1.56  0.3043 0.5983312  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

block  1 715.56  715.56 139.3791 7.093e-06 ***  

A      1  68.06   68.06 13.2574 0.0082753 **  

B      1  0.06   0.06  0.0122 0.9152401  

A:B    1  0.56   0.56  0.1096 0.7503276  

C      1 232.56  232.56 45.2991 0.0002698 ***  

A:C    1  0.06   0.06  0.0122 0.9152401  

B:C    1  7.56   7.56  1.4730 0.2642229  

A:B:C  1  1.56   1.56  0.3043 0.5983312  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  

  

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

block  1 715.56  715.56 139.3791 7.093e-06 ***  

A      1  68.06   68.06 13.2574 0.0082753 **  

B      1  0.06   0.06  0.0122 0.9152401  

A:B    1  0.56   0.56  0.1096 0.7503276  

C      1 232.56  232.56 45.2991 0.0002698 ***  

A:C    1  0.06   0.06  0.0122 0.9152401  

B:C    1  7.56   7.56  1.4730 0.2642229  

A:B:C  1  1.56   1.56  0.3043 0.5983312  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.4.2 p273

(136) MODEL

```
v2p273 = read.table("C:/G/Rt/Kemp/v2p273.txt", head=TRUE)
v2p273 = af(v2p273, c("block", "A", "B", "C"))
GLM(y ~ block + A*B*C + block:A:B:C, v2p273) # OK
```

```
$ANOVA
Response : y
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	15	2245.0	149.665	129.44	8.427e-14 ***
RESIDUALS	16	18.5	1.156		
CORRECTED TOTAL	31	2263.5			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	y	Mean	Coef	Var	R-square	Adj R-sq
1.075291	25.78125	4.170824	0.9918267	0.9841642		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	1498.78	1498.78	1296.2432	< 2.2e-16 ***
A	1	132.03	132.03	114.1892	1.083e-08 ***
B	1	0.03	0.03	0.0270	0.87148
A:B	1	1.53	1.53	1.3243	0.26673
C	1	504.03	504.03	435.9189	4.926e-13 ***
A:C	1	0.78	0.78	0.6757	0.42316
B:C	1	3.78	3.78	3.2703	0.08938 .
A:B:C	1	2.53	2.53	2.1892	0.15840
block:A:B:C	7	101.47	14.50	12.5367	1.965e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	1498.78	1498.78	1296.2432	< 2.2e-16 ***
A	1	132.03	132.03	114.1892	1.083e-08 ***
B	1	0.03	0.03	0.0270	0.87148
A:B	1	1.53	1.53	1.3243	0.26673
C	1	504.03	504.03	435.9189	4.926e-13 ***
A:C	1	0.78	0.78	0.6757	0.42316
B:C	1	3.78	3.78	3.2703	0.08938 .
A:B:C	1	2.53	2.53	2.1892	0.15840
block:A:B:C	7	101.47	14.50	12.5367	1.965e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	1498.78	1498.78	1296.2432	< 2.2e-16 ***
A	1	132.03	132.03	114.1892	1.083e-08 ***
B	1	0.03	0.03	0.0270	0.87148
A:B	1	1.53	1.53	1.3243	0.26673
C	1	504.03	504.03	435.9189	4.926e-13 ***
A:C	1	0.78	0.78	0.6757	0.42316
B:C	1	3.78	3.78	3.2703	0.08938 .

```

A:B:C      1    2.53    2.53    2.1892   0.15840
block:A:B:C 7  101.47   14.50   12.5367  1.965e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.5 Chapter 8

9.5.1 p304

(137) MODEL

```

v2p304 = read.table("C:/G/Rt/Kemp/v2p304.txt", head=TRUE)
v2p304 = af(v2p304, c("rep", "block", "A", "B", "C"))
GLM(y ~ rep + block %in% rep + A*B*C - A:B:C, v2p304) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      9 699.06  77.674  248.56 5.096e-07 ***
RESIDUALS   6   1.88   0.312
CORRECTED TOTAL 15 700.94
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  y Mean Coef Var R-square  Adj R-sq
0.559017 23.0625 2.423922 0.997325 0.9933125

```

```

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 390.06  390.06 1248.2 3.428e-08 ***
rep:block 2   8.12   4.06   13.0 0.0065918 **
A        1 18.06  18.06   57.8 0.0002696 ***
B        1 175.56  175.56  561.8 3.702e-07 ***
A:B      1   0.06   0.06    0.2 0.6704121
C        1 68.06  68.06  217.8 6.083e-06 ***
A:C      1   0.06   0.06    0.2 0.6704121
B:C      1 39.06  39.06  125.0 3.056e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 390.06  390.06 1248.2 3.428e-08 ***
rep:block 2   8.12   4.06   13.0 0.0065918 **
A        1 18.06  18.06   57.8 0.0002696 ***
B        1 175.56  175.56  561.8 3.702e-07 ***
A:B      1   0.06   0.06    0.2 0.6704121
C        1 68.06  68.06  217.8 6.083e-06 ***

```

```

A:C      1  0.06   0.06    0.2 0.6704121
B:C      1 39.06  39.06  125.0 3.056e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
  Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 390.06 390.06 1248.2 3.428e-08 ***
rep:block 2   8.12   4.06   13.0 0.0065918 **
A       1 18.06  18.06   57.8 0.0002696 ***
B       1 175.56 175.56  561.8 3.702e-07 ***
A:B     1  0.06   0.06    0.2 0.6704121
C       1 68.06  68.06  217.8 6.083e-06 ***
A:C     1  0.06   0.06    0.2 0.6704121
B:C     1 39.06  39.06  125.0 3.056e-05 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.5.2 p309

(138) MODEL

```
GLM(y ~ rep*A*B*C, v2p304) # OK
```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      15 700.94 46.729
RESIDUALS   0   0.00
CORRECTED TOTAL 15 700.94

$Fitness
Root MSE  y Mean Coef Var R-square
NA 23.0625      NA        1

```

```

$`Type I` 
  Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 390.06 390.06
A       1 18.06  18.06
rep:A    1  0.06   0.06
B       1 175.56 175.56
rep:B    1  1.56   1.56
A:B     1  0.06   0.06
rep:A:B  1  0.06   0.06
C       1 68.06  68.06
rep:C    1  0.06   0.06
A:C     1  0.06   0.06
rep:A:C  1  0.06   0.06
B:C     1 39.06  39.06

```

```

rep:B:C    1   0.06   0.06
A:B:C      1   7.56   7.56
rep:A:B:C  1   0.56   0.56

```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	1	390.06	390.06		
A	1	18.06	18.06		
rep:A	1	0.06	0.06		
B	1	175.56	175.56		
rep:B	1	1.56	1.56		
A:B	1	0.06	0.06		
rep:A:B	1	0.06	0.06		
C	1	68.06	68.06		
rep:C	1	0.06	0.06		
A:C	1	0.06	0.06		
rep:A:C	1	0.06	0.06		
B:C	1	39.06	39.06		
rep:B:C	1	0.06	0.06		
A:B:C	1	7.56	7.56		
rep:A:B:C	1	0.56	0.56		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rep	1	390.06	390.06		
A	1	18.06	18.06		
rep:A	1	0.06	0.06		
B	1	175.56	175.56		
rep:B	1	1.56	1.56		
A:B	1	0.06	0.06		
rep:A:B	1	0.06	0.06		
C	1	68.06	68.06		
rep:C	1	0.06	0.06		
A:C	1	0.06	0.06		
rep:A:C	1	0.06	0.06		
B:C	1	39.06	39.06		
rep:B:C	1	0.06	0.06		
A:B:C	1	7.56	7.56		
rep:A:B:C	1	0.56	0.56		

9.6 Chapter 9

9.6.1 p343

(139) MODEL

```

v2p343 = read.table("C:/G/Rt/Kemp/v2p343.txt", head=TRUE)
v2p343 = af(v2p343, c("rep", "block", "A", "B", "C"))
GLM(y ~ rep + block %in% rep + A*B*C - A:B:C, v2p343) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       17 1889.8 111.167 14.659 0.001608 **
RESIDUALS     6   45.5   7.583
CORRECTED TOTAL 23 1935.3
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE      y Mean Coef Var R-square Adj R-sq
2.753785 21.66667 12.70978 0.9764898 0.9098777

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep       2 1537.33  768.67 101.3626 2.375e-05 ***
rep:block 9 127.00   14.11   1.8608  0.23163
A         1  36.00   36.00   4.7473  0.07218 .
B         1  36.00   36.00   4.7473  0.07218 .
A:B       1  12.25   12.25   1.6154  0.25079
C         1  56.25   56.25   7.4176  0.03448 *
A:C       1  81.00   81.00  10.6813  0.01707 *
B:C       1   4.00    4.00   0.5275  0.49502
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep       2 1537.33  768.67 101.3626 2.375e-05 ***
rep:block 9 119.83   13.31   1.7558  0.25388
A         1  36.00   36.00   4.7473  0.07218 .
B         1  36.00   36.00   4.7473  0.07218 .
A:B       1  12.25   12.25   1.6154  0.25079
C         1  56.25   56.25   7.4176  0.03448 *
A:C       1  81.00   81.00  10.6813  0.01707 *
B:C       1   4.00    4.00   0.5275  0.49502
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep       2 1537.33  768.67 101.3626 2.375e-05 ***
rep:block 9 119.83   13.31   1.7558  0.25388
A         1  36.00   36.00   4.7473  0.07218 .
B         1  36.00   36.00   4.7473  0.07218 .
A:B       1  12.25   12.25   1.6154  0.25079

```

```

C           1   56.25   56.25   7.4176  0.03448 *
A:C         1   81.00   81.00  10.6813  0.01707 *
B:C         1    4.00    4.00   0.5275  0.49502
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.6.2 p348

(140) MODEL

```
GLM(y ~ rep + A*B*C + block %in% rep, v2p343) # OK
```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      17 1889.8 111.167 14.659 0.001608 **
RESIDUALS     6   45.5   7.583
CORRECTED TOTAL 23 1935.3
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE   y Mean Coef Var R-square Adj R-sq
2.753785 21.66667 12.70978 0.9764898 0.9098777

```

```

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 1537.33  768.67 101.3626 2.375e-05 ***
A        1   88.17   88.17  11.6264  0.01432 *
B        1   37.50   37.50   4.9451  0.06785 .
A:B      1    2.67    2.67   0.3516  0.57484
C        1   66.67   66.67   8.7912  0.02512 *
A:C      1   37.50   37.50   4.9451  0.06785 .
B:C      1    0.17    0.17   0.0220  0.88700
A:B:C    1   24.00   24.00   3.1648  0.12555
rep:block 8   95.83   11.98   1.5797  0.29730
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 1537.33  768.67 101.3626 2.375e-05 ***
A        1   36.00   36.00   4.7473  0.07218 .
B        1   36.00   36.00   4.7473  0.07218 .
A:B      1   12.25   12.25   1.6154  0.25079
C        1   56.25   56.25   7.4176  0.03448 *
A:C      1   81.00   81.00  10.6813  0.01707 *
B:C      1    4.00    4.00   0.5275  0.49502
A:B:C    0

```

```

rep:block 8 95.83 11.98 1.5797 0.29730
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value    Pr(>F)
rep      2 1537.33 768.67 101.3626 2.375e-05 ***
A        1   36.00   36.00   4.7473  0.07218 .
B        1   36.00   36.00   4.7473  0.07218 .
A:B      1   12.25   12.25   1.6154  0.25079
C        1   56.25   56.25   7.4176  0.03448 *
A:C      1   81.00   81.00  10.6813  0.01707 *
B:C      1     4.00     4.00   0.5275  0.49502
A:B:C    0
rep:block 8 95.83 11.98 1.5797 0.29730
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.6.3 p353

(141) MODEL

```

v2p353 = read.table("C:/G/Rt/Kemp/v2p353.txt", head=TRUE)
v2p353 = af(v2p353, c("rep", "block", "A", "B", "C", "D"))
GLM(y ~ rep + rep:block + A*B*C*D - A:B:C:D, v2p353) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      21 7132.2 339.63 56.022 9.795e-08 ***
RESIDUALS   10   60.6    6.06
CORRECTED TOTAL 31 7192.9
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  y Mean Coef Var R-square Adj R-sq
2.462214 37.1875 6.621081 0.9915715 0.9738717

```

```

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
rep       1 5940.5 5940.5 979.8763 2.600e-11 ***
rep:block 6  777.4   129.6  21.3711 3.675e-05 ***
A         1   171.1   171.1  28.2268 0.0003412 ***
B         1    18.0    18.0   2.9691 0.1155937
A:B       1     1.6     1.6   0.2577 0.6226914
C         1   120.1   120.1  19.8144 0.0012326 **
A:C       1     0.6     0.6   0.0928 0.7669127

```

```

B:C      1   2.0   2.0   0.3299 0.5784103
A:B:C    1   4.5   4.5   0.7423 0.4091189
D        1   6.1   6.1   1.0103 0.3385304
A:D      1   1.1   1.1   0.1856 0.6757693
B:D      1   5.1   5.1   0.8351 0.3823203
A:B:D    1   0.5   0.5   0.0825 0.7798349
C:D      1   1.6   1.6   0.2577 0.6226914
A:C:D    1  10.1  10.1  1.6701 0.2253083
B:C:D    1  72.0  72.0  11.8763 0.0062660 **

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq  F value    Pr(>F)
rep      1 5940.5 5940.5 979.8763 2.6e-11 ***
rep:block 6  406.9   67.8 11.1856 0.0006129 ***  

A        1  171.1  171.1 28.2268 0.0003412 ***  

B        1   18.0   18.0  2.9691 0.1155937  

A:B     1   1.6   1.6   0.2577 0.6226914  

C        1  120.1  120.1 19.8144 0.0012326 **  

A:C     1   0.6   0.6   0.0928 0.7669127  

B:C     1   2.0   2.0   0.3299 0.5784103  

A:B:C   1   4.5   4.5   0.7423 0.4091189  

D        1   6.1   6.1   1.0103 0.3385304  

A:D      1   1.1   1.1   0.1856 0.6757693  

B:D      1   5.1   5.1   0.8351 0.3823203  

A:B:D   1   0.5   0.5   0.0825 0.7798349  

C:D      1   1.6   1.6   0.2577 0.6226914  

A:C:D   1  10.1  10.1  1.6701 0.2253083  

B:C:D   1  72.0  72.0  11.8763 0.0062660 **

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df Sum Sq Mean Sq  F value    Pr(>F)
rep      1 5940.5 5940.5 979.8763 2.6e-11 ***
rep:block 6  406.9   67.8 11.1856 0.0006129 ***  

A        1  171.1  171.1 28.2268 0.0003412 ***  

B        1   18.0   18.0  2.9691 0.1155937  

A:B     1   1.6   1.6   0.2577 0.6226914  

C        1  120.1  120.1 19.8144 0.0012326 **  

A:C     1   0.6   0.6   0.0928 0.7669127  

B:C     1   2.0   2.0   0.3299 0.5784103  

A:B:C   1   4.5   4.5   0.7423 0.4091189  

D        1   6.1   6.1   1.0103 0.3385304  

A:D      1   1.1   1.1   0.1856 0.6757693  

B:D      1   5.1   5.1   0.8351 0.3823203  

A:B:D   1   0.5   0.5   0.0825 0.7798349

```

```

C:D      1    1.6     1.6   0.2577 0.6226914
A:C:D    1   10.1    10.1   1.6701 0.2253083
B:C:D    1   72.0    72.0  11.8763 0.0062660 **

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.7 Chapter 10

9.7.1 p388

(142) MODEL

```

v2p388 = read.table("C:/G/Rt/Kemp/v2p388.txt", head=TRUE)
v2p388 = af(v2p388, c("rep", "block", "A", "B"))
GLM(y ~ rep + A*B + rep:block, v2p388) # OK

```

\$ANOVA

```

Response : y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      11 1136.8 103.343 124.01 3.698e-06 ***
RESIDUALS   6    5.0   0.833
CORRECTED TOTAL 17 1141.8
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```

Root MSE    y Mean Coef Var R-square Adj R-sq
0.9128709 26.11111 3.496101 0.9956209 0.9875924

```

\$`Type I`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 410.89  410.89 493.0667 5.455e-07 ***
A        2 228.11  114.06 136.8667 9.868e-06 ***
B        2   3.44    1.72   2.0667  0.207585
A:B      4 464.22  116.06 139.2667 4.801e-06 ***
rep:block 2  30.11   15.06 18.0667  0.002888 **
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
rep      1 410.89  410.89 493.0667 5.455e-07 ***
A        2 228.11  114.06 136.8667 9.868e-06 ***
B        2   3.44    1.72   2.0667  0.207585
A:B      2  18.78    9.39  11.2667  0.009298 **
rep:block 2  30.11   15.06 18.0667  0.002888 **
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  
CAUTION: Singularity Exists !  

      Df Sum Sq Mean Sq F value Pr(>F)  
rep      1 410.89 410.89 493.0667 5.455e-07 ***  
A         2 228.11 114.06 136.8667 9.868e-06 ***  
B         2   3.44    1.72    2.0667  0.207585  
A:B       2  18.78    9.39   11.2667  0.009298 **  
rep:block 2  30.11   15.06   18.0667  0.002888 **  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.8 Chapter 14

9.8.1 p570

(143) MODEL

```
v2p570 = read.table("C:/G/Rt/Kemp/v2p570.txt", head=TRUE)
v2p570 = af(v2p570, c("A", "B", "C", "D"))
GLM(Y ~ A + B + C + D + A:B + A:C + A:D + B:C + B:D + C:D, v2p570) # OK
```

```
$ANOVA
Response : Y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL     8 22.222  2.7778
RESIDUALS 0  0.000
CORRECTED TOTAL 8 22.222
```

```
$Fitness
Root MSE  Y Mean Coef Var R-square
      NA 6.555556      NA        1
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  
A      2  2.8889  1.4444  
B      2  2.8889  1.4444  
C      2  1.5556  0.7778  
D      2 14.8889  7.4444  
A:B    0  
A:C    0  
A:D    0  
B:C    0  
B:D    0  
C:D    0
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  
A      0  
B      0
```

```

C      0
D      0
A:B    0
A:C    0
A:D    0
B:C    0
B:D    0
C:D    0

$`Type III`  

CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value Pr(>F)
A      0
B      0
C      0
D      0
A:B    0
A:C    0
A:D    0
B:C    0
B:D    0
C:D    0

```

9.8.2 p578

(144) MODEL

```

v2p578 = read.table("C:/G/Rt/Kemp/v2p578.txt", head=TRUE)
v2p578 = af(v2p578, 1:11)
GLM(Y ~ A + B + C + D + E + F + G + H + J + K + L, v2p578) # OK

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL       11     575   52.273
RESIDUALS    0      0
CORRECTED TOTAL 11     575

```

```

$Fitness
Root MSE Y Mean Coef Var R-square
NA     25.5      NA        1

```

```

$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)
A 1 3.000 3.000
B 1 27.000 27.000
C 1 12.000 12.000
D 1 16.333 16.333
E 1 176.333 176.333

```

```

F 1 133.333 133.333
G 1 1.333 1.333
H 1 21.333 21.333
J 1 108.000 108.000
K 1 1.333 1.333
L 1 75.000 75.000

```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	3.000	3.000		
B	1	27.000	27.000		
C	1	12.000	12.000		
D	1	16.333	16.333		
E	1	176.333	176.333		
F	1	133.333	133.333		
G	1	1.333	1.333		
H	1	21.333	21.333		
J	1	108.000	108.000		
K	1	1.333	1.333		
L	1	75.000	75.000		

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	3.000	3.000		
B	1	27.000	27.000		
C	1	12.000	12.000		
D	1	16.333	16.333		
E	1	176.333	176.333		
F	1	133.333	133.333		
G	1	1.333	1.333		
H	1	21.333	21.333		
J	1	108.000	108.000		
K	1	1.333	1.333		
L	1	75.000	75.000		

(145) MODEL

```
GLM(Y ~ E*F + E*J + F*J + E*L + F*L + J*L, v2p578) # OK
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	10	574.5	57.45	114.9	0.07249 .
RESIDUALS	1	0.5	0.50		
CORRECTED TOTAL	11	575.0			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root	MSE	Y	Mean	Coef	Var	R-square	Adj R-sq
0.7071068	25.5	2.772968	0.9991304	0.9904348			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
E	1	176.333	176.333	352.6667	0.03387 *						
F	1	133.333	133.333	266.6667	0.03894 *						
E:F	1	65.333	65.333	130.6667	0.05555 .						
J	1	66.667	66.667	133.3333	0.05500 .						
E:J	1	2.667	2.667	5.3333	0.26015						
F:J	1	112.667	112.667	225.3333	0.04235 *						
L	1	10.800	10.800	21.6000	0.13492						
E:L	1	5.486	5.486	10.9714	0.18666						
F:L	1	0.176	0.176	0.3516	0.65925						
J:L	1	1.038	1.038	2.0769	0.38618						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'..'	0.1	' '	1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
E	1	61.633	61.633	123.2667	0.05719 .						
F	1	75.208	75.208	150.4167	0.05179 .						
E:F	1	9.346	9.346	18.6923	0.14470						
J	1	54.675	54.675	109.3500	0.06069 .						
E:J	1	0.115	0.115	0.2308	0.71490						
F:J	1	72.115	72.115	144.2308	0.05289 .						
L	1	10.800	10.800	21.6000	0.13492						
E:L	1	5.654	5.654	11.3077	0.18402						
F:L	1	0.115	0.115	0.2308	0.71490						
J:L	1	1.038	1.038	2.0769	0.38618						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'..'	0.1	' '	1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
E	1	61.038	61.038	122.0769	0.05746 .						
F	1	61.038	61.038	122.0769	0.05746 .						
E:F	1	9.346	9.346	18.6923	0.14470						
J	1	61.038	61.038	122.0769	0.05746 .						
E:J	1	0.115	0.115	0.2308	0.71490						
F:J	1	72.115	72.115	144.2308	0.05289 .						
L	1	9.346	9.346	18.6923	0.14470						
E:L	1	5.654	5.654	11.3077	0.18402						
F:L	1	0.115	0.115	0.2308	0.71490						
J:L	1	1.038	1.038	2.0769	0.38618						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'..'	0.1	' '	1

9.9 Chapter 16

9.9.1 p619

(146) MODEL

```
v2p619 = read.table("C:/G/Rt/Kemp/v2p619.txt", head=TRUE)
v2p619 = af(v2p619, c("A", "B", "C"))
GLM(y ~ A + B + C + A:B, v2p619) # OK
```

\$ANOVA

```
Response : y
             Df Sum Sq Mean Sq F value Pr(>F)
MODEL          4 31.429  7.8571
RESIDUALS      2  0.000  0.0000
CORRECTED TOTAL 6 31.429
```

\$Fitness

Root MSE	y	Mean Coef	Var	R-square	Adj R-sq
0	10.78571	0	1	1	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	13.7619	13.7619	Inf	< 2.2e-16 ***
B	1	1.6667	1.6667	Inf	< 2.2e-16 ***
C	1	10.0000	10.0000	Inf	< 2.2e-16 ***
A:B	1	6.0000	6.0000	Inf	< 2.2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	19.6	19.6	Inf	< 2.2e-16 ***
B	1	3.6	3.6	Inf	< 2.2e-16 ***
C	1	13.5	13.5	Inf	< 2.2e-16 ***
A:B	1	6.0	6.0	Inf	< 2.2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	24.0	24.0	Inf	< 2.2e-16 ***
B	1	6.0	6.0	Inf	< 2.2e-16 ***
C	1	13.5	13.5	Inf	< 2.2e-16 ***
A:B	1	6.0	6.0	Inf	< 2.2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(147) MODEL

```

GLM(y ~ A + B + C + A:C, v2p619) # OK

$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      4 26.0952  6.5238  2.4464 0.3106
RESIDUALS   2  5.3333  2.6667
CORRECTED TOTAL 6 31.4286

```

```

$Fitness
Root MSE    y Mean Coef Var R-square  Adj R-sq
1.632993 10.78571 15.14033 0.830303 0.4909091

```

```

$`Type I`
      Df  Sum Sq Mean Sq F value Pr(>F)
A      1 13.7619 13.7619  5.1607 0.1511
B      1  1.6667  1.6667  0.6250 0.5120
C      1 10.0000 10.0000  3.7500 0.1924
A:C    1  0.6667  0.6667  0.2500 0.6667

```

```

$`Type II`
      Df  Sum Sq Mean Sq F value Pr(>F)
A      1 19.6000 19.6000  7.35 0.1134
B      1  2.6667  2.6667  1.00 0.4226
C      1 10.0000 10.0000  3.75 0.1924
A:C    1  0.6667  0.6667  0.25 0.6667

```

```

$`Type III`
      Df  Sum Sq Mean Sq F value Pr(>F)
A      1 16.6667 16.6667  6.2500 0.1296
B      1  2.6667  2.6667  1.0000 0.4226
C      1  8.1667  8.1667  3.0625 0.2222
A:C    1  0.6667  0.6667  0.2500 0.6667

```

(148) MODEL

```

GLM(y ~ A + B + C + B:C, v2p619) # OK

```

```

$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL      4 26.0952  6.5238  2.4464 0.3106
RESIDUALS   2  5.3333  2.6667
CORRECTED TOTAL 6 31.4286

```

```

$Fitness
Root MSE    y Mean Coef Var R-square  Adj R-sq
1.632993 10.78571 15.14033 0.830303 0.4909091

```

```
$`Type I`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 13.7619 13.7619 5.1607 0.1511  
B 1 1.6667 1.6667 0.6250 0.5120  
C 1 10.0000 10.0000 3.7500 0.1924  
B:C 1 0.6667 0.6667 0.2500 0.6667
```

```
$`Type II`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 16.6667 16.6667 6.25 0.1296  
B 1 3.6000 3.6000 1.35 0.3652  
C 1 10.0000 10.0000 3.75 0.1924  
B:C 1 0.6667 0.6667 0.25 0.6667
```

```
$`Type III`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 16.6667 16.6667 6.2500 0.1296  
B 1 2.6667 2.6667 1.0000 0.4226  
C 1 8.1667 8.1667 3.0625 0.2222  
B:C 1 0.6667 0.6667 0.2500 0.6667
```

9.9.2 p626

(149) MODEL

```
v2p626 = read.table("C:/G/Rt/Kemp/v2p626.txt", head=TRUE)  
v2p626 = af(v2p626, c("A", "B", "C"))  
GLM(y ~ A + B + C + A:B, v2p626) # OK
```

```
$ANOVA  
Response : y  
Df Sum Sq Mean Sq F value Pr(>F)  
MODEL 4 42.092 10.5231 22.002 0.04395 *  
RESIDUALS 2 0.957 0.4783  
CORRECTED TOTAL 6 43.049  
---  
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  
Root MSE y Mean Coef Var R-square Adj R-sq  
0.6915708 11.12243 6.217804 0.9777801 0.9333402
```

```
$`Type I`  
Df Sum Sq Mean Sq F value Pr(>F)  
A 1 16.2088 16.2088 33.890 0.02826 *  
B 1 4.8150 4.8150 10.068 0.08662 .  
C 1 15.7339 15.7339 32.898 0.02908 *  
A:B 1 5.3346 5.3346 11.154 0.07916 .  
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	25.4131	25.4131	53.136	0.01830 *
B	1	8.6630	8.6630	18.113	0.05102 .
C	1	19.5193	19.5193	40.812	0.02364 *
A:B	1	5.3346	5.3346	11.154	0.07916 .

```
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	29.7950	29.7950	62.297	0.01568 *
B	1	11.7460	11.7460	24.559	0.03839 *
C	1	19.5193	19.5193	40.812	0.02364 *
A:B	1	5.3346	5.3346	11.154	0.07916 .

```
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
(150) MODEL
```

```
GLM(y ~ A + B + C + A:C, v2p626) # OK
```

```
$ANOVA
```

```
Response : y
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	4	39.229	9.8072	5.1346	0.1696
RESIDUALS	2	3.820	1.9100		
CORRECTED TOTAL	6	43.049			

```
$Fitness
```

Root MSE	y	Mean Coef	Var	R-square	Adj R-sq
1.382033	11.12243	12.42564	0.9112627	0.733788	

```
$`Type I`
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	16.2088	16.2088	8.4862	0.1004
B	1	4.8150	4.8150	2.5209	0.2533
C	1	15.7339	15.7339	8.2376	0.1030
A:C	1	2.4711	2.4711	1.2937	0.3733

```
$`Type II`
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	25.4131	25.4131	13.3052	0.06762 .
B	1	6.0361	6.0361	3.1602	0.21743
C	1	15.7339	15.7339	8.2376	0.10298
A:C	1	2.4711	2.4711	1.2937	0.37327

```
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A   1 20.1428 20.1428 10.5459 0.08317 .  

B   1  6.0361  6.0361  3.1602 0.21743  

C   1 11.8863 11.8863  6.2232 0.13007  

A:C 1  2.4711  2.4711  1.2937 0.37327  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(151) MODEL  

GLM(y ~ A + B + C + B:C, v2p626) # OK

$ANOVA  

Response : y  

  Df Sum Sq Mean Sq F value Pr(>F)  

MODEL          4 37.340  9.3349  3.2701 0.2477  

RESIDUALS      2  5.709  2.8546  

CORRECTED TOTAL 6 43.049

$Fitness  

  Root MSE    y Mean Coef Var R-square Adj R-sq  

  1.689558 11.12243 15.19055 0.8673781 0.6021342

$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A   1 16.2088 16.2088  5.6781 0.1400  

B   1  4.8150  4.8150  1.6867 0.3236  

C   1 15.7339 15.7339  5.5118 0.1434  

B:C 1  0.5819  0.5819  0.2038 0.6959

$`Type II`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A   1 21.9995 21.9995  7.7067 0.1090  

B   1  8.6630  8.6630  3.0347 0.2236  

C   1 15.7339 15.7339  5.5118 0.1434  

B:C 1  0.5819  0.5819  0.2038 0.6959

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A   1 21.9995 21.9995  7.7067 0.1090  

B   1  7.0709  7.0709  2.4770 0.2562  

C   1 13.3221 13.3221  4.6669 0.1633  

B:C 1  0.5819  0.5819  0.2038 0.6959

```

9.10 Chapter 17

9.10.1 p642

(152) MODEL

```
v2p642 = read.table("C:/G/Rt/Kemp/v2p642.txt", head=TRUE)
v2p642 = af(v2p642, 2:11)
GLM(Y ~ A + B + C + D + E + F + G, v2p642) # OK
```

\$ANOVA

Response : Y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	7	11.0	1.57143	1.6688	0.1646
RESIDUALS	24	22.6	0.94167		
CORRECTED TOTAL	31	33.6			

\$Fitness

Root	MSE	Y	Mean	Coef	Var	R-square	Adj	R-sq
0.9703951	2.25	43.12867	0.327381	0.1312004				

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	5.7800	5.7800	6.1381	0.02066 *
B	1	0.1800	0.1800	0.1912	0.66587
C	1	0.1250	0.1250	0.1327	0.71879
D	1	2.5312	2.5312	2.6881	0.11415
E	1	0.6613	0.6613	0.7022	0.41031
F	1	0.0112	0.0112	0.0119	0.91387
G	1	1.7113	1.7113	1.8173	0.19023

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	5.7800	5.7800	6.1381	0.02066 *
B	1	0.1800	0.1800	0.1912	0.66587
C	1	0.1250	0.1250	0.1327	0.71879
D	1	2.5312	2.5312	2.6881	0.11415
E	1	0.6613	0.6613	0.7022	0.41031
F	1	0.0112	0.0112	0.0119	0.91387
G	1	1.7113	1.7113	1.8173	0.19023

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	5.7800	5.7800	6.1381	0.02066 *
B	1	0.1800	0.1800	0.1912	0.66587

```

C 1 0.1250 0.1250 0.1327 0.71879
D 1 2.5312 2.5312 2.6881 0.11415
E 1 0.6613 0.6613 0.7022 0.41031
F 1 0.0112 0.0112 0.0119 0.91387
G 1 1.7113 1.7113 1.8173 0.19023
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(153) MODEL

GLM(log(S) ~ A + B + C + D + E + F + G, v2p642) # OK

$ANOVA
Response : log(S)
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7 266.43 38.062
RESIDUALS      24   0.00   0.000
CORRECTED TOTAL 31 266.43

$Fitness
Root MSE log(S) Mean Coef Var R-square Adj R-sq
          0    -2.233358        0       1       1

$`Type I`
            Df Sum Sq Mean Sq F value     Pr(>F)
A 1 1.511 1.511 Inf < 2.2e-16 ***
B 1 0.600 0.600 Inf < 2.2e-16 ***
C 1 0.284 0.284 Inf < 2.2e-16 ***
D 1 0.384 0.384 Inf < 2.2e-16 ***
E 1 0.741 0.741 Inf < 2.2e-16 ***
F 1 261.783 261.783 Inf < 2.2e-16 ***
G 1 1.127 1.127 Inf < 2.2e-16 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
            Df Sum Sq Mean Sq F value     Pr(>F)
A 1 1.511 1.511 Inf < 2.2e-16 ***
B 1 0.600 0.600 Inf < 2.2e-16 ***
C 1 0.284 0.284 Inf < 2.2e-16 ***
D 1 0.384 0.384 Inf < 2.2e-16 ***
E 1 0.741 0.741 Inf < 2.2e-16 ***
F 1 261.783 261.783 Inf < 2.2e-16 ***
G 1 1.127 1.127 Inf < 2.2e-16 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
            Df Sum Sq Mean Sq F value     Pr(>F)

```

```

A 1 1.511 1.511      Inf < 2.2e-16 ***
B 1 0.600 0.600      Inf < 2.2e-16 ***
C 1 0.284 0.284      Inf < 2.2e-16 ***
D 1 0.384 0.384      Inf < 2.2e-16 ***
E 1 0.741 0.741      Inf < 2.2e-16 ***
F 1 261.783 261.783   Inf < 2.2e-16 ***
G 1 1.127 1.127      Inf < 2.2e-16 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.11 Chapter 19

9.11.1 p700

(154) MODEL

```

v2p700 = read.table("C:/G/Rt/Kemp/v2p700.txt", head=TRUE)
v2p700 = af(v2p700, 2:5)
GLM(Y ~ P + S + T + C, v2p700) # OK

```

\$ANOVA

```

Response : Y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       12 378.80 31.5670  57.256 0.003319 **
RESIDUALS    3   1.65  0.5513
CORRECTED TOTAL 15 380.46
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```

Root MSE  Y Mean Coef Var R-square Adj R-sq
0.7425182 19.6375 3.781124 0.9956526 0.978263

```

\$`Type I`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
P 3 53.888 17.963 32.580 0.008646 **
S 3 154.508 51.503 93.414 0.001845 **
T 3 149.848 49.949 90.597 0.001930 **
C 3 20.561  6.854 12.431 0.033708 *
---
```

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```

          Df Sum Sq Mean Sq F value    Pr(>F)
P 2 2.220  1.110 2.0133 0.278974
S 3 111.966 37.322 67.6941 0.002969 **
T 3 161.828 53.943 97.8403 0.001722 **
C 3 20.561  6.854 12.4311 0.033708 *
---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
  Df Sum Sq Mean Sq F value    Pr(>F)
P   2   2.220   1.110   2.0133 0.278974
S   3 111.966   37.322  67.6941 0.002969 **
T   3 161.828   53.943  97.8403 0.001722 **
C   3   20.561   6.854  12.4311 0.033708 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

9.11.2 p703

(155) MODEL

```

v2p703 = read.table("C:/G/Rt/Kemp/v2p703.txt", head=TRUE)
v2p703$C = ifelse(v2p703$C == 0, 4, v2p703$C)
v2p703 = af(v2p703, 2:5)
GLM(Y ~ P + S + T + C, v2p703) # OK

```

```

$ANOVA
Response : Y
  Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          13 385.18 29.6293  21.766 0.0005673 ***
RESIDUALS       6   8.17  1.3613
CORRECTED TOTAL 19 393.35
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
  Root MSE Y Mean Coef Var R-square Adj R-sq
  1.166726 19.46 5.99551 0.9792359 0.9342472

```

```

$`Type I`
  Df Sum Sq Mean Sq F value    Pr(>F)
P   4   56.408  14.102 10.3596 0.0073255 **
S   3  119.260   39.753 29.2036 0.0005620 ***
T   3  190.430   63.477 46.6312 0.0001498 ***
C   3   19.083   6.361  4.6728 0.0518237 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`
  Df Sum Sq Mean Sq F value    Pr(>F)
P   4   52.288  13.072  9.6028 0.0088641 **
S   3  167.414   55.805 40.9952 0.0002163 ***
T   3  190.430   63.477 46.6312 0.0001498 ***
C   3   19.083   6.361  4.6728 0.0518237 .

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

Df	Sum Sq	Mean Sq	F value	Pr(>F)
P	4	52.288	13.072	9.6028 0.0088641 **
S	3	167.414	55.805	40.9952 0.0002163 ***
T	3	190.430	63.477	46.6312 0.0001498 ***
C	3	19.083	6.361	4.6728 0.0518237 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10 Lawson - DAE with SAS

Reference

- Lawson J. Design and Analysis of Experiments with SAS. Taylor and Francis Group. 2010.

```
require(daewr)
```

10.1 Chapter 2

10.1.1 p22

(156) MODEL

```
GLM(height ~ time, bread) # OK
```

\$ANOVA

Response : height

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	2	21.573	10.7865	4.6022	0.042 *
RESIDUALS	9	21.094	2.3438		
CORRECTED TOTAL	11	42.667			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	MSE	height	Mean	Coef	Var	R-square	Adj R-sq
1.530931	7.333333	20.87633	0.5056152	0.395752			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
time	2	21.573	10.787	4.6022	0.042 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
time	2	21.573	10.787	4.6022	0.042 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
time	2	21.573	10.787	4.6022	0.042 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10.1.2 p32

(157) MODEL

```

GLM(height^(1 - 1.294869) ~ time, bread) # OK

$ANOVA
Response : height^(1 - 1.294869)
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL    2 0.0130560 0.0065280 5.9356 0.02271 *
RESIDUALS 9 0.0098983 0.0010998
CORRECTED TOTAL 11 0.0229544
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE height^(1 - 1.294869) Mean Coef Var R-square Adj R-sq
0.03316344           0.5629811 5.890685 0.5687825 0.4729564

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
time  2 0.013056 0.006528 5.9356 0.02271 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
time  2 0.013056 0.006528 5.9356 0.02271 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
time  2 0.013056 0.006528 5.9356 0.02271 *
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.1.3 p42

(158) MODEL

```
GLM(yield ~ treat, sugarbeet) # OK
```

```

$ANOVA
Response : yield
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL    3 291.00 97.002   45.9 1.718e-07 ***
RESIDUALS 14 29.59  2.113
CORRECTED TOTAL 17 320.59
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE yield Mean Coef Var R-square Adj R-sq
1.453727 45.68333 3.182182 0.9077128 0.8879369

$`Type I` 
Df Sum Sq Mean Sq F value    Pr(>F)
treat  3     291   97.002    45.9 1.718e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
Df Sum Sq Mean Sq F value    Pr(>F)
treat  3     291   97.002    45.9 1.718e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
Df Sum Sq Mean Sq F value    Pr(>F)
treat  3     291   97.002    45.9 1.718e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2 Chapter 3

10.2.1 p63

(159) MODEL

```
GLM(CO ~ Eth + Ratio + Eth:Ratio, C0data) # OK
```

```

$ANOVA
Response : CO
Df Sum Sq Mean Sq F value    Pr(>F)
MODEL          8 1654.0 206.750  40.016 3.861e-06 ***
RESIDUALS      9    46.5    5.167
CORRECTED TOTAL 17 1700.5
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE CO Mean Coef Var R-square Adj R-sq
2.27303 72.83333 3.120865 0.9726551 0.9483485

```

```

$`Type I` 
Df Sum Sq Mean Sq F value    Pr(>F)
Eth        2     324   162.0  31.355 8.790e-05 ***
Ratio      2     652   326.0  63.097 5.067e-06 ***
Eth:Ratio  4     678   169.5  32.806 2.240e-05 ***
---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth       2     324    162.0  31.355 8.790e-05 ***  

Ratio     2     652    326.0  63.097 5.067e-06 ***  

Eth:Ratio 4     678    169.5  32.806 2.240e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth       2     324    162.0  31.355 8.790e-05 ***  

Ratio     2     652    326.0  63.097 5.067e-06 ***  

Eth:Ratio 4     678    169.5  32.806 2.240e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(160) MODEL  

GLM(CO ~ Ratio + Eth + Ratio:Eth, C0data) # OK

$ANOVA  

Response : CO  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL      8 1654.0 206.750  40.016 3.861e-06 ***  

RESIDUALS   9    46.5    5.167  

CORRECTED TOTAL 17 1700.5  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness  

Root MSE  CO Mean Coef Var  R-square  Adj R-sq  

  2.27303 72.83333 3.120865 0.9726551 0.9483485

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Ratio     2     652    326.0  63.097 5.067e-06 ***  

Eth       2     324    162.0  31.355 8.790e-05 ***  

Ratio:Eth 4     678    169.5  32.806 2.240e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Ratio     2     652    326.0  63.097 5.067e-06 ***  

Eth       2     324    162.0  31.355 8.790e-05 ***  

Ratio:Eth 4     678    169.5  32.806 2.240e-05 ***  

---
```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Ratio      2     652    326.0  63.097 5.067e-06 ***  

Eth        2     324    162.0  31.355 8.790e-05 ***  

Ratio:Eth  4     678    169.5  32.806 2.240e-05 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2.2 p74

(161) MODEL

```
GLM(CO ~ Eth + Ratio + Eth:Ratio, C0data[-18,]) # OK
```

```

$ANOVA  

Response : CO  

      Df Sum Sq Mean Sq F value    Pr(>F)  

MODEL       8 1423.0 177.879  31.978 2.749e-05 ***  

RESIDUALS   8    44.5    5.563  

CORRECTED TOTAL 16 1467.5  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

Root MSE  CO Mean Coef Var  R-square  Adj R-sq  

2.358495 73.70588 3.199874 0.9696769 0.9393539

```

```

$`Type I`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth        2 472.66 236.33  42.486 5.482e-05 ***  

Ratio      2 395.33 197.66  35.535 0.0001048 ***  

Eth:Ratio  4 555.04 138.76  24.945 0.0001427 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth        2 398.26 199.13  35.799 0.0001020 ***  

Ratio      2 395.33 197.66  35.535 0.0001048 ***  

Eth:Ratio  4 555.04 138.76  24.945 0.0001427 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

Eth        2 319.45 159.73  28.715 0.0002235 ***  

Ratio      2 511.45 255.73  45.973 4.105e-05 ***

```

```

Eth:Ratio 4 555.04 138.76 24.945 0.0001427 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.2.3 p91

(162) MODEL

```

volt$XA = (as.numeric(as.character(volt$A)) - 27)/5
volt$XB = (as.numeric(as.character(volt$B)) - 2.75)/2.25
volt$XC = (as.numeric(as.character(volt$C)) - 2.75)/2.25
GLM(y ~ XA + XB + XC + XA:XB + XA:XC + XB:XC + XA:XB:XC, volt) # OK

```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	7	8843.4	1263.35	3.8686	0.0385 *
RESIDUALS	8	2612.5	326.56		
CORRECTED TOTAL	15	11455.9			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	MSE	y	Mean	Coef	Var	R-square	Adj R-sq
18.07104	668.5625	2.702969	0.7719523	0.5724106			

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
XA	1	4522.6	4522.6	13.8490	0.005859 **
XB	1	14.1	14.1	0.0431	0.840793
XC	1	473.1	473.1	1.4486	0.263154
XA:XB	1	715.6	715.6	2.1912	0.177071
XA:XC	1	2525.1	2525.1	7.7322	0.023899 *
XB:XC	1	52.6	52.6	0.1610	0.698780
XA:XB:XC	1	540.6	540.6	1.6553	0.234218

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
XA	1	4522.6	4522.6	13.8490	0.005859 **
XB	1	14.1	14.1	0.0431	0.840793
XC	1	473.1	473.1	1.4486	0.263154
XA:XB	1	715.6	715.6	2.1912	0.177071
XA:XC	1	2525.1	2525.1	7.7322	0.023899 *
XB:XC	1	52.6	52.6	0.1610	0.698780
XA:XB:XC	1	540.6	540.6	1.6553	0.234218

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
$`Type III`  

      Df Sum Sq Mean Sq F value    Pr(>F)  

XA        1 4522.6  4522.6 13.8490 0.005859 **  

XB        1   14.1    14.1  0.0431 0.840793  

XC        1   473.1    473.1  1.4486 0.263154  

XA:XB     1   715.6    715.6  2.1912 0.177071  

XA:XC     1 2525.1  2525.1  7.7322 0.023899 *  

XB:XC     1    52.6    52.6  0.1610 0.698780  

XA:XB:XC  1   540.6    540.6  1.6553 0.234218  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.2.4 p97

(163) MODEL

```
chem2 = af(chem, c("A","B","C","D"))  

GLM(y ~ A*B*C*D, chem2) # OK
```

```
$ANOVA  

Response : y  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL       15 6369.4  424.63  

RESIDUALS    0    0.0  

CORRECTED TOTAL 15 6369.4
```

```
$Fitness  

Root MSE  y Mean Coef Var R-square  

      NA 62.3125      NA      1
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

A        1   637.6    637.6  

B        1 5076.6  5076.6  

A:B      1   451.6    451.6  

C        1    0.6     0.6  

A:C      1   10.6    10.6  

B:C      1    1.6     1.6  

A:B:C    1    0.6     0.6  

D        1    7.6     7.6  

A:D      1   68.1    68.1  

B:D      1    0.1     0.1  

A:B:D    1    7.6     7.6  

C:D      1    7.6     7.6  

A:C:D    1   95.1    95.1  

B:C:D    1    3.1     3.1  

A:B:C:D 1    1.6     1.6
```

```
$`Type II`  
          Df Sum Sq Mean Sq F value Pr(>F)  
A           1   637.6   637.6  
B           1 5076.6 5076.6  
A:B         1   451.6   451.6  
C           1     0.6     0.6  
A:C         1    10.6    10.6  
B:C         1     1.6     1.6  
A:B:C       1     0.6     0.6  
D           1     7.6     7.6  
A:D         1   68.1    68.1  
B:D         1     0.1     0.1  
A:B:D       1     7.6     7.6  
C:D         1     7.6     7.6  
A:C:D       1   95.1    95.1  
B:C:D       1     3.1     3.1  
A:B:C:D    1     1.6     1.6
```

```
$`Type III`  
          Df Sum Sq Mean Sq F value Pr(>F)  
A           1   637.6   637.6  
B           1 5076.6 5076.6  
A:B         1   451.6   451.6  
C           1     0.6     0.6  
A:C         1    10.6    10.6  
B:C         1     1.6     1.6  
A:B:C       1     0.6     0.6  
D           1     7.6     7.6  
A:D         1   68.1    68.1  
B:D         1     0.1     0.1  
A:B:D       1     7.6     7.6  
C:D         1     7.6     7.6  
A:C:D       1   95.1    95.1  
B:C:D       1     3.1     3.1  
A:B:C:D    1     1.6     1.6
```

10.2.5 p104

(164) MODEL

```
GLM(y ~ A*B*C*D, BoxM) # OK
```

```
$ANOVA  
Response : y  
          Df Sum Sq Mean Sq F value Pr(>F)  
MODEL      15   207.1   13.807  
RESIDUALS   0     0.0  
CORRECTED TOTAL 15   207.1
```

```
$Fitness
Root MSE y Mean Coef Var R-square
      NA 48.245       NA        1
```

```
$`Type I` 
      Df Sum Sq Mean Sq F value Pr(>F)
A       1   2.560   2.560
B       1 71.234  71.234
A:B     1   3.312   3.312
C       1 55.056  55.056
A:C     1 24.800  24.800
B:C     1   2.560   2.560
A:B:C   1   5.760   5.760
D       1   4.080   4.080
A:D     1   1.346   1.346
B:D     1   5.570   5.570
A:B:D   1   2.074   2.074
C:D     1   8.880   8.880
A:C:D   1   0.640   0.640
B:C:D   1   9.986   9.986
A:B:C:D 1   9.242   9.242
```

```
$`Type II` 
      Df Sum Sq Mean Sq F value Pr(>F)
A       1   2.560   2.560
B       1 71.234  71.234
A:B     1   3.312   3.312
C       1 55.056  55.056
A:C     1 24.800  24.800
B:C     1   2.560   2.560
A:B:C   1   5.760   5.760
D       1   4.080   4.080
A:D     1   1.346   1.346
B:D     1   5.570   5.570
A:B:D   1   2.074   2.074
C:D     1   8.880   8.880
A:C:D   1   0.640   0.640
B:C:D   1   9.986   9.986
A:B:C:D 1   9.242   9.242
```

```
$`Type III` 
      Df Sum Sq Mean Sq F value Pr(>F)
A       1   2.560   2.560
B       1 71.234  71.234
A:B     1   3.312   3.312
C       1 55.056  55.056
A:C     1 24.800  24.800
B:C     1   2.560   2.560
```

A:B:C	1	5.760	5.760
D	1	4.080	4.080
A:D	1	1.346	1.346
B:D	1	5.570	5.570
A:B:D	1	2.074	2.074
C:D	1	8.880	8.880
A:C:D	1	0.640	0.640
B:C:D	1	9.986	9.986
A:B:C:D	1	9.242	9.242

10.3 Chapter 4

10.3.1 p122

(165) MODEL

```
GLM(rate ~ rat + dose, drug) # OK
```

\$ANOVA

Response : rate

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	13	2.12867	0.163744	19.613	1.59e-12 ***
RESIDUALS	36	0.30055	0.008349		
CORRECTED TOTAL	49	2.42922			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	rate	Mean Coef	Var	R-square	Adj R-sq
0.09137104	0.9142	9.994644	0.8762762	0.8315982	

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rat	9	1.66846	0.18538	22.205	3.749e-12 ***
dose	4	0.46021	0.11505	13.781	6.535e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rat	9	1.66846	0.18538	22.205	3.749e-12 ***
dose	4	0.46021	0.11505	13.781	6.535e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
rat	9	1.66846	0.18538	22.205	3.749e-12 ***
dose	4	0.46021	0.11505	13.781	6.535e-07 ***

```
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.3.2 p127

(166) MODEL

```
GLM(y ~ block + treat + strain + treat:strain, bha) # OK
```

\$ANOVA

Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	8	543.22	67.902	26.203	0.0001507 ***
RESIDUALS	7	18.14	2.591		
CORRECTED TOTAL	15	561.36			

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

Root MSE	MSE	y	Mean Coef	Var	R-square	Adj R-sq
1.609791	12.9875	12.39493	0.9676855	0.9307546		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	47.61	47.61	18.3721	0.003627 **
treat	1	422.30	422.30	162.9613	4.194e-06 ***
strain	3	32.96	10.99	4.2399	0.052741 .
treat:strain	3	40.34	13.45	5.1892	0.033685 *

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	47.61	47.61	18.3721	0.003627 **
treat	1	422.30	422.30	162.9613	4.194e-06 ***
strain	3	32.96	10.99	4.2399	0.052741 .
treat:strain	3	40.34	13.45	5.1892	0.033685 *

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
block	1	47.61	47.61	18.3721	0.003627 **
treat	1	422.30	422.30	162.9613	4.194e-06 ***
strain	3	32.96	10.99	4.2399	0.052741 .
treat:strain	3	40.34	13.45	5.1892	0.033685 *

```
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.3.3 p129

(167) MODEL

```
GLM(cdistance ~ id + teehgt, rcb) # OK
```

\$ANOVA

Response : cdistance

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	10	126465	12646.5	161.72	< 2.2e-16 ***
RESIDUALS	124	9697	78.2		
CORRECTED TOTAL	134	136162			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	MSE	cdistance	Mean Coef	Var	R-square	Adj R-sq
8.8431	176.3778	5.013727	0.9287846	0.9230414		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
id	8	124741	15593	199.394	< 2.2e-16 ***
teehgt	2	1724	862	11.023	3.926e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
id	8	124741	15593	199.394	< 2.2e-16 ***
teehgt	2	1724	862	11.023	3.926e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
id	8	124741	15593	199.394	< 2.2e-16 ***
teehgt	2	1724	862	11.023	3.926e-05 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

10.3.4 p136

(168) MODEL

```
GLM(AUC ~ Subject + Period + Treat, bioequiv) # OK
```

\$ANOVA

Response : AUC

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	6	174461	29077	0.1315	0.9774

```

RESIDUALS      2 442158  221079
CORRECTED TOTAL 8 616618

$Fitness
Root MSE AUC Mean Coef Var R-square Adj R-sq
470.1902 1141.556 41.18855 0.2829314 -1.868274

```

```

$`Type I` 
  Df Sum Sq Mean Sq F value Pr(>F)
Subject  2 114264   57132  0.2584 0.7946
Period    2  45196   22598  0.1022 0.9073
Treat     2 15000    7500  0.0339 0.9672

```

```

$`Type II` 
  Df Sum Sq Mean Sq F value Pr(>F)
Subject  2 114264   57132  0.2584 0.7946
Period    2  45196   22598  0.1022 0.9073
Treat     2 15000    7500  0.0339 0.9672

```

```

$`Type III` 
  Df Sum Sq Mean Sq F value Pr(>F)
Subject  2 114264   57132  0.2584 0.7946
Period    2  45196   22598  0.1022 0.9073
Treat     2 15000    7500  0.0339 0.9672

```

10.4 Chapter 5

10.4.1 p152

(169) MODEL

```
GLM(conc ~ lab, Apo) # OK
```

```

$ANOVA
Response : conc
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      3 0.092233 0.0307444 42.107 4.009e-10 ***
RESIDUALS  26 0.018984 0.0007302
CORRECTED TOTAL 29 0.111217
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE conc Mean Coef Var R-square Adj R-sq
0.02702142 1.141567 2.367047 0.8293064 0.809611

```

```

$`Type I` 
  Df Sum Sq Mean Sq F value Pr(>F)
lab  3 0.092233 0.030744 42.107 4.009e-10 ***

```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
  Df  Sum Sq  Mean Sq F value    Pr(>F)
lab  3 0.092233 0.030744  42.107 4.009e-10 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
  Df  Sum Sq  Mean Sq F value    Pr(>F)
lab  3 0.092233 0.030744  42.107 4.009e-10 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.4.2 p181

(170) MODEL

```
GLM(residue ~ form + tech + form:tech + plot:form:tech, pesticide) # OK
```

```
$ANOVA
Response : residue
  Df  Sum Sq  Mean Sq F value    Pr(>F)
MODEL      7 0.036857 0.0052653 11.804 0.001187 **
RESIDUALS  8 0.003569 0.0004461
CORRECTED TOTAL 15 0.040426
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
  Root MSE residue Mean Coef Var  R-square Adj R-sq
0.02112019    0.3165625 6.671729 0.9117275 0.834489
```

```
$`Type I`
  Df  Sum Sq  Mean Sq F value    Pr(>F)
form       1 0.000018 0.000018  0.0405   0.84554
tech       1 0.032310 0.032310 72.4339 2.789e-05 ***
form:tech  1 0.002186 0.002186  4.8997   0.05776 .
form:tech:plot 4 0.002344 0.000586  1.3136   0.34317
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
  Df  Sum Sq  Mean Sq F value    Pr(>F)
form       1 0.000018 0.000018  0.0405   0.84554
tech       1 0.032310 0.032310 72.4339 2.789e-05 ***
form:tech  1 0.002186 0.002186  4.8997   0.05776 .
form:tech:plot 4 0.002344 0.000586  1.3136   0.34317
```

```

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

form       1 0.000018 0.000018 0.0405 0.84554  

tech       1 0.032310 0.032310 72.4339 2.789e-05 ***  

form:tech   1 0.002186 0.002186 4.8997 0.05776 .  

form:tech:plot 4 0.002344 0.000586 1.3136 0.34317  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.5 Chapter 7

10.5.1 p260

(171) MODEL

```
GLM(score ~ recipe + panelist, taste) # OK
```

```
$ANOVA  

Response : score  

      Df Sum Sq Mean Sq F value Pr(>F)  

MODEL      14 28.458 2.03274 2.661 0.0719 .  

RESIDUALS    9  6.875 0.76389  

CORRECTED TOTAL 23 35.333  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness  

  Root MSE score Mean Coef Var R-square Adj R-sq  

0.8740074 5.833333 14.98298 0.8054245 0.5027516
```

```
$`Type I`  

      Df Sum Sq Mean Sq F value Pr(>F)  

recipe     3 21.0000  7.000  9.1636 0.004246 **  

panelist  11  7.4583  0.678  0.8876 0.581099  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`  

      Df Sum Sq Mean Sq F value Pr(>F)  

recipe     3 9.1250 3.04167 3.9818 0.04649 *  

panelist  11 7.4583 0.67803 0.8876 0.58110  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)
```

```

recipe      3 9.1250 3.04167  3.9818 0.04649 *
panelist   11 7.4583 0.67803  0.8876 0.58110
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.5.2 p262

(172) MODEL

```
GLM(pressure ~ Block + Treatment, BPmonitor) # OK
```

```
$ANOVA
Response : pressure
          Df Sum Sq Mean Sq F value Pr(>F)
MODEL       8 321.00 40.125 4.4174 0.1245
RESIDUALS    3 27.25  9.083
CORRECTED TOTAL 11 348.25
```

```
$Fitness
Root MSE pressure Mean Coef Var R-square Adj R-sq
3.013857           77.75 3.876343 0.9217516 0.7130893
```

```
$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
Block       5 73.75 14.750 1.6239 0.36606
Treatment   3 247.25 82.417 9.0734 0.05149 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
Block       5 83.25 16.650 1.8330 0.32772
Treatment   3 247.25 82.417 9.0734 0.05149 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III`
          Df Sum Sq Mean Sq F value Pr(>F)
Block       5 83.25 16.650 1.8330 0.32772
Treatment   3 247.25 82.417 9.0734 0.05149 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.5.3 p276

(173) MODEL

```
GLM(weight ~ Blocks + A + B + C + D + E + F + G + H, Bff) # OK
```

```
$ANOVA
```

```

Response : weight
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL          15 158.37 10.558
RESIDUALS       0   0.00
CORRECTED TOTAL 15 158.37

$Fitness
Root MSE weight Mean Coef Var R-square
      NA     5.925625      NA        1

$`Type I`
      Df Sum Sq Mean Sq F value Pr(>F)
Blocks  7 30.567  4.367
A       1 21.879 21.879
B       1  8.338  8.338
C       1  6.213  6.213
D       1 12.870 12.870
E       1  0.098  0.098
F       1  1.260  1.260
G       1 71.868 71.868
H       1  5.279  5.279

$`Type II`
      Df Sum Sq Mean Sq F value Pr(>F)
Blocks  7 30.567  4.367
A       1 21.879 21.879
B       1  8.338  8.338
C       1  6.213  6.213
D       1 12.870 12.870
E       1  0.098  0.098
F       1  1.260  1.260
G       1 71.868 71.868
H       1  5.279  5.279

$`Type III`
      Df Sum Sq Mean Sq F value Pr(>F)
Blocks  7 30.567  4.367
A       1 21.879 21.879
B       1  8.338  8.338
C       1  6.213  6.213
D       1 12.870 12.870
E       1  0.098  0.098
F       1  1.260  1.260
G       1 71.868 71.868
H       1  5.279  5.279

```

10.6 Chapter 8

10.6.1 p315

(174) MODEL

```
GLM(ys ~ Block + A*B + Block:A:B + C*D + A:C + A:D + B:C + B:D + A:B:C + A:B:D +  
     A:C:D + B:C:D + A:B:C:D, sausage) # OK
```

\$ANOVA

Response : ys

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	19	0.064059	0.0033715	14.134	1.74e-05 ***
RESIDUALS	12	0.002862	0.0002385		
CORRECTED TOTAL	31	0.066922			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$Fitness

Root MSE	ys	Mean	Coef	Var	R-square	Adj R-sq
0.01544479	2.023438	0.7632948	0.9572262	0.8895011		

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Block	1	0.000903	0.000903	3.7860	0.075482 .
A	1	0.045753	0.045753	191.8035	9.647e-09 ***
B	1	0.002628	0.002628	11.0175	0.006119 **
A:B	1	0.001128	0.001128	4.7293	0.050371 .
Block:A:B	3	0.005484	0.001828	7.6638	0.004007 **
C	1	0.003828	0.003828	16.0480	0.001743 **
D	1	0.000528	0.000528	2.2140	0.162566
C:D	1	0.000253	0.000253	1.0611	0.323272
A:C	1	0.000153	0.000153	0.6419	0.438593
A:D	1	0.000903	0.000903	3.7860	0.075482 .
B:C	1	0.000078	0.000078	0.3275	0.577693
B:D	1	0.000253	0.000253	1.0611	0.323272
A:B:C	1	0.001378	0.001378	5.7773	0.033299 *
A:B:D	1	0.000703	0.000703	2.9476	0.111680
A:C:D	1	0.000028	0.000028	0.1179	0.737260
B:C:D	1	0.000028	0.000028	0.1179	0.737260
A:B:C:D	1	0.000028	0.000028	0.1179	0.737260

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
Block	1	0.000903	0.000903	3.7860	0.075482 .
A	1	0.045753	0.045753	191.8035	9.647e-09 ***
B	1	0.002628	0.002628	11.0175	0.006119 **

```

A:B      1 0.001128 0.001128   4.7293  0.050371 .
Block:A:B 3 0.005484 0.001828   7.6638  0.004007 **
C       1 0.003828 0.003828  16.0480  0.001743 **
D       1 0.000528 0.000528   2.2140  0.162566
C:D     1 0.000253 0.000253   1.0611  0.323272
A:C     1 0.000153 0.000153   0.6419  0.438593
A:D     1 0.000903 0.000903   3.7860  0.075482 .
B:C     1 0.000078 0.000078   0.3275  0.577693
B:D     1 0.000253 0.000253   1.0611  0.323272
A:B:C   1 0.001378 0.001378   5.7773  0.033299 *
A:B:D   1 0.000703 0.000703   2.9476  0.111680
A:C:D   1 0.000028 0.000028   0.1179  0.737260
B:C:D   1 0.000028 0.000028   0.1179  0.737260
A:B:C:D 1 0.000028 0.000028   0.1179  0.737260
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type III`  

      Df  Sum Sq  Mean Sq F value    Pr(>F)  

Block     1 0.000903 0.000903   3.7860  0.075482 .  

A        1 0.045753 0.045753 191.8035 9.647e-09 ***  

B        1 0.002628 0.002628  11.0175  0.006119 **  

A:B      1 0.001128 0.001128   4.7293  0.050371 .  

Block:A:B 3 0.005484 0.001828   7.6638  0.004007 **  

C       1 0.003828 0.003828  16.0480  0.001743 **  

D       1 0.000528 0.000528   2.2140  0.162566  

C:D     1 0.000253 0.000253   1.0611  0.323272  

A:C     1 0.000153 0.000153   0.6419  0.438593  

A:D     1 0.000903 0.000903   3.7860  0.075482 .  

B:C     1 0.000078 0.000078   0.3275  0.577693  

B:D     1 0.000253 0.000253   1.0611  0.323272  

A:B:C   1 0.001378 0.001378   5.7773  0.033299 *  

A:B:D   1 0.000703 0.000703   2.9476  0.111680  

A:C:D   1 0.000028 0.000028   0.1179  0.737260  

B:C:D   1 0.000028 0.000028   0.1179  0.737260  

A:B:C:D 1 0.000028 0.000028   0.1179  0.737260
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.6.2 p320

(175) MODEL

```
GLM(y ~ A*B*C*D+E, plasma) # OK
```

```
$ANOVA  

Response : y  

      Df  Sum Sq  Mean Sq F value Pr(>F)  

MODEL  31 6672.9   215.26
```

RESIDUALS 0 0.0
 CORRECTED TOTAL 31 6672.9

\$Fitness
 Root MSE y Mean Coef Var R-square
 NA 40.98125 NA 1

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	1118.65	1118.65		
B	1	142.81	142.81		
A:B	1	141.96	141.96		
C	1	91.80	91.80		
A:C	1	70.81	70.81		
B:C	1	5.78	5.78		
A:B:C	1	65.55	65.55		
D	1	1824.08	1824.08		
A:D	1	2194.53	2194.53		
B:D	1	87.78	87.78		
A:B:D	1	87.12	87.12		
C:D	1	22.45	22.45		
A:C:D	1	42.78	42.78		
B:C:D	1	12.25	12.25		
A:B:C:D	1	375.38	375.38		
E	1	78.75	78.75		
A:E	1	278.48	278.48		
B:E	1	0.72	0.72		
A:B:E	1	0.10	0.10		
C:E	1	0.15	0.15		
A:C:E	1	0.24	0.24		
B:C:E	1	6.48	6.48		
A:B:C:E	1	1.53	1.53		
D:E	1	8.40	8.40		
A:D:E	1	5.28	5.28		
B:D:E	1	0.28	0.28		
A:B:D:E	1	0.60	0.60		
C:D:E	1	0.85	0.85		
A:C:D:E	1	0.55	0.55		
B:C:D:E	1	6.30	6.30		
A:B:C:D:E	1	0.50	0.50		

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	1118.65	1118.65		
B	1	142.81	142.81		
A:B	1	141.96	141.96		
C	1	91.80	91.80		
A:C	1	70.81	70.81		

B:C	1	5.78	5.78
A:B:C	1	65.55	65.55
D	1	1824.08	1824.08
A:D	1	2194.53	2194.53
B:D	1	87.78	87.78
A:B:D	1	87.12	87.12
C:D	1	22.45	22.45
A:C:D	1	42.78	42.78
B:C:D	1	12.25	12.25
A:B:C:D	1	375.38	375.38
E	1	78.75	78.75
A:E	1	278.48	278.48
B:E	1	0.72	0.72
A:B:E	1	0.10	0.10
C:E	1	0.15	0.15
A:C:E	1	0.24	0.24
B:C:E	1	6.48	6.48
A:B:C:E	1	1.53	1.53
D:E	1	8.40	8.40
A:D:E	1	5.28	5.28
B:D:E	1	0.28	0.28
A:B:D:E	1	0.60	0.60
C:D:E	1	0.85	0.85
A:C:D:E	1	0.55	0.55
B:C:D:E	1	6.30	6.30
A:B:C:D:E	1	0.50	0.50

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	1118.64	1118.64		
B	1	142.80	142.80		
A:B	1	141.96	141.96		
C	1	91.80	91.80		
A:C	1	70.81	70.81		
B:C	1	5.78	5.78		
A:B:C	1	65.55	65.55		
D	1	1824.08	1824.08		
A:D	1	2194.53	2194.53		
B:D	1	87.78	87.78		
A:B:D	1	87.12	87.12		
C:D	1	22.45	22.45		
A:C:D	1	42.78	42.78		
B:C:D	1	12.25	12.25		
A:B:C:D	1	375.38	375.38		
E	1	78.75	78.75		
A:E	1	278.48	278.48		
B:E	1	0.72	0.72		
A:B:E	1	0.10	0.10		

C:E	1	0.15	0.15
A:C:E	1	0.24	0.24
B:C:E	1	6.48	6.48
A:B:C:E	1	1.53	1.53
D:E	1	8.40	8.40
A:D:E	1	5.28	5.28
B:D:E	1	0.28	0.28
A:B:D:E	1	0.60	0.60
C:D:E	1	0.85	0.85
A:C:D:E	1	0.55	0.55
B:C:D:E	1	6.30	6.30
A:B:C:D:E	1	0.50	0.50

10.6.3 p335

(176) MODEL

```

gear$A = as.numeric(as.character(gear$A))
gear$B = as.numeric(as.character(gear$B))
gear$C = as.numeric(as.character(gear$C))
gear$P = as.numeric(as.character(gear$P))
gear$Q = as.numeric(as.character(gear$Q))
REG(y ~ A*B*C + P + Q + A:P + A:Q + B:P + B:Q + C:P + C:Q, gear) # OK

```

Warning in qt(0.5 + conf.level/2, Res0[, "Df"]): NaNs produced

\$ANOVA
 Response : y

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	15	1104.6	73.641		
RESIDUALS	0	0.0			
CORRECTED TOTAL	15	1104.6			

\$Fitness

Root MSE	y	Mean	Coef	Var	R-square	PRESS	R2pred
NA	15.40625	NA	1	NaN	NaN		

\$Coefficients

	Estimate	Std. Error	Df	Lower CL	Upper CL	t value	Pr(> t)
(Intercept)	15.4062	0	0				
A	-4.9062	0	0				
B	-0.1562	0	0				
A:B	0.5312	0	0				
C	3.9688	0	0				
A:C	2.9062	0	0				
B:C	0.4062	0	0				
A:B:C	0.5938	0	0				
P	-2.3438	0	0				
Q	-3.4062	0	0				

A:P	-0.9062	0
A:Q	-0.3438	0
B:P	1.0938	0
B:Q	0.1562	0
C:P	-0.2812	0
C:Q	0.7812	0

10.7 Chapter 9

10.7.1 p349

(177) MODEL

```
GLM(pl ~ Subject + Period + Treat, antifungal) # OK
```

```
$ANOVA
Response : pl
      Df  Sum Sq Mean Sq F value Pr(>F)
MODEL       18 118.558  6.5866 1.4435 0.2388
RESIDUALS    15  68.444  4.5630
CORRECTED TOTAL 33 187.002
```

```
$Fitness
Root MSE  pl Mean Coef Var  R-square  Adj R-sq
2.136109 13.15882 16.23328 0.6339915 0.1947814
```

```
$`Type I`
      Df  Sum Sq Mean Sq F value Pr(>F)
Subject  16 114.642  7.1651  1.5703 0.1942
Period    1   0.922  0.9224  0.2021 0.6594
Treat     1   2.993  2.9932  0.6560 0.4306
```

```
$`Type II`
      Df  Sum Sq Mean Sq F value Pr(>F)
Subject  16 114.642  7.1651  1.5703 0.1942
Period    1   0.734  0.7344  0.1609 0.6939
Treat     1   2.993  2.9932  0.6560 0.4306
```

```
$`Type III`
      Df  Sum Sq Mean Sq F value Pr(>F)
Subject  16 114.642  7.1651  1.5703 0.1942
Period    1   0.734  0.7344  0.1609 0.6939
Treat     1   2.993  2.9932  0.6560 0.4306
```

10.7.2 p355

(178) MODEL

```
GLM(y ~ Group + Subject:Group + Period + Treat + Carry, bioequiv) # OK
```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL        39 417852 10714.1   20.367 < 2.2e-16 ***
RESIDUALS     68 35772   526.1
CORRECTED TOTAL 107 453624
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    y Mean Coef Var R-square Adj R-sq
22.93611 101.3846 22.62287 0.9211408 0.8759128

$`Type I` 
      Df Sum Sq Mean Sq F value    Pr(>F)
Group        1 43335  43335 82.3763  2.46e-13 ***
Group:Subject 34 370970  10911 20.7406 < 2.2e-16 ***
Period       2    287     143  0.2723   0.7624
Treat         1   2209    2209  4.1993   0.0443 *
Carry         1   1051    1051  1.9970   0.1622
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
      Df Sum Sq Mean Sq F value    Pr(>F)
Group        1 32616  32616 61.9998 3.712e-11 ***
Group:Subject 34 370970  10911 20.7406 < 2.2e-16 ***
Period       1    38     38  0.0724   0.7888
Treat         1   2209    2209  4.1993   0.0443 *
Carry         1   1051    1051  1.9970   0.1622
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value    Pr(>F)
Group        1 32616  32616 61.9998 3.712e-11 ***
Group:Subject 34 370970  10911 20.7406 < 2.2e-16 ***
Period       1    38     38  0.0724   0.7888
Treat         1   2209    2209  4.1993   0.0443 *
Carry         1   1051    1051  1.9970   0.1622
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
(179) MODEL
GLM(y ~ Subject + Period + Treat + Carry, bioequiv) # OK

```

\$ANOVA

```

Response : y
          Df Sum Sq Mean Sq F value    Pr(>F)
MODEL      39 417852 10714.1 20.367 < 2.2e-16 ***
RESIDUALS   68 35772   526.1
CORRECTED TOTAL 107 453624
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE    y Mean Coef Var R-square Adj R-sq
22.93611 101.3846 22.62287 0.9211408 0.8759128

$`Type I`
          Df Sum Sq Mean Sq F value Pr(>F)
Subject  35 414306 11837.3 22.5016 <2e-16 ***
Period    2     287    143.3  0.2723 0.7624
Treat     1    2209   2209.1  4.1993 0.0443 *
Carry     1    1051   1050.6  1.9970 0.1622
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
          Df Sum Sq Mean Sq F value Pr(>F)
Subject  35 403586 11531.0 21.9194 <2e-16 ***
Period    1     38    38.1  0.0724 0.7888
Treat     1    2209   2209.1  4.1993 0.0443 *
Carry     1    1051   1050.6  1.9970 0.1622
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
          Df Sum Sq Mean Sq F value Pr(>F)
Subject  35 403586 11531.0 21.9194 <2e-16 ***
Period    1     38    38.1  0.0724 0.7888
Treat     1    2209   2209.1  4.1993 0.0443 *
Carry     1    1051   1050.6  1.9970 0.1622
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.7.3 p361

(180) MODEL

```
GLM(Time ~ Subject + Period + Treat + Carry, chipman) # OK
```

```
$ANOVA
Response : Time
          Df Sum Sq Mean Sq F value    Pr(>F)
```

```

MODEL           17 28.0757 1.65151  64.421 1.139e-12 ***
RESIDUALS      18  0.4615 0.02564
CORRECTED TOTAL 35 28.5372
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE Time Mean Coef Var  R-square Adj R-sq
  0.1601128 6.250556 2.561577 0.9838299 0.9685581

$`Type I` 
  Df  Sum Sq Mean Sq F value    Pr(>F)
Subject 11 24.2084 2.20076 85.8462 3.157e-13 ***
Period   2  3.2065 1.60325 62.5388 7.894e-09 ***
Treat    2  0.4276 0.21382  8.3406  0.002733 **
Carry    2  0.2332 0.11660  4.5484  0.025188 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II` 
  Df  Sum Sq Mean Sq F value    Pr(>F)
Subject 11 24.2547 2.20497 86.0105 3.104e-13 ***
Period   1  0.0018 0.00184  0.0717 0.7919554
Treat    2  0.6392 0.31958 12.4661 0.0004003 ***
Carry    2  0.2332 0.11660  4.5484  0.0251881 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III` 
CAUTION: Singularity Exists !
  Df  Sum Sq Mean Sq F value    Pr(>F)
Subject 11 24.2547 2.20497 86.0105 3.104e-13 ***
Period   1  0.0018 0.00184  0.0717 0.7919554
Treat    2  0.6392 0.31958 12.4661 0.0004003 ***
Carry    2  0.2332 0.11660  4.5484  0.0251881 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.7.4 p372

(181) MODEL

```

residue$lc1 = log(residue$X1)
residue$lc2 = log(residue$X2)
residue$lc3 = log(residue$X3)
residue$lc4 = log(residue$X4)
residue$lc5 = log(residue$X5)
residue$sp = 7*residue$lc2+ 14*residue$lc3 + 30*residue$lc4 + 60*residue$lc5

```

```

residue$sm = residue$lc1 + residue$lc2+ residue$lc3 + residue$lc4 + residue$lc5
residue$num = 5*residue$sp - 111*residue$sm
residue$den = 5*4745 - 111^2
residue$k = residue$num/residue$den
residue$HL = -log(2)/residue$k
residue$logHL = log(residue$HL)
GLM(logHL ~ temp*moisture*soil, residue) # OK

$ANOVA
Response : logHL
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       7 7.5133 1.07332 13.543 0.0007329 ***
RESIDUALS   8 0.6340 0.07925
CORRECTED TOTAL 15 8.1473
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
Root MSE logHL Mean Coef Var R-square Adj R-sq
0.2815174 4.875155 5.774532 0.9221806 0.8540886

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
temp          1 6.0503 6.0503 76.3427 2.303e-05 ***
moisture      1 0.9521 0.9521 12.0134 0.008492 **
temp:moisture 1 0.0013 0.0013 0.0162 0.901779
soil          1 0.4098 0.4098 5.1712 0.052559 .
temp:soil     1 0.0086 0.0086 0.1081 0.750753
moisture:soil 1 0.0860 0.0860 1.0855 0.327921
temp:moisture:soil 1 0.0051 0.0051 0.0648 0.805427
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
temp          1 6.0503 6.0503 76.3427 2.303e-05 ***
moisture      1 0.9521 0.9521 12.0134 0.008492 **
temp:moisture 1 0.0013 0.0013 0.0162 0.901779
soil          1 0.4098 0.4098 5.1712 0.052559 .
temp:soil     1 0.0086 0.0086 0.1081 0.750753
moisture:soil 1 0.0860 0.0860 1.0855 0.327921
temp:moisture:soil 1 0.0051 0.0051 0.0648 0.805427
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
      Df Sum Sq Mean Sq F value    Pr(>F)

```

```

temp          1 6.0503  6.0503 76.3427 2.303e-05 ***
moisture      1 0.9521  0.9521 12.0134  0.008492 **
temp:moisture 1 0.0013  0.0013  0.0162  0.901779
soil          1 0.4098  0.4098  5.1712  0.052559 .
temp:soil     1 0.0086  0.0086  0.1081  0.750753
moisture:soil 1 0.0860  0.0860  1.0855  0.327921
temp:moisture:soil 1 0.0051  0.0051  0.0648  0.805427
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.8 Chapter 11

10.8.1 p461

(182) MODEL

```
GLM(y ~ x1 + x2 + x1:x2 + x1:x3 + x2:x3, pest) # OK
```

```
$ANOVA
Response : y
      Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL      5 275.642  55.128 160.38 4.631e-07 ***
RESIDUALS   7   2.406   0.344
CORRECTED TOTAL 12 278.048
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
  Root MSE    y Mean Coef Var  R-square Adj R-sq
  0.5862902 52.63077 1.113968 0.9913463 0.985165
```

```
$`Type I` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
x1      1  83.402  83.402 242.6351 1.086e-06 ***
x2      1 161.734 161.734 470.5191 1.116e-07 ***
x1:x2  1   0.246   0.246   0.7169 0.4251627
x1:x3  1  15.663  15.663  45.5660 0.0002649 ***
x2:x3  1  14.596  14.596  42.4614 0.0003291 ***
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II` 
      Df  Sum Sq Mean Sq F value    Pr(>F)
x1      1 215.951 215.951 628.246 4.105e-08 ***
x2      1 175.256 175.256 509.855 8.458e-08 ***
x1:x2  1   0.025   0.025   0.072 0.7961658
x1:x3  1  14.539  14.539  42.298 0.0003330 ***
x2:x3  1  14.596  14.596  42.461 0.0003291 ***
---

```

```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

x1     1 178.372 178.372 518.922 7.958e-08 ***  

x2     1 145.518 145.518 423.341 1.608e-07 ***  

x1:x2  1   0.025   0.025   0.072 0.7961658  

x1:x3  1 14.539  14.539  42.298 0.0003330 ***  

x2:x3  1 14.596  14.596  42.461 0.0003291 ***  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.8.2 p469

(183) MODEL

```
GLM(y ~ x1 + x2 + x1:x2 + x1:x3 + x2:x3 + x1:x2:x3, polvdat) # OK
```

```

$ANOVA  

Response : y  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL       6 12.5313 2.08854 37.056 0.0005473 ***  

RESIDUALS   5  0.2818 0.05636  

CORRECTED TOTAL 11 12.8131  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness  

  Root MSE   y Mean Coef Var R-square Adj R-sq  

  0.2374067 5.406667   4.391 0.9780061 0.9516133

```

```

$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

x1     1 5.4668  5.4668 96.9942 0.0001839 ***  

x2     1 0.3660  0.3660  6.4944 0.0513654 .  

x1:x2   1 4.6897  4.6897 83.2068 0.0002652 ***  

x1:x3   1 1.2450  1.2450 22.0887 0.0053378 **  

x2:x3   1 0.4707  0.4707  8.3509 0.0341949 *  

x1:x2:x3 1 0.2931  0.2931  5.2004 0.0714991 .  

---  

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$`Type II`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

x1     1 0.0184  0.0184  0.3265 0.5924707  

x2     1 0.2419  0.2419  4.2911 0.0930613 .  

x1:x2   1 3.8824  3.8824 68.8834 0.0004147 ***  

x1:x3   1 1.4383  1.4383 25.5196 0.0039276 **  

x2:x3   1 0.4707  0.4707  8.3509 0.0341949 *

```

```

x1:x2:x3 1 0.2931 0.2931 5.2004 0.0714991 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

      Df Sum Sq Mean Sq F value Pr(>F)  

x1       1 0.25744 0.25744 4.5677 0.08562 .  

x2       1 0.12956 0.12956 2.2987 0.18992  

x1:x2    1 0.65909 0.65909 11.6939 0.01885 *  

x1:x3    1 0.26323 0.26323 4.6704 0.08307 .  

x2:x3    1 0.12999 0.12999 2.3063 0.18931  

x1:x2:x3 1 0.29310 0.29310 5.2004 0.07150 .  

---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.8.3 p482

(184) MODEL

```

REG(y ~ x1 + x2 + x3 + x1:x2 + x1:x3 + x2:x3 + x1:z1 + x2:z1 + x3:z1 +
     x1:x2:z1 + x1:x3:z1 + x2:x3:z1 + x1:z2 + x2:z2 + x3:z2 +
     x1:x2:z2 + x1:x3:z2 + x2:x3:z2 + x1:z1:z2 + x2:z1:z2 + x3:z1:z2 +
     x1:x2:z1:z2 + x1:x3:z1:z2 + x2:x3:z1:z2 - 1, MPV) # OK

```

```

$ANOVA
Response : y
      Df Sum Sq Mean Sq F value Pr(>F)
MODEL      24 535997257 22333219 96.728 1.142e-09 ***
RESIDUALS   11 2539743   230886
UNCORRECTED TOTAL 35 538537000
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

$Fitness
Root MSE  y Mean Coef Var R-square Adj R-sq PRESS R2pred
480.5057 3582.857 13.41124 0.995284 0.9849945 51495197 0.9043795

```

```

$Coefficients
      Estimate Std. Error Df Lower CL Upper CL t value Pr(>|t|)  

x1       346948    294197 11  -300575   994471  1.1793 0.2631550  

x2        8223      490 11     7144    9301 16.7869 3.467e-09 ***  

x3       1656      459 11     646    2665  3.6104 0.0040950 **  

x1:x2    -414463   312262 11  -1101748   272822 -1.3273 0.2113017  

x1:x3    -334747   311426 11  -1020190   350696 -1.0749 0.3054382  

x2:x3    -6476      1199 11    -9114   -3838 -5.4032 0.0002156 ***  

x1:z1    103044    328922 11   -620909   826997  0.3133 0.7599297  

x2:z1    -2241      548 11    -3446   -1036 -4.0924 0.0017824 **  

x3:z1     823       513 11    -305    1952  1.6056 0.1366709  

x1:x2:z1 -64013    349120 11   -832421   704395 -0.1834 0.8578546

```

```

x1:x3:z1      -123730    348184 11  -890079    642618 -0.3554 0.7290412
x2:x3:z1       4659      1340 11   1709     7608  3.4765 0.0051806 ** 
x1:z2          244320    328922 11  -479632    968273 0.7428 0.4731733
x2:z2          886       548 11   -319     2092  1.6187 0.1338108
x3:z2          86        513 11  -1043     1214  0.1670 0.8704301
x1:x2:z2      -266052    349120 11  -1034460   502356 -0.7621 0.4620497
x1:x3:z2      -253151    348184 11  -1019500   513198 -0.7271 0.4823761
x2:x3:z2      -1822      1340 11  -4771     1128 -1.3593 0.2012686
x1:z1:z2      259038    328922 11  -464915    982990 0.7875 0.4476062
x2:z1:z2      -137       548 11  -1342     1068 -0.2500 0.8071853
x3:z1:z2       100       513 11  -1028     1229  0.1955 0.8485983
x1:x2:z1:z2  -269527    349120 11  -1037935   498881 -0.7720 0.4563702
x1:x3:z1:z2  -269249    348184 11  -1035597   497100 -0.7733 0.4556454
x2:x3:z1:z2  -328       1340 11  -3278     2621 -0.2448 0.8111141
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.9 Chapter 12

10.9.1 p513

(185) MODEL

```
GLM(ybar ~ A + B + C + D + E + F + G, tile) # OK
```

```
$ANOVA
Response : ybar
           Df  Sum Sq Mean Sq F value Pr(>F)
MODEL        7 0.68737 0.098196
RESIDUALS    0 0.00000
CORRECTED TOTAL 7 0.68737
```

```
$Fitness
Root MSE ybar Mean Coef Var R-square
      NA 0.7424626      NA      1
```

```
$`Type I` 
           Df  Sum Sq Mean Sq F value Pr(>F)
A 1 0.04984 0.04984
B 1 0.01992 0.01992
C 1 0.51534 0.51534
D 1 0.01532 0.01532
E 1 0.05965 0.05965
F 1 0.00879 0.00879
G 1 0.01851 0.01851
```

```
$`Type II` 
           Df  Sum Sq Mean Sq F value Pr(>F)
A 1 0.04984 0.04984
```

```

B 1 0.01992 0.01992
C 1 0.51534 0.51534
D 1 0.01532 0.01532
E 1 0.05965 0.05965
F 1 0.00879 0.00879
G 1 0.01851 0.01851

$`Type III`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 0.04984 0.04984
B 1 0.01992 0.01992
C 1 0.51534 0.51534
D 1 0.01532 0.01532
E 1 0.05965 0.05965
F 1 0.00879 0.00879
G 1 0.01851 0.01851

```

(186) MODEL

```
GLM(lns2 ~ A + B + C + D + E + F + G, tile) # OK
```

```

$ANOVA
Response : lns2
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7 12.305  1.7578
RESIDUALS      0  0.000
CORRECTED TOTAL 7 12.305

```

```

$Fitness
Root MSE lns2 Mean Coef Var R-square
      NA -2.623421      NA         1

```

```

$`Type I`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 1.6436  1.6436
B 1 0.3109  0.3109
C 1 7.1858  7.1858
D 1 2.3199  2.3199
E 1 0.0248  0.0248
F 1 0.7379  0.7379
G 1 0.0820  0.0820

```

```

$`Type II`
  Df Sum Sq Mean Sq F value Pr(>F)
A 1 1.6436  1.6436
B 1 0.3109  0.3109
C 1 7.1858  7.1858
D 1 2.3199  2.3199
E 1 0.0248  0.0248

```

```

F 1 0.7379 0.7379
G 1 0.0820 0.0820

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A 1 1.6436 1.6436  

B 1 0.3109 0.3109  

C 1 7.1858 7.1858  

D 1 2.3199 2.3199  

E 1 0.0248 0.0248  

F 1 0.7379 0.7379  

G 1 0.0820 0.0820

```

10.9.2 p521

(187) MODEL

```

strng = reshape(tile,
  direction = "long",
  varying = list(c("y1", "y2")),
  v.names = "y",
  idvar = c("A", "B", "C", "D", "E", "F", "G"),
  timevar = "H",
  times = c(-1, 1))
GLM(y ~ A/H + B/H + C/H + D/H + E/H + F/H + G/H, strng) # OK

```

```

$ANOVA
Response : y
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      14 1.65427 0.11816 0.1433 0.9807
RESIDUALS   1 0.82473 0.82473
CORRECTED TOTAL 15 2.47901

```

```

$Fitness
  Root MSE     y Mean Coef Var R-square  Adj R-sq
  0.9081486 0.7424626 122.3157 0.667313 -3.990305

```

```

$`Type I`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A 1 0.09968 0.09968 0.1209 0.7870  

A:H 1 0.04015 0.04015 0.0487 0.8618  

B 1 0.03984 0.03984 0.0483 0.8623  

H:B 1 0.00043 0.00043 0.0005 0.9854  

C 1 1.03069 1.03069 1.2497 0.4646  

H:C 1 0.15307 0.15307 0.1856 0.7410  

D 1 0.03064 0.03064 0.0372 0.8788  

H:D 1 0.04690 0.04690 0.0569 0.8510  

E 1 0.11929 0.11929 0.1446 0.7686  

H:E 1 0.01883 0.01883 0.0228 0.9045

```

```

F      1 0.01758 0.01758  0.0213  0.9077
H:F    1 0.01384 0.01384  0.0168  0.9180
G      1 0.03702 0.03702  0.0449  0.8671
H:G    1 0.00632 0.00632  0.0077  0.9444

```

\$`Type II`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	0.09968	0.09968	0.1209	0.7870
A:H	1	0.04015	0.04015	0.0487	0.8618
B	1	0.03984	0.03984	0.0483	0.8623
H:B	1	0.00043	0.00043	0.0005	0.9854
C	1	1.03069	1.03069	1.2497	0.4646
H:C	1	0.15307	0.15307	0.1856	0.7410
D	1	0.03064	0.03064	0.0372	0.8788
H:D	1	0.04690	0.04690	0.0569	0.8510
E	1	0.11929	0.11929	0.1446	0.7686
H:E	1	0.01883	0.01883	0.0228	0.9045
F	1	0.01758	0.01758	0.0213	0.9077
H:F	1	0.01384	0.01384	0.0168	0.9180
G	1	0.03702	0.03702	0.0449	0.8671
H:G	1	0.00632	0.00632	0.0077	0.9444

\$`Type III`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
A	1	0.09968	0.09968	0.1209	0.7870
A:H	1	0.04015	0.04015	0.0487	0.8618
B	1	0.03984	0.03984	0.0483	0.8623
H:B	1	0.00043	0.00043	0.0005	0.9854
C	1	1.03069	1.03069	1.2497	0.4646
H:C	1	0.15307	0.15307	0.1856	0.7410
D	1	0.03064	0.03064	0.0372	0.8788
H:D	1	0.04690	0.04690	0.0569	0.8510
E	1	0.11929	0.11929	0.1446	0.7686
H:E	1	0.01883	0.01883	0.0228	0.9045
F	1	0.01758	0.01758	0.0213	0.9077
H:F	1	0.01384	0.01384	0.0168	0.9180
G	1	0.03702	0.03702	0.0449	0.8671
H:G	1	0.00632	0.00632	0.0077	0.9444

10.9.3 p525

(188) MODEL

```

prod2 = af(prodstd, 1:7)
GLM(Pof ~ A + B + C + D + E + F + G + A:G + A:E:F + B:E:G + C:E:G + C:E:G:F +
     D:E + D:F, prod2) # OK

```

\$ANOVA

Response : Pof

	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
MODEL	47	769.49	16.3721	5.1667	2.737e-05 ***						
RESIDUALS	24	76.05	3.1688								
CORRECTED TOTAL	71	845.54									

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'. '	0.1	' '	1
\$Fitness											
Root MSE	MSE	Pof	Mean	Coef	Var	R-square	Adj	R-sq			
1.780098	19.73194	9.021403	0.9100571	0.7339189							
\$`Type I`											
	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
A	2	50.577	25.288	7.9806	0.0022023 **						
B	2	13.384	6.692	2.1118	0.1429491						
C	2	68.594	34.297	10.8234	0.0004463 ***						
D	2	23.674	11.837	3.7355	0.0386914 *						
E	1	275.733	275.733	87.0165	1.878e-09 ***						
F	1	161.700	161.700	51.0296	2.204e-07 ***						
G	1	1.051	1.051	0.3318	0.5699896						
A:G	2	26.567	13.284	4.1921	0.0274494 *						
A:E:F	7	28.404	4.058	1.2806	0.3013844						
B:E:G	7	22.453	3.208	1.0123	0.4475160						
C:E:G	6	35.546	5.924	1.8696	0.1277692						
C:E:F:G	10	24.607	2.461	0.7766	0.6500534						
D:E	2	21.745	10.873	3.4312	0.0489076 *						
D:F	2	15.450	7.725	2.4379	0.1086730						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'. '	0.1	' '	1
\$`Type II`											
	Df	Sum Sq	Mean Sq	F value	Pr(>F)						
A	2	50.577	25.288	7.9806	0.0022023 **						
B	2	13.384	6.692	2.1118	0.1429491						
C	2	68.594	34.297	10.8234	0.0004463 ***						
D	2	23.674	11.837	3.7355	0.0386914 *						
E	1	275.733	275.733	87.0165	1.878e-09 ***						
F	1	161.700	161.700	51.0296	2.204e-07 ***						
G	1	1.051	1.051	0.3318	0.5699896						
A:G	2	26.567	13.284	4.1921	0.0274494 *						
A:E:F	6	24.623	4.104	1.2951	0.2970196						
B:E:G	6	19.770	3.295	1.0398	0.4246194						
C:E:G	6	35.546	5.924	1.8696	0.1277692						
C:E:F:G	10	24.607	2.461	0.7766	0.6500534						
D:E	2	21.745	10.873	3.4312	0.0489076 *						
D:F	2	15.450	7.725	2.4379	0.1086730						

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'. '	0.1	' '	1

```
$`Type III`  

CAUTION: Singularity Exists !  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

A       2   50.577  25.288  7.9806 0.0022023 **  

B       2   13.384   6.692  2.1118 0.1429491  

C       2   68.594  34.297 10.8234 0.0004463 ***  

D       2   23.674  11.837  3.7355 0.0386914 *  

E       1 275.733 275.733 87.0165 1.878e-09 ***  

F       1 161.700 161.700 51.0296 2.204e-07 ***  

G       1     1.051   1.051  0.3318 0.5699896  

A:G     2   26.567  13.284  4.1921 0.0274494 *  

A:E:F   6   24.623   4.104  1.2951 0.2970196  

B:E:G   6   19.770   3.295  1.0398 0.4246194  

C:E:G   6   35.546   5.924  1.8696 0.1277692  

C:E:F:G 10   24.607   2.461  0.7766 0.6500534  

D:E     2   21.745  10.873  3.4312 0.0489076 *  

D:F     2   15.450   7.725  2.4379 0.1086730  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

10.9.4 p532

(189) MODEL

```
GLM(torque ~ A + B + C + D + E + A:B + A:C + A:D + A:E, Smotor) # OK
```

```
$ANOVA  

Response : torque  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

MODEL      15 0.0112217 0.00074811 102.2 0.009731 **  

RESIDUALS   2 0.0000146 0.00000732  

CORRECTED TOTAL 17 0.0112363  

---  

Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

Root	MSE	torque	Mean	Coef	Var	R-square	Adj R-sq
0.002705567	0.2572743	1.051627	0.9986971	0.988925			

```
$`Type I`  

      Df  Sum Sq Mean Sq F value    Pr(>F)  

A       1 0.0039545 0.0039545 540.2187 0.001846 **  

B       2 0.0003817 0.0001909  26.0732 0.036937 *  

C       2 0.0057241 0.0028620 390.9837 0.002551 **  

D       2 0.0000265 0.0000133   1.8104 0.355820  

E       1 0.0000984 0.0000984  13.4406 0.067009 .  

A:B     2 0.0010068 0.0005034  68.7668 0.014333 *  

A:C     2 0.0000031 0.0000016   0.2134 0.824110
```

```

A:D 2 0.0000009 0.0000004 0.0599 0.943521
A:E 1 0.0000258 0.0000258 3.5198 0.201458
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A 1 0.0039545 0.0039545 540.2187 0.001846 **  

B 2 0.0003817 0.0001909 26.0732 0.036937 *  

C 2 0.0032014 0.0016007 218.6753 0.004552 **  

D 2 0.0000268 0.0000134 1.8319 0.353123  

E 1 0.0000423 0.0000423 5.7744 0.138172  

A:B 2 0.0010068 0.0005034 68.7668 0.014333 *  

A:C 2 0.0000031 0.0000016 0.2134 0.824110  

A:D 2 0.0000052 0.0000026 0.3536 0.738760  

A:E 1 0.0000258 0.0000258 3.5198 0.201458
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`  

  Df Sum Sq Mean Sq F value Pr(>F)  

A 1 0.0034241 0.0034241 467.7636 0.002131 **  

B 2 0.0003817 0.0001909 26.0732 0.036937 *  

C 2 0.0032014 0.0016007 218.6753 0.004552 **  

D 2 0.0000268 0.0000134 1.8319 0.353123  

E 1 0.0000423 0.0000423 5.7744 0.138172  

A:B 2 0.0010068 0.0005034 68.7668 0.014333 *  

A:C 2 0.0000031 0.0000016 0.2134 0.824110  

A:D 2 0.0000052 0.0000026 0.3536 0.738760  

A:E 1 0.0000258 0.0000258 3.5198 0.201458
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.9.5 p535

(190) MODEL

```
GLM(shrinkage ~ A + B + C + D + E + F + G + A:B + A:C + A:D + A:E + A:F + A:G +
     B:D, inject) # OK
```

```
$ANOVA
Response : shrinkage
  Df Sum Sq Mean Sq F value Pr(>F)
MODEL      14 6659.4 475.67 129.08 1.97e-05 ***
RESIDUALS   5   18.4    3.68
CORRECTED TOTAL 19 6677.8
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$Fitness
Root MSE shrinkage Mean Coef Var R-square Adj R-sq
1.919635          27.1 7.083525 0.9972409 0.9895153
```

```
$`Type I` 
   Df Sum Sq Mean Sq  F value    Pr(>F)
A     1  770.1  770.1  208.9722 2.858e-05 ***
B     1 5076.6 5076.6 1377.6289 2.674e-07 ***
C     1     3.1     3.1    0.8311  0.403773
D     1     7.6     7.6    2.0522  0.211416
E     1     0.6     0.6    0.1526  0.712112
F     1     0.6     0.6    0.1526  0.712112
G     1   95.1   95.1   25.7972  0.003837 **
A:B   1  564.1  564.1  153.0699 6.112e-05 ***
A:C   1   10.6   10.6   2.8664  0.151230
A:D   1  115.6  115.6   31.3602  0.002508 **
A:E   1   14.1   14.1   3.8161  0.108185
A:F   1     1.6     1.6    0.4240  0.543677
A:G   1     0.1     0.1    0.0170  0.901459
B:D   1     0.1     0.1    0.0170  0.901459
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type II` 
   Df Sum Sq Mean Sq  F value    Pr(>F)
A     1  770.1  770.1  208.9722 2.858e-05 ***
B     1 5076.6 5076.6 1377.6289 2.674e-07 ***
C     1     3.1     3.1    0.8311  0.403773
D     1     7.6     7.6    2.0522  0.211416
E     1     0.6     0.6    0.1526  0.712112
F     1     0.6     0.6    0.1526  0.712112
G     1   95.1   95.1   25.7972  0.003837 **
A:B   1  564.1  564.1  153.0699 6.112e-05 ***
A:C   1   10.6   10.6   2.8664  0.151230
A:D   1  115.6  115.6   31.3602  0.002508 **
A:E   1   14.1   14.1   3.8161  0.108185
A:F   1     1.6     1.6    0.4240  0.543677
A:G   1     0.1     0.1    0.0170  0.901459
B:D   1     0.1     0.1    0.0170  0.901459
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
$`Type III` 
   Df Sum Sq Mean Sq  F value    Pr(>F)
A     1  770.1  770.1  208.9722 2.858e-05 ***
B     1 5076.6 5076.6 1377.6289 2.674e-07 ***
C     1     3.1     3.1    0.8311  0.403773
D     1     7.6     7.6    2.0522  0.211416
```

```

E    1    0.6    0.6    0.1526  0.712112
F    1    0.6    0.6    0.1526  0.712112
G    1    95.1   95.1   25.7972  0.003837 ***
A:B  1    564.1  564.1  153.0699 6.112e-05 ***
A:C  1    10.6   10.6   2.8664   0.151230
A:D  1    115.6  115.6  31.3602  0.002508 **
A:E  1    14.1   14.1   3.8161   0.108185
A:F  1    1.6    1.6    0.4240   0.543677
A:G  1    0.1    0.1    0.0170   0.901459
B:D  1    0.1    0.1    0.0170   0.901459
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

10.9.6 p539

(191) MODEL

```

eptax = cbind(eptaxr[1:16,], y2=eptaxr[17:32,9], y3=eptaxr[33:48,9],
               y5=eptaxr[49:64,9])
eptax$ybar = (eptax$y + eptax$y2 + eptax$y3 + eptax$y5)/4
GLM(ybar ~ A + B + C + D + E + F + G + H + A:B + A:C + A:D + A:E + A:F + A:G +
     A:H, eptax) # OK

```

```

$ANOVA
Response : ybar
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          15 2.8452 0.18968
RESIDUALS       0 0.0000
CORRECTED TOTAL 15 2.8452

```

```

$Fitness
Root MSE ybar Mean Coef Var R-square
      NA 14.36122      NA        1

```

```

`Type I`
            Df Sum Sq Mean Sq F value Pr(>F)
A      1 0.02686 0.02686
B      1 0.00042 0.00042
C      1 0.06306 0.06306
D      1 2.49443 2.49443
E      1 0.00304 0.00304
F      1 0.03209 0.03209
G      1 0.02954 0.02954
H      1 0.12879 0.12879
A:B   1 0.00047 0.00047
A:C   1 0.03218 0.03218
A:D   1 0.01185 0.01185
A:E   1 0.00380 0.00380
A:F   1 0.01674 0.01674

```

A:G 1 0.00186 0.00186
A:H 1 0.00012 0.00012

\$`Type II`
Df Sum Sq Mean Sq F value Pr(>F)
A 1 0.02686 0.02686
B 1 0.00042 0.00042
C 1 0.06306 0.06306
D 1 2.49443 2.49443
E 1 0.00304 0.00304
F 1 0.03209 0.03209
G 1 0.02954 0.02954
H 1 0.12879 0.12879
A:B 1 0.00047 0.00047
A:C 1 0.03218 0.03218
A:D 1 0.01185 0.01185
A:E 1 0.00380 0.00380
A:F 1 0.01674 0.01674
A:G 1 0.00186 0.00186
A:H 1 0.00012 0.00012

\$`Type III`
Df Sum Sq Mean Sq F value Pr(>F)
A 1 0.02686 0.02686
B 1 0.00042 0.00042
C 1 0.06306 0.06306
D 1 2.49443 2.49443
E 1 0.00304 0.00304
F 1 0.03209 0.03209
G 1 0.02954 0.02954
H 1 0.12879 0.12879
A:B 1 0.00047 0.00047
A:C 1 0.03218 0.03218
A:D 1 0.01185 0.01185
A:E 1 0.00380 0.00380
A:F 1 0.01674 0.01674
A:G 1 0.00186 0.00186
A:H 1 0.00012 0.00012

11 Searle - Linear Models 2e

Reference

- Searle SR, Gruber MHJ. Linear Models 2e, Kindle Edition. John Wiley & Sons Inc. 2016.

11.1 7.2 (p390, 59%)

(192) MODEL

```
weight = c(8,13,9,12,7,11,6,12,12,14,9,7,14,16,10,14,11,13)
treatment = c("ta","ta","ta","ta","ta","tb","tb","tb","tb","tc","tc","tc",
             "tc","tc","tc")
variety = c("va","va","va","vc","vd","vd","va","vb","vb","vb","vb","vc",
           "vc","vd","vd","vd")
d1 = data.frame(weight, treatment, variety)
GLM(weight ~ treatment*variety, d1)
```

\$ANOVA

```
Response : weight
            Df Sum Sq Mean Sq F value Pr(>F)
MODEL          7   82   11.714  2.0918  0.14
RESIDUALS      10   56    5.600
CORRECTED TOTAL 17  138
```

\$Fitness

```
Root MSE weight Mean Coef Var R-square Adj R-sq
2.366432           11 21.51302 0.5942029 0.3101449
```

\$`Type I`

```
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2 10.500   5.250  0.9375 0.42348
variety        3 36.786  12.262  2.1896 0.15232
treatment:variety  2 34.714  17.357  3.0995 0.08965 .
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2   9.486  4.7429  0.8469 0.45731
variety        3 36.786 12.2619  2.1896 0.15232
treatment:variety  2 34.714 17.3571  3.0995 0.08965 .
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```
            Df Sum Sq Mean Sq F value Pr(>F)
treatment      2 12.471  6.2353  1.1134 0.36595
variety        3 34.872 11.6240  2.0757 0.16719
```

```

treatment:variety 2 34.714 17.3571 3.0995 0.08965 .
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts = c("contr.sum", "contr.poly"))
Anova(lm(weight ~ treatment*variety, d1), type=3, singular.ok=TRUE) # NOT OK

Note: model has aliased coefficients
      sums of squares computed by model comparison

Anova Table (Type III tests)

Response: weight
          Sum Sq Df F values Pr(>F)
treatment      0.000  0
variety        0.000  0
treatment:variety 34.714  2   3.0995 0.08965 .
Residuals     56.000 10
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

11.2 7.2 (p393, 60%)

(193) MODEL

```

percent = c(31,33,44,36,38,26,37,59,42,42,34,42,28,39,36,32,38,42,36,22,42,46,
          26,37,43)
refinery = c(rep("g",9),rep("n",8),rep("s",8))
process = as.factor(c(1,1,1,1,1,1,2,2,2,1,1,1,2,2,2,2,1,1,1,2,2,2,2,2))
source0 = c("t","t","t","t","o","m","t","t","o","m","i","i","i","t","o","m","m",
           "t","o","i","o","o","m","i","i")
d2 = data.frame(percent, refinery, process, source=source0)
GLM(percent ~ refinery*source, d2)

```

\$ANOVA

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
MODEL	10	442.56	44.256	0.6361	0.7616
RESIDUALS	14	974.00	69.571		
CORRECTED TOTAL	24	1416.56			

\$Fitness

Root MSE	percent	Mean Coef	Var	R-square	Adj R-sq
8.340949		37.24	22.39782	0.3124188	-0.1787106

\$`Type I`

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
refinery	2	20.963	10.481	0.1507	0.8615
source	3	266.124	88.708	1.2751	0.3212
refinery:source	5	155.474	31.095	0.4469	0.8086

```
$`Type II`  
Df Sum Sq Mean Sq F value Pr(>F)  
refinery 2 25.535 12.767 0.1835 0.8343  
source 3 266.124 88.708 1.2751 0.3212  
refinery:source 5 155.474 31.095 0.4469 0.8086  
  
$`Type III`  
Df Sum Sq Mean Sq F value Pr(>F)  
refinery 2 10.766 5.383 0.0774 0.9259  
source 3 282.633 94.211 1.3542 0.2972  
refinery:source 5 155.474 31.095 0.4469 0.8086  
  
options(contrasts=c("contr.sum", "contr.poly"))  
Anova(lm(percent ~ refinery*source, d2), type=3, singular.ok=TRUE) # NOT OK
```

Note: model has aliased coefficients
 sums of squares computed by model comparison

Anova Table (Type III tests)

Response: percent

	Sum Sq	Df	F values	Pr(>F)
refinery	2.52	1	0.0362	0.8518
source	268.19	2	1.9275	0.1822
refinery:source	155.47	5	0.4469	0.8086
Residuals	974.00	14		

12 Web site examples

12.1 <https://github.com/djnavarro/psyr>

(194) MODEL

```
d21 = read.csv("http://r.acr.kr/psyr/coffee.csv")
GLM(babble ~ sugar*milk - 1, d21)

$ANOVA
Response : babble
      Df Sum Sq Mean Sq F value    Pr(>F)
MODEL       6 472.54 78.756 298.84 2.39e-12 ***
RESIDUALS   12   3.16   0.264
UNCORRECTED TOTAL 18 475.70
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$Fitness
  Root MSE babble Mean Coef Var R-square Adj R-sq
0.5133631   5.066667 10.13217 0.9933519 0.9900279

$`Type I`
      Df Sum Sq Mean Sq F value    Pr(>F)
sugar      3 465.64 155.213 588.9486 2.756e-13 ***
milk       1   0.96   0.956   3.6279  0.081061 .
sugar:milk 2   5.94   2.972  11.2769  0.001754 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type II`
      Df Sum Sq Mean Sq F value    Pr(>F)
sugar      2 3.0696 1.53482  5.8238 0.017075 *
milk       1 0.9561 0.95611  3.6279 0.081061 .
sugar:milk 2 5.9439 2.97193 11.2769 0.001754 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$`Type III`
CAUTION: Singularity Exists !
      Df Sum Sq Mean Sq F value    Pr(>F)
sugar      2 2.1318  1.0659  4.0446 0.045426 *
milk       1 1.0041  1.0041  3.8102 0.074672 .
sugar:milk 2 5.9439  2.9719 11.2769 0.001754 **
---
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
options(contrasts=c("contr.sum", "contr.poly"))
r21 = lm(babble ~ sugar*milk - 1, d21)
```

```
anova(r21) # Type I SS OK
```

Analysis of Variance Table

Response: babble

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
sugar	3	465.64	155.213	588.9486	2.756e-13 ***
milk	1	0.96	0.956	3.6279	0.081061 .
sugar:milk	2	5.94	2.972	11.2769	0.001754 **
Residuals	12	3.16	0.264		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
Anova(r21, type=2) # NOT OK
```

Anova Table (Type II tests)

Response: babble

	Sum Sq	Df	F value	Pr(>F)
sugar	453.76	3	573.9233	3.214e-13 ***
milk	0.96	1	3.6279	0.081061 .
sugar:milk	5.94	2	11.2769	0.001754 **
Residuals	3.16	12		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
Anova(r21, type=3) # NOT OK
```

Anova Table (Type III tests)

Response: babble

	Sum Sq	Df	F value	Pr(>F)
sugar	454.77	3	575.1970	3.172e-13 ***
milk	1.00	1	3.8102	0.074672 .
sugar:milk	5.94	2	11.2769	0.001754 **
Residuals	3.16	12		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

13 Bioequivalence (BE) data example

(195) MODEL

```
GLM(log(CMAX) ~ SEQ/SUBJ + PRD + TRT, BEdata) # a BE dataset in sasLM package
```

\$ANOVA

```
Response : log(CMAX)
            Df  Sum Sq Mean Sq F value    Pr(>F)
MODEL          48 23.1924 0.48317  5.6278 4.395e-08 ***
RESIDUALS      42  3.6059 0.08585
CORRECTED TOTAL 90 26.7983
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$Fitness

```
Root MSE log(CMAX) Mean Coef Var R-square Adj R-sq
0.2930098      6.071036 4.826355 0.8654428 0.7116631
```

\$`Type I`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.6454 0.64544  7.5178  0.008938 **
SEQ:SUBJ 45 22.4395 0.49866  5.8081 3.359e-08 ***
PRD     1  0.0969 0.09686  1.1281  0.294242
TRT     1  0.0106 0.01057  0.1231  0.727410
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type II`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.6440 0.64395  7.5005  0.009011 **
SEQ:SUBJ 45 22.5232 0.50052  5.8298 3.173e-08 ***
PRD     1  0.0996 0.09958  1.1599  0.287632
TRT     1  0.0106 0.01057  0.1231  0.727410
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

\$`Type III`

```
Df  Sum Sq Mean Sq F value    Pr(>F)
SEQ     1  0.3368 0.33679  3.9228  0.05421 .
SEQ:SUBJ 45 22.5232 0.50052  5.8298 3.173e-08 ***
PRD     1  0.0996 0.09958  1.1599  0.28763
TRT     1  0.0106 0.01057  0.1231  0.72741
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
options(contrasts=c("contr.sum", "contr.poly"))
Anova(lm(log(CMAX) ~ SEQ/SUBJ + PRD + TRT, BEdata), type=3, singular.ok=TRUE)
```

Note: model has aliased coefficients
sums of squares computed by model comparison

Anova Table (Type III tests)

Response: log(CMAX)

	Sum Sq	Df	F values	Pr(>F)
SEQ	0.0000	0		
PRD	0.0996	1	1.1599	0.2876
TRT	0.0106	1	0.1231	0.7274
SEQ:SUBJ	22.5232	45	5.8298	3.173e-08 ***
Residuals	3.6059	42		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1				

14 Test Summary

Package	Version	Total Count	Identical to SAS	Different from SAS
sasLM	0.10.3	195	195 (100%)	0 (0%)
car	3.1.2	195	173 (89%)	22 (11%)

All of the results by sasLM 0.10.3 were practically identical to those of SAS.

Last digit difference by 1 is resulted from the round-to-even number way of R rounding function.

If you are uncertain about the equivalence of the ‘sasLM’ to ‘SAS,’ you can check these examples using ‘SAS onDemand’ for free.

If you have any question, please mail to the author, Kyun-Seop Bae k@acr.kr.

15 Session Information

```
R version 4.3.3 (2024-02-29 ucrt)
Platform: x86_64-w64-mingw32/x64 (64-bit)
Running under: Windows 10 x64 (build 19044)
```

```
Matrix products: default
```

```
locale:
```

```
[1] LC_COLLATE=Korean_Korea.utf8  LC_CTYPE=Korean_Korea.utf8
[3] LC_MONETARY=Korean_Korea.utf8 LC_NUMERIC=C
[5] LC_TIME=Korean_Korea.utf8
```

```
time zone: Asia/Seoul
```

```
tzcode source: internal
```

```
attached base packages:
```

```
[1] stats      graphics   grDevices utils      datasets  methods   base
```

```
other attached packages:
```

```
[1] daewr_1.2-11    car_3.1-2       carData_3.0-5   sasLM_0.10.3   mvtnorm_1.2-4
[6] rmarkdown_2.25
```

```
loaded via a namespace (and not attached):
```

```
[1] digest_0.6.34    fastmap_1.1.1    xfun_0.41        abind_1.4-5
[5] knitr_1.45       htmltools_0.5.7  tinytex_0.49    cli_3.6.2
[9] compiler_4.3.3   tools_4.3.3     evaluate_0.23   yaml_2.3.8
[13] rlang_1.1.3     MASS_7.3-60.0.1
```