

Research Article

Refinements of Kantorovich Inequality for Hermitian Matrices

Feixiang Chen

School of Mathematics and Statistics, Chongqing Three Gorges University, Wanzhou, Chongqing 404000, China

Correspondence should be addressed to Feixiang Chen, cfx2002@126.com

Received 28 August 2012; Accepted 5 November 2012

Academic Editor: K. C. Sivakumar

Copyright © 2012 Feixiang Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Some new Kantorovich-type inequalities for Hermitian matrix are proposed in this paper. We consider what happens to these inequalities when the positive definite matrix is allowed to be invertible and provides refinements of the classical results.

1. Introduction and Preliminaries

We first state the well-known Kantorovich inequality for a positive definite Hermite matrix (see [1, 2]), let $A \in M_n$ be a positive definite Hermitian matrix with real eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Then

$$1 \leq x^* A x x^* A^{-1} x \leq \frac{(\lambda_1 + \lambda_n)^2}{4\lambda_1 \lambda_n}, \quad (1.1)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$, where A^* denotes the conjugate transpose of matrix A . A matrix $A \in M_n$ is Hermitian if $A = A^*$. An equivalent form of this result is incorporated in

$$0 \leq x^* A x x^* A^{-1} x - 1 \leq \frac{(\lambda_n - \lambda_1)^2}{4\lambda_1 \lambda_n}, \quad (1.2)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$.

Attributed to Kantorovich, the inequality has built up a considerable literature. This typically comprises generalizations. Examples are [3–5] for matrix versions. Operator

versions are developed in [6, 7]. Multivariate versions have been useful in statistics to assess the robustness of least squares, see [8, 9] and the references therein.

Due to the important applications of the original Kantorovich inequality for matrices [10] in Statistics [8, 11, 12] and Numerical Analysis [13, 14], any new inequality of this type will have a flow of consequences in the areas of applications.

Motivated by the interest in both pure and applied mathematics outlined above we establish in this paper some improvements of Kantorovich inequalities. The classical Kantorovich-type inequalities are modified to apply not only to positive definite but also to invertible Hermitian matrices. As natural tools in deriving the new results, the recent Grüss-type inequalities for vectors in inner product in [6, 15–19] are utilized.

To simplify the proof, we first introduce some lemmas.

2. Lemmas

Let B be a Hermitian matrix with real eigenvalues $\mu_1 \leq \mu_2 \leq \dots \leq \mu_n$, if $A - B$ is positive semidefinite, we write

$$A \geq B, \quad (2.1)$$

that is, $\lambda_i \geq \mu_i$, $i = 1, 2, \dots, n$. On \mathbb{C}^n , we have the standard inner product defined by $\langle x, y \rangle = \sum_{i=1}^n x_i \bar{y}_i$, where $x = (x_1, \dots, x_n)^* \in \mathbb{C}^n$ and $y = (y_1, \dots, y_n)^* \in \mathbb{C}^n$.

Lemma 2.1. *Let a, b, c , and d be real numbers, then one has the following inequality:*

$$(a^2 - b^2)(c^2 - d^2) \leq (ac - bd)^2. \quad (2.2)$$

Lemma 2.2. *Let A and B be Hermitian matrices, if $AB = BA$, then*

$$AB \leq \frac{(A + B)^2}{4}. \quad (2.3)$$

Lemma 2.3. *Let $A \geq 0$, $B \geq 0$, if $AB = BA$, then*

$$AB \geq 0. \quad (2.4)$$

3. Some Results

The following lemmas can be obtained from [16–19] by replacing Hilbert space $(H, \langle \cdot, \cdot \rangle)$ with inner product spaces \mathbb{C}^n , so we omit the details.

Lemma 3.1. *Let u, v , and e be vectors in \mathbb{C}^n , and $\|e\| = 1$. If α, β, δ , and γ are real or complex numbers such that*

$$\operatorname{Re}\langle \beta e - u, u - \alpha e \rangle \geq 0, \quad \operatorname{Re}\langle \delta e - v, v - \gamma e \rangle \geq 0, \quad (3.1)$$

then

$$|\langle u, v \rangle - \langle u, e \rangle \langle e, v \rangle| \leq \frac{1}{4}(\beta - \alpha)(\delta - \gamma) - [\operatorname{Re}(\beta e - u, u - \alpha e) \operatorname{Re}(\delta e - v, v - \gamma e)]^{1/2}. \quad (3.2)$$

Lemma 3.2. *With the assumptions in Lemma 3.1, one has*

$$|\langle u, v \rangle - \langle u, e \rangle \langle e, v \rangle| \leq \frac{1}{4}(\beta - \alpha)(\gamma - \delta) - \left| \langle u, e \rangle - \frac{\alpha + \beta}{2} \right| \left| \langle v, e \rangle - \frac{\gamma + \delta}{2} \right|. \quad (3.3)$$

Lemma 3.3. *With the assumptions in Lemma 3.1, if $\operatorname{Re}(\beta \bar{\alpha}) \geq 0$, $\operatorname{Re}(\delta \bar{\gamma}) \geq 0$, one has*

$$|\langle u, v \rangle - \langle u, e \rangle \langle e, v \rangle| \leq \frac{|\beta - \alpha| |\delta - \gamma|}{4[\operatorname{Re}(\beta \bar{\alpha}) \operatorname{Re}(\delta \bar{\gamma})]^{1/2}} |\langle u, e \rangle \langle e, v \rangle|. \quad (3.4)$$

Lemma 3.4. *With the assumptions in Lemma 3.3, one has*

$$\begin{aligned} & |\langle u, v \rangle - \langle u, e \rangle \langle e, v \rangle| \\ & \leq \left\{ \left(|\alpha + \beta| - 2[\operatorname{Re}(\beta \bar{\alpha})]^{1/2} \right) \left(|\delta + \gamma| - 2[\operatorname{Re}(\delta \bar{\gamma})]^{1/2} \right) \right\}^{1/2} [|\langle u, e \rangle \langle e, v \rangle|]^{1/2}. \end{aligned} \quad (3.5)$$

4. New Kantorovich Inequalities for Hermitian Matrices

For a Hermitian matrix A , as in [6], we define the following transform:

$$C(A) = (\lambda_n I - A)(A - \lambda_1 I). \quad (4.1)$$

When A is invertible, if $\lambda_1 \lambda_n > 0$, then,

$$C(A^{-1}) = \left(\frac{1}{\lambda_1} I - A^{-1} \right) \left(A^{-1} - \frac{1}{\lambda_n} I \right). \quad (4.2)$$

Otherwise, $\lambda_1 \lambda_n < 0$, then,

$$C(A^{-1}) = \left(\frac{1}{\lambda_{k+1}} I - A^{-1} \right) \left(A^{-1} - \frac{1}{\lambda_k} I \right), \quad (4.3)$$

where

$$\lambda_1 \leq \dots \leq \lambda_k < 0 < \lambda_{k+1} \leq \dots \leq \lambda_n. \quad (4.4)$$

From Lemma 2.3 we can conclude that $C(A) \geq 0$ and $C(A^{-1}) \geq 0$.

For two Hermitian matrices A and B , and $x \in \mathbb{C}^n$, $\|x\| = 1$, we define the following functional:

$$G(A, B; x) = \langle Ax, Bx \rangle - \langle Ax, x \rangle \langle x, Bx \rangle. \quad (4.5)$$

When $A = B$, we denote

$$G(A; x) = \|Ax\|^2 - \langle Ax, x \rangle^2, \quad (4.6)$$

for $x \in \mathbb{C}^n$, $\|x\| = 1$.

Lemma 4.1. *With notations above, and for $x \in \mathbb{C}^n$, $\|x\| = 1$, then*

$$0 \leq \langle C(A)x, x \rangle \leq \frac{(\lambda_n - \lambda_1)^2}{4}, \quad (4.7)$$

if $\lambda_n \lambda_1 > 0$,

$$0 \leq \langle C(A^{-1})x, x \rangle \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)^2}, \quad (4.8)$$

if $\lambda_n \lambda_1 < 0$,

$$0 \leq \langle C(A^{-1})x, x \rangle \leq \frac{(\lambda_{k+1} - \lambda_k)^2}{4(\lambda_{k+1} \lambda_k)^2}. \quad (4.9)$$

Proof. From $C(A) \geq 0$, then

$$\langle C(A)x, x \rangle \geq 0. \quad (4.10)$$

While, from Lemma 2.2, we can get

$$C(A) = (\lambda_n I - A)(A - \lambda_1 I) \leq \frac{(\lambda_n - \lambda_1)^2}{4} I. \quad (4.11)$$

Then $\langle C(A)x, x \rangle \leq (\lambda_n - \lambda_1)^2/4$ is straightforward. The proof for $C(A^{-1})$ is similar. \square

Lemma 4.2. *With notations above, and for $x \in \mathbb{C}^n$, $\|x\| = 1$, then*

$$\left| x^* A x x^* A^{-1} x - 1 \right|^2 \leq G(A; x) G(A^{-1}; x). \quad (4.12)$$

Proof. Thus,

$$\begin{aligned} \left| x^* A x x^* A^{-1} x - 1 \right|^2 &= \left| x^* \left((x^* A^{-1} x) I - A^{-1} \right) \left((x^* A x) I - A \right) x \right|^2 \\ &\leq \left\| \left((x^* A^{-1} x) I - A^{-1} \right) x \right\|^2 \left\| \left((x^* A x) I - A \right) x \right\|^2, \end{aligned} \quad (4.13)$$

while

$$\begin{aligned} \left\| \left((x^* A x) I - A \right) x \right\|^2 &= x^* \left((x^* A x)^2 I - 2(x^* A x) A + A^2 \right) x \\ &= x^* A^2 x - (x^* A x)^2 \\ &= \|Ax\|^2 - \langle Ax, x \rangle^2 \\ &= G(A; x). \end{aligned} \quad (4.14)$$

Similarly, we can get $\left\| \left((x^* A^{-1} x) I - A^{-1} \right) x \right\|^2 = G(A^{-1}; x)$, then we complete the proof. \square

Theorem 4.3. Let A, B be two Hermitian matrices, and $C(A) \geq 0, C(B) \geq 0$ are defined as above, then

$$\begin{aligned} |G(A, B; x)| &\leq \frac{1}{4} (\lambda_n - \lambda_1) (\mu_n - \mu_1) - [\langle C(A)x, x \rangle \langle C(B)x, x \rangle]^{1/2}, \\ |G(A, B; x)| &\leq \frac{1}{4} (\lambda_n - \lambda_1) (\mu_n - \mu_1) - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} \right) x, x \right\rangle \left\langle \left(B - \frac{\mu_1 + \mu_n}{2} \right) x, x \right\rangle \right|, \end{aligned} \quad (4.15)$$

for any $x \in \mathbb{C}^n, \|x\| = 1$.

If $\lambda_1 \lambda_n > 0, \mu_1 \mu_n > 0$, then

$$\begin{aligned} |G(A, B; x)| &\leq \frac{(\lambda_n - \lambda_1) (\mu_n - \mu_1)}{4 [(\lambda_n \lambda_1) (\mu_n \mu_1)]^{1/2}} |\langle Ax, x \rangle \langle Bx, x \rangle|, \\ |G(A, B; x)| &\leq \left\{ \left(|\lambda_1 + \lambda_n| - 2(\lambda_1 \lambda_n)^{1/2} \right) \left(|\mu_1 + \mu_n| - 2(\mu_1 \mu_n)^{1/2} \right) \right\}^{1/2} [|\langle Ax, x \rangle \langle Bx, x \rangle|]^{1/2}, \end{aligned} \quad (4.16)$$

for any $x \in \mathbb{C}^n, \|x\| = 1$.

Proof. The proof follows by Lemmas 3.1, 3.2, 3.3, and 3.4 on choosing $u = Ax, v = Bx$, and $e = x, \beta = \lambda_n, \alpha = \lambda_1, \delta = \mu_n$, and $\gamma = \mu_1, x \in \mathbb{C}^n, \|x\| = 1$, respectively. \square

Corollary 4.4. Let A be a Hermitian matrices, and $C(A) \geq 0$ is defined as above, then

$$|G(A; x)| \leq \frac{1}{4}(\lambda_n - \lambda_1)^2 - \langle C(A)x, x \rangle, \quad (4.17)$$

$$|G(A; x)| \leq \frac{1}{4}(\lambda_n - \lambda_1)^2 - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right|^2, \quad (4.18)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$.
If $\lambda_1 \lambda_n > 0$, then

$$|G(A; x)| \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)} |\langle Ax, x \rangle|^2, \quad (4.19)$$

$$|G(A; x)| \leq \left(|\lambda_1 + \lambda_n| - 2\sqrt{\lambda_1 \lambda_n} \right) |\langle Ax, x \rangle|, \quad (4.20)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$.

Proof. The proof follows by Theorem 4.3 on choosing $A = B$, respectively. \square

Corollary 4.5. Let A be a Hermitian matrices and $C(A^{-1}) \geq 0$ is defined as above, then one has the following.

If $\lambda_1 \lambda_n > 0$, then

$$|G(A^{-1}; x)| \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)^2} - \langle C(A^{-1})x, x \rangle, \quad (4.21)$$

$$|G(A^{-1}; x)| \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)^2} - \left| \left\langle \left(A^{-1} - \frac{\lambda_1 + \lambda_n}{2\lambda_n \lambda_1} I \right) x, x \right\rangle \right|^2, \quad (4.22)$$

$$|G(A^{-1}; x)| \leq \frac{(\lambda_n - \lambda_1)^2}{4\lambda_n \lambda_1} |\langle A^{-1}x, x \rangle|^2, \quad (4.23)$$

$$|G(A^{-1}; x)| \leq \left(\frac{|\lambda_1 + \lambda_n|}{\lambda_1 \lambda_n} - 2\frac{1}{\sqrt{\lambda_1 \lambda_n}} \right) |\langle A^{-1}x, x \rangle|, \quad (4.24)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$.
If $\lambda_1 \lambda_n < 0$, then

$$|G(A^{-1}; x)| \leq \frac{(\lambda_k - \lambda_{k+1})^2}{4(\lambda_k \lambda_{k+1})^2} - \langle C(A^{-1})x, x \rangle, \quad (4.25)$$

where $C(A^{-1}) = ((1/\lambda_{k+1})I - A^{-1})(A^{-1} - (1/\lambda_k)I)$, and

$$|G(A^{-1}; x)| \leq \frac{(\lambda_k - \lambda_{k+1})^2}{4(\lambda_k \lambda_{k+1})^2} - \left| \left\langle \left(A^{-1} - \frac{\lambda_{k+1} + \lambda_k}{2\lambda_k \lambda_{k+1}} I \right) x, x \right\rangle \right|^2, \quad (4.26)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$.

Proof. The proof follows by Corollary 4.4 by replacing A with A^{-1} , respectively. \square

Theorem 4.6. Let A be an $n \times n$ invertible Hermitian matrix with real eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$, then one has the following.

If $\lambda_1 \lambda_n > 0$, then

$$|x^* A x x^* A^{-1} x - 1| \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)} - \sqrt{\langle C(A)x, x \rangle \langle C(A^{-1})x, x \rangle}, \quad (4.27)$$

$$|x^* A x x^* A^{-1} x - 1| \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)} - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right| \left| \left\langle \left(A^{-1} - \frac{\lambda_1 + \lambda_n}{2\lambda_n \lambda_1} I \right) x, x \right\rangle \right|, \quad (4.28)$$

$$|x^* A x x^* A^{-1} x - 1| \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)^2} \langle Ax, x \rangle \langle A^{-1}x, x \rangle, \quad (4.29)$$

$$|x^* A x x^* A^{-1} x - 1| \leq \frac{(\sqrt{|\lambda_n|} - \sqrt{|\lambda_1|})^2}{\sqrt{\lambda_n \lambda_1}} \sqrt{\langle Ax, x \rangle \langle A^{-1}x, x \rangle}, \quad (4.30)$$

for any $x \in \mathbb{C}^n$, $\|x\| = 1$.

If $\lambda_1 \lambda_n < 0$, then

$$|x^* A x x^* A^{-1} x - 1| \leq \frac{(\lambda_n - \lambda_1)(\lambda_{k+1} - \lambda_k)}{4|\lambda_k \lambda_{k+1}|} - \sqrt{\langle C(A)x, x \rangle \langle C(A^{-1})x, x \rangle}, \quad (4.31)$$

where $C(A^{-1}) = ((1/\lambda_{k+1})I - A^{-1})(A^{-1} - (1/\lambda_k)I)$,

$$|x^* A x x^* A^{-1} x - 1| \leq \frac{(\lambda_n - \lambda_1)(\lambda_{k+1} - \lambda_k)}{4|\lambda_k \lambda_{k+1}|} - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right| \left| \left\langle \left(A^{-1} - \frac{\lambda_k + \lambda_{k+1}}{2\lambda_1 \lambda_{k+1}} I \right) x, x \right\rangle \right|. \quad (4.32)$$

Proof. Considering

$$|x^* A x x^* A^{-1} x - 1|^2 \leq G(A; x)G(A^{-1}; x). \quad (4.33)$$

When $\lambda_1\lambda_n > 0$, from (4.17) and (4.21), we get

$$\left| x^* A x x^* A^{-1} x - 1 \right|^2 \leq \left\{ \frac{1}{4} (\lambda_n - \lambda_1)^2 - \langle C(A)x, x \rangle \right\} \left\{ \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n\lambda_1)^2} - \langle C(A^{-1})x, x \rangle \right\}. \quad (4.34)$$

From $(a^2 - b^2)(c^2 - d^2) \leq (ac - bd)^2$, we have

$$\left| x^* A x x^* A^{-1} x - 1 \right|^2 \leq \left\{ \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n\lambda_1)} - \sqrt{[\langle C(A)x, x \rangle \langle C(A^{-1})x, x \rangle]} \right\}^2, \quad (4.35)$$

then, the conclusion (4.27) holds.

Similarly, from (4.18) and (4.22), we get

$$\begin{aligned} \left| x^* A x x^* A^{-1} x - 1 \right|^2 &\leq \left\{ \frac{1}{4} (\lambda_n - \lambda_1)^2 - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right|^2 \right\} \\ &\quad \times \left\{ \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n\lambda_1)^2} - \left| \left\langle \left(A^{-1} - \frac{\lambda_1 + \lambda_n}{2\lambda_n\lambda_1} I \right) x, x \right\rangle \right|^2 \right\} \\ &\leq \left\{ \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n\lambda_1)} - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right| \left| \left\langle \left(A^{-1} - \frac{\lambda_1 + \lambda_n}{2\lambda_n\lambda_1} I \right) x, x \right\rangle \right| \right\}^2, \end{aligned} \quad (4.36)$$

then, the conclusion (4.28) holds.

From (4.19) and (4.23), we get

$$\left| x^* A x x^* A^{-1} x - 1 \right|^2 \leq \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n\lambda_1)} |\langle Ax, x \rangle|^2 \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n\lambda_1)} |\langle A^{-1}x, x \rangle|^2, \quad (4.37)$$

then, the conclusion (4.29) holds.

From (4.20) and (4.24), we get

$$\left| x^* A x x^* A^{-1} x - 1 \right|^2 \leq (|\lambda_1 + \lambda_n| - 2\sqrt{\lambda_1\lambda_n}) |\langle Ax, x \rangle| \left(\frac{|\lambda_1 + \lambda_n|}{\lambda_1\lambda_n} - 2\frac{1}{\sqrt{\lambda_1\lambda_n}} \right) |\langle A^{-1}x, x \rangle|, \quad (4.38)$$

then, the conclusion (4.30) holds.

When $\lambda_1\lambda_n < 0$, from (4.17) and (4.25), we get

$$\left| x^* A x x^* A^{-1} x - 1 \right|^2 \leq \left\{ \frac{1}{4} (\lambda_n - \lambda_1)^2 - \langle C(A)x, x \rangle \right\} \left\{ \frac{(\lambda_k - \lambda_{k+1})^2}{4(\lambda_k\lambda_{k+1})^2} - \langle C(A^{-1})x, x \rangle \right\}, \quad (4.39)$$

then, the conclusion (4.31) holds.

From (4.18) and (4.26), we get

$$\begin{aligned}
|x^*Ax x^*A^{-1}x - 1|^2 &\leq \left\{ \frac{1}{4}(\lambda_n - \lambda_1)^2 - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right|^2 \right\} \\
&\quad \times \left\{ \frac{(\lambda_k - \lambda_{k+1})^2}{4(\lambda_k \lambda_{k+1})^2} - \left| \left\langle \left(A^{-1} - \frac{\lambda_{k+1} + \lambda_k}{2\lambda_k \lambda_{k+1}} I \right) x, x \right\rangle \right|^2 \right\} \\
&\leq \left\{ \frac{(\lambda_n - \lambda_1)(\lambda_{k+1} - \lambda_k)}{4|\lambda_k \lambda_{k+1}|} \right. \\
&\quad \left. - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right| \left| \left\langle \left(A^{-1} - \frac{\lambda_k + \lambda_{k+1}}{2\lambda_1 \lambda_{k+1}} I \right) x, x \right\rangle \right| \right\}^2,
\end{aligned} \tag{4.40}$$

then, the conclusion (4.32) holds. \square

Corollary 4.7. *With the notations above, for any $x \in \mathbb{C}^n$, $\|x\| = 1$, one lets*

$$\begin{aligned}
|G_1(A; x)| &= \frac{1}{4}(\lambda_n - \lambda_1)^2 - \langle C(A)x, x \rangle, \\
|G_2(A; x)| &= \frac{1}{4}(\lambda_n - \lambda_1)^2 - \left| \left\langle \left(A - \frac{\lambda_1 + \lambda_n}{2} I \right) x, x \right\rangle \right|^2.
\end{aligned} \tag{4.41}$$

If $\lambda_1 \lambda_n > 0$, one lets

$$\begin{aligned}
|G_3(A; x)| &= \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)} |\langle Ax, x \rangle|^2, \\
|G_4(A; x)| &= \left(|\lambda_1 + \lambda_n| - 2\sqrt{\lambda_1 \lambda_n} \right) |\langle Ax, x \rangle|.
\end{aligned} \tag{4.42}$$

If $\lambda_1 \lambda_n > 0$

$$\begin{aligned}
|G^1(A^{-1}; x)| &= \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)^2} - \langle C(A^{-1})x, x \rangle, \\
|G^2(A^{-1}; x)| &= \frac{(\lambda_n - \lambda_1)^2}{4(\lambda_n \lambda_1)^2} - \left| \left\langle \left(A^{-1} - \frac{\lambda_1 + \lambda_n}{2\lambda_n \lambda_1} I \right) x, x \right\rangle \right|^2, \\
|G^3(A^{-1}; x)| &= \frac{(\lambda_n - \lambda_1)^2}{4\lambda_n \lambda_1} \left| \langle A^{-1}x, x \rangle \right|^2, \\
|G^4(A^{-1}; x)| &= \left(\frac{|\lambda_1 + \lambda_n|}{\lambda_1 \lambda_n} - 2\frac{1}{\sqrt{\lambda_1 \lambda_n}} \right) \left| \langle A^{-1}x, x \rangle \right|.
\end{aligned} \tag{4.43}$$

If $\lambda_1\lambda_n < 0$, then

$$\left| G^5(A^{-1}; x) \right| = \frac{(\lambda_k - \lambda_{k+1})^2}{4(\lambda_k\lambda_{k+1})^2} - \langle C(A^{-1})x, x \rangle, \quad (4.44)$$

where $C(A^{-1}) = ((1/\lambda_{k+1})I - A^{-1})(A^{-1} - (1/\lambda_k)I)$, and

$$\left| G^6(A^{-1}; x) \right| = \frac{(\lambda_k - \lambda_{k+1})^2}{4(\lambda_k\lambda_{k+1})^2} - \left| \left\langle \left(A^{-1} - \frac{\lambda_{k+1} + \lambda_k}{2\lambda_k\lambda_{k+1}} I \right) x, x \right\rangle \right|^2. \quad (4.45)$$

Then, one has the following.

If $\lambda_1\lambda_n > 0$,

$$\left| x^* A x x^* A^{-1} x - 1 \right| \leq \sqrt{G^*(A; x) G^*(A^{-1}; x)}, \quad (4.46)$$

where

$$\begin{aligned} G^*(A; x) &= \min\{G_1(A; x), G_2(A; x), G_3(A; x), G_4(A; x)\}, \\ G^*(A; x) &= \min\{G^1(A^{-1}; x), G^2(A^{-1}; x), G^3(A^{-1}; x), G^4(A^{-1}; x)\}. \end{aligned} \quad (4.47)$$

If $\lambda_1\lambda_n < 0$

$$\left| x^* A x x^* A^{-1} x - 1 \right| \leq \sqrt{G^\circ(A; x) G^\bullet(A^{-1}; x)}, \quad (4.48)$$

where

$$\begin{aligned} G^\circ(A; x) &= \min\{G_1(A; x), G_2(A; x)\}, \\ G^\bullet(A; x) &= \min\{G^5(A^{-1}; x), G^6(A^{-1}; x)\}. \end{aligned} \quad (4.49)$$

Proof. The proof follows from that the conclusions in Corollaries 4.4 and 4.5 are independent. \square

Remark 4.8. It is easy to see that if $\lambda_1 > 0$, $\lambda_n > 0$, our result coincides with the inequality of operator versions in [6]. So we conclude that our results give an improvement of the Kantorovich inequality [6] that applies to all invertible Hermite matrices.

5. Conclusion

In this paper, we introduce some new Kantorovich-type inequalities for the invertible Hermitian matrices. Inequalities (4.27) and (4.31) are the same as [4], but our proof is simple. In Theorem 4.6, if $\lambda_1 > 0$, $\lambda_n > 0$, the results are similar to the well-known Kantorovich-type inequalities for operators in [6]. Moreover, for any invertible Hermitian matrix, there exists a similar inequality.

Acknowledgments

The authors are grateful to the associate editor and two anonymous referees whose comments have helped to improve the final version of the present work. This work was supported by the Natural Science Foundation of Chongqing Municipal Education Commission (no. KJ091104).

References

- [1] R. A. Horn and C. R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, UK, 1985.
- [2] A. S. Householder, *The Theory of Matrices in Numerical Analysis*, Blaisdell, New York, NY, USA, 1964.
- [3] S. Liu and H. Neudecker, "Several matrix Kantorovich-type inequalities," *Journal of Mathematical Analysis and Applications*, vol. 197, no. 1, pp. 23–26, 1996.
- [4] A. W. Marshall and I. Olkin, "Matrix versions of the Cauchy and Kantorovich inequalities," *Aequationes Mathematicae*, vol. 40, no. 1, pp. 89–93, 1990.
- [5] J. K. Baksalary and S. Puntanen, "Generalized matrix versions of the Cauchy-Schwarz and Kantorovich inequalities," *Aequationes Mathematicae*, vol. 41, no. 1, pp. 103–110, 1991.
- [6] S. S. Dragomir, "New inequalities of the Kantorovich type for bounded linear operators in Hilbert spaces," *Linear Algebra and its Applications*, vol. 428, no. 11-12, pp. 2750–2760, 2008.
- [7] P. G. Spain, "Operator versions of the Kantorovich inequality," *Proceedings of the American Mathematical Society*, vol. 124, no. 9, pp. 2813–2819, 1996.
- [8] S. Liu and H. Neudecker, "Kantorovich inequalities and efficiency comparisons for several classes of estimators in linear models," *Statistica Neerlandica*, vol. 51, no. 3, pp. 345–355, 1997.
- [9] P. Bloomfield and G. S. Watson, "The inefficiency of least squares," *Biometrika*, vol. 62, pp. 121–128, 1975.
- [10] L. V. Kantorovič, "Functional analysis and applied mathematics," *Uspekhi Matematicheskikh Nauk*, vol. 3, no. 6(28), pp. 89–185, 1948 (Russian).
- [11] C. G. Khatri and C. R. Rao, "Some extensions of the Kantorovich inequality and statistical applications," *Journal of Multivariate Analysis*, vol. 11, no. 4, pp. 498–505, 1981.
- [12] C. T. Lin, "Extrema of quadratic forms and statistical applications," *Communications in Statistics A*, vol. 13, no. 12, pp. 1517–1520, 1984.
- [13] A. Galántai, "A study of Auchmuty's error estimate," *Computers & Mathematics with Applications*, vol. 42, no. 8-9, pp. 1093–1102, 2001.
- [14] P. D. Robinson and A. J. Wathen, "Variational bounds on the entries of the inverse of a matrix," *IMA Journal of Numerical Analysis*, vol. 12, no. 4, pp. 463–486, 1992.
- [15] S. S. Dragomir, "A generalization of Grüss's inequality in inner product spaces and applications," *Journal of Mathematical Analysis and Applications*, vol. 237, no. 1, pp. 74–82, 1999.
- [16] S. S. Dragomir, "Some Grüss type inequalities in inner product spaces," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 2, article 42, 2003.
- [17] S. S. Dragomir, "On Bessel and Grüss inequalities for orthonormal families in inner product spaces," *Bulletin of the Australian Mathematical Society*, vol. 69, no. 2, pp. 327–340, 2004.
- [18] S. S. Dragomir, "Reverses of Schwarz, triangle and Bessel inequalities in inner product spaces," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 5, no. 3, article 76, 2004.
- [19] S. S. Dragomir, "Reverses of the Schwarz inequality generalising a Klamkin-McLenaghan result," *Bulletin of the Australian Mathematical Society*, vol. 73, no. 1, pp. 69–78, 2006.
- [20] Z. Liu, L. Lu, and K. Wang, "On several matrix Kantorovich-type inequalities," *Journal of Inequalities and Applications*, vol. 2010, Article ID 571629, 5 pages, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

