

## Research Article

# Bernstein Widths of Some Classes of Functions Defined by a Self-Adjoint Operator

**Guo Feng**

*School of Mathematics and Information Engineering, Taizhou University, Taizhou 317000, China*

Correspondence should be addressed to Guo Feng, [gfgeng@tzc.edu.cn](mailto:gfgeng@tzc.edu.cn)

Received 9 May 2011; Accepted 2 August 2011

Academic Editor: Kai Diethelm

Copyright © 2012 Guo Feng. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the classes of periodic functions with formal self-adjoint linear differential operators  $W_p(\mathcal{L}_r)$ , which include the classical Sobolev class as its special case. Using the iterative method of Buslaev, with the help of the spectrum of linear differential equations, we determine the exact values of Bernstein width of the classes  $W_p(\mathcal{L}_r)$  in the space  $L_q$  for  $1 < p \leq q < \infty$ .

## 1. Introduction and Main Result

Let  $\mathbb{C}$ ,  $\mathbb{R}$ ,  $\mathbb{Z}$ ,  $\mathbb{N}$ , and  $\mathbb{N}^+$  be the sets of all complex numbers, real numbers, integers, nonnegative integers, and positive integers, respectively.

Let  $\mathbb{T}$  be the unit circle realized as the interval  $[0, 2\pi]$  with the points 0 and  $2\pi$  identified, and as usual, let  $L_q := L_q[0, 2\pi]$  be the classical Lebesgue integral space of  $2\pi$ -periodic real-valued functions with the usual norm  $\|\cdot\|_q$ ,  $1 \leq q \leq \infty$ . Denote by  $\widetilde{W}_p^r$  the Sobolev space of functions  $x(\cdot)$  on  $\mathbb{T}$  such that the  $(r-1)$ st derivative  $x^{(r-1)}(\cdot)$  is absolutely continuous on  $\mathbb{T}$  and  $x^{(r)}(\cdot) \in L_p$ ,  $r \in \mathbb{N}$ . The corresponding Sobolev class is the set

$$W_p^r := \left\{ x(\cdot) : x \in \widetilde{W}_p^r, \left\| x^{(r)}(\cdot) \right\|_p \leq 1 \right\}. \quad (1.1)$$

Tikhomirov [1] introduced the notion of Bernstein width of a centrally symmetric set  $C$  in a normed space  $X$ . It is defined by the formula

$$b_n(C, X) := \sup_L \sup \{ \lambda \geq 0 : L \cap \lambda BX \subset C \}, \quad (1.2)$$

where  $BX$  is the unit ball of  $X$  and the outer supremum is taken over all subspaces  $L \subset X$  such that  $\dim L \geq n + 1$ ,  $n \in \mathbb{N}$ .

In particular, Tikhomirov posed the problem of finding the exact value of  $b_n(C; X)$ , where  $C = W_p^r$  and  $X = L_q$ ,  $1 \leq p, q \leq \infty$ . He also obtained the first results [1] for  $p = q = \infty$  and  $n = 2k - 1$ . Pinkus [2] found  $b_{2n-1}(W_p^r; L_q)$ , where  $p = q = 1$ . Later, Magaril-Il'yaev [3] obtained the exact value of  $b_{2n-1}(W_p^r; L_q)$  for  $1 < p = q < \infty$ . The latest contribution to these fields is due to Buslaev et al. [4] who found the exact values of  $b_{2n-1}(W_p^r; L_q)$  for all  $1 < p \leq q < \infty$ .

Let

$$\mathcal{L}_r(D) = D^r + a_{r-1}D^{r-1} + \dots + a_1D + a_0, \quad D = \frac{d}{dt}, \quad (1.3)$$

be an arbitrary linear differential operator of order  $r$  with constant real coefficients  $a_0, a_1, \dots, a_{r-1}$ . Denote by  $p_r$  the characteristic polynomial of  $\mathcal{L}_r(D)$ . The linear differential operator  $\mathcal{L}_r(D)$  will be called formal self-adjoint if  $p_r(-t) = (-1)^r p_r(t)$  for each  $t \in \mathbb{C}$ .

We define the function classes  $W_p(\mathcal{L}_r)$  as follows:

$$W_p(\mathcal{L}_r) = \left\{ x(\cdot) : x^{r-1} \in AC_{2\pi}, \|\mathcal{L}_r(D)x(\cdot)\|_p \leq 1 \right\}, \quad (1.4)$$

where  $1 \leq p \leq \infty$ .

In this paper, we consider some classes of periodic functions with formal self-adjoint linear differential operators  $W_p(\mathcal{L}_r)$ , which include the classical Sobolev class as its special case. Using the iterative method of Buslaev and Tikhomirov [5], with the help of the spectrum of linear differential equations, we determine the exact values of Bernstein width of the classes  $W_p(\mathcal{L}_r)$  in the space  $L_q$  for  $1 < p \leq q < \infty$ . The results of Buslaev et al. [4] are extended to the classes (1.4) defined by differential operators (1.3).

We define  $Q_q$  to be the nonlinear transformation

$$(Q_q f)(t) := |f(t)|^{q-1} \operatorname{sign} f(t). \quad (1.5)$$

The main result of this paper is the following.

**Theorem 1.1.** *Let  $\mathcal{L}_r(D)$  be an arbitrary formal self-adjoint linear differential operator given by (1.3), and  $n, r \in \mathbb{N}$ ,  $1 < p \leq q < \infty$ . Then*

$$b_{2n-1}(W_p(\mathcal{L}_r); L_q) = \lambda_{2n} := \lambda_{2n}(p, q, \mathcal{L}_r), \quad (1.6)$$

where  $\lambda_{2n}$  is that eigenvalue  $\lambda$  of the boundary value problem

$$\begin{aligned} \mathcal{L}_r(D)y(t) &= (-1)^r \lambda^{-q} (Q_q x)(t), \\ y(t) &= (Q_p \mathcal{L}_r(D)x)(t), \\ x^{(j)}(0) &= x^{(j)}(2\pi), \quad y^{(j)}(0) = y^{(j)}(2\pi), \quad j = 0, 1, \dots, n-1, \end{aligned} \quad (1.7)$$

for which the corresponding eigenfunction  $x(\cdot) = x_{2n}(\cdot)$  has only  $2n$  simple zeros on  $\mathbb{T}$  and is normalized by the condition  $\|\mathcal{L}_r(D)x(\cdot)\|_p = 1$ .

## 2. Proof of the Theorem

First we introduce some notations and formulate auxiliary statements.

Let  $\mathcal{L}_r(D)$  be an arbitrary linear differential operator (1.3). Denote the  $2\pi$ -periodic kernel of  $\mathcal{L}_r(D)$  by

$$\text{Ker } \mathcal{L}_r(D) = \{x(\cdot) \in C^r(\mathbb{T}) : \mathcal{L}_r(D)x(t) \equiv 0\}. \quad (2.1)$$

Let  $\mu(0 \leq \mu \leq r)$  be the dimension of  $\text{Ker } \mathcal{L}_r(D)$  and  $\{\varphi_1, \dots, \varphi_\mu\}$  an arbitrary basis in  $\text{Ker } \mathcal{L}_r(D)$ .

$Z_c(f)$  denotes the number of zeros of  $f$  on a period, counting multiplicity, and  $S_c(f)$  is the cyclic sign change count for a piecewise continuous,  $2\pi$ -Periodic function  $f$  [2]. Following,  $(x(\cdot), \lambda)$  is called the spectral pair of (1.7) if the function  $x(\cdot)$  is normalized by the condition  $\|\mathcal{L}_r(D)x(\cdot)\|_p = 1$ . The set of all spectral pairs is denoted by  $\text{SP}(p, q, \mathcal{L}_r)$ . Define the spectral classes  $\text{SP}_{2k}(p, q, \mathcal{L}_r)$  as

$$\text{SP}_{2k}(p, q, \mathcal{L}_r) = \{(x(\cdot), \lambda) \in \text{SP}(p, q, \mathcal{L}_r) : S_c(x(\cdot)) = 2k\}. \quad (2.2)$$

Let  $\widehat{x}_{2n}(\cdot)$  be the solution of the extremal problem as follows:

$$\begin{aligned} & \int_0^{\pi/2n} |X(t)|^q dt \longrightarrow \sup, \\ & \int_0^{\pi/2n} |\mathcal{L}_r(D)X(t)|^p dt \leq 1, \\ & x^{(k)}\left(\frac{\left((\pi/2n) + (-1)^{k+1}(\pi/2n)\right)}{2}\right) = 0, \quad k = 0, 1, \dots, n-1 \end{aligned} \quad (2.3)$$

and the function  $x_{2n}(\cdot)$  is such that  $x_{2n}(t) = -x_{2n}(t - \pi/n)$  for all  $t \in \mathbb{T}$

$$x_{2n}(t) := \begin{cases} \widehat{x}_{2n}(t), & 0 \leq t \leq \frac{\pi}{2n}, \\ \widehat{x}_{2n}\left(\frac{\pi}{n} - t\right), & \frac{\pi}{2n} < t \leq \frac{\pi}{n}. \end{cases} \quad (2.4)$$

Let us extend periodically the function  $x_{2n}(t)$  onto  $\mathbb{R}$  and normalize the obtained function as it is required in the definition of spectral pairs. From what has been done above, we get a function  $x_{2n}(t)$  belonging to  $\text{SP}_{2n}(p, q, \mathcal{L}_r)$ . Furthermore, by [6], which any other function from  $\text{SP}_{2n}(p, q, \mathcal{L}_r)$  differs from  $x_{2n}(\cdot)$  only in the sign and in a shift of its argument, and there exists a number  $N \in \mathbb{N}^+$  such that for every  $n \geq N$ , all zeros of  $x_{2n}(\cdot)$  are simple, equidistant

with a step equal to  $\pi/n$ , and  $S_c(x_{2n}) = S_c(\mathcal{L}_r(D)x_{2n}) = 2n$ . We denote the set of zeros (= sign variations) of  $\mathcal{L}_r(D)x_{2n}$  on the period by  $Q_{2n} = (\tau_1, \dots, \tau_{2n})$ . Let

$$G_r(t) = \frac{1}{2\pi} \sum_{k \notin \Lambda} \frac{e^{ikt}}{p_r(ik)}, \quad (2.5)$$

where  $\Lambda = \{k \in \mathbb{Z} : p_r(ik) = 0\}$  and  $i$  is the imaginary unit.

The  $2\pi$ -periodic G-splines are defined as elements of the linear space

$$S(Q_{2n}, G_r) = \text{span}\{\varphi_1(t), \dots, \varphi_\mu(t), G_r(t - \tau_1), \dots, G_r(t - \tau_{2n})\}. \quad (2.6)$$

As was proved in [7], if  $n \geq N$ , then  $\dim S(Q_{2n}, G_r) = 2n$ .

We assume (shifting  $x(\cdot)$  if necessary) that  $\mathcal{L}_r(D)\hat{x}_{2n}(\cdot)$  is positive on  $(-\pi, \pi + \pi/n)$ . Let  $L_{2n} := L_{2n}(r, p, q)$  denote the space of functions of the form

$$x(t) = \sum_{j=1}^{\mu} a_j \varphi_j(t) + \frac{1}{\pi} \int_{\mathbb{T}} G_r(t - \tau) \left( \sum_{i=1}^{2n} b_i y_i(\tau) \right) d\tau, \quad (2.7)$$

where  $a_1, \dots, a_\mu, b_1, \dots, b_{2n} \in \mathbb{R}$ ,  $\sum_{i=1}^{2n} b_i = 0$ ,  $y_i(\cdot) = \chi_i(\cdot) \mathcal{L}_r(D)x_{2n}(\cdot - (i-1)\pi/n)$ , and  $\chi_i(\cdot)$  is the characteristic function of the interval  $\Delta_i := [-\pi + (i-1)\pi/n, -\pi + i\pi/n]$ ,  $1 \leq i \leq 2n$ . Obviously,  $\dim L_{2n} = 2n$  and  $L_{2n} \subset W_p(\mathcal{L}_r)$ .

Let us now consider exact estimate of Bernstein  $n$ -width. This was introduced in [1]. We reformulate the definition for a linear operator  $P$  mapping  $X$  to  $Y$ .

*Definition 2.1* (see [2, page 149]). Let  $P \in L(X, Y)$ . Then the Bernstein  $n$ -width is defined by

$$b_n(P(X), Y) = \sup_{X_{n+1}} \inf_{\substack{Px \in X_{n+1} \\ Px \neq 0}} \frac{\|Px\|_Y}{\|x\|_X}, \quad (2.8)$$

where  $X_{n+1}$  is any subspace of  $\text{span}\{Px : x \in X\}$  of dimension  $\geq n+1$ .

## 2.1. Lower Estimate of Bernstein $n$ -Width

Consider the extremal problem

$$\frac{\|x(\cdot)\|_q^q}{\|\mathcal{L}_r(D)x(\cdot)\|_p^p} \longrightarrow \inf, \quad x(\cdot) \in L_{2n} \quad (2.9)$$

and denote the value of this problem by  $\alpha^q$ . Let us show that  $\alpha \geq \lambda_n$ ; this will imply the desired lower bound for  $b_{2n-1}$ . Let  $x(\cdot) \in L_{2n}$ , then

$$\|\mathcal{L}_r(D)x(\cdot)\|_p^p = \sum_{i=1}^{2n} \int_{\Delta_i} \left| \sum_{i=1}^{2n} b_i y_i(t) \right|^p dt = \sum_{i=1}^{2n} \int_{\Delta_i} |b_i|^p |\mathcal{L}_r(D)x_n(t)|^p dt = \frac{1}{2n} \sum_{i=1}^{2n} |b_i|^p \quad (2.10)$$

and by setting

$$z_i(\cdot) := \frac{1}{\mathcal{I}} \int_{\mathbb{T}} G_r(\cdot - \tau) y_i(\tau) d\tau, \quad i = 1, 2, \dots, 2n, \quad (2.11)$$

we reduce problem (2.9) to the form

$$\frac{\left\| \sum_{j=1}^{\mu} a_j \varphi_j(\cdot) + \sum_{i=1}^{2n} b_i z_i(\cdot) \right\|_q^q}{1/2n \sum_{i=1}^{2n} |b_i|^p} \rightarrow \inf, \quad a_1, \dots, a_{\mu}, b_1, \dots, b_{2n} \in \mathbb{R}. \quad (2.12)$$

This is a smooth finite-dimensional problem. It has a solution  $(\bar{a}_1, \dots, \bar{a}_{\mu}, \bar{b}_1, \dots, \bar{b}_{2n})$  and,  $(\bar{b}_1, \dots, \bar{b}_{2n}) \neq \vec{0}$ . According to the Lagrange multiplier rule, there exists a  $\eta \in \mathbb{R}$  such that the derivatives of the function  $(a_1, \dots, a_{\mu}, b_1, \dots, b_{2n}) \rightarrow g(a_1, \dots, a_{\mu}, b_1, \dots, b_{2n}) + \eta(b_1 + b_2 + \dots + b_{2n})$  (where  $g(\cdot)$  is the function being minimized in (2.12)) with respect to  $a_1, \dots, a_{\mu}, b_1, \dots, b_{2n}$  at the point  $(\bar{a}_1, \dots, \bar{a}_{\mu}, \bar{b}_1, \dots, \bar{b}_{2n})$  are equal to zero. This leads to the relations

$$\int_{\mathbb{T}} \varphi_j(t) (Q_q \bar{x})(t) dt = 0, \quad j = 1, \dots, \mu, \quad (2.13)$$

$$\int_{\mathbb{T}} z_i(t) (Q_q \bar{x})(t) dt = \frac{1}{2n} \frac{\|\bar{x}(\cdot)\|_q^q}{\|\mathcal{L}_r(D)\bar{x}(\cdot)\|_p^p} Q_p \bar{b}_i, \quad i = 1, \dots, 2n, \quad (2.14)$$

where  $\bar{x}(\cdot) = \sum_{j=1}^{\mu} \bar{a}_j \varphi_j(t) + \sum_{i=1}^{2n} \bar{b}_i z_i(\cdot)$ .

We remark that  $g(a_1, \dots, a_{\mu}, b_1, \dots, b_{2n}) = g(da_1, \dots, da_{\mu}, db_1, \dots, db_{2n})$  for any  $d \neq 0$ , and hence the vector  $(d\bar{a}_1, \dots, d\bar{a}_{\mu}, d\bar{b}_1, \dots, d\bar{b}_{2n})$  is also a solution of (2.12). Thus, it can be assumed that  $|\bar{b}_i| \leq 1, i = 1, \dots, 2n$  and  $\bar{b}_{i_0} = (-1)^{i_0+1}$  for some  $i_0, 1 \leq i_0 \leq 2n$ .

Let

$$\tilde{x}_{2n}(t) = \sum_{j=1}^{\mu} a_j^* \varphi_j(t) + \sum_{i=1}^{2n} (-1)^{i+1} z_i(t) \quad (2.15)$$

and  $\tilde{x}_{2n}$  satisfies (1.7). Let  $a^* = (a_1^*, \dots, a_{2n}^*)$  and let  $b^* = (1, -1, \dots, 1, -1) \in \mathbb{R}^{2n}$ . It follows from the definitions of  $\tilde{x}_{2n}(\cdot)$  and  $\bar{x}(\cdot)$  that

$$\mathcal{L}_r(D)\tilde{x}_{2n}(t) - \mathcal{L}_r(D)\bar{x}(t) = \sum_{\substack{i=1 \\ i \neq i_0}}^{2n} \left( (-1)^{i+1} - \bar{b}_i \right) \chi_i(t) \mathcal{L}_r(D)x_{2n} \left( t - \frac{(i-1)\mathcal{T}}{n} \right), \quad (2.16)$$

and hence  $S_c(\mathcal{L}_r(D)\tilde{x}_{2n}(\cdot), \mathcal{L}_r(D)\bar{x}(\cdot))$  has at most  $2n - 2$  sign changes. Then, by Rolle's theorem,  $S_c(\mathcal{L}_r(D)\tilde{x}_{2n}(\cdot) - \mathcal{L}_r(D)\bar{x}(\cdot)) \leq 2n - 2$ . For any  $a, b \in \mathbb{R}$ ,  $\text{sign}(a+b) = \text{sign}(Q_p a + Q_p b)$ ; therefore

$$S_c((Q_q \tilde{x}_{2n})(\cdot) - (Q_q \bar{x})(\cdot)) \leq 2n - 2. \quad (2.17)$$

In addition, since  $\tilde{x}_{2n}$  is  $2\pi$ -periodic solution of the linear differential equation,  $\mathcal{L}_r(D)y(t) = (-1)^r \lambda^{-q} (Q_q x)(t)$  and  $\varphi_j(t) \in \text{Ker } \mathcal{L}_r(D)$ . Then, by [8, page 94], we have

$$\int_{\mathbb{T}} \varphi_j(t) (Q_q \tilde{x})(t) dt = 0, \quad j = 1, \dots, \mu. \quad (2.18)$$

If we now multiply both sides of (2.15) by  $(Q_q \tilde{x}_{2n})(t)$  and integrate over the interval  $\Delta_i$ ,  $1 \leq i \leq 2n$ , we get

$$\int_{\Delta_i} z_i(t) (Q_q \tilde{x}_{2n})(t) dt = (-1)^{i+1} \int_{\Delta_i} |\tilde{x}_{2n}(t)|^q dt = (-1)^{i+1} \frac{\lambda_{2n}^q}{2n}, \quad (2.19)$$

due to  $\int_{\mathbb{T}} z_i(t) (Q_q \tilde{x}_{2n})(t) dt = \int_{\Delta_i} z_i(t) (Q_q \tilde{x}_{2n})(t) dt$ . Therefore, we have

$$\int_{\mathbb{T}} z_i(t) (Q_q \tilde{x}_{2n})(t) dt = (-1)^{i+1} \frac{\lambda_{2n}^q}{2n}, \quad i = 1, \dots, 2n. \quad (2.20)$$

Changing the order of integration and using (2.14) and (2.20), we get that

$$\begin{aligned} & \int_{\Delta_i} \mathcal{L}_r(D)x_{2n} \left( t - \frac{(i-1)\pi}{n} \right) \left( \frac{1}{\pi} \int_{\mathbb{T}} G_r(t-\tau) ((Q_q \tilde{x}_{2n})(\tau) - (Q_q \bar{x})(\tau)) d\tau \right) dt \\ &= \int_{\mathbb{T}} z_i(t) ((Q_q \tilde{x}_{2n})(t) - (Q_q \bar{x})(t)) dt = \frac{(-1)^r}{2n} \left( (-1)^{i+1} \lambda_{2n}^q - \frac{\|x(\cdot)\|_q^q}{\|(\mathcal{L}_r(D)\bar{x})(\cdot)\|_p^p} Q_p \bar{b}_i \right) \\ &= \frac{(-1)^r}{2n} \left( (-1)^{i+1} \lambda_{2n}^q - \alpha^q \|(\mathcal{L}_r(D)\bar{x})(\cdot)\|^{q-p} Q_p \bar{b}_i \right), \quad i = 1, \dots, 2n. \end{aligned} \quad (2.21)$$

Denote by  $f(\cdot)$  the factor multiply  $\mathcal{L}_r(D)x_{2n}(t - (i-1)\pi/n)$  in the integral in the left-hand side of this equality. Since  $\|(\mathcal{L}_r(D)\bar{x})(\cdot)\|_p \leq 1$  and hence  $\|(\mathcal{L}_r(D)\bar{x})(\cdot)\|_p^{q-p} \leq 1$  for  $p \leq q$ , if we assume that  $\lambda_{2n} > \alpha$ , then we arrive at the relations

$$\text{sign} \int_{\Delta_i} \mathcal{L}_r(D)x_{2n} \left( t - \frac{(i-1)\pi}{n} \right) f(\cdot) dt = (-1)^{r+i+1}, \quad i = 1, \dots, 2n. \quad (2.22)$$

Suppose for definiteness that  $\mathcal{L}_r(D)x_{2n}(t - (i-1)\pi/n) > 0$  interior to  $\Delta_i$ ,  $i = 1, \dots, 2n$ . Then it follows from (2.22) that there are points  $t_i \in \Delta_i$  such that  $\text{sign } f(t_i) = (-1)^{i+1}$ ,  $i = 1, \dots, 2n$ , that is,  $S_c(f(\cdot)) \geq 2n - 1$ . But  $f(\cdot)$  is periodic, and hence  $S_c(f(\cdot)) \geq 2n$ ; therefore,  $S_c(\mathcal{L}_r(D)f(\cdot)) \geq 2n$ . Further,  $\mathcal{L}_r(D)f(\cdot) = (Q_q \tilde{x}_{2n})(t) - (Q_q \bar{x})(t)$ , that is,  $S_c((Q_q \tilde{x}_{2n})(t) - (Q_q \bar{x})(t)) \geq 2n$ .

We have arrived at a contradiction to (2.17), and hence  $\lambda_{2n} \leq \alpha$ . Thus  $b_{2n-1}(W_p(\mathcal{L}_r); L_q) \geq \lambda_{2n}$ .

## 2.2. Upper Estimate of Bernstein $n$ -Width

Assume the contrary:  $b_{2n-1}(W_p(\mathcal{L}_r); L_q) > \lambda_{2n}$ , ( $1 < p \leq q < \infty$ ). Then, by definition, there exists a linearly independent system of  $2n$  functions  $L_{2n} := \text{span}\{f_1, \dots, f_{2n}\} \subset L_q$  and number  $\gamma > \lambda_{2n}$  such that  $L_{2n} \cap \gamma S(L_q) \subseteq \mathcal{L}_r(D)$ , or equivalently,

$$\min_{x(\cdot) \in L_{2n}} \frac{\|x(\cdot)\|_q}{\|\mathcal{L}_r(D)x(\cdot)\|_p} \geq \gamma > \lambda_{2n}. \quad (2.23)$$

Let us assign a vector  $c \in \mathbb{R}^{2n}$  to each function  $x(\cdot) \in L_{2n}$  by the following rule:

$$x(\cdot) \longrightarrow c = (c_1, \dots, c_{2n}) \in \mathbb{R}^{2n}, \quad \text{where } x(\cdot) = \sum_{j=1}^{2n} c_j f_j(\cdot). \quad (2.24)$$

Then inequality (2.23) acquires the form

$$\min_{c \in \mathbb{R}^{2n} \setminus \{0\}} \frac{\left\| \sum_{j=1}^{2n} c_j f_j(\cdot) \right\|_q}{\left\| \sum_{j=1}^{2n} c_j \mathcal{L}_r(D) f_j(\cdot) \right\|_p} \geq \gamma > \lambda_{2n}. \quad (2.25)$$

Let  $c_0 = 0$ . Consider the sphere  $S^{2n-1}$  in the space  $\mathbb{R}^{2n}$  with radius  $2\pi$ , that is,

$$S^{2n-1} := \left\{ c : c = (c_1, \dots, c_{2n}) \in \mathbb{R}^{2n}, \|c\| = \sum_{j=1}^{2n} |c_j| = 2\pi \right\}. \quad (2.26)$$

To every vector  $c \in \mathbb{R}^{2n}$  we assign function  $u(t, c)$  defined by

$$u(t, c) = \begin{cases} (2\pi)^{-1/p} \text{sign } c_k, & \text{for } t \in (t_{k-1}, t_k), \quad k = 1, \dots, 2n, \\ 0, & \text{for } t = t_k, \quad k = 1, \dots, 2n-1, \end{cases} \quad (2.27)$$

where  $t_0 = 0$ ,  $t_k = \sum_{i=1}^k |c_i|$ ,  $k = 1, \dots, 2n$ , and the extended  $2\pi$ -periodically onto  $\mathbb{R}$ .

An analog of the Buslaev iteration process [5] is constructed in the following way: the function  $x(t, c)$  is found as a periodic solution of the linear differential equation  $\mathcal{L}_r(D)x_0 = u$ ; then the periodic functions  $\{x_k(t, c)\}_{k \in \mathbb{N}^+}$  are successively determined from the differential equations

$$\begin{aligned} \mathcal{L}_r(D)x_k(t) &= (Q_p y_k)(t), \\ \mathcal{L}_r(D)y_k(t) &= (-1)^r \mu_{k-1}^{-q} (Q_q x_{k-1})(t), \end{aligned} \quad (2.28)$$

where  $p' = p/(p - 1)$ , and the constants  $\{\mu_k : k = 0, \dots\}$  are uniquely determined by the conditions

$$\|\mathcal{L}_r(D)x_k\|_p = 1, \quad (Q_q x_k)(t) \perp \text{Ker } \mathcal{L}_r(D), \quad (Q_{p'} y_k)(t) \perp \text{Ker } \mathcal{L}_r(D). \quad (2.29)$$

By analogy with the reasoning in [5], we can prove the following assertions.

- (i) The iteration procedure (2.28)-(2.29) is well defined; the sequences  $\{\mu_k\}_{k \in \mathbb{N}}$  are monotone nondecreasing and converge to an eigenvalue  $\lambda(c) > 0$  of the problem (1.7).
- (ii) The sequence  $\{x_k(\cdot, c)\}_{k \in \mathbb{N}}$  has a subsequence that is convergent to an eigenfunction  $x(\cdot, c)$  of the problem (1.7), with  $\lambda(c) = \|x(\cdot, c)\|_p$ .
- (iii) For any  $k \in \mathbb{N}$  there exists a  $\hat{c} \in S^{2n-1}$  such that  $x_k(\cdot, \hat{c})$  has at least  $2n$  zeros ( $Z_c(x_k(\cdot, \hat{c})) \geq 2n$ ) on  $\mathbb{T}$ .
- (iv) In the set of spectral pairs  $(\lambda(c), x(\cdot, c))$ , there exists a pair  $(\lambda(\hat{c}), x(\cdot, \hat{c}))$  such that  $S_c(x(\cdot, \hat{c})) = 2N \geq 2n$ .

Items (i) and (ii) can be proved in the same way as Lemmas 1 and 2 of [5, Sections 6 and 10]. Item (iii) follows from the Borsuk theorem [10], which states that there exists a  $\hat{c} \in S^{2n-1}$  such that  $Z_c(x_k(\cdot, \hat{c})) \geq 2n - 1$ , but since the function  $x_k(\cdot, \hat{c})$  is periodic, we actually have  $Z_c(x_k(\cdot, \hat{c})) \geq 2n$ . Finally, in item (iv), by (ii) and (iii),  $Z_c(x(\cdot, \hat{c})) \geq 2n$ . In view of  $x(\cdot, \hat{c})$  zeros are simple; therefore,  $S_c(x(\cdot, \hat{c})) \geq 2n$ .

Note that [8] the linear differential equation  $\mathcal{L}_r(D)f = g$  has a  $2\pi$ -periodic solution if and only if  $\int_{\mathbb{T}} g(t)v(t)dt = 0$ , where  $v(\cdot) \in \text{Ker } \mathcal{L}_r(D)$  and  $g$  is an integrable  $2\pi$ -periodic function. Using the method similar to [5, 11], it is not difficult to show that spectral pairs of (1.7) are unique and spectral value  $\lambda_n$  is monotone decreasing for  $n$ ; it follows that

$$\lambda(\hat{c}) = \lambda_{2N} \leq \lambda_{2n}. \quad (2.30)$$

Therefore, by virtue of items (i), (ii), and (2.30), we obtain

$$\begin{aligned} \min_{c \in \mathbb{R}^{2n} \setminus \{0\}} \frac{\left\| \sum_{j=1}^{2n} c_j f_j(\cdot) \right\|_q}{\left\| \sum_{j=1}^{2n} c_j \mathcal{L}_r(D) f_j(\cdot) \right\|_p} &\leq \frac{\left\| \sum_{j=1}^{2n} \hat{c}_j f_j(\cdot) \right\|_q}{\left\| \sum_{j=1}^{2n} \hat{c}_j \mathcal{L}_r(D) f_j(\cdot) \right\|_p} \\ &\leq \frac{\|x_k(\cdot, \hat{c})\|_q}{\|\mathcal{L}_r(D)x_k(\cdot, \hat{c})\|_p} \leq \lambda(\hat{c}) \leq \lambda_{2n}, \end{aligned} \quad (2.31)$$

which contradicts (2.25). Hence  $b_{2n-1}(W_p(\mathcal{L}_r); L_q) \leq \lambda_{2n}$ . Thus, the upper bound is proved. This completes the proof of the theorem.

## Acknowledgments

The author would like to thank Professor Kai Diethelm and the anonymous referees for their valuable comments, remarks, and suggestions which greatly help us to improve the



presentation of this paper and make it more readable. Project Supported by the Natural Science Foundation of China (Grant no. 10671019) and Scientific Research fund of Zhejiang Provincial Education Department (Grant no. 20070509).

## References

- [1] V. M. Tikhomirov, *Some Questions in Approximation Theory*, Izdat. Moskov. Univ., Moscow, Russia, 1976.
- [2] A. Pinkus, *n-Widths in Approximation Theory*, Springer, New York, NY, USA, 1985.
- [3] G. G. Magaril-Il'yaev, "Mean dimension, widths, and optimal recovery of Sobolev classes of functions on the line," *Mathematics of the USSR—Sbornik*, vol. 74, no. 2, pp. 381–403, 1993.
- [4] A. P. Buslaev, G. G. Magaril-Il'yaev, and Nguen T'en Nam, "Exact values of Bernstein widths for Sobolev classes of periodic functions," *Matematicheskie Zametki*, vol. 58, no. 1, pp. 139–143, 1995 (Russian).
- [5] A. P. Buslaev and V. M. Tikhomirov, "Spectra of nonlinear differential equations and widths of Sobolev classes," *Mathematics of the USSR—Sbornik*, vol. 71, no. 2, pp. 427–446, 1992.
- [6] S. I. Novikov, "Exact values of widths for some classes of periodic functions," *The East Journal on Approximations*, vol. 4, no. 1, pp. 35–54, 1998.
- [7] Nguen Thi Thien Hoa, *Optimal quadrature formulae and methods for recovery on function classes defined by variation diminishing convolutions*, Candidate's Dissertation, Moscow State University, Moscow, Russia, 1985.
- [8] V. A. Jakubovitch and V. I. Starzhinski, *Linear Differential Equations with Periodic Coefficients and Its Applications*, Nauka, Moscow, Russia, 1972.
- [9] A. Pinkus, "*n*-widths of Sobolev spaces in  $L^p$ ," *Constructive Approximation*, vol. 1, no. 1, pp. 15–62, 1985.
- [10] K. Borsuk, "Drei Sätze über die *n*-dimensionale euklidische Sphäre," *Fundamenta Mathematicae*, vol. 20, pp. 177–190, 1933.
- [11] A. P. Buslaev and V. M. Tikhomirov, "Some problems of nonlinear analysis and approximation theory," *Soviet Mathematics—Doklady*, vol. 283, no. 1, pp. 13–18, 1985.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

