



# A STUDY OF THE REAL HARDY INEQUALITY

**MOHAMMAD SABABHEH**

Department of Science and Arts  
Princess Sumaya University for Technology  
Amman 11941 Jordan  
EMail: [sababbeh@psut.edu.jo](mailto:sababbeh@psut.edu.jo)

*Received:* 11 November, 2008

*Accepted:* 20 October, 2009

*Communicated by:* [L. Pick](#)

*2000 AMS Sub. Class.:* 42A16, 42A05, 42B30.

*Key words:* Hardy's inequality, inequalities involving Fourier transforms.

*Abstract:* We show that some Hardy-type inequalities on the circle can be proved to be true on the real line. Namely, we discuss the idea of getting Hardy inequalities on the real line by the use of corresponding inequalities on the circle. In the last section, we prove the truth of a certain open problem under some restrictions.

---

Real Hardy Inequality  
Mohammad Sababbeh  
vol. 10, iss. 4, art. 104, 2009

---

[Title Page](#)

[Contents](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

Page 1 of 18

[Go Back](#)

[Full Screen](#)


[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756

# Contents

1	Introduction	3
2	Setup	5
3	From The Circle To The Line	7
4	Another Hardy Inequality	10



---

**Real Hardy Inequality**

Mohammad Sababbeh

vol. 10, iss. 4, art. 104, 2009

---

Title Page

Contents



Page 2 of 18


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Title Page

Contents



Page 3 of 18

Go Back

Full Screen

Close

## 1. Introduction

When McGehee, Pigno and Smith [4] proved the Littlewood conjecture, many questions regarding the best possible generalization of Hardy's inequality were asked. The longstanding question is: Does there exist a constant  $c > 0$  such that

$$(1.1) \quad \sum_{n=1}^{\infty} \frac{|\hat{f}(n)|}{n} \leq c \|f\|_1 + c \sum_{n=1}^{\infty} \frac{|\hat{f}(-n)|}{n}$$

for all  $f \in L^1(\mathbb{T})$ ? The truth of this inequality is an open problem. Many attempts were made to answer this question and many partial results were obtained. We refer the reader to [2], [3], [6] and [7] for some partial results.

Almost all articles in the literature treat Hardy-type inequalities on the circle and a very few articles treat them on the real line.

In [8] it was proved that a constant  $c > 0$  exists such that for all

$$f \in \left\{ g \in L^1(\mathbb{R}) : \int_{-\infty}^x g(t) dt \in L^1(\mathbb{R}) \right\}$$


we have

$$\int_0^{\infty} \frac{|\hat{f}(\xi)|^2}{\xi} d\xi \leq c \|f\|_1^2 + c \int_0^{\infty} \frac{|\hat{f}(-\xi)|^2}{\xi} d\xi.$$

Although this is not the first proved Hardy inequality on the real line, its proof is the first proof which uses the construction of a bounded function on  $\mathbb{R}$  whose Fourier coefficients have some desired decay properties.

We have two main goals in this article. The first is to prove Hardy inequalities on the real line using well known inequalities on the circle and the second is to prove a real Hardy inequality on the real line which is related to the open problem (1.1).

Proving a Hardy inequality usually involves quite a difficult construction and this is because of the way we prove such inequalities. Again we refer the reader to [2], [4], [6], [7] and [8] for more information on how and why we construct bounded functions with desired Fourier coefficients.



---

### Real Hardy Inequality

Mohammad Sababheh

vol. 10, iss. 4, art. 104, 2009

---

[Title Page](#)

[Contents](#)



Page 4 of 18


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



Title Page

Contents



Page 5 of 18

Go Back

Full Screen

Close

## 2. Setup

Let  $L^1$  denote the space of all integrable functions (equivalent classes) defined on  $\mathbb{R}$ . For  $f \in L^1$ , we define the Fourier transform of  $f$  to be

$$\hat{f}(\xi) = \int_{\mathbb{R}} f(x)e^{-ix\xi} d\xi.$$

If  $\hat{f} \in L^1$  then the inversion formula for  $f$  holds:

$$f(x) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\xi)e^{i\xi x} d\xi.$$

If  $f \in L^2$  then the Fourier transform of  $f$  is defined to be the  $L^2$ - limit:

$$\hat{f}(\xi) = \lim_{n \rightarrow \infty} \int_{-n}^n f(x)e^{-i\xi x} dx.$$

In this case  $\hat{f} \in L^2$  and


$$f(x) = \lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{-n}^n \hat{f}(\xi)e^{ix\xi} d\xi, \quad \text{in the } L^2 \text{ sense.}$$

The Plancherel theorem then says:

$$\|f\|_2 = \frac{1}{\sqrt{2\pi}} \|\hat{f}\|_2.$$

**Lemma 2.1.** For  $f, g \in L^2$  we have

$$\int_{\mathbb{R}} f(x)\overline{g(x)} dx = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\xi)\overline{\hat{g}(\xi)} d\xi.$$



Title Page

Contents



Page 6 of 18

Go Back

Full Screen

Close

We refer the reader to any standard book in analysis for more on these concepts, see for example [5].

Observe that if  $f \in L^1$  is such that  $\hat{f}$  is compactly supported, then  $\hat{f} \in L^2$  and hence  $f \in L^2$ .

Now, given  $f \in L^1(\mathbb{R})$ , define


$$(2.1) \quad \varphi_N(t) = 2\pi \sum_{j=-\infty}^{\infty} f_N(t + 2\pi j), \quad \text{where } f_N(x) = Nf(Nx).$$

Then we have

$$(2.2) \quad \varphi_N \in L^1(\mathbb{T}), \hat{\varphi}_N(n) = \hat{f}\left(\frac{n}{N}\right) \quad \text{and} \quad \lim_{N \rightarrow \infty} \|\varphi_N\|_{L^1(\mathbb{T})} = \|f\|_{L^1(\mathbb{R})}.$$

For a discussion of this idea we refer the reader to [1, p. 160-162].

Thus, the equations in (2.1) and (2.2) enable us to move from a function integrable on the line to a function integrable on the circle. This idea will be used efficiently in the next section to prove some Hardy-type inequalities on the real line using only a corresponding inequality on the circle.



Title Page

Contents



Page 7 of 18

Go Back

Full Screen

Close

### 3. From The Circle To The Line

In this section we discuss how one can use a Hardy inequality on the circle to obtain an inequality on the real line.

Recall that Hardy's inequality on the circle states that a constant  $C > 0$  exists such that for all  $f \in L^1(\mathbb{T})$  with  $\hat{f}(n) = 0, n < 0$  we have

$$\sum_{n=1}^{\infty} \frac{|\hat{f}(n)|}{n} \leq C \|f\|_1.$$

As a matter of notation, let  $H^1(\mathbb{R})$  be defined by

$$H^1(\mathbb{R}) = \left\{ f \in L^1(\mathbb{R}) : \hat{f}(\xi) = 0, \forall \xi < 0 \right\}.$$

In the following theorem we use the above stated Hardy's inequality to get the well known Hardy inequality on the line. We should remark that proving such an inequality (on the line) is a difficult task, but transforming the problem from the circle to the line greatly simplifies the proof.


**Theorem 3.1.** *There exists an absolute constant  $C > 0$  such that*

$$\int_0^{\infty} \frac{|\hat{f}(\xi)|}{\xi} \leq C \|f\|_1$$

for all  $f \in H^1(\mathbb{R})$ .

*Proof.* Let  $f \in H^1(\mathbb{R})$  be arbitrary and let, for  $N \in \mathbb{N}$ ,  $\varphi_N$  (as in (2.1) and (2.2)) be such that:

$$\varphi_N \in L^1(\mathbb{T}), \quad \lim_{N \rightarrow \infty} \|\varphi_N\|_{L^1(\mathbb{T})} = \|f\|_{L^1(\mathbb{R})} \quad \text{and} \quad \hat{\varphi}_N(n) = \hat{f}(n/N).$$



Title Page

Contents

◀ ▶

◀ ▶

Page 8 of 18

Go Back

Full Screen

Close

Clearly  $\hat{\varphi}_N(n) = 0, \forall n < 0$ , hence Hardy's inequality applies and we have

$$\sum_{n=1}^{\infty} \frac{|\hat{\varphi}_N(n)|}{n} \leq C \|\varphi_N\|_{L^1(\mathbb{T})}.$$

However, this implies that

$$\sum_{n=1}^N \frac{1}{N} \frac{|\hat{f}(n/N)|}{n/N} \leq C \|\varphi_N\|_{L^1(\mathbb{T})}.$$

We take the limit as  $N \rightarrow \infty$  to get

$$(3.1) \quad \int_0^1 \frac{|\hat{f}(\xi)|}{\xi} \leq C \|f\|_1.$$

Thus, we have shown (3.1) for any function in  $H^1(\mathbb{R})$ . Now we proceed to prove the inequality stated in the theorem. That is, we would like to replace the upper limit of the above integral by  $\infty$ . For this, let  $M \in \mathbb{N}$  be arbitrary and put  $h(x) = f(x/M)$ . Then

$$\|h\|_1 = M \|f\|_1 \quad \text{and} \quad \hat{h}(\xi) = M \hat{f}(M\xi).$$

Now apply (3.1) on  $h$  to obtain


$$\int_0^1 \frac{|\hat{h}(\xi)|}{\xi} \leq C \|h\|_1 \Rightarrow \int_0^1 \frac{|\hat{f}(M\xi)|}{\xi} d\xi \leq C \|f\|_1.$$

Put  $M\xi = t$  to get

$$(3.2) \quad \int_0^M \frac{|\hat{f}(\xi)|}{\xi} d\xi \leq C \|f\|_1.$$

Now, letting  $M$  tend to  $\infty$ , we obtain the result. □





Title Page

Contents



Page 9 of 18

Go Back

Full Screen

Close

In fact, the idea of this proof can be used to prove many Hardy inequalities on the real line!

It is proved that, see for example [3], for all functions  $f \in L^1(\mathbb{T})$  we have


$$\sum_{n=1}^{\infty} \frac{|\hat{f}(n)|^2}{n} \leq C \|f\|_1^2 + C \sum_{n=1}^{\infty} \frac{|\hat{f}(-n)|^2}{n}.$$

Using an argument similar to that of Theorem 3.1 we can prove:

**Theorem 3.2.** *There exists a constant  $C > 0$  such that for all functions  $f \in L^1(\mathbb{R})$  we have*

$$\int_0^{\infty} \frac{|\hat{f}(\xi)|^2}{\xi} d\xi \leq C \|f\|_1^2 + C \int_0^{\infty} \frac{|\hat{f}(-\xi)|^2}{\xi} d\xi.$$

This modifies the result proved in [8].



[Title Page](#)

[Contents](#)



Page 10 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

## 4. Another Hardy Inequality

In this section we prove (1.1) on the real line under a certain condition on the signs of the Fourier coefficients.

We should remark here that the method used to prove this inequality is standard and all recent articles use this idea; we need a bounded function whose Fourier coefficients obey some desired decay conditions.

One last remark before proceeding, although the given proof is for a real Hardy inequality, we can imitate the given steps to prove a similar inequality on the circle.

For  $j \geq 1$  put

$$f_j(x) = \frac{1}{4^j} \int_{4^{j-1}}^{4^j} e^{ix\xi} d\xi, \quad x \in \mathbb{R}.$$

Then we have our first lemma:

**Lemma 4.1.** *Let  $f_j$  be as above, then*


$$\hat{f}_j(\xi) = \begin{cases} \frac{2\pi}{4^j}, & 4^{j-1} < \xi < 4^j, \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* Let

$$g_j(\xi) = \begin{cases} \frac{2\pi}{4^j}, & 4^{j-1} < \xi < 4^j, \\ 0, & \text{otherwise,} \end{cases}$$

then  $g_j \in L^2$ . Therefore,  $g_j$  is the Fourier transform of some function  $h_j \in L^2$  and

$$\begin{aligned} h_j(x) &= \frac{1}{2\pi} \int_{\mathbb{R}} g(\xi) e^{ix\xi} d\xi \\ &= f(x). \end{aligned}$$



Title Page

Contents



Page 11 of 18

Go Back

Full Screen

Close

However,  $\hat{h}_j = g_j$ , which implies  $\hat{f}_j = g_j$  as claimed. □

Observe that the above proof implies that  $f_j \in L^2$  and

**Lemma 4.2.** For  $f_j$  as above, we have

$$\|f_j\|_2 = \frac{\sqrt{6\pi}}{4^{j/2}}.$$

*Proof.* Since  $f_j \in L^2$  we have


$$\begin{aligned} \|f_j\|_2 &= \frac{1}{\sqrt{2\pi}} \|\hat{f}_j\|_2 \\ &= \frac{1}{\sqrt{2\pi}} \frac{2\pi}{4^j} \left( \int_{4^{j-1}}^{4^j} d\xi \right)^{\frac{1}{2}} \\ &= \frac{\sqrt{6\pi}}{4^{j/2}}. \end{aligned}$$

□

**Lemma 4.3.** For  $f_j$  as above and for  $M \in \mathbb{N}$ , let

$$F_M(x) = \sum_{j=1}^M \left| f_j(x) - \overline{f_j(x)} \right|,$$

then  $\|F_M\|_\infty \leq C$  where  $C$  is some absolute constant (independent of  $M$ ).



Title Page

Contents



Page 12 of 18

Go Back

Full Screen

Close

*Proof.* Observe first that

$$\begin{aligned}
 \left| f_j(x) - \overline{f_j(x)} \right| &= \frac{2}{4^j} \left| \int_{4^{j-1}}^{4^j} \sin(x\xi) d\xi \right| \\
 &= \frac{2}{4^j} \left| \frac{\cos(4^j x) - \cos(4^{j-1} x)}{x} \right| \\
 &= \frac{2}{4^j} \left| \frac{2 \sin^2(4^{j-1} x/2) [-1 + 16 \cos^2(4^{j-1} x) \cos^2(4^{j-1} x/2)]}{x} \right| \\
 &\leq \frac{60 \sin^2(4^{j-1} x/2)}{4^j |x|}.
 \end{aligned}$$

Note that  $F_M$  is an even function, so it suffices to consider only the case  $x > 0$ .

Now, fix  $x > 0$  and observe that


$$\begin{aligned}
 F_M(x) &\leq 60 \sum_{j=1}^{\infty} \frac{1}{4^j} \frac{\sin^2(4^{j-1} x/2)}{x} \\
 &= 60 \left( \sum_{4^{-j} \leq x} \frac{1}{4^j} \frac{\sin^2(4^{j-1} x/2)}{x} + \sum_{4^{-j} > x} \frac{1}{4^j} \frac{\sin^2(4^{j-1} x/2)}{x} \right),
 \end{aligned}$$

where, by convention, the second sum is zero if  $4^{-j} \leq x$  for all  $j \geq 1$ .

Now

$$\sum_{4^{-j} \leq x} \frac{1}{4^j} \frac{\sin^2(4^{j-1} x/2)}{x} \leq \frac{1}{x} \sum_{j=k_x}^{\infty} \frac{1}{4^j},$$

where  $k_x$  is the smallest positive integer such that  $4^{-j} \leq x$  for all  $j \geq k_x$ .



[Title Page](#)

[Contents](#)



Page 13 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

Consequently

$$\begin{aligned} \sum_{4^{-j} \leq x} \frac{1}{4^j} \frac{\sin^2(4^{j-1}x/2)}{x} &\leq \frac{1}{x} \frac{4}{3} \frac{1}{4^{k_x}} \\ &\leq \frac{1}{x} \frac{4}{3} x = \frac{4}{3}. \end{aligned}$$


On the other hand, if  $\sum_{4^{-j} > x} \frac{1}{4^j} \frac{\sin^2(4^{j-1}x/2)}{x}$  is not zero, we get

$$\begin{aligned} \sum_{4^{-j} > x} \frac{1}{4^j} \frac{\sin^2(4^{j-1}x/2)}{x} &= \sum_{j=1}^{\log_4(1/x)} \frac{1}{4^j} \frac{\sin^2(4^{j-1}x/2)}{x} \\ &\leq \sum_{j=1}^{\log_4(1/x)} \frac{1}{4^j x} \left( \frac{1}{2} 4^{j-1} x \right)^2 \\ &= \frac{1}{64} x \sum_{j=1}^{\log_4(1/x)} 4^j \\ &\leq \frac{1}{64} x \frac{4^{\log_4(1/x)} - 4}{3} \\ &\leq \frac{1}{192}. \end{aligned}$$

Thus, we have

$$F_M(x) \leq 60 \left( \frac{4}{3} + \frac{1}{192} \right) := C.$$

□



Title Page

Contents



Page 14 of 18

Go Back

Full Screen

Close

Now let  $X$  be the set of all  $f \in L^1$  such that the sign of  $\hat{f}(\xi)$  is constant in the block  $(4^{j-1}, 4^j)$ . Then we have our main result:

**Theorem 4.4.** *There is an absolute constant  $K > 0$  such that*

$$(4.1) \quad \int_1^\infty \frac{|\hat{f}(\xi)|}{\xi} d\xi \leq K \|f\|_1 + K \int_1^\infty \frac{|\hat{f}(-\xi)|}{\xi} d\xi$$

for all  $f \in X$ .


*Proof.* Let  $f \in X$  be such that  $\hat{f}$  is compactly supported, let  $f_j$  be as above and let  $M \in \mathbb{N}$  be such that the support of  $\hat{f}$  is contained in  $[-M, M]$ . Denote the sign of  $\hat{f}$  in the block  $(4^{j-1}, 4^j)$  by  $\sigma_j$  and put

$$F(x) = \sum_{j=1}^M \sigma_j \left( f_j(x) - \overline{f_j(x)} \right).$$

Then  $\|F\|_\infty \leq C$ , where  $C$  is the constant of Lemma 4.3. Moreover,  $\hat{F}(\xi) = \frac{2\pi}{4^j} \sigma_j$ , where  $j$  is the unique index such that  $\xi \in [4^{j-1}, 4^j] \cup [-4^j, -4^{j-1}]$  if  $-4^M \leq \xi \leq 4^M$  and  $\hat{F}(\xi) = 0$  otherwise.

Moreover, since  $\hat{f}$  is of compact support, we have  $f \in L^2$ . Now we apply a standard duality argument:

$$\begin{aligned} C \|f\|_1 &\geq \left| \int_{\mathbb{R}} f(x) \overline{F(x)} dx \right| \\ &= \frac{1}{2\pi} \left| \int_{\mathbb{R}} \hat{f}(\xi) \overline{\hat{F}(\xi)} d\xi \right| \quad (\text{see Lemma 2.1}) \end{aligned}$$



Title Page

Contents



Page 15 of 18

Go Back

Full Screen

Close

$$\begin{aligned} &\geq \frac{1}{2\pi} \left| \sum_{j=1}^{\infty} \int_{4^{j-1}}^{4^j} \hat{f}(\xi) \overline{\hat{F}(\xi)} d\xi \right| - \frac{1}{2\pi} \left| \int_{-1}^1 \hat{f}(\xi) \overline{\hat{F}(\xi)} d\xi \right| \\ &\quad - \frac{1}{2\pi} \left| \sum_{j=1}^{\infty} \int_{4^{j-1}}^{4^j} \hat{f}(-\xi) \overline{\hat{F}(-\xi)} d\xi \right|. \end{aligned}$$

That is,

$$(4.2) \quad \frac{1}{2\pi} \left| \sum_{j=1}^{\infty} \int_{4^{j-1}}^{4^j} \hat{f}(\xi) \overline{\hat{F}(\xi)} d\xi \right| \leq C \|f\|_1 + \frac{1}{2\pi} \left| \int_{-1}^1 \hat{f}(\xi) \overline{\hat{F}(\xi)} d\xi \right| + \frac{1}{2\pi} \left| \sum_{j=1}^{\infty} \int_{4^{j-1}}^{4^j} \hat{f}(-\xi) \overline{\hat{F}(-\xi)} d\xi \right|.$$

However, when  $\xi \in (4^{j-1}, 4^j) \cup (-4^j, -4^{j-1})$ , we have  $\hat{F}(\xi) = \frac{2\pi}{4^j} \sigma_j$ , where  $\sigma_j$  is the sign of  $\hat{f}$  in  $(4^j, 4^{j-1})$ . Thus, when  $\xi \in (4^{j-1}, 4^j)$  we have  $\hat{f}(\xi) \overline{\hat{F}(\xi)} = |\hat{f}(\xi)|$ . Moreover

$$\begin{aligned} \left| \int_{-1}^1 \hat{f}(\xi) \overline{\hat{F}(\xi)} d\xi \right| &\leq \|f\|_1 \int_{-1}^1 |\hat{F}(\xi)| d\xi \\ &\leq \|f\|_1 \left( \int_{-1}^1 d\xi \right)^{\frac{1}{2}} \left( \int_{-1}^1 |\hat{F}(\xi)|^2 d\xi \right)^{\frac{1}{2}} \\ &\leq \sqrt{2} \|f\|_1 \|\hat{F}\|_2 \\ &= \sqrt{2} \|f\|_1 \sqrt{2\pi} \|F\|_2 \end{aligned}$$

$$\begin{aligned} &\leq 2\sqrt{\pi}\|f\|_1 \times 2 \sum_{j=1}^{\infty} \|f_j\|_2 \\ &= 4\sqrt{6\pi}\|f\|_1, \end{aligned}$$

where we have used the facts

$$F(x) = \sum_{j=1}^M \sigma_j \left( f_j(x) - \overline{f_j(x)} \right) \quad \text{and} \quad \|f_j\|_2 = \sqrt{6\pi}4^{-j/2}.$$

Consequently, (4.2) becomes

$$(4.3) \quad \sum_{j=1}^{\infty} \int_{4^{j-1}}^{4^j} \frac{|\hat{f}(\xi)|}{4^j} d\xi \leq C\|f\|_1 + 2\sqrt{6}\|f\|_1 + \sum_{j=1}^{\infty} \int_{4^{j-1}}^{4^j} \frac{|\hat{f}(-\xi)|}{4^j} d\xi.$$

However, when  $\xi \in [4^{j-1}, 4^j]$  we have


$$\frac{1}{4^j} \leq \frac{1}{\xi} \quad \text{and} \quad \frac{1}{4^j} \geq \frac{1}{4\xi}.$$

Therefore, (4.3) boils down to

$$\int_1^{\infty} \frac{|\hat{f}(\xi)|}{\xi} \leq K\|f\|_1 + K \int_1^{\infty} \frac{|\hat{f}(-\xi)|}{\xi} d\xi,$$

where  $K = 4(C + 2\sqrt{6})$  and where  $C$  is the constant of Lemma 4.3.

This completes the proof for  $f \in X$  with the property that  $\hat{f}$  is compactly supported. Now for general  $f \in X$ , consider the convolution  $f * K_\lambda$  where  $K_\lambda$  is the Fejer Kernel of order  $\lambda$ . A standard limiting process yields the result.  $\square$



Real Hardy Inequality

Mohammad Sababheh

vol. 10, iss. 4, art. 104, 2009

Title Page

Contents



Page 16 of 18

Go Back


Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756





## Remark:

1. We note that the above proof can be adopted to prove that: There is an absolute constant  $C' > 0$  such that for any function  $f \in L^1(\mathbb{T})$  whose Fourier coefficients have the same sign on the block  $[4^{j-1}, 4^j)$  we have

$$\sum_{n=1}^{\infty} \frac{|\hat{f}(n)|}{n} \leq C' \|f\|_1 + C' \sum_{n=1}^{\infty} \frac{|\hat{f}(-n)|}{n}.$$

2. Observe that the condition that  $\hat{f}$  has the same sign on the block  $(4^{j-1}, 4^j)$  is flexible. This is because functions in  $L^p$  are in fact equivalent classes. Therefore, even if  $\hat{f}$  obeys our condition but for a set of measure zero, then we may modify our choice by changing the values of  $\hat{f}$  so that the new  $\hat{f}$  satisfies our condition.

We should remark that this process does not effect the proof above.

---

Real Hardy Inequality

Mohammad Sababbeh

vol. 10, iss. 4, art. 104, 2009

---

Title Page

Contents



Page 17 of 18


Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756



## References

- [1] Y. KATZNELSON, *An Introduction To Harmonic Analysis*, John Wiley and Sons, Inc., 1968.
- [2] I. KLEMES, A note on Hardy's inequality, *Canad. Math. Bull.*, **36**(4) (1993), 442–448.
- [3] P. KOOSIS, *Conference On Harmonic Analysis in Honour of Antoni Zygmund*, Volume 2, Wadsworth International Group, Belmont, California, A Division of Wadsworth, Inc., (1981), 740–748.
- [4] O.C. McGEHEE, L. PIGNO AND B. SMITH, Hardy's inequality and the  $L^1$  norm of exponential sums, *Annals of Math.*, **113** (1981), 613–618.
- [5] W. RUDIN, *Real and Complex Analysis*, 3rd edition, McGraw-Hill Book Co., New York, 1987.
- [6] M. SABABHEH, Two-sided probabilistic versions of Hardy's inequality, *J. Fourier Anal. and Applics.*, (2007), 577–587.
- [7] M. SABABHEH, On an argument of Körner and Hardy's inequality, *Analysis Mathematica*, **34** (2008), 51–57.
- [8] M. SABABHEH, Hardy-type inequalities on the real line, *J. Inequal. in Pure and Appl. Math.*, **9**(3) (2008), Art. 72. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=1009>].

Real Hardy Inequality

Mohammad Sababheh

vol. 10, iss. 4, art. 104, 2009

Title Page

Contents



Page 18 of 18

Go Back

Full Screen

Close

journal of **inequalities**  
in pure and applied  
mathematics

issn: 1443-5756